

LE MATEMATICHE

Vol. LXX (2015) – Fasc. I, pp. 3–20

doi: 10.4418/2015.70.1.1

HERMITE-HADAMARD TYPE FRACTIONAL INTEGRAL INEQUALITIES FOR GEOMETRIC-GEOMETRIC CONVEX FUNCTIONS

SHENGDA LIU - JINRONG WANG

By utilizing two fractional integral identities and elementary inequalities via geometric-geometric (*GG* for short) convex functions, we derive new type Hermite-Hadamard inequalities involving Hadamard fractional integrals. Some applications to special means of real numbers are given.

1. Introduction

Fractional calculus originally appeared in the letter between L'Hospital and Leibniz. Since 1695, Riemann, Liouville, Caputo, Hadamard and other famous mathematicians paid attention to study such a branch of mathematical analysis. Meanwhile, fractional calculus have been widely applied to the fields of electricity, biology, economics and signal and image processing [1–8].

The classical Hermite-Hadamard inequality was firstly discovered by Hermite in 1881 in the journal *Mathesis*, which provides a lower and an upper estimations for the integral average of any convex function defined on a compact

Entrato in redazione: 30 dicembre 2013

AMS 2010 Subject Classification: 26A33, 26A51, 26D15.

Keywords: Hermite-Hadamard inequalities, Hadamard fractional integrals, *GG*-convex functions. This work is supported by the National Natural Science Foundation of China (11201091), Key Projects of Science and Technology Research in the Chinese Ministry of Education (211169), Key Support Subject (Applied Mathematics) and Key project on the reforms of teaching contents and course system of Guizhou Normal College.

interval, involving the midpoint and the endpoints of the domain. Very recently, many authors pay attention to study Hermite-Hadamard type inequalities involving Riemann-Liouville and Hadamard fractional integrals. For more recent results which generalize, improve, and extend the classical Hermite-Hadamard inequality, one can see [9–23] and the references therein.

For $f \in L[a, b]$, the Hadamard fractional integrals [1] $J_{a+}^{\alpha} f$ and $J_{b-}^{\alpha} f$ of order $\alpha \in R^+$ with $a \geq 0$ are defined by

$$({}_H J_{a+}^{\alpha} f)(x) = \frac{1}{\Gamma(\alpha)} \int_a^x \left(\ln \frac{x}{t} \right)^{\alpha-1} f(t) \frac{dt}{t}, \quad (0 < a < x \leq b),$$

and

$$({}_H J_{b-}^{\alpha} f)(x) = \frac{1}{\Gamma(\alpha)} \int_x^b \left(\ln \frac{t}{x} \right)^{\alpha-1} f(t) \frac{dt}{t}, \quad (0 < a \leq x < b),$$

where $\Gamma(\cdot)$ is the Gamma function.

Recently, Wang et al. [17, 18] established the following two powerful fractional integral identities involving Hadamard fractional integrals.

Lemma 1.1 (see [17]). *Let $f : [a, b] \rightarrow R$ be a differentiable mapping on (a, b) with $0 < a < b$. If $f' \in L[a, b]$, then the following equality for fractional integrals holds*

$$\begin{aligned} & \frac{(\ln x - \ln a)^{\alpha} + (\ln b - \ln x)^{\alpha}}{\ln b - \ln a} f(x) - \frac{\Gamma(\alpha + 1)}{\ln b - \ln a} [{}_H J_{x-}^{\alpha} f(a) + {}_H J_{x+}^{\alpha} f(b)] \\ &= \frac{(\ln x - \ln a)^{\alpha+1}}{\ln b - \ln a} \int_0^1 t^{\alpha} e^{t \ln x + (1-t) \ln a} f'(e^{t \ln x + (1-t) \ln a}) dt \\ & - \frac{(\ln b - \ln x)^{\alpha+1}}{\ln b - \ln a} \int_0^1 t^{\alpha} e^{t \ln x + (1-t) \ln b} f'(e^{t \ln x + (1-t) \ln b}) dt \\ &= \frac{(\ln x - \ln a)^{\alpha+1}}{\ln b - \ln a} \int_0^1 t^{\alpha} x^t a^{1-t} f'(x^t a^{1-t}) dt \\ & - \frac{(\ln b - \ln x)^{\alpha+1}}{\ln b - \ln a} \int_0^1 t^{\alpha} x^t b^{1-t} f'(x^t a^{1-t}) dt, \end{aligned}$$

for any $x \in (a, b)$.

Lemma 1.2 (see [18]). *Let $f : [a, b] \rightarrow R$ be a differentiable mapping on (a, b) with $0 < a < b$. If $f' \in L[a, b]$, then the following equality for fractional integrals holds:*

$$\begin{aligned} & \frac{\Gamma(\alpha + 1)}{2(\ln b - \ln a)^{\alpha}} [{}_H J_{a+}^{\alpha} f(b) + {}_H J_{b-}^{\alpha} f(a)] - f(\sqrt{ab}) = \\ &= \frac{\ln b - \ln a}{2} \left[\int_0^1 k e^{t \ln a + (1-t) \ln b} f'(e^{t \ln a + (1-t) \ln b}) dt \right. \end{aligned}$$

$$\begin{aligned}
& - \int_0^1 [(1-t)^\alpha - t^\alpha] e^{t \ln a + (1-t) \ln b} f'(e^{t \ln a + (1-t) \ln b}) dt \Big] \\
& = \frac{\ln b - \ln a}{2} \left[\int_0^1 k a^t b^{1-t} f'(a^t b^{1-t}) dt - \int_0^1 [(1-t)^\alpha - t^\alpha] a^t b^{1-t} f'(a^t b^{1-t}) dt \right],
\end{aligned}$$

where

$$k = \begin{cases} 1, & 0 \leq t < \frac{1}{2}, \\ -1, & \frac{1}{2} \leq t < 1. \end{cases}$$

Meanwhile, Wang et al. [17] introduced a new classes of functions satisfying “*s-e*-condition”.

Definition 1.3. A function $f : I \subset (0, \infty) \rightarrow R$ is said to satisfy *s-e*-condition if

$$f(e^{\lambda x + (1-\lambda)y}) \leq \lambda^s f(e^x) + (1-\lambda)^s f(e^y)$$

for all $x, y \in I$, $\lambda \in [0, 1]$ and for some fixed $s \in (0, 1]$.

With the help of the new concept of “*s-e*-condition”, monotonicity and Lemma 1.1 and Lemma 1.2, some new Hermite-Hadamard inequalities involving Hadamard fractional integrals are established [17].

Motivated by our previous works [17, 18], we do not use the concept of “*s-e*-condition” and use the concept of *GG*-convex functions (see Definition 2.1), which maybe more suitable for solving such problem in some sense. We will use Lemma 1.1, Lemma 1.2 and elementary equalities and inequalities via *GG*-convex functions to derive new type Hermite-Hadamard inequalities involving Hadamard fractional integrals.

2. Definitions, elementary equalities and inequalities

Definition 2.1 (see [24, 25]). Let $f : I \subseteq R^+ \rightarrow R^+$. A function f is said to be *GG*-convex on I if for every $x, y \in I$ and $\lambda \in [0, 1]$, we have

$$f(x^\lambda y^{1-\lambda}) \leq [f(x)]^\lambda [f(y)]^{1-\lambda}. \quad (1)$$

Remark 2.2. By the arithmetic-geometric mean inequality, we have

$$[f(x)]^\lambda [f(y)]^{1-\lambda} \leq \lambda f(x) + (1-\lambda)f(y). \quad (2)$$

Linking (1) and (2), we obtain

$$f(x^\lambda y^{1-\lambda}) \leq \lambda f(x) + (1-\lambda)f(y),$$

which appears in the standard definition of geometric-arithmetic (GA for short) convex function [24]. So *GG*-convex function is GA-convex function.

Remark 2.3 (see [26]). The functions $f(x) = \cosh x$ and $g(x) = e^x$ are GG-convex on $(0, +\infty)$.

It follows Lemma 2.1 and Lemma 2.2 in [14, 17] that we give two singular integral equalities.

Lemma 2.4. For $\alpha > 0$ and $k > 0$, we have

$$I_f(\alpha, x, y) = \int_0^1 t^{\alpha-1} T_f^t(x, y) dt = T_f(x, y) \sum_{i=1}^{\infty} (-1)^{i-1} \frac{(\ln T_f(x, y))^{i-1}}{(\alpha)_i} < +\infty$$

where T is defined by

$$T_f(x, y) = \frac{x|f'(x)|}{y|f'(y)|},$$

and

$$(\alpha)_i = \alpha(\alpha+1)(\alpha+2) \cdots (\alpha+i-1).$$

Lemma 2.5. For $\alpha > 0$ and $k > 0, z > 0$, we have

$$I_f(a, x, y, z) = \int_0^z t^{\alpha-1} T_f^t(x, y) dt = z^\alpha T_f^z(x, y) \sum_{i=0}^{\infty} \frac{(-z \ln T_f(x, y))^{i-1}}{(\alpha)_i} < +\infty.$$

The following two inequalities are also needed.

Lemma 2.6 (see [14]). For $A > B > 0$, it holds

$$\begin{aligned} (A - B)^\theta &\leq A^\theta - B^\theta \quad \text{when } \theta \geq 1, \\ (A - B)^\theta &\geq A^\theta - B^\theta \quad \text{when } 0 < \theta \leq 1. \end{aligned}$$

Lemma 2.7 (see [19]). For $t \in [0, 1]$, we have

$$\begin{aligned} (1-t)^n &\leq 2^{1-n} - t^n \quad \text{for } n \in [0, 1], \\ (1-t)^n &\geq 2^{1-n} - t^n \quad \text{for } n \in [1, \infty). \end{aligned}$$

3. Main results

In this section, we will use the results via GG-convex functions in Section 2 to derive our main results in this paper.

Theorem 3.1. Let $f : [0, b] \rightarrow R$ be a differentiable mapping. If $|f'|$ is measurable and $|f'|$ is GG-convex on $[0, b]$ for some fixed $\alpha \in (0, \infty)$, $0 \leq a < b$, then the following inequality for fractional integrals holds:

$$\left| \frac{(\ln x - \ln a)^\alpha + (\ln b - \ln x)^\alpha}{\ln b - \ln a} f(x) - \frac{\Gamma(\alpha+1)}{\ln b - \ln a} [{}_H J_{x^-}^\alpha f(a) + {}_H J_{x^+}^\alpha f(b)] \right|$$

$$\begin{aligned} &\leq \frac{(\ln x - \ln a)^{\alpha+1}}{\ln b - \ln a} \cdot a \cdot |f'(a)| \cdot T_f(x, a) \sum_{i=1}^{\infty} (-1)^{i-1} \frac{(\ln T_f(x, a))^{i-1}}{(\alpha+1)_i} \\ &+ \frac{(\ln b - \ln x)^{\alpha+1}}{\ln b - \ln a} \cdot b \cdot |f'(b)| \cdot T_f(x, b) \sum_{i=1}^{\infty} (-1)^{i-1} \frac{(\ln T_f(x, b))^{i-1}}{(\alpha+1)_i}, \end{aligned}$$

for any $x \in (a, b)$.

Proof. By using Definition 2.1 and Lemma 2.4, we have

$$\begin{aligned} \int_0^1 t^\alpha x^t a^{1-t} |f'(x^t a^{1-t})| dt &\leq \int_0^1 t^\alpha x^t a^{1-t} |f'(x)|^t |f'(a)|^{1-t} dt \\ &= a |f'(a)| I_f(\alpha+1, x, a). \end{aligned}$$

Similarly, one has

$$\int_0^1 t^\alpha x^t b^{1-t} |f'(x^t b^{1-t})| dt = b |f'(b)| I_f(\alpha+1, x, b).$$

By using Lemma 1.1, we obtain

$$\begin{aligned} &\left| \frac{(\ln x - \ln a)^\alpha + (\ln b - \ln x)^\alpha}{\ln b - \ln a} f(x) - \frac{\Gamma(\alpha+1)}{\ln b - \ln a} [{}_H J_{x^-}^\alpha f(a) + {}_H J_{x^+}^\alpha f(b)] \right| \\ &= \left| \frac{(\ln x - \ln a)^{\alpha+1}}{\ln b - \ln a} \int_0^1 t^\alpha x^t a^{1-t} f'(x^t a^{1-t}) dt \right. \\ &\quad \left. - \frac{(\ln b - \ln x)^{\alpha+1}}{\ln b - \ln a} \int_0^1 t^\alpha x^t b^{1-t} f'(x^t a^{1-t}) dt \right| \\ &\leq \frac{(\ln x - \ln a)^{\alpha+1}}{\ln b - \ln a} \int_0^1 t^\alpha x^t a^{1-t} |f'(x^t a^{1-t})| dt \\ &\quad + \frac{(\ln b - \ln x)^{\alpha+1}}{\ln b - \ln a} \int_0^1 t^\alpha x^t b^{1-t} |f'(x^t b^{1-t})| dt \\ &\leq \frac{(\ln x - \ln a)^{\alpha+1}}{\ln b - \ln a} a |f'(a)| I_f(\alpha+1, x, a) \\ &\quad + \frac{(\ln b - \ln x)^{\alpha+1}}{\ln b - \ln a} b |f'(b)| I_f(\alpha+1, x, b) \\ &= \frac{(\ln x - \ln a)^{\alpha+1}}{\ln b - \ln a} \cdot a \cdot |f'(a)| \cdot T_f(x, a) \sum_{i=1}^{\infty} (-1)^{i-1} \frac{(\ln T_f(x, a))^{i-1}}{(\alpha+1)_i} \\ &\quad + \frac{(\ln b - \ln x)^{\alpha+1}}{\ln b - \ln a} \cdot b \cdot |f'(b)| \cdot T_f(x, b) \sum_{i=1}^{\infty} (-1)^{i-1} \frac{(\ln T_f(x, b))^{i-1}}{(\alpha+1)_i}. \end{aligned}$$

The proof is completed. \square

Theorem 3.2. Let $f : [0, b] \rightarrow R$ be a differentiable mapping and $1 < q < \infty$. If $|f'|^q$ is measurable and $|f'|^q$ is GG-convex on $[0, b]$ for some fixed $\alpha \in (0, \infty)$, $0 \leq a < b$, then the following inequality for fractional integrals holds:

$$\begin{aligned} & \left| \frac{(\ln x - \ln a)^\alpha + (\ln b - \ln x)^\alpha}{\ln b - \ln a} f(x) - \frac{\Gamma(\alpha + 1)}{\ln b - \ln a} [{}_H J_{x^-}^\alpha f(a) + {}_H J_{x^+}^\alpha f(b)] \right| \\ & \leq \frac{(\ln x - \ln a)^{\alpha+1}}{\ln b - \ln a} \cdot a \cdot |f'(a)| \cdot \left(\frac{1}{\alpha p + 1} \right)^{\frac{1}{p}} \cdot \left(\frac{T_f^q(x, a) - 1}{q \ln(T_f(x, a))} \right)^{\frac{1}{q}} \\ & \quad + \frac{(\ln b - \ln x)^{\alpha+1}}{\ln b - \ln a} \cdot b \cdot |f'(b)| \cdot \left(\frac{1}{\alpha p + 1} \right)^{\frac{1}{p}} \cdot \left(\frac{T_f^q(x, b) - 1}{q \ln(T_f(x, b))} \right)^{\frac{1}{q}}, \end{aligned}$$

where $\frac{1}{p} + \frac{1}{q} = 1$.

Proof. By using Definition 2.1, Hölder inequality, we obtain

$$\begin{aligned} & \int_0^1 t^\alpha x^t a^{1-t} |f'(x^t a^{1-t})| dt \\ & \leq \left(\int_0^1 t^{\alpha p} dt \right)^{\frac{1}{p}} \left(\int_0^1 (x^t a^{1-t} |f'(x^t a^{1-t})|)^q dt \right)^{\frac{1}{q}} \\ & \leq \left(\frac{1}{\alpha p + 1} \right)^{\frac{1}{p}} \left(\int_0^1 x^{qt} a^{q(1-t)} |f'(x)|^{qt} |f'(a)|^{q(1-t)} dt \right)^{\frac{1}{q}} \\ & \leq \left(\frac{1}{\alpha p + 1} \right)^{\frac{1}{p}} \cdot a \cdot |f'(a)| \cdot \left(\int_0^1 T_f^q(x, a) dt \right)^{\frac{1}{q}} \\ & \leq \left(\frac{1}{\alpha p + 1} \right)^{\frac{1}{p}} \cdot a \cdot |f'(a)| \cdot \left(\frac{T_f^q(x, a) - 1}{q \ln(T_f(x, a))} \right)^{\frac{1}{q}}. \end{aligned}$$

Similarly, one has

$$\int_0^1 t^\alpha x^t b^{1-t} |f'(x^t b^{1-t})| dt \leq \left(\frac{1}{\alpha p + 1} \right)^{\frac{1}{p}} \cdot b \cdot |f'(b)| \cdot \left(\frac{T_f^q(x, b) - 1}{q \ln(T_f(x, b))} \right)^{\frac{1}{q}}.$$

By using Lemma 1.1, we have

$$\begin{aligned} & \left| \frac{(\ln x - \ln a)^\alpha + (\ln b - \ln x)^\alpha}{\ln b - \ln a} f(x) - \frac{\Gamma(\alpha + 1)}{\ln b - \ln a} [{}_H J_{x^-}^\alpha f(a) + {}_H J_{x^+}^\alpha f(b)] \right| \\ & = \left| \frac{(\ln x - \ln a)^{\alpha+1}}{\ln b - \ln a} \int_0^1 t^\alpha x^t a^{1-t} f'(x^t a^{1-t}) dt - \right. \end{aligned}$$

$$\begin{aligned}
& - \frac{(\ln b - \ln x)^{\alpha+1}}{\ln b - \ln a} \left| \int_0^1 t^\alpha x^t b^{1-t} f'(x^t b^{1-t}) dt \right| \\
& \leq \frac{(\ln x - \ln a)^{\alpha+1}}{\ln b - \ln a} \int_0^1 t^\alpha x^t a^{1-t} |f'(x^t a^{1-t})| dt \\
& + \frac{(\ln b - \ln x)^{\alpha+1}}{\ln b - \ln a} \int_0^1 t^\alpha x^t b^{1-t} |f'(x^t b^{1-t})| dt \\
& \leq \frac{(\ln x - \ln a)^{\alpha+1}}{\ln b - \ln a} \cdot a \cdot |f'(a)| \cdot \left(\frac{1}{\alpha p + 1} \right)^{\frac{1}{p}} \cdot \left(\frac{T_f^q(x, a) - 1}{q \ln(T_f(x, a))} \right)^{\frac{1}{q}} \\
& + \frac{(\ln b - \ln x)^{\alpha+1}}{\ln b - \ln a} \cdot b \cdot |f'(b)| \cdot \left(\frac{1}{\alpha p + 1} \right)^{\frac{1}{p}} \cdot \left(\frac{T_f^q(x, b) - 1}{q \ln(T_f(x, b))} \right)^{\frac{1}{q}}.
\end{aligned}$$

The proof is done. \square

Theorem 3.3. *Let $f : [0, b] \rightarrow R$ be a differentiable mapping. If $|f'|$ is measurable and $|f'|$ is GG-convex on $[0, b]$ for some fixed $\alpha \in (0, \infty)$, $0 \leq a < b$, then the following inequality for fractional integrals holds:*

$$\begin{aligned}
& \left| \frac{\Gamma(\alpha + 1)}{2(\ln b - \ln a)^\alpha} [{}_H J_{b^-}^\alpha f(a) + {}_H J_{a^+}^\alpha f(b)] - f(\sqrt{ab}) \right| \\
& \leq \max \left\{ \frac{\ln b - \ln a}{2} b |f'(b)| \left[\frac{2T_f(a, b) - 1 + (2^{1-\alpha} + 1)T_f^{\frac{1}{2}}(a, b)}{\ln(T_f(a, b))} \right. \right. \\
& \quad \left. \left. - 4I_f(\alpha + 1, a, b, \frac{1}{2}) + 2I_f(\alpha + 1, a, b, a) \right], \right. \\
& \quad \left. \frac{\ln b - \ln a}{2} b |f'(b)| \left[\frac{2T_f^{\frac{1}{2}}(a, b) - 1}{\ln(T_f(a, b))} - 4I_f(\alpha + 1, a, b, \frac{1}{2}) \right. \right. \\
& \quad \left. \left. + 2I_f(\alpha + 1, a, b, a) \right] \right\}.
\end{aligned}$$

Proof. To achieve our aim, we divide our proof into two cases.

Case 1: $\alpha \in (0, 1)$. By using Definition 2.1, Lemma 1.2, Lemma 2.4, Lemma 2.5, Lemma 2.6, Lemma 2.7 via Hölder inequality, we have

$$\begin{aligned}
& \left| \frac{\Gamma(\alpha + 1)}{2(\ln b - \ln a)^\alpha} [{}_H J_{b^-}^\alpha f(a) + {}_H J_{a^+}^\alpha f(b)] - f(\sqrt{ab}) \right| \\
& = \left| \frac{\ln b - \ln a}{2} \left[\int_0^1 k a^t b^{1-t} f'(a^t b^{1-t}) dt \right. \right. \\
& \quad \left. \left. - \int_0^1 [(1-t)^\alpha - t^\alpha] a^t b^{1-t} f'(a^t b^{1-t}) dt \right] \right| \leq
\end{aligned}$$

$$\begin{aligned}
&\leq \frac{\ln b - \ln a}{2} \left[\int_0^1 a^t b^{1-t} |f'(a^t b^{1-t})| dt \right. \\
&+ \left. \int_0^1 |(1-t)^\alpha - t^\alpha| \cdot a^t b^{1-t} \cdot |f'(a^t b^{1-t})| dt \right] \\
&\leq \frac{\ln b - \ln a}{2} \left[\int_0^1 a^t b^{1-t} |f'(a)|^t |f'(b)|^{1-t} dt \right. \\
&+ \left. \int_0^1 |(1-t)^\alpha - t^\alpha| a^t b^{1-t} |f'(a)|^t |f'(b)|^{1-t} dt \right] \\
&= \frac{\ln b - \ln a}{2} \left[b |f'(b)| \int_0^1 T_f^t(a, b) dt + b |f'(b)| \int_0^1 |(1-t)^\alpha - t^\alpha| T_f^t(a, b) dt \right] \\
&= \frac{\ln b - \ln a}{2} b |f'(b)| \left[\int_0^1 T_f^t(a, b) dt + \int_0^1 |(1-t)^\alpha - t^\alpha| T_f^t(a, b) dt \right] \\
&= \frac{\ln b - \ln a}{2} b |f'(b)| \left[\int_0^1 T_f^t(a, b) dt + \int_0^{\frac{1}{2}} ((1-t)^\alpha - t^\alpha) T_f^t(a, b) dt \right. \\
&+ \left. \int_{\frac{1}{2}}^1 (-(1-t)^\alpha + t^\alpha) T_f^t(a, b) dt \right] \\
&\leq \frac{\ln b - \ln a}{2} b |f'(b)| \left[\int_0^1 T_f^t(a, b) dt + \int_0^{\frac{1}{2}} (1-t)^\alpha T_f^t(a, b) dt \right. \\
&- \left. \int_0^{\frac{1}{2}} t^\alpha T_f^t(a, b) dt - \int_{\frac{1}{2}}^1 (1-t)^\alpha T_f^t(a, b) dt + \int_{\frac{1}{2}}^1 t^\alpha T_f^t(a, b) dt \right] \\
&\leq \frac{\ln b - \ln a}{2} b |f'(b)| \left[\int_0^1 T_f^t(a, b) dt + \int_0^{\frac{1}{2}} (2^{1-\alpha} - t^\alpha) T_f^t(a, b) dt \right. \\
&- \left. \int_0^{\frac{1}{2}} t^\alpha T_f^t(a, b) dt - \int_{\frac{1}{2}}^1 (1-t^\alpha) T_f^t(a, b) dt + \int_{\frac{1}{2}}^1 t^\alpha T_f^t(a, b) dt \right] \\
&= \frac{\ln b - \ln a}{2} b |f'(b)| \left[\int_0^1 T_f^t(a, b) dt \right. \\
&+ \left. \int_0^{\frac{1}{2}} (2^{1-\alpha} - 2t^\alpha) T_f^t(a, b) dt + \int_{\frac{1}{2}}^1 (2t^\alpha - 1) T_f^t(a, b) dt \right] \\
&= \frac{\ln b - \ln a}{2} b |f'(b)| \left[\int_0^1 T_f^t(a, b) dt + \int_0^{\frac{1}{2}} 2^{1-\alpha} T_f^t(a, b) dt \right. \\
&- \left. 2 \int_0^{\frac{1}{2}} t^\alpha T_f^t(a, b) dt - \int_{\frac{1}{2}}^1 T_f^t(a, b) dt + 2 \int_{\frac{1}{2}}^1 t^\alpha T_f^t(a, b) dt \right] \\
&= \frac{\ln b - \ln a}{2} b |f'(b)| \left[\int_0^1 T_f(a, b) dt + 2^{1-\alpha} \int_0^{\frac{1}{2}} T_f(a, b) dt \right. \\
&- \left. \int_{\frac{1}{2}}^1 T_f(a, b) dt - 4 \int_0^{\frac{1}{2}} t^\alpha T_f(a, b) dt + 2 \int_0^1 t^\alpha T_f(a, b) dt \right] =
\end{aligned}$$

$$\begin{aligned}
&= \frac{\ln b - \ln a}{2} b |f'(b)| \left[\frac{2T_f(a, b) - 1 + (2^{1-\alpha} + 1)T_f^{\frac{1}{2}}(a, b)}{\ln(T_f(a, b))} \right. \\
&\quad \left. - 4I_f(\alpha + 1, a, b, \frac{1}{2}) + 2I_f(\alpha + 1, a, b, a) \right].
\end{aligned}$$

Case 2: $\alpha \in [1, \infty)$. By using Definition 2.1, Lemma 1.2, Lemma 2.4, Lemma 2.6, Lemma 2.7, Hölder inequality and Lemma 2.5 again, we have

$$\begin{aligned}
&\left| \frac{\Gamma(\alpha + 1)}{2(\ln b - \ln a)^\alpha} [{}_H J_{b^-}^\alpha f(a) + {}_H J_{a^+}^\alpha f(b)] - f(\sqrt{ab}) \right| \\
&= \left| \frac{\ln b - \ln a}{2} \left[\int_0^1 k a^t b^{1-t} f'(a^t b^{1-t}) dt \right. \right. \\
&\quad \left. \left. - \int_0^1 [(1-t)^\alpha - t^\alpha] a^t b^{1-t} f'(a^t b^{1-t}) dt \right] \right| \\
&\leq \frac{\ln b - \ln a}{2} \left[\int_0^1 a^t b^{1-t} |f'(a^t b^{1-t})| dt \right. \\
&\quad \left. + \int_0^1 |(1-t)^\alpha - t^\alpha| a^t b^{1-t} |f'(a^t b^{1-t})| dt \right] \\
&\leq \frac{\ln b - \ln a}{2} \left[\int_0^1 a^t b^{1-t} |f'(a)|^t |f'(b)|^{1-t} dt \right. \\
&\quad \left. + \int_0^1 |(1-t)^\alpha - t^\alpha| a^t b^{1-t} |f'(a)|^t |f'(b)|^{1-t} dt \right] \\
&= \frac{\ln b - \ln a}{2} \left[b |f'(b)| \int_0^1 T_f^t(a, b) dt \right. \\
&\quad \left. + b |f'(b)| \int_0^1 |(1-t)^\alpha - t^\alpha| T_f^t(a, b) dt \right] \\
&= \frac{\ln b - \ln a}{2} b |f'(b)| \left[\int_0^1 T_f^t(a, b) dt + \int_0^1 |(1-t)^\alpha - t^\alpha| T_f^t(a, b) dt \right] \\
&= \frac{\ln b - \ln a}{2} b |f'(b)| \left[\int_0^1 T_f^t(a, b) dt + \int_0^{\frac{1}{2}} ((1-t)^\alpha - t^\alpha) T_f^t(a, b) dt \right. \\
&\quad \left. + \int_{\frac{1}{2}}^1 (-(1-t)^\alpha + t^\alpha) T_f^t(a, b) dt \right] \\
&= \frac{\ln b - \ln a}{2} b |f'(b)| \left[\int_0^1 T_f^t(a, b) dt + \int_0^{\frac{1}{2}} (1-t)^\alpha T_f^t(a, b) dt \right. \\
&\quad \left. - \int_0^{\frac{1}{2}} t^\alpha T_f^t(a, b) dt - \int_{\frac{1}{2}}^1 (1-t)^\alpha T_f^t(a, b) dt + \int_{\frac{1}{2}}^1 t^\alpha T_f^t(a, b) dt \right] \\
&\leq \frac{\ln b - \ln a}{2} b |f'(b)| \left[\int_0^1 T_f^t(a, b) dt + \int_0^{\frac{1}{2}} (1-t)^\alpha T_f^t(a, b) dt - \right.
\end{aligned}$$

$$\begin{aligned}
& - \int_0^{\frac{1}{2}} t^\alpha T_f^t(a, b) dt - \int_{\frac{1}{2}}^1 (2^{1-\alpha} - t^\alpha) T_f^t(a, b) dt + \int_{\frac{1}{2}}^1 t^\alpha T_f^t(a, b) dt \Big] \\
& = \frac{\ln b - \ln a}{2} b \left[\int_0^1 T_f^t(a, b) dt + \int_0^{\frac{1}{2}} (1 - 2t^\alpha) T_f^t(a, b) dt \right. \\
& \quad \left. + \int_{\frac{1}{2}}^1 (-1 + 2t^\alpha) T_f^t(a, b) dt \right] \\
& = \frac{\ln b - \ln a}{2} b |f'(b)| \left[\int_0^1 T_f^t(a, b) dt + \int_0^{\frac{1}{2}} T_f^t(a, b) dt - \int_{\frac{1}{2}}^1 T_f^t(a, b) dt \right. \\
& \quad \left. - 2 \int_0^{\frac{1}{2}} t^\alpha T_f^t(a, b) dt + 2 \int_0^1 t^\alpha T_f^t(a, b) dt - 2 \int_0^{\frac{1}{2}} t^\alpha T_f^t(a, b) dt \right] \\
& = \frac{\ln b - \ln a}{2} b |f'(b)| \left[\int_0^1 T_f^t(a, b) dt + \int_0^{\frac{1}{2}} T_f^t(a, b) dt \right. \\
& \quad \left. - \int_{\frac{1}{2}}^1 T_f^t(a, b) dt - 4 \int_0^{\frac{1}{2}} t^\alpha T_f^t(a, b) dt + 2 \int_0^1 t^\alpha T_f^t(a, b) dt \right] \\
& = \frac{\ln b - \ln a}{2} b |f'(b)| \left[\frac{2T_f^{\frac{1}{2}}(a, b) - 1}{\ln(T_f(a, b))} - 4I_f(\alpha + 1, a, b, \frac{1}{2}) + 2I_f(\alpha + 1, a, b, a) \right].
\end{aligned}$$

The proof is done. \square

Corollary 3.4. *Under the conditions of Theorem 3.3, if $x|f'(x)|$ is decreasing, then*

$$\begin{aligned}
& \left| \frac{\Gamma(\alpha + 1)}{2(\ln b - \ln a)^\alpha} [{}_H J_b^\alpha f(a) + {}_H J_{a^+}^\alpha f(b)] - f(\sqrt{ab}) \right| \leq \\
& \leq \max \left\{ \frac{\ln b - \ln a}{2} b |f'(b)| \left[\frac{2T_f(a, b) - 1 + (2^{1-\alpha} + 1)T_f^{\frac{1}{2}}(a, b)}{\ln(T_f(a, b))} \right. \right. \\
& \quad \left. \left. - 4I_f(\alpha + 1, a, b, \frac{1}{2}) + 2\frac{T_f(a, b)}{\alpha + 1} \right], \right. \\
& \quad \left. \frac{\ln b - \ln a}{2} b |f'(b)| \left[\frac{2T_f^{\frac{1}{2}}(a, b) - 1}{\ln(T_f(a, b))} - 4I_f(\alpha + 1, a, b, \frac{1}{2}) + 2\frac{T_f(a, b)}{\alpha + 1} \right] \right\}.
\end{aligned}$$

Proof. Note that $T_f^t(a, b) \leq T_f(a, b)$ for $t \in [0, 1]$ gives

$$\int_0^1 t^\alpha T_f^t(a, b) dt \leq \int_0^1 t^\alpha T_f(a, b) dt = \frac{T_f(a, b)}{\alpha + 1}.$$

Then, the result can be derived immediately. \square

Theorem 3.5. *Let $f : [0, b] \rightarrow R$ be a differentiable mapping and $1 < q < \infty$. If $|f'|^q$ is measurable and $|f'|^q$ is GG-convex on $[0, b]$ for some fixed $\alpha \in (0, \infty)$, $0 \leq a < b$, then the following inequality for fractional integrals holds*

$$\begin{aligned} & \left| \frac{\Gamma(\alpha + 1)}{2(\ln b - \ln a)^\alpha} [{}_H J_b^\alpha f(a) + {}_H J_a^\alpha f(b)] - f(\sqrt{ab}) \right| \\ & \leq \max \left\{ \frac{\ln b - \ln a}{2} \cdot b \cdot |f'(b)| \cdot \left(\frac{(1 + 2^{1-\alpha})^p}{2} - \frac{2^p(1 - 2^{-p\alpha})}{p\alpha + 1} \right)^{\frac{1}{p}} \right. \\ & \quad \cdot \left(\frac{T_f^q(a, b)}{q \ln T_f(a, b)} \right)^{\frac{1}{q}}, \frac{\ln b - \ln a}{2} \cdot b \cdot |f'(b)| \cdot \left(2^p - \frac{2^{p+1} - 2^{p(1-\alpha)}}{p\alpha + 1} \right)^{\frac{1}{p}} \\ & \quad \cdot \left. \left(\frac{T_f^q(a, b)}{q \ln T_f(a, b)} \right)^{\frac{1}{q}} \right\}, \end{aligned}$$

where $\frac{1}{p} + \frac{1}{q} = 1$.

Proof. To achieve our aim, we divide into two cases.

Case 1: $\alpha \in (0, 1)$. For

$$k = \begin{cases} 1, & 0 \leq t < \frac{1}{2}, \\ -1, & \frac{1}{2} \leq t < 1. \end{cases}$$

By using Lemma 2.6, Lemma 2.7, we obtain

$$\begin{aligned} & \int_0^1 |k - (1-t)^\alpha + t^\alpha|^p dt \\ & = \int_0^{\frac{1}{2}} (1 - (1-t)^\alpha + t^\alpha)^p dt + \int_{\frac{1}{2}}^1 (1 - t^\alpha + (1-t)^\alpha)^p dt \\ & \leq \int_0^{\frac{1}{2}} [(1+t^\alpha) - (1-t^\alpha)]^p dt + \int_{\frac{1}{2}}^1 [1 - t^\alpha + 2^{1-\alpha} - t^\alpha]^p dt \\ & = \int_0^{\frac{1}{2}} 2^p t^{p\alpha} dt + \int_{\frac{1}{2}}^1 [(1 + 2^{1-\alpha})^p - 2^p t^{p\alpha}] dt \\ & = \frac{(1 + 2^{1-\alpha})^p}{2} - \frac{2^p(1 - 2^{-p\alpha})}{p\alpha + 1}. \end{aligned}$$

Case 2: $\alpha \in [1, \infty)$. By using Lemma 2.6 and Lemma 2.7, we obtain

$$\int_0^1 |k - (1-t)^\alpha + t^\alpha|^p dt =$$

$$\begin{aligned}
&= \int_0^{\frac{1}{2}} (1 - (1-t)^\alpha + t^\alpha)^p dt + \int_{\frac{1}{2}}^1 (1 - t^\alpha + (1-t)^\alpha)^p dt \\
&= 2 \int_{\frac{1}{2}}^1 (1 - t^\alpha + (1-t)^\alpha)^p dt \leq 2 \int_{\frac{1}{2}}^1 (1 - t^\alpha + 1 - t^\alpha)^p dt \\
&\leq 2^{p+1} \int_{\frac{1}{2}}^1 (1 - t^\alpha)^p dt \leq 2^{p+1} \int_{\frac{1}{2}}^1 (1 - t^{\alpha p}) dt \\
&= 2^{p+1} \left(\frac{1}{2} - \int_{\frac{1}{2}}^1 t^{p\alpha} dt \right) \leq 2^p - \frac{2^{p+1} - 2^{p(1-\alpha)}}{p\alpha + 1}.
\end{aligned}$$

Since $|f'(x)|^q$ is GG -convex on $[0, b]$, from Definition 2.1, we drive

$$\begin{aligned}
\int_0^1 (a^t b^{1-t} |f'(a^t b^{1-t})|)^q dt &\leq \int_0^1 (a^t b^{1-t})^q |f'(a)|^{tq} |f'(b)|^{q(1-t)} dt \\
&= (b \cdot |f'(b)|)^q \int_0^1 T_f^q(a, b) dt \\
&= (b \cdot |f'(b)|)^q \frac{T_f^q(a, b)}{q \ln T_f(a, b)}.
\end{aligned}$$

Therefore, by Lemma 2.7 and Hölder inequality, we have

$$\begin{aligned}
&\left| \frac{\Gamma(\alpha + 1)}{2(\ln b - \ln a)^\alpha} [{}_H J_{b^-}^\alpha f(a) + {}_H J_{a^+}^\alpha f(b)] - f(\sqrt{ab}) \right| \\
&= \left| \frac{\ln b - \ln a}{2} \left[\int_0^1 (k - (1-t)^\alpha + t^\alpha) a^t b^{1-t} f'(a^t b^{1-t}) dt \right] \right| \\
&\leq \frac{\ln b - \ln a}{2} \int_0^1 |k - (1-t)^\alpha + t^\alpha| \cdot a^t b^{1-t} \cdot |f'(a^t b^{1-t})| dt \\
&\leq \frac{\ln b - \ln a}{2} \left(\int_0^1 |k - (1-t)^\alpha + t^\alpha|^p dt \right)^{\frac{1}{p}} \left(\int_0^1 (a^t b^{1-t} |f'(a^t b^{1-t})|)^q dt \right)^{\frac{1}{q}} \\
&\leq \max \left\{ \frac{\ln b - \ln a}{2} \cdot b \cdot |f'(b)| \cdot \left(\frac{(1 + 2^{1-\alpha})^p}{2} - \frac{2^p(1 - 2^{-p\alpha})}{p\alpha + 1} \right)^{\frac{1}{p}} \right. \\
&\quad \cdot \left(\frac{T_f^q(a, b)}{q \ln T_f(a, b)} \right)^{\frac{1}{q}}, \frac{\ln b - \ln a}{2} \cdot b \cdot |f'(b)| \cdot \left(2^p - \frac{2^{p+1} - 2^{p(1-\alpha)}}{p\alpha + 1} \right)^{\frac{1}{p}} \\
&\quad \left. \cdot \left(\frac{T_f^q(a, b)}{q \ln T_f(a, b)} \right)^{\frac{1}{q}} \right\}.
\end{aligned}$$

The proof is done. □

4. Applications to special means

Consider the following special means (see [27]) for arbitrary real numbers $x, y, x \neq y$ as follows:

- (i) $A(x, y) = \frac{x+y}{2}, x, y \in R.$
- (ii) $H(x, y) = \frac{2}{\frac{1}{x} + \frac{1}{y}}, x, y \in R \setminus \{0\}.$
- (iii) $G(x, y) = \sqrt{xy}.$
- (iv) $L(x, y) = \frac{y-x}{\ln|y| - \ln|x|}, |x| \neq |y|, xy \neq 0.$
- (v) $L_n(x, y) = \left[\frac{y^{n+1} - x^{n+1}}{(n+1)(y-x)} \right]^{\frac{1}{n}}, n \in Z \setminus \{-1, 0\}, x, y \in R, x \neq y.$

Using the results obtained in Section 3, we give some applications to special means of real numbers.

Proposition 4.1. *Let $a, b \in R^+ \setminus \{0\}, 0 \leq a < b, x \in [0, b].$ Then*

$$\begin{aligned} \left| A(x, x) - L(a, b) \right| &\leq \frac{(2 \ln x - \ln a - \ln b - 2)x + a + b}{\ln b - \ln a}, \\ \left| A(x, x) - L(a, b) \right| &\leq \frac{(\ln x - \ln a)^2}{\ln b - \ln a} a \left(\frac{1}{p+1} \right)^{\frac{1}{p}} \left(\frac{\left(\frac{x}{a}\right)^q - 1}{q(\ln x - \ln a)} \right)^{\frac{1}{q}} \\ &\quad + \frac{(\ln b - \ln x)^2}{\ln b - \ln a} b \left(\frac{1}{p+1} \right)^{\frac{1}{p}} \left(\frac{\left(\frac{x}{b}\right)^q - 1}{q(\ln x - \ln b)} \right)^{\frac{1}{q}}, \\ \left| L(a, b) - [A(a, b)H(a, b)]^{\frac{1}{2}} \right| &\leq a - b - \frac{(\sqrt{a} + \sqrt{b})^2}{\ln a - \ln b}, \\ \left| L(a, b) - [A(a, b)H(a, b)]^{\frac{1}{2}} \right| &\leq \frac{a(\ln b - \ln a)}{2(q(\ln a - \ln b))^{\frac{1}{q}}} \left(2^p - \frac{2^{p+1} - 1}{p+1} \right)^{\frac{1}{p}}. \end{aligned}$$

Proof. Applying Theorems 3.1, 3.2, 3.3, and 3.5 for $f(x) = x$ and $\alpha = 1$, one can obtain the results immediately. □

Proposition 4.2. *Let $a, b \in R^+ \setminus \{0\}, 0 \leq a < b, x \in [0, b], n \geq 2.$ Then*

$$\begin{aligned} \left| A(x^n, x^n) - L(a, b)L_{n-1}^{n-1}(a, b) \right| &\leq \left(1 - \frac{2}{n(\ln b - \ln a)} \right) x^n - \frac{a^n + b^n}{n(\ln b - \ln a)}, \\ \left| A(x^n, x^n) - L(a, b)L_{n-1}^{n-1}(a, b) \right| &\leq \frac{n \left(\frac{1}{p+1}\right)^{\frac{1}{p}}}{\ln b - \ln a} \left[\left(\frac{(\ln x - \ln a)^{2q-1} (x^{qn} - a^{qn})}{qn} \right)^{\frac{1}{q}} \right. \\ &\quad \left. + \left(\frac{(\ln x - \ln b)^{2q-1} (x^{qn} - b^{qn})}{qn} \right)^{\frac{1}{q}} \right], \end{aligned}$$

$$\left| L(a^n, b^n) - (H(a, b)A(a, b))^{\frac{n}{2}} \right| \leq -nb^n \left(\left(\frac{a}{b} \right)^{\frac{n}{2}} - 1 + \frac{2 \left(\frac{a}{b} \right)^{\frac{n}{2}} - \left(\frac{a}{b} \right)^n - 1}{n^2(\ln a - \ln b)} \right),$$

$$\left| L(a^n, b^n) - (H(a, b)A(a, b))^{\frac{n}{2}} \right| \leq \frac{\ln b - \ln a}{2(qn(\ln a - \ln b))^{\frac{1}{q}}} na^n \left(2^p - \frac{2^{p+1} - 1}{p+1} \right)^{\frac{1}{p}}.$$

Proof. Applying Theorems 3.1, 3.2, 3.3, and 3.5 for $f(x) = x^n$ and $\alpha = 1$, one can obtain the results immediately. \square

Proposition 4.3. *Let $a, b \in \mathbb{R}^+ \setminus \{0\}, 0 \leq a < b, x \in [0, b], n \geq 2$. Then*

$$\left| A\left(\frac{1}{x}, \frac{1}{x}\right) - \frac{1}{2}L\left(\frac{1}{a}, \frac{1}{b}\right) \right| \leq \frac{2(\ln a + \ln b) - 4\ln x - 4}{x(\ln b - \ln a)} - \frac{2}{a(\ln b - \ln a)},$$

$$\left| A\left(\frac{1}{x}, \frac{1}{x}\right) - \frac{1}{2}L\left(\frac{1}{a}, \frac{1}{b}\right) \right| \leq \frac{2(\ln x - \ln a)^2}{a(\ln b - \ln a)} \left(\frac{1}{p+1} \right)^{\frac{1}{p}} \left(\frac{\left(\frac{a}{x}\right)^q - 1}{q(\ln a - \ln x)} \right)^{\frac{1}{q}}$$

$$+ \frac{2(\ln b - \ln x)^2}{b(\ln b - \ln a)} \left(\frac{1}{p+1} \right)^{\frac{1}{p}} \left(\frac{\left(\frac{b}{x}\right)^q - 1}{q(\ln b - \ln x)} \right)^{\frac{1}{q}},$$

$$\left| L\left(\frac{1}{b}, \frac{1}{a}\right) - G\left(\frac{1}{a}, \frac{1}{b}\right) \right| \leq -\frac{1}{2b} \left(-1 + \frac{b}{a} + \frac{2\left(\frac{b}{a}\right)^{\frac{1}{2}} - \frac{b}{a} - 1}{\ln b - \ln a} \right),$$

$$\left| L\left(\frac{1}{b}, \frac{1}{a}\right) - G\left(\frac{1}{a}, \frac{1}{b}\right) \right| \leq \frac{\ln b - \ln a}{4a(q(\ln b - \ln a))^{\frac{1}{q}}} \left(2^p - \frac{2^{p+1} - 1}{p+1} \right)^{\frac{1}{p}}.$$

Proof. Applying Theorems 3.1, 3.2, 3.3, and 3.5 for $f(x) = \frac{1}{x}$ and $\alpha = 1$, one can obtain the results immediately. \square

Proposition 4.4. *Let $a, b \in \mathbb{R}^+ \setminus \{0\}, 0 \leq a < b, x \in [0, b], n \geq 2$. Then*

$$\left| A(x, x) - L(b^{-1}, a^{-1}) \right|$$

$$\leq \frac{(2\ln x + \ln a + \ln b - 2)x + a^{-1} + b^{-1}}{\ln b - \ln a},$$

$$\left| A(x, x) - L(b^{-1}, a^{-1}) \right|$$

$$\leq \frac{(\ln x + \ln b)^2}{\ln b - \ln a} b^{-1} \left(\frac{1}{p+1} \right)^{\frac{1}{p}} \left(\frac{(xb)^q - 1}{q(\ln x + \ln b)} \right)^{\frac{1}{q}}$$

$$+ \frac{(\ln a + \ln x)^2}{\ln b - \ln a} a^{-1} \left(\frac{1}{p+1} \right)^{\frac{1}{p}} \left(\frac{(xa)^q - 1}{q(\ln x + \ln a)} \right)^{\frac{1}{q}},$$

$$\begin{aligned}
& \left| L(b^{-1}, a^{-1}) - [A(b^{-1}, a^{-1})H(b^{-1}, a^{-1})]^{\frac{1}{2}} \right| \\
& \leq -a^{-1} + b^{-1} - \frac{(\sqrt{a^{-1}} + \sqrt{b^{-1}})^2}{\ln a - \ln b}, \\
& \left| L(b^{-1}, a^{-1}) - [A(b^{-1}, a^{-1})H(b^{-1}, a^{-1})]^{\frac{1}{2}} \right| \\
& \leq \frac{\ln b - \ln a}{2} b^{-1} \left(2^p - \frac{2^{p+1} - 1}{p+1} \right)^{\frac{1}{p}} \left(\frac{1}{q(\ln a - \ln b)} \right)^{\frac{1}{q}}. \\
& \left| A(x^n, x^n) - L(b^{-1}, a^{-1})L_{n-1}^{n-1}(b^{-1}, a^{-1}) \right| \\
& \leq \left(1 - \frac{2}{n(\ln b - \ln a)} \right) x^n - \frac{a^{-n} + b^{-n}}{n(\ln b - \ln a)}, \\
& \left| A(x^n, x^n) - L(b^{-1}, a^{-1})L_{n-1}^{n-1}(b^{-1}, a^{-1}) \right| \\
& \leq \frac{n \left(\frac{1}{p+1} \right)^{\frac{1}{p}}}{\ln b - \ln a} \left[\left((\ln x + \ln b)^{2q-1} (x^{qn} - b^{-qn}) \cdot \frac{1}{qn} \right)^{\frac{1}{q}} \right. \\
& \quad \left. + \left((\ln x + \ln a)^{2q-1} (x^{qn} - a^{-qn}) \cdot \frac{1}{qn} \right)^{\frac{1}{q}} \right], \\
& \left| L(b^{-n}, a^{-n}) - (H(b^{-1}, a^{-1})A(b^{-1}, a^{-1}))^{\frac{n}{2}} \right| \\
& \leq -na^{-n} \left(\left(\frac{a}{b} \right)^{\frac{n}{2}} - 1 + \frac{2 \left(\frac{a}{b} \right)^{\frac{n}{2}} - \left(\frac{a}{b} \right)^n - 1}{n^2(\ln a - \ln b)} \right), \\
& \left| L(b^{-n}, a^{-n}) - (H(b^{-1}, a^{-1})A(b^{-1}, a^{-1}))^{\frac{n}{2}} \right| \\
& \leq \frac{\ln b - \ln a}{2} nb^{-n} \left(2^p - \frac{2^{p+1} - 1}{p+1} \right)^{\frac{1}{p}} \left(\frac{1}{qn(\ln a - \ln b)} \right)^{\frac{1}{q}}. \\
& \left| A\left(\frac{1}{x}, \frac{1}{x}\right) - \frac{1}{2}L(b, a) \right| \leq \frac{-2(\ln a + \ln b) - 4\ln x - 4}{x(\ln b - \ln a)} - \frac{2b}{\ln b - \ln a}, \\
& \left| A\left(\frac{1}{x}, \frac{1}{x}\right) - \frac{1}{2}L(b, a) \right| \leq \frac{2b(\ln x + \ln b)^2}{(\ln b - \ln a)(p+1)^{\frac{1}{p}}} \left(\frac{\left(\frac{1}{xb}\right)^q - 1}{q(-\ln b - \ln x)} \right)^{\frac{1}{q}} \\
& \quad + \frac{2a(\ln a + \ln x)^2}{(\ln b - \ln a)(p+1)^{\frac{1}{p}}} \left(\frac{\left(\frac{1}{xa}\right)^q - 1}{q(-\ln a - \ln x)} \right)^{\frac{1}{q}},
\end{aligned}$$

$$\left| L(a, b) - G(b, a) \right| \leq -2a \left(-1 + \frac{b}{a} + \frac{2 \left(\frac{b}{a} \right)^{\frac{1}{2}} - \frac{b}{a} - 1}{\ln b - \ln a} \right),$$

$$\left| L(a, b) - G(b, a) \right| \leq \frac{b(\ln b - \ln a)}{4(q(\ln b - \ln a))^{\frac{1}{q}}} \left(2^p - \frac{2^{p+1} - 1}{p+1} \right)^{\frac{1}{p}}.$$

Proof. Making the substitutions $a \rightarrow b^{-1}$, $b \rightarrow a^{-1}$ in the Proposition 4.1, Proposition 4.2 and Proposition 4.3, one can obtain desired inequalities respectively. \square

5. Acknowledgement.

We acknowledge the valuable comments and suggestions of the anonymous referee and editor.

REFERENCES

- [1] A. A. Kilbas - H. M. Srivastava - J. J. Trujillo, *Theory and applications of fractional differential equations*, Elsevier Science B.V., 2006.
- [2] D. Baleanu - J. A. T. Machado - A. C. J. Luo, *Fractional dynamics and control*, Springer, 2012.
- [3] K. Diethelm, *The analysis of fractional differential equations*, Lecture Notes in Mathematics, 2010.
- [4] V. Lakshmikantham - S. Leela - J. V. Devi, *Theory of fractional dynamic systems*, Cambridge Scientific Publishers, 2009.
- [5] K. S. Miller - B. Ross, *An introduction to the fractional calculus and differential equations*, John Wiley, 1993.
- [6] M. W. Michalski, *Derivatives of noninteger order and their applications*, Dissertationes Mathematicae, CCCXXVIII, Inst. Math., Polish Acad. Sci., 1993.
- [7] I. Podlubny, *Fractional differential equations*, Academic Press, 1999.
- [8] V. E. Tarasov, *Fractional dynamics. Application of fractional calculus to dynamics of particles, fields and media*, Nonlinear Physical Science. Springer, Heidelberg; HEP, Beijing, 2010.
- [9] M. Z. Sarikaya - E. Set - H. Yaldiz - N. Başak, *Hermite-Hadamard's inequalities for fractional integrals and related fractional inequalities*, Math. Comput. Model. 57 (2013), 2403–2407.
- [10] C. Zhu - M. Fečkan J. Wang, *Fractional integral inequalities for differentiable convex mappings and applications to special means and a midpoint formula*, J. Appl. Math. Statistics Inform. 8 (2012), 21–28.

- [11] M. A. Latif - S. S. Dragomir - A. E. Matouk, *New inequalities of Ostrowski type for co-ordinated convex functions via fractional integrals*, J. Frac. Calc. Appl. 2 (2012), 1–15.
- [12] J. Wang - X. Li - C. Zhu, *Refinements of Hermite-Hadamard type inequalities involving fractional integrals*, Bull. Belg. Math. Soc. Simon Stevin 20 (2013), 655–666.
- [13] E. Set, *New inequalities of Ostrowski type for mappings whose derivatives are s -convex in the second sense via fractional integrals*, Comput. Math. Appl. 63 (2012), 1147–1154.
- [14] J. Wang - J. Deng - M. Fečkan, *Hermite-Hadamard type inequalities for r -convex functions via Riemann-Liouville fractional integrals*, Ukrainian Math. J. 65 (2013), 193–211.
- [15] Y. Zhang - J. Wang, *On some new Hermite-Hadamard inequalities involving Riemann-Liouville fractional integrals*, J. Inequal. Appl. 2013 (2013):220, 1–27.
- [16] J. Wang - X. Li - M. Fečkan - Y. Zhou, *Hermite-Hadamard-type inequalities for Riemann-Liouville fractional integrals via two kinds of convexity*, Appl. Anal. 92 (2013), 2241–2253.
- [17] J. Wang - J. Deng - M. Fečkan, *Exploring s - e -condition and applications to some Ostrowski type inequalities via Hadamard fractional integrals*, Math. Slovaca, (2013), in press.
- [18] J. Wang - C. Zhu - Y. Zhou, *New generalized Hermite-Hadamard type inequalities and applications to special means*, J. Inequal. Appl. 2013 (2013):325, 1–15.
- [19] J. Deng - J. Wang, *Fractional Hermite-Hadamard inequalities for (α, m) -logarithmically convex functions*, J. Inequal. Appl. 2013 (2013):364, 1–11.
- [20] Y. Liao - J. Deng - J. Wang, *Riemann-Liouville fractional Hermite-Hadamard inequalities. Part I: for once differentiable geometric-arithmetically s -convex functions*, J. Inequal. Appl. 2013 (2013):443, 1–13.
- [21] Y. Liao - J. Deng - J. Wang, *Riemann-Liouville fractional Hermite-Hadamard inequalities. Part II: for twice differentiable geometric-arithmetically s -convex functions*, J. Inequal. Appl. 2013 (2013):517, 1–13.
- [22] I. İşcan, *New general integral inequalities for quasi-geometrically convex functions via fractional integrals*, J. Inequal. Appl. 2013 (2013):491, 1–15.
- [23] I. İşcan, *Generalization of different type integral inequalities for s -convex functions via fractional integrals*, Appl. Anal. 6 (2014), 1846–1862.
- [24] C. P. Niculescu, *Convexity according to means*, Math. Inequal. Appl. 6 (2003), 571–579.
- [25] R. A. Satnoianu, *Improved GA-convexity inequalities*, J. Inequal. Pure Appl. Math. 3 (2002), Article 82, 1–6.
- [26] G. D. Anderson - M. K. Vamanamurthy - M. Vuorinen, *Generalized convexity and inequalities*, J. Math. Anal. Appl. 335 (2007), 1294–1308.

- [27] C. E. M. Pearce - J. E. Pečarić, *Inequalities for differentiable mappings with application to special means and quadrature formula*, Appl. Math. Lett. 13 (2000), 51–55.

SHENGDA LIU

*Department of Mathematics, College of Science
Guizhou University, Guiyang
Guizhou 550025, P.R.China
e-mail: thinksheng@foxmail.com*

JINRONG WANG

*Department of Mathematics, College of Science
Guizhou University, Guiyang
Guizhou 550025, P.R.China
School of Mathematics and Computer Science
Guizhou Normal College, Guiyang
Guizhou 550018, P.R. China
e-mail: sci.jrwang@gzu.edu.cn*