

LE MATEMATICHE

Vol. LXIX (2014) – Fasc. I, pp. 169–178

doi: 10.4418/2014.69.1.12

SOME PROPERTIES OF SKEW HURWITZ SERIES

A. M. HASSANEIN - MOHAMED A. FARAHAT

In this paper we show that, if R is a ring and σ an endomorphism of R , then the *skew Hurwitz series ring* $T = (HR, \sigma)$ is an n -clean ring if and only if R is an n -clean ring. Moreover, if R is an integral domain and a torsion-free \mathbb{Z} -module, then $T = (HR, \sigma)$ is a Prüfer domain if and only if R is a field. Also, we investigate when the ring $T = (HR, \sigma)$ is $g(x)$ -clean, $(n, g(x))$ -clean and a Neat ring.

1. Introduction

Throughout this paper R is an associative ring with identity 1, $U(R)$ its group of units, $Id(R)$ its set of idempotents and $C(R)$ its center and σ an endomorphism of the ring R .

In a series of papers ([15], [16], [17]) Keigher demonstrated that the ring HR of *Hurwitz series* over a commutative ring R with identity has many interesting applications in differential algebra.

Some properties which are shared between R and HR have been studied by Keigher [17], Zhongkui [24], Hassanein, et al in [12, 13], Benhissi [1, 2] and Ghanem [5].

The concept of Hurwitz series was extended by Hassanein in [11] to the ring of *skew Hurwitz series* as follows: the elements of $T = (HR, \sigma)$, the ring of *skew Hurwitz series*, are the ordinary functions $f: \mathbb{N} \rightarrow R$ with component wise

Entrato in redazione: 8 marzo 2013

AMS 2010 Subject Classification: Primary 16E50; Secondary 16U99, 16S70.

Keywords: Clean rings, n -clean rings, $g(x)$ -clean rings, $(n, g(x))$ -clean rings, Prüfer domain.

addition and the following operation of multiplication: For each two functions $f, g \in T = (HR, \sigma)$,

$$(fg)(n) = \sum_{k=0}^n \binom{n}{k} f(k) \sigma^k(g(n-k)).$$

Define the mappings $h_n: \mathbb{N} \rightarrow R$ via $h_n(n-1) = 1$ and $h_n(m) = 0$ for each $m \neq n-1$ in \mathbb{N} . And $h'_r: \mathbb{N} \rightarrow R$ via $h'_r(0) = r$ and $h'_r(n) = 0$ for each $0 \neq n$ in \mathbb{N} and $r \in R$. It can be easily shown that $T = (HR, \sigma)$ is a ring with identity h_1 , defined by $h_1: \mathbb{N} \rightarrow R$ via $h_1(0) = 1$ and $h_1(n) = 0$ for each $n \neq 0$ in \mathbb{N} and $1 \in R$.

There is a ring homomorphism $\lambda_R: R \rightarrow T = (HR, \sigma)$ defined for any $r \in R$ by $\lambda_R(r) = h'_r$. So, the ring R is canonically embedded as a subring of T via $r \in R \mapsto h'_r \in T$. Note also that there is a ring homomorphism $\varepsilon_R: T = (HR, \sigma) \rightarrow R$ defined for any $f \in T = (HR, \sigma)$ by $\varepsilon_R(f) = f(0)$. Clearly, $\varepsilon_R \circ \lambda_R = \text{id}_R$.

Let $\text{supp}(f)$ denote the support of $f \in T = (HR, \sigma)$, i.e.,

$$\text{supp}(f) = \{i \in \mathbb{N} \mid 0 \neq f(i) \in R\},$$

$\pi(f)$ denote the minimal element in $\text{supp}(f)$. See [10] for more details.

Recently, Hassanein [10, 11, 14], Handam [9] and Yu-juan, et al [23] studied the transfer of some algebraic properties between R and $T = (HR, \sigma)$.

The motivation of this paper is to show that and extend the results in [5] to the ring $T = (HR, \sigma)$ of skew Hurwitz series over the ring R . Neat skew Hurwitz rings are also considered.

2. n -clean skew Hurwitz ring.

An element $r \in R$ is called *clean* if it can be expressed as a sum of an idempotent and a unit in R . This definition was introduced by Nicholson [19].

According to Xiao and Tong [21], an element x of a ring R is called n -clean, where n is a positive integer, if $x = e + u_1 + u_2 + \dots + u_n$ where $e \in \text{Id}(R)$ and $u_i \in U(R)$; $i = 1, 2, \dots, n$. The ring R is called n -clean if every element of R is n -clean for some fixed positive integer n .

We need the following construction. Let R be a ring and let ${}_R V_R$ be an R -bimodule. Then the ideal extension $I(R; V)$ of R by V is defined to be the additive abelian group $I(R; V) = R \oplus V$ with multiplication given as follows: for all $v, w \in V$ and $r, s \in R$, we get,

$$(r, v)(s, w) = (rs, rw + vs + vw).$$

Note that if S is a ring and $S = R \oplus A$, where R is a subring of S and A is a two sided ideal of S , then $S \cong I(R; A)$.

Proposition 2.1. *Let R be a ring and σ an endomorphism of the ring R , then:*

1) $A = \{f \in T \mid f(0) = 0\}$ is a two sided ideal of T .

2) For each two sided σ -ideal I of R we have $H_I = \{h'_r \in T \mid r \in I\}$ is a two sided ideal in T and

$$(HR, \sigma) / (H_I + A) \cong (H(R/I), \sigma).$$

In particular, if I is a maximal σ -ideal of R , then $H_I + A$ is a maximal σ -ideal of T .

Proof. The proof of (1) is clear and that of (2) follows from Proposition 3.2 in [10]. □

Proposition 2.2 ([11]). *Let R be a ring and $\sigma \in \text{End}(R)$. Then $T = (HR, \sigma) \cong I(R; A)$, where $A = \{f \in T \mid f(0) = 0\}$ is a two sided ideal of T .*

In the following Theorem shows us how the n -clean property shared between R and $T = (HR, \sigma)$.

Theorem 2.3. *Let R be a ring and $\sigma \in \text{End}(R)$. Then $T = (HR, \sigma)$ is an n -clean ring if and only if R is an n -clean ring.*

Proof. Since $\langle h_2 \rangle = Th_2 = \{fh_2 \mid f \in T\}$ is an ideal of T and clearly $(fh_2)(0) = 0$, by Proposition 2.2, we have $T \cong I(R; \langle h_2 \rangle)$. Since $R \cong T / \langle h_2 \rangle$, by Proposition 2.4 in [21], we conclude that if $T = (HR, \sigma)$ is an n -clean ring, then its homomorphic image R is.

Conversely, suppose that R is an n -clean ring and $f \in T$, hence $f(0) \in R$, therefore we can write

$$f(0) = e + u_1 + u_2 + \dots + u_n,$$

where $e \in \text{Id}(R)$ and $u_i \in U(R); i = 1, 2, \dots, n$. Then

$$f = h'_e + g + h'_{u_2} + \dots + h'_{u_n}$$

where $g \in T$ defined by

$$g(0) = u_1 \text{ and } g(n) = f(n) \text{ for each } n \geq 1.$$

Since $g(0) = u_1$ is a unit in R , then, by Proposition 2.2 in [10], g is a unit in T . Also, we can easily check that $h'_{u_2}, \dots, h'_{u_n} \in U(T); i = 2, \dots, n$ and $h'_e \in \text{Id}(T)$. Thus, we conclude that $T = (HR, \sigma)$ is an n -clean ring. □

Taking $\sigma = \text{id}_R$, the identity automorphism on R , we get the next result

Corollary 2.4. *Let R be a ring, then the ring of Hurwitz series HR is an n -clean ring if and only if R is an n -clean ring.*

The previous corollary generalizes the following result due to Ghanem [5].

Theorem 2.5. *Suppose R is a commutative ring and n is a positive integer. Then HR is an n -clean ring if and only if R is an n -clean ring.*

3. $g_H(x)$ -clean skew Hurwitz ring.

Camilo and Simon in [3] introduced the $g(x)$ -clean ring for a polynomial $g(x) \in C(R)[x]$. A ring R is said to be $g(x)$ -clean if every element of R is a sum of a unit and a root of the polynomial $g(x)$. Nicholson and Zhou in [20] showed that $\text{End}({}_R M)$ is a $g(x)$ -clean where ${}_R M$ is a semisimple left R -module and $g(x) \in (x-a)(x-b)C(R)[x]$ where $a, b \in C(R)$ and $b, b-a \in U(R)$. Fan and Yang [4] investigated $g(x)$ -clean rings and obtained several important results. Clearly, any clean ring is n -clean and $g(x)$ -clean. The following example shows us that the converse need not be true:

Example 3.1 (Example 3.1, [22]). Let G be a cyclic group of order 3, then the group ring $\mathbb{Z}_{(7)}G$ is not clean, while Theorem 2.3, in [21], illustrates that $\mathbb{Z}_{(7)}G$ is a 2-clean ring. Hence, n -clean ring need not be clean.

Next, we give a characterization of $g_H(x)$ -clean of skew Hurwitz series rings.

Theorem 3.2. *Let R be a ring, $\sigma \in \text{End}(R)$ and $g(x) = a_0 + a_1x + \dots + a_mx^m \in C(R)[x]$. Then the ring R is $g(x)$ -clean if and only if $T = (HR, \sigma)$ is $g_H(x)$ -clean, where*

$$g_H(x) = h'_{a_0} + h'_{a_1}x + \dots + h'_{a_m}x^m \in C(T)[x].$$

Proof. Suppose R is a $g(x)$ -clean ring and $f \in T$. Hence $f(0) = u + s$ where $u \in U(R)$ and $g(s) = 0$. Therefore, $f = v + h'_s$ where $v \in T$ defined by $v(0) = u$ and $v(n) = f(n)$ for each $n \geq 1$. Since $v(0) = u$ is a unit of R , then v is a unit of T , by Proposition 2.2, in [10]. Clearly, h'_s is a root of the polynomial $g_H(x) \in C(T)[x]$. Therefore, T is a $g_H(x)$ -clean ring.

Conversely, suppose T is a $g_H(x)$ -clean ring and $r \in R$, then $\lambda_R(r) \in T$. Hence $\lambda_R(r) = f + q$ where $f \in U(T)$ and $g_H(q) = 0$. Therefore $\varepsilon_R(f) \in U(R)$, by Proposition 2.2, in [10], and $g(\varepsilon_R(q)) = 0$. Moreover, $r = \varepsilon_R(f) + \varepsilon_R(q)$. So, R is a $g(x)$ -clean ring. \square

Taking $\sigma = \text{id}_R$, the identity automorphism on R , we get the next result

Corollary 3.3. *Let R be a ring and $g(x) \in C(R)[x]$. The ring of Hurwitz series HR is a $g_H(x)$ -clean ring if and only if R is a $g(x)$ -clean ring.*

The previous corollary generalizes the following result due to Ghanem [5].

Theorem 3.4. *Suppose R is a commutative ring and $g(x) \in C(R)[x]$. The ring of Hurwitz series HR is a $g_H(x)$ -clean ring if and only if R is a $g(x)$ -clean ring.*

4. $(n, g_H(x))$ -clean skew Hurwitz ring.

In [8], Handam extended the definition of $g(x)$ -clean ring to obtain a larger class of rings, call it $(n, g(x))$ -clean. A ring R is said to be $(n, g(x))$ -clean if every element of R can be written as a sum of a root of the polynomial $g(x)$ and n -units. The following two examples are due to Handam in [8]:

Example 4.1. Let R be the ring of all 3×3 upper triangular matrices over \mathbb{Z}_2 . Since

$$\begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} + \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} + \begin{pmatrix} 0 & 1 & 1 \\ 0 & 0 & 1 \\ 0 & 0 & 1 \end{pmatrix},$$

where $\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$ are units in R and

$$\begin{pmatrix} 0 & 1 & 1 \\ 0 & 0 & 1 \\ 0 & 0 & 1 \end{pmatrix}^2 + \begin{pmatrix} 0 & 1 & 1 \\ 0 & 0 & 1 \\ 0 & 0 & 1 \end{pmatrix}^3 = 0.$$

Hence, $\begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 1 \end{pmatrix}$ is a $(2, x^2 + x^3)$ -clean element.

Clearly, clean rings are $(1, x^2 - x)$ -clean rings, n -clean rings are $(n, x^2 - x)$ -clean rings and $g(x)$ -clean rings are $(1, g(x))$ -clean rings. Thus, the classes of n -clean and $g(x)$ -clean rings are proper subclasses of $(n, g(x))$ -clean rings.

Example 4.2. Let G be a cyclic group of order 3, then the group ring $\mathbb{Z}_{(7)}G$ is not clean, by [7], while Theorem 2.3, in [21], illustrates that $\mathbb{Z}_{(7)}G$ is a 2-clean ring. Hence, n -clean ring need not be clean. So, $\mathbb{Z}_{(7)}G$ is a $(2, x^2 - x)$ -clean ring which is not a $(1, x^2 - x)$ -clean ring. Thus we obtain an example which is a $(2, x^2 - x)$ -clean ring but not a $(x^2 - x)$ -clean ring.

Propositions 2.9 and 2.10, in [8], tell us the following: if R is an $(n, g(x))$ -clean ring, then the power series ring $R[[x]]$ is an $(n, g(x))$ -clean ring but its subring $R[x]$ is not an $(n, g(x))$ -clean ring.

In the following we give the necessary and sufficient condition for the skew Hurwitz series ring $T = (HR, \sigma)$ to be $(n, g_H(x))$ -clean ring:

Theorem 4.3. *Let R be a ring, $\sigma \in \text{End}(R)$, n a positive integer and $g(x)$ a fixed polynomial in $C(R)[x]$. Then $T = (HR, \sigma)$ is an $(n, g_H(x))$ -clean ring if and only if R is an $(n, g(x))$ -clean ring.*

Proof. Since $\langle h_2 \rangle = Th_2 = \{fh_2 \mid f \in T\}$ is an ideal of T and, by Proposition 2.2, we have $T \cong I(R; \langle h_2 \rangle) = R \oplus \langle h_2 \rangle$. If $T = (HR, \sigma)$ is an $(n, g_H(x))$ -clean ring, then $R \cong T / \langle h_2 \rangle$ is an $(n, g(x))$ -clean ring, by Proposition 2.8 in [8].

Conversely, suppose that R is an $(n, g(x))$ -clean ring and $f \in T$, hence $f(0) \in R$. Write

$$f(0) = s + u_1 + u_2 + \dots + u_n,$$

where $u_i \in U(R)$; $i = 1, 2, \dots, n$ and $g(s) = 0$.

Then

$$f = h'_s + v + h'_{u_2} + \dots + h'_{u_n}$$

where $v \in T$ defined by

$$v(0) = u_1 \text{ and } v(n) = f(n) \text{ for each } n \geq 1.$$

Since $v(0) = u_1$ is a unit in R , then, by Proposition 2.2 in [10], v is a unit in T . Also, we can easily check that $h'_{u_2}, \dots, h'_{u_n} \in U(T)$; $i = 2, \dots, n$ and $g(h'_s) = 0$. Thus, we conclude that $T = (HR, \sigma)$ is an $(n, g_H(x))$ -clean ring. \square

Taking $\sigma = \text{id}_R$, the identity automorphism on R , we get the next result

Corollary 4.4. *Let R be a ring, n a positive integer and $g(x)$ be a fixed polynomial in $C(R)[x]$. Then the ring of Hurwitz series HR is an $(n, g_H(x))$ -clean ring if and only if R is an $(n, g(x))$ -clean ring.*

The previous corollary generalizes the following result due to Ghanem [5].

Theorem 4.5. *Suppose R is a commutative ring, n a positive integer and $g(x)$ be a fixed polynomial in $C(R)[x]$. Then HR is an $(n, g_H(x))$ -clean ring if and only if R is an $(n, g(x))$ -clean ring.*

5. Neat skew Hurwitz ring.

One of the fundamental properties of a clean ring is that every homomorphic image of a clean ring is clean. McGovern [18] defined a neat ring to be: the ring in which every proper homomorphic image is clean. Clearly, every clean ring is a neat ring but the converse need not be true, for example any nonlocal PID is a neat ring but is not clean.

In the following we give the necessary and sufficient condition for the skew Hurwitz series ring $T = (HR, \sigma)$ to be a neat ring:

Theorem 5.1. *Let R be a ring and $\sigma \in \text{End}(R)$. Then:*

- 1) $T = (HR, \sigma)$ is a neat ring if and only if R is a clean ring.
- 2) $T = (HR, \sigma)$ is a neat ring if and only if it is a clean ring.

Proof. 1) Since $\langle h_2 \rangle = Th_2 = \{fh_2 \mid f \in T\}$ is a two-sided ideal of T , we have $T \cong I(R; \langle h_2 \rangle)$, by Proposition 2.2, if $T = (HR, \sigma)$ is a neat ring, then $R \cong T / \langle h_2 \rangle$ is a clean ring. The converse direction is clear.

The conclusion (2) follows from (1) and Theorem 2.3. □

6. Prüfer domain of skew Hurwitz ring.

A commutative ring R is called Prüfer if every finitely generated ideal is invertible. An invertible ideal $A = \langle a_1, a_2, \dots, a_m \rangle$ has the property that $A^n = \langle a_1^n, a_2^n, \dots, a_m^n \rangle$ for each $n \in \mathbb{N}$. Thus it is clear that the Prüfer ring satisfies the following condition, if $a, b \in R$ and at least one of a and b is regular, then $ab \in \langle a^2, b^2 \rangle$. In [6], Gilmer called the ring satisfies the above condition a P -ring.

Throughout, unless otherwise stated, we assume that R is a commutative ring with identity 1 and D is an integral domain.

Proposition 6.1. *Suppose that R is a ring and $\sigma \in \text{End}(R)$. If $T = (HR, \sigma)$ is a P -ring, then R is a von-Neumann regular ring.*

Proof. Assume T is a P -ring. Let $0 \neq r \in R$ be a regular element $1 \neq n \in \mathbb{N}$, whence h_n is a regular element of T and $h'_r h_n \in \langle h'_r, h_n \rangle^2 = \langle h'^2_r, h^2_n \rangle = \langle h'^2_{r^2}, h^2_n \rangle$. Hence $h'_r h_n = h'^2_{r^2} f + h^2_n g$ for some $f, g \in T = (HR, \sigma)$. Since

$$\pi(h'^2_n g) = \pi(h^2_n) + \pi(g) = 2n - 2 + \pi(g)$$

and $\pi(h'_r h_n) = n - 1$, therefore, $r = (h'_r h_n)(n - 1) = (h'^2_{r^2} f)(n - 1) = r^2 f(n - 1) \in r^2 R$. Since R is a commutative ring, then R is a von-Neumann regular ring. □

Proposition 6.2. $T = (HR, \sigma)$ is an integral domain if and only if R is an integral domain and a torsion-free \mathbb{Z} -module.

Proof. Let $T = (HR, \sigma)$ be an integral domain. Since R has a natural embedding in T , then clearly R is an integral domain. Now suppose that the ring R is a torsion-free \mathbb{Z} -module, then there is a positive integer m , such that $m1 = 0$. Now, we have

$$(h_2 h_m)(m) = \binom{1+m-1}{1} h_2(1) \sigma(h_m(m-1)) = m1 = 0,$$

which implies that $h_2 h_{m-1} = 0$, a contradiction with the assumption that $T = (HR, \sigma)$ is an integral domain, so we conclude that R is a torsion-free \mathbb{Z} -module. The converse direction is clear. \square

Theorem 6.3. Let D be an integral domain and a torsion-free \mathbb{Z} -module. Then $T = (HR, \sigma)$ is a Prüfer domain if and only if D is a field.

Proof. Using the same argument in the proof of Proposition 6.1, it can be easily shown that $d \in d^2 D$. Since D is an integral domain, then d is invertible and D must be a field.

Conversely, assume that D is a field, then, by Proposition 2.2, every element in the subset $J = \langle h_2 \rangle = Th_2 = \{fh_2 \mid f \in T\}$ satisfies $(fh_2)(0) = 0$, so J is a two sided ideal of T . We can easily check that J is the only non-zero maximal ideal of T and the other ideal are principal in the form $J_n = \langle h_n \rangle = Th_n$ for each $n \geq 3$. Hence T is a principal ideal domain, in particular, T is a Prüfer domain. \square

Acknowledgements

The authors wish to express their sincere thanks to the *referee* for his/her helpful comments and valuable remarks which improved the results of this paper.

REFERENCES

- [1] A. Benhissi, *Ideal structure of Hurwitz series rings*, Contributions to Algebra and Geometry 48 (1) (2007), 251–256.
- [2] A. Benhissi - F. Koja, *Basic properties of Hurwitz series rings*, Ricerche mat., DOI 10.1007/s11587-012-0128-2.
- [3] V. Camilo - J.J. Simon, *The Nicholson-Varadarajan theorem on clean linear transformations*, Glasgow Math. J. 44 (2002), 365–369.

- [4] L. Fan - X. Yang, *On rings whose elements are the sum of a unit and a root of a fixed polynomial*, Comm. Algebra 36 (1) (2008), 855–861.
- [5] M. Ghanem, *Some properties of Hurwitz series ring*, Int. Math. Forum 40 (6) (2011), 1973–1981.
- [6] R. Gilmer, *Commutative semigroup rings*, University of Chicago Press, Chicago, 1984.
- [7] J. Han - W. K. Nicholson, *Extension of clean rings*, Comm. Algebra 29 (6) (2001), 2589–2595.
- [8] A. H. Handam, *$(n, g(x))$ -clean rings*, Int. Math. Forum 21 (4) (2009), 1007–1011.
- [9] A. H. Handam, *On f -clean rings and f -clean elements*, Proyecciones Journal of Mathematics 30 (2) (2011), 277–284.
- [10] A. M. Hassanein, *Clean rings of skew Hurwitz series*, Le Matematiche 62 (1) (2007), 47–54.
- [11] A. M. Hassanein, *On uniquely clean skew Hurwitz series*, South-east Bull. of Math. 35 (2012), 5–10.
- [12] A. M. Hassanein - R. M. Salem - M. A. Farahat, *Quasi-Baer and Quasi-Baer-* of Hurwitz series*, Proc. Math. and Phys. Soc. Egypt 86 (1) (2008), 35–43.
- [13] A. M. Hassanein - R. M. Salem - M. A. Farahat, *Noncommutative clean rings of Hurwitz series*, Proc. Math. and Phys. Soc. Egypt 86 (1) (2008), 45–51.
- [14] A. M. Hassanein - R. M. Salem, *Skew Hurwitz series of Baer and PP-rings*, J. of Adv. Research in pure Math. 3 (3) (2011), 61–69.
- [15] W. F. Keigher, *Adjunctions and comonads in differential algebra*, Pacific. J. Math 248 (1975), 99–112.
- [16] W. F. Keigher, *On the ring of Hurwitz series*, Comm. Algebra 25 (6) (1997), 1845–1859.
- [17] W. F. Keigher - F. L. Pritchard, *Hurwitz series as formal functions*, J. Pure Appl. Algebra 146 (2000), 291–304.
- [18] W. Wm. McGovern, *Neat rings*, J. Pure Appl. Algebra 205 (2) (2006), 243–265.
- [19] W. K. Nicholson, *Lifting idempotents and exchange rings*, Trans. Amer. Math. Soc. 229 (1977), 269–278.
- [20] W. K. Nicholson - Y. Zhou, *Endomorphisms that are sum of a unit and a root of a fixed polynomial*, Canadian Math. Bull. 49 (2006), 265–269.
- [21] G. Xiao - W. Tong, *n -clean rings and weakly unit stable rings*, Comm. Algebra 33 (5) (2003), 1501–1517.
- [22] Y. Q. Ye, *Semiclean rings*, Comm. Algebra 31 (11) (2003), 5609–5625.
- [23] J. Yu-Juan - Z. Shen-Gui, *Traingular matrix representations of rings of skew Hurwitz series*, J. of Shandong University (Natural Science) 46 (2) (2011), 105–109.
- [24] L. Zhongkui, *Hermite and PS- rings of Hurwitz series*, Comm. Algebra 28(1) (2000), 299–305.

A. M. HASSANEIN
Department of Mathematics
Faculty of Science,
Al-Azhar University, Cairo, Egypt.
e-mail: mhassaneien_05@yahoo.com

MOHAMED A. FARAHAT
Department of Mathematics
Faculty of Science,
Al-Azhar University, Cairo, Egypt.
e-mail: m_farahat79@yahoo.com