

LE MATEMATICHE

Vol. LXXI (2016) – Fasc. I, pp. 89–97

doi: 10.4418/2016.71.1.7

COEFFICIENT BOUNDS FOR A GENERAL SUBCLASS OF BI-UNIVALENT FUNCTIONS

ŞAHSENE ALTINKAYA - SIBEL YALÇIN

In the present investigation, we introduce and investigate a new subclass of the function class Σ of bi-univalent functions defined in the open unit disc. We find estimates on the coefficients $|a_2|$ and $|a_3|$ for functions in the function class $S_{\Sigma}(n, h, \lambda)$. The results presented in this paper improve or generalize the recent works of *Jothibasu* [13] and other authors.

1. Introduction and Definitions

Let A denote the class of analytic functions in the unit disk

$$U = \{z \in \mathbb{C} : |z| < 1\}$$

that have the form

$$f(z) = z + \sum_{n=2}^{\infty} a_n z^n \quad (1)$$

and let S be the class of all functions from A which are univalent in U .

The Koebe one-quarter theorem [8] states that the image of U under every function f from S contains a disk of radius $\frac{1}{4}$. Thus every such univalent function has an inverse f^{-1} which satisfies

$$f^{-1}(f(z)) = z, \quad (z \in U)$$

Entrato in redazione: 14 febbraio 2015

AMS 2010 Subject Classification: 30C45, 30C50.

Keywords: Analytic and univalent functions, bi-univalent functions, coefficient bounds.

and

$$f(f^{-1}(w)) = w, \left(|w| < r_0(f), r_0(f) \geq \frac{1}{4} \right),$$

where

$$f^{-1}(w) = w - a_2 w^2 + (2a_2^2 - a_3) w^3 - (5a_2^3 - 5a_2 a_3 + a_4) w^4 + \dots$$

A function $f(z) \in A$ is said to be bi-univalent in U if both $f(z)$ and $f^{-1}(z)$ are univalent in U .

Let Σ denote the class of bi-univalent functions defined in the unit disk U . For a brief history and interesting examples of functions in the class Σ ; see [3].

The research into Σ was started by Lewin ([15]). It focused on problems connected with coefficients and obtained the bound for the second coefficient. Several authors have subsequently studied similar problems in this direction (see [4], [19]). Thus, following Brannan and Taha [3], a function $f(z) \in A$ is said to be in the class $S_{\Sigma}^*(\alpha)$ of strongly bi-starlike functions of order α ($0 < \alpha \leq 1$) if each of the following conditions is satisfied:

$$f \in \Sigma, \left| \arg \left(\frac{zf'(z)}{f(z)} \right) \right| < \frac{\alpha\pi}{2} \quad (0 < \alpha \leq 1, z \in U)$$

and

$$\left| \arg \left(\frac{wg'(w)}{g(w)} \right) \right| < \frac{\alpha\pi}{2} \quad (0 < \alpha \leq 1, w \in U)$$

and is said to be in the class $K_{\Sigma}(\alpha)$ of strongly bi-convex functions of order α ($0 < \alpha \leq 1$) if each of the following conditions is satisfied:

$$f \in \Sigma, \left| \arg \left(1 + \frac{zf'(z)}{f(z)} \right) \right| < \frac{\alpha\pi}{2} \quad (0 < \alpha \leq 1, z \in U)$$

and

$$\left| \arg \left(1 + \frac{wg'(w)}{g(w)} \right) \right| < \frac{\alpha\pi}{2} \quad (0 < \alpha \leq 1, w \in U)$$

where g is the extension of f^{-1} to U . The classes $S_{\Sigma}^*(\alpha)$ and $K_{\Sigma}(\alpha)$ of bi-starlike functions of order α and bi-convex functions of order α , corresponding to the function classes $S^*(\alpha)$ and $K(\alpha)$, were also introduced analogously. For each of the function classes $S_{\Sigma}^*(\alpha)$ and $K_{\Sigma}(\alpha)$, they found non-sharp estimates on the initial coefficients. Recently, Srivastava et al. [22] introduced and investigated subclasses of the bi-univalent functions and obtained bounds for the

initial coefficients; it was followed by such works as those by Murugunsundaramoorthy et al. [18], Frasin and Aouf [9], Çağlar et al. [6] and others (see, for example, [1], [7], [14], [16], [17], [20], [24]).

Not much is known about the bounds on the general coefficient $|a_n|$ for $n \geq 4$. In the literature, the only a few works determining the general coefficient bounds $|a_n|$ for the analytic bi-univalent functions ([2], [5], [10], [11], [12]). The coefficient estimate problem for each of $|a_n|$ ($n \in \mathbb{N} \setminus \{1, 2\}$; $\mathbb{N} = \{1, 2, 3, \dots\}$) is still an open problem.

Let $f \in A$. We define the differential operator D^n , $n \in \mathbb{N}_0 = \mathbb{N} \cup \{0\}$ by (see [21])

$$\begin{aligned} D^0 f(z) &= f(z); \\ D^1 f(z) &= Df(z) = zf'(z); \\ &\vdots \\ D^n f(z) &= D^1(D^{n-1}f(z)). \end{aligned}$$

In this paper, by using the method [23] different from that used by other authors, we obtain bounds for the coefficients $|a_2|$ and $|a_3|$ for the subclasses of bi-univalent functions considered Jothibasud and get more accurate estimates than that given in [13].

Definition 1.1. Let the functions $h, p : U \rightarrow \mathbb{C}$ be so constrained that

$$\min \{Re(h(z)), Re(p(z))\} > 0$$

and

$$h(0) = p(0) = 1.$$

Definition 1.2. A function $f \in \Sigma$ is said to be $S_\Sigma(n, h, \lambda)$, $n \in \mathbb{N}_0$ and $0 \leq \lambda < 1$, if the following conditions are satisfied:

$$\frac{D^{n+1}f(z)}{(1-\lambda)D^n f(z) + \lambda D^{n+1}f(z)} \in h(U) \quad (z \in U) \tag{2}$$

and

$$\frac{D^{n+1}g(w)}{(1-\lambda)D^n g(w) + \lambda D^{n+1}g(w)} \in p(U) \quad (w \in U) \tag{3}$$

where $g(w) = f^{-1}(w)$.

2. Coefficient Estimates

Theorem 2.1. Let f given by (1) be in the class $S_{\Sigma}(n, h, \lambda)$, $n \in \mathbb{N}_0$ and $0 \leq \lambda < 1$. Then

$$|a_2| \leq \min \left\{ \sqrt{\frac{|h'(0)|^2 + |p'(0)|^2}{2(1-\lambda)^2 4^n}}, \sqrt{\frac{|h''(0)| + |p''(0)|}{4|(\lambda^2 - 1)4^n + 2(1-\lambda)3^n}} \right\} \quad (4)$$

and

$$|a_3| \leq \min \left\{ \frac{|h'(0)|^2 + |p'(0)|^2}{2(1-\lambda)^2 4^n} + \frac{|h''(0)| + |p''(0)|}{8(1-\lambda)3^n}, \frac{[4(1-\lambda)3^n + (\lambda^2 - 1)4^n] |h''(0)| + (1-\lambda^2)4^n |p''(0)|}{8|(\lambda^2 - 1)4^n + 2(1-\lambda)3^n| (1-\lambda)3^n} \right\}. \quad (5)$$

Proof. Let $f \in S_{\Sigma}(n, h, \lambda)$ and $0 \leq \lambda < 1$. It follows from (2) and (3) that

$$\frac{D^{n+1}f(z)}{(1-\lambda)D^n f(z) + \lambda D^{n+1}f(z)} = h(z) \quad (6)$$

and

$$\frac{D^{n+1}g(w)}{(1-\lambda)D^n g(w) + \lambda D^{n+1}g(w)} = p(w), \quad (7)$$

where $h(z)$ and $p(w)$ satisfy the conditions of Definiton 1. Furthermore, the functions $h(z)$ and $p(w)$ have the following Taylor-Maclaurin series expansions:

$$h(z) = 1 + h_1 z + h_2 z^2 + \dots$$

and

$$p(w) = 1 + p_1 w + p_2 w^2 + \dots,$$

respectively. It follows from (6) and (7) that

$$(1-\lambda)2^n a_2 = h_1, \quad (8)$$

$$(\lambda^2 - 1)4^n a_2^2 + 2(1-\lambda)3^n a_3 = h_2, \quad (9)$$

and

$$-(1-\lambda)2^n a_2 = p_1, \quad (10)$$

$$2(1-\lambda)3^n (2a_2^2 - a_3) + (\lambda^2 - 1)4^n a_2^2 = p_2. \quad (11)$$

From (8) and (10) we obtain

$$h_1 = -p_1, \quad (12)$$

and

$$2(1-\lambda)^2 4^n a_2^2 = h_1^2 + p_1^2. \tag{13}$$

By adding (9) to (11), we find that

$$[2(\lambda^2 - 1)4^n + 4(1-\lambda)3^n] a_2^2 = h_2 + p_2, \tag{14}$$

which gives us the desired estimate on $|a_2|$ as asserted in (4).

Next, in order to find the bound on $|a_3|$, by subtracting (11) from (9), we obtain

$$4(1-\lambda)3^n a_3 - 4(1-\lambda)3^n a_2^2 = h_2 - p_2. \tag{15}$$

Then, in view of (13), (14) and (15), it follows that

$$a_3 = \frac{h_1^2 + p_1^2}{2(1-\lambda)^2 4^n} + \frac{h_2 - p_2}{4(1-\lambda)3^n}$$

and

$$a_3 = \frac{h_2 + p_2}{2(\lambda^2 - 1)4^n + 4(1-\lambda)3^n} + \frac{h_2 - p_2}{4(1-\lambda)3^n}.$$

as claimed. This completes the proof of Theorem 2.1. □

3. Corollaries and Consequences

By setting $n = 0$ and $\lambda = 0$ in Theorem 2.1, we get Corollary 3.1 below.

Corollary 3.1. *Let the function $f(z)$ given by (1) be in the class $S_\Sigma(h)$ ($0 \leq \lambda < 1$). Then*

$$|a_2| \leq \min \left\{ \sqrt{\frac{|h'(0)|^2 + |p'(0)|^2}{2}}, \sqrt{\frac{|h''(0)| + |p''(0)|}{4}} \right\}$$

and

$$|a_3| \leq \min \left\{ \frac{|h'(0)|^2 + |p'(0)|^2}{2} + \frac{|h''(0)| + |p''(0)|}{8}, \frac{3|h''(0)| + |p''(0)|}{8} \right\}.$$

By setting $n = 1$ and $\lambda = 0$ in Theorem 2.1, we get Corollary 3.2 below.

Corollary 3.2. *Let the function $f(z)$ given by (1) be in the class $K_\Sigma(h)$ ($0 \leq \lambda < 1$). Then*

$$|a_2| \leq \min \left\{ \sqrt{\frac{|h'(0)|^2 + |p'(0)|^2}{8}}, \sqrt{\frac{|h''(0)| + |p''(0)|}{8}} \right\}$$

and

$$|a_3| \leq \min \left\{ \frac{|h'(0)|^2 + |p'(0)|^2}{8} + \frac{|h''(0)| + |p''(0)|}{24}, \frac{2|h''(0)| + |p''(0)|}{12} \right\}.$$

Corollary 3.3. *If*

$$h(z) = p(z) = \left(\frac{1+z}{1-z} \right)^\alpha = 1 + 2\alpha z + 2\alpha^2 z^2 + \dots \quad (0 < \alpha \leq 1),$$

then inequalities (4) and (5) become

$$|a_2| \leq \min \left\{ \frac{2\alpha}{(1-\lambda)2^n}, \sqrt{\frac{2}{|(\lambda^2-1)4^n + 2(1-\lambda)3^n|}} \alpha \right\}$$

and

$$|a_3| \leq \min \left\{ \frac{4\alpha^2}{(1-\lambda)^2 4^n} + \frac{\alpha^2}{(1-\lambda)3^n}, \frac{2\alpha^2}{|(\lambda^2-1)4^n + 2(1-\lambda)3^n|} \right\}.$$

Remark 3.4. Putting $n = 0$ and $\lambda = 0$ in Corollary 3.3, we obtain the following corollary.

Corollary 3.5 (See also [23]). *Let the function $f(z)$ given by (1) be in the class $S_\Sigma^*(\alpha)$. Then*

$$|a_2| \leq \sqrt{2}\alpha$$

and

$$|a_3| \leq 2\alpha^2.$$

The estimates on the coefficient $|a_2|$ and $|a_3|$ of Corollary 3.5 are improvement of the estimates obtained in [13] and [18].

Remark 3.6. Putting $n = 1$ and $\lambda = 0$ in Corollary 3.3, we obtain the following corollary.

Corollary 3.7. *Let the function $f(z)$ given by (1) be in the class $K_\Sigma(\alpha)$. Then*

$$|a_2| \leq \alpha$$

and

$$|a_3| \leq \alpha^2.$$

The estimate on the coefficient $|a_3|$ of Corollary 3.7 is improvement of the estimates obtained in [13].

Corollary 3.8. *If*

$$h(z) = p(z) = \frac{1 + (1 - 2\beta)z}{1 - z} = 1 + 2(1 - \beta)z + 2(1 - \beta)z^2 + \dots \quad (0 \leq \beta < 1),$$

then inequalities (4) and (5) become

$$|a_2| \leq \min \left\{ \frac{2(1 - \beta)}{(1 - \lambda)2^n}, \sqrt{\frac{2(1 - \beta)}{|(\lambda^2 - 1)4^n + 2(1 - \lambda)3^n|}} \right\}.$$

and

$$|a_3| \leq \min \left\{ \frac{4(1 - \beta)^2}{(1 - \lambda)^2 4^n} + \frac{1 - \beta}{(1 - \lambda)3^n}, \frac{2(1 - \beta)}{|(\lambda^2 - 1)4^n + 2(1 - \lambda)3^n|} \right\}.$$

Remark 3.9. Putting $n = 0$ and $\lambda = 0$ in Corollary 3.8, we obtain the following corollary.

Corollary 3.10. *Let the function $f(z)$ given by (1) be in the class $K_\Sigma(\beta)$. Then*

$$|a_2| \leq \sqrt{2(1 - \beta)}$$

and

$$|a_3| \leq 2(1 - \beta).$$

The estimate on the coefficient $|a_3|$ of Corollary 3.10 is improvement of the estimates obtained in [13] and [18].

Remark 3.11. Further, taking $n = 1$ and $\lambda = 0$ in Corollary 3.8, we obtain the following corollary.

Corollary 3.12. *Let the function $f(z)$ given by (1) be in the class $K_\Sigma(\beta)$. Then*

$$|a_2| \leq \sqrt{1 - \beta}$$

and

$$|a_3| \leq (1 - \beta).$$

The estimate on the coefficient $|a_3|$ of Corollary 3.12 is improvement of the estimates obtained in [3] and [13].

REFERENCES

- [1] Ş. Altinkaya - S. Yalçın, *Initial coefficient bounds for a general class of bi-univalent functions*, International Journal of Analysis, Article ID 867871, (2014), 4 pp.
- [2] Ş. Altinkaya - S. Yalçın, *Coefficient bounds for a subclass of bi-univalent functions*, TWMS Journal of Pure and Applied Mathematics, in press.
- [3] D. A. Brannan - T. S. Taha, *On some classes of bi-univalent functions*, Studia Universitatis Babeş-Bolyai. Mathematica, 31 (2) (1986), 70–77.
- [4] D. A. Brannan - J. G. Clunie, *Aspects of contemporary complex analysis*, Proceedings of the NATO Advanced Study Institute held at University of Durham: July 1-20, 1979. Academic Press, New York 1980.
- [5] S. Bulut, *Faber polynomial coefficient estimates for a comprehensive subclass of analytic bi-univalent functions*, C. R. Acad. Sci. Paris, Ser. I, in press.
- [6] M. Caglar - H. Orhan - N. Yagmur, *Coefficient bounds for new subclasses of bi-univalent functions*, Filomat 27 (2013), 1165–1171.
- [7] O. Crişan, *Coefficient estimates certain subclasses of bi-univalent functions*, Gen. Math. Notes 16 (2) (2013), 93–102.
- [8] P.L. Duren, *Univalent Functions*, Grundlehren der Mathematischen Wissenschaften 259, Springer, New York, USA, 1983.
- [9] B. A. Frasin - M. K. Aouf, *New subclasses of bi-univalent functions*, Applied Mathematics Letters 24 (2011), 1569–1573.
- [10] S. G. Hamidi - J. M. Jahangiri, *Faber polynomial coefficient estimates for analytic bi-close-to-convex functions*, C. R. Acad. Sci. Paris Ser.I 352 (1) (2014), 17–20.
- [11] J. M. Jahangiri - S. G. Hamidi, *Coefficient estimates for certain classes of bi-univalent functions*, Int. J. Math. Math. Sci., ArticleID 190560, (2013), 4 pp.
- [12] J. M. Jahangiri - S. G. Hamidi - S. A. Halim, *Coefficients of bi-univalent functions with positive real part derivatives*, Bull. Malays. Math. Soc., in press, <http://math.usm.my/bulletin/pdf/acceptedpapers/2013-04-050-R1.pdf>.
- [13] J. Jothibasu, *Certain subclasses of bi-univalent functions defined by Salagean Operator*, Electron. J. Math. Anal. Appl. 3 (1) (2015), 150–157.
- [14] B. Srutha Keerthi - B. Raja, *Coefficient inequality for certain new subclasses of analytic bi-univalent functions*, Theoretical Mathematics and Applications 3 (1) (2013), 1–10.
- [15] M. Lewin, *On a coefficient problem for bi-univalent functions*, Proceeding of the American Mathematical Society 18 (1967), 63–68.
- [16] X. F. Li - A. P. Wang, *Two new subclasses of bi-univalent functions*, Internat. Math. Forum 7 (2012), 1495–1504.
- [17] N. Magesh - J. Yamini, *Coefficient bounds for a certain subclass of bi-univalent functions*, International Mathematical Forum 8 (27) (2013), 1337–1344.
- [18] G. Murugunsundaramoorthy - N. Magesh - V. Prameela, *Coefficient bounds for certain subclasses of bi-univalent functions*, Abstr. Appl. Anal., Article ID

- 573017, (2013), 1–3.
- [19] E. Netanyahu, *The minimal distance of the image boundary from the origin and the second coefficient of a univalent function in $|z| < 1$* , Archive for Rational Mechanics and Analysis 32 (1969), 100–112.
- [20] S. Porwal - M. Darus, *On a new subclass of bi-univalent functions*, J. Egypt. Math. Soc. 21 (3) (2013), 190–193.
- [21] G. S. Salagean, *Subclasses of univalent functions*, in: *Complex Analysis*, Fifth Romanian Finish Seminar, Bucharest, 1 (1983), 362–372.
- [22] H. M. Srivastava - A. K. Mishra - P. Gochhayat, *Certain subclasses of analytic and bi-univalent functions*, Applied Mathematics Letters 23 (10) (2010), 1188–1192.
- [23] H. M. Srivastava - S. Bulut - M. Çağlar - N. Yağmur, *Coefficient estimates for a general subclass of analytic and bi-univalent functions*, Filomat 27 (5) (2013), 831–842.
- [24] Q. H. Xu - Y. C. Gui - H. M. Srivastava, *Coefficient estimates for a certain subclass of analytic and bi-univalent functions*, Applied Mathematics Letters 25 (2012), 990–994.

ŞAHSENE ALTINKAYA

Department of Mathematics

Faculty of Arts and Science

Uludag University

e-mail: sahsene@uludag.edu.tr

SIBEL YALÇIN

Department of Mathematics

Faculty of Arts and Science

Uludag University

e-mail: syalcin@uludag.edu.tr