



# Effect of yoga practices on Hs-CRP in Indian railway engine drivers of metropolis

Shete Sanjay. U, Kulkarni Dattatraya. D and Thakur Ghanshyam Singh

Scientific Research Department, Kaivalyadhama, S.M.Y.M. Samiti, Lonavla, Pune, Maharashtra, India.

## Abstract

**Objectives:** To examine the effect of yoga training on high sensitivity C reactive protein (hs-CRP) and lipid profile levels in railway engine drivers working in metropolis.

**Methods:** Male drivers of Indian railways, age ranged from 30 to 42 yrs with no known medical disorders, were randomized to yoga group (n=16) and control group (n=16). At the baseline and after completion of one month yoga training both the groups were assessed for hs-CRP and lipid profile levels. The yoga group practiced in a set of yoga techniques for 1 hr. daily in the morning along with daily routine work, while control group engaged in daily routine work only.

**Results:** Statistically significant reduction ( $p < 0.01$ ) was evident in both hs-CRP and serum total cholesterol.

**Conclusion:** Yoga based lifestyle modifications could contribute to prevention of coronary artery disease.

**Keywords:** hs-CRP, CAD, Lipid profiles, Yoga, Railway engine drivers.

## INTRODUCTION

The process of inflammation is an immunological response evoked by the body injury or infection. The past clinical and population research studies suggest the importance of inflammation in progression of atherosclerosis [1]. This is the process in which fatty deposits build up in the inner lining of arteries. C-reactive protein is an acute phase protein which increases during systemic inflammation. In fact, it has been found that the *high sensitivity C reactive protein* (hs-CRP) is a reliable indicator of inflammatory atherosclerosis among subjects with or without cardiovascular disease (CVD) [2-8].

There is evidence that physical activity may modify the inflammatory process. In fact, the effect of regular exercise has shown inverse association on levels of inflammatory markers [9]. Rohde et al. reported that healthy men who exercise more than one time a week had lower mean CRP than the men who did not exercise at least once a week [10]. In addition, some of the CVD risk factors, in particular those relating to inflammation and hemostasis, are also modified favorably with physical activity [11-16]. This likely represents an additional mechanistic pathway through which physical activity decreases CVD risk. Ridker et al. found that aspirin use decreased risk of myocardial infarction and ischemic stroke in apparently healthy men, primarily in those with the highest levels of C-reactive protein [5]. This suggests that even a moderate reduction in inflammation may be protective.

The profession of railway engine driver, especially in metropolitan cities in India, is very hectic and full of uncertainty that influence more stress due to night work and long irregular working

hours, which may increase risk of cardiovascular diseases. Some of the earlier investigations have revealed an increased incidence of myocardial infarction among male railway engine drivers that may be due to their continuous exposure to electro-magnetic field [17, 18].

Research has suggested that inflammatory markers, such as high-sensitivity C-reactive protein (hs-CRP), provide an alternative method for assessment of cardiovascular risk [19-21]. Furthermore, previous research reports suggest that yoga based lifestyle modifications help in regression of coronary lesions [22-24]. Hence, the main objective of this study was to examine the effect of yoga training on high sensitivity CRP (hs-CRP) and lipid profile level in railway engine drivers.

## MATERIALS AND METHODS

**Subjects :** After approval of experimental procedures by the Institutional Ethics Committee, informed consent was obtained from thirty two male engine-drivers working in Indian railways at Mumbai region, who had no yoga practice background [age: 30-42 yr]. The participants were examined by a medical officer and those with known coronary disease were excluded from this study.

**Experimental design:** The subjects were randomly divided into two groups viz., yoga group (n = 16) and control group (n=16). Primarily, baseline concentrations of serum lipid profile and hs-CRP were taken from all the selected subjects of both the groups. The subjects of yoga group were then underwent a training of yoga practices under the overall supervision of yoga expert, whereas the comparable control group did not. The training was imparted to the yoga group daily one hour in the morning including Sundays and holidays for a total period of one month. However, both the groups participated in their regular lifestyle activities and duties assigned by the railways authorities. They were also advised to avoid non-vegetarian food. After completion of the experiment for one month, the testing of serum biochemistry was repeated.

**Yoga Practices:** The yoga group practiced a set of yoga techniques in the form of *asana* (postures) and *pranayama* (breathing techniques). The supine position asanas were *ardh-*

Received: Nov 26, 2011; Revised: Jan 03, 2012; Accepted: Feb 15, 2012.

\*Corresponding Author

Shete Sanjay. U  
Scientific Research Department, Kaivalyadhama, S.M.Y.M. Samiti, Lonavla,  
Pune, Maharashtra, India.

Tel: +91-9860594828; Fax: +91- 2114-271983  
Email: shete.sanjay@gmail.com

halasana (halplough) viparita karani (inverted pose), matsyasana (fish pose), naukasana (boat pose), setubandhasana (bridge pose), and supta vajrasana (reclining adamant pose). The prone position asanas included were bhujangasana (cobra pose), ardashalabhasana (half locust pose), shalabhasana (locust pose), and dhanurasana (bow pose). The sitting position asanas were vakrasana (twisted pose), gomukhasana (cow face pose), paschimatanasana (forward bending pose), ardha ushtrasana (half camel pose) while standing position asanas were tadasana (mountain pose), chakrasana (wheel pose), utkatasana (chair pose) vrikshasana (tree pose). The pranayama practices for this experiment were anulom vilom and bhramari. Each session of yoga practices was concluded with om chanting. The duration of each asana (posture) was ranged from 2 to 3 minutes depending upon the improvement in performance, whereas for practice of pranayama the duration was from 2-5 minutes.

**Serum biochemistry:** Serum samples were separated from the collected blood by using Vacutainer blood-collection tubes (Becton Dickinson) with the centrifugation at 1,000 g for 10 min after the blood was allowed to clot at room temperature for 30 min. Serum hs-CRP activity was measured by using Calbiotech (USA) enzyme immunoassay kit on ELISA plate reader (Bio-Rad 680, Bio-Rad PW 40, USA), where the sensitivity limit was 0.2 µg/ml. Further, biochemical assay kits as prescribed for the analyzer Statfax-2000 (Awareness technology, USA) were used to measure Serum total Cholesterol, Triglycerides HDL, and LDL concentration.

**Statistics:** Since the primary outcome-variables were lipid profile and hs-CRP, the collected data were analyzed for evaluating

mean and standard deviation; whereas within group comparisons were performed using paired t-tests, while between-group comparisons were performed using independent t-tests.

**RESULTS**

The result of within group comparison revealed that the yoga group showed a significant decrease in total cholesterol (Tc), triglyceride (Tg) and hs-CRP (t= 6.4, p<0.01; t= 10.42, p<0.01; t= 27.87, p<0.01) (Table 1), whereas the control group revealed no change in Tc (t= 0.28 p >0.05) and in hs-CRP (t= 1.58, p>0.05) respectively, but significant increase in Tg (t=19.66, p<0.01) was evident. However, a significant decrease in LDL were observed in yoga (t= 5.21, p<0.01) and control groups (t= 4.79, p<0.01). Further, no change was evident in HDL in both yoga and control groups (t=0.48, p>0.05, t=0.33, p>0.05). This indicates yoga practice helps to reduce lipid profile to normal level and decreased CRP that reflects yoga reduced inflammation.

Further, the results between the group confirmed that the yoga group had significantly lower level of Tc, Tg and hs-CRP as compared to control group (t=1.62 p< 0.05, t=1.83, p< 0.05, t=1.96, p< 0.05). Trend of reduction in LDL was evident among the subjects of both the yoga and control groups; however, no statistically significant difference was evident between these groups (t=1.04, p>0.05) (Table 1). In case of HDL, trend of improvement was seen in both the yoga and control groups; however, no statistically difference between the groups was seen (t=0.39, p>0.05).

Table 1. Comparison of the baseline and final values (end of one month) of serum lipids and hs-CRP levels recorded at the end of the yoga program

Variable	Control Group			Yoga Group			Control Vs Yoga (t-value)
	Baseline (M±SD)	Final (M±SD)	t-value	Baseline (M±SD)	Final (M±SD)	t-value	
Total cholesterol (mg/dL)	175.06 (±18.23)	174.56 (±17.86)	0.28	191.81 (±20.38)	179.375 (±16.59)	6.4**	1.62*
Triglycerides (mg/dL)	118.56 (±12.45)	141.87 (±13.55)	19.66**	149.81 (±13.76)	134.18 (±13.34)	10.42**	1.83*
HDL (mg/dL)	51.75 (±6.23)	53.52 (±5.98)	0.48	51.36 (±5.77)	53.093 (±5.48)	0.33	0.39
LDL (mg/dL)	99.59 (±8.23)	94.81 (±8.56)	4.79**	99.95 (±8.67)	97.93 (±9.08)	5.21**	1.04
hsCRP(microgram/ml)	2.40 (±0.22)	2.37 (±0.18)	1.58	2.35 (±0.20)	1.87 (±0.19)	27.87**	1.96*
*p<0.05, **p<0.01							

**DISCUSSION**

The result of this randomized control trial of one month yoga training could reduce total cholesterol, triglycerides, low density lipoprotein (LDL) and hs-CRP in railway engine drivers. However, there was no significant improvement in high density lipoprotein (HDL).

In the present study participants were railway engine drivers who are believed to be under stress. Previous study on railway

engine drivers shows prevalence of high cardiovascular risk factors among these professionals [25]. Further, the occupational stress can lead to progression of coronary atherosclerosis but it can be prevented by appropriate lifestyle modification. Elevated hsCRP has consistently provided incremental prognostic value for cardiovascular risk prediction beyond traditional risk factor assessment [26-31]. Weight loss and physical activity can lower hsCRP levels, and lifestyle changes are first-line therapy to lower CVD risk and regression of atherosclerosis [32, 33]. In this perspective Indian

traditional yoga practices were found beneficial for reducing stress and inflammation because one randomized trial suggested positive benefits for inflammation in heart patients [34]. There is ample of evidence suggesting positive effect of yoga lifestyle in cardiovascular disease [35, 36, 22, 24]. In our study results shows reduction in hsCRP and lipoproteins and as the atherosclerotic process is characterized by inflammation one alternative explanation would be that regular practice of yoga, which offer protection against atherosclerosis, indirectly offers protection against vascular inflammation and hence, systemic low grade inflammation. This may be the mechanism by which yoga practices could have helped to reduce CRP levels. Other mechanisms linking exercise to lower inflammation levels may involve antioxidant effect of exercise. There is evidence that yoga practices can significantly elevate antioxidant defense [37].

Finally, the reduction in the lipoproteins and CRP, even though, is less in magnitude but achieved by a very simple and inexpensive intervention. This is of importance, because railway engine drivers are at higher risk of developing cardiovascular disease. Although lipoproteins and hsCRP are the tools for estimating coronary artery disease in healthy adults, this study has some limitations. It does not include other well established risk factors, such as BMI, alcohol intake, levels of physical activity or the presence of or absence of parental history of coronary artery disease. Though this study was conducted on a small sample, the results appeared are promising. However, future study on larger population is required to confirm the beneficial effects of yoga. Since, our study was limited to measures of hsCRP and lipoproteins other prospective studies estimating specific cytokines related to inflammation will be required to elucidate role of yoga in the prevention and regression of CAD.

## CONCLUSION

The present study has demonstrated that yoga practices are associated with regression of inflammatory process by reducing CRP levels in a representative sample of apparently healthy middle aged railway engine drivers. Since, elevated levels of CRP and other markers have been shown to important predictors of coronary atherosclerosis, current study implies that yoga practices seems to be of lower risk in controlling coronary atherosclerosis by reducing inflammation. Further studies that examine yoga as a prospective predictor of inflammation in general population sample are needed to definitively establish whether yoga truly prevents or reduces inflammation by assessing various other inflammatory markers.

## ACKNOWLEDGEMENT

The investigators are grateful to Swami Maheshananda (Chairman), O.P.Tiwari (Secretary) and Subodh Tiwari (Administrator) of Kaivalyadhama SMYM Samiti, Lonavla (India) for their inspiration and financial support to complete this project. They are also thankful to Dr. Desai, ADR, Scientific Research Department, Kaivalyadhama SMYM Samiti, Lonavla (India) for encouragement to complete this project. Further, suggestions rendered by Dr.T.K.Bera, Former Asst. Director of Research of the department to upgrade the manuscript are thankfully acknowledged.

## REFERENCES

- [1] Heinrich, J., H. Schulte, R. Schonfeld, E. Kohler, and G. Assmann. 1995. Association of variables of coagulation, fibrinolysis and acute-phase with atherosclerosis in coronary and peripheral arteries and those arteries supplying the brain. *Thromb Haemost.* 73:374–379.
- [2] Tracy, R. P., R. N. Lemaitre, B. M. Psaty, D. G. Ives, R. W. Evans, M. Cushman, E. N. Meilahn, and L. H. Kuller. 1997. Relationship of C-reactive protein to risk of cardiovascular disease in the elderly: results from the Cardiovascular Health Study and the Rural Health Promotion Project. *Arterioscler Thromb Vasc Biol.* 17:1121–1127.
- [3] Liuzzo, G., L. M. Biasucci, J. R. Gallimore, R. L. Grillo, A. G. Rebuzzi, M. B. Pepys, and A. Maseri. 1994. The prognostic value of C-reactive protein and serum amyloid a protein in severe unstable angina. *N Engl J Med.* 331:417–424.
- [4] Thompson, S. G., J. Kienast, S. D. Pyke, F. Haverkate, and J. C. van de Loo. 1995. Hemostatic factors and the risk of myocardial infarction or sudden death in patients with angina pectoris: European Concerted Action on Thrombosis and Disabilities Angina Pectoris Study Group. *N Engl J Med.* 332:635–641.
- [5] Ridker, P. M., M. Cushman, M. J. Stampfer, R. P. Tracy, and C. H. Hennekens. 1997. Inflammation, aspirin, and the risk of cardiovascular disease in apparently healthy men. *N Engl J Med.* 336:973–979.
- [6] Ridker, P. M., R. J. Glynn, and C. H. Hennekens. 1998. C-reactive protein adds to the predictive value of total and HDL cholesterol in determining risk of first myocardial infarction. *Circulation.* 97:2007–2011.
- [7] Haverkate, F., S. G. Thompson, S. D. Pyke, J. R. Gallimore, and M. B. Pepys. 1997. Production of C-reactive protein and risk of coronary events in stable and unstable angina: European Concerted Action on Thrombosis and Disabilities Angina Pectoris Study Group. *Lancet.* 349:462–466.
- [8] Kuller, L. H., R. P. Tracy, J. Shaten, and E. N. Meilahn. 1996. Relation of C-reactive protein and coronary heart disease in the MRFIT nested case-control study: Multiple Risk Factor Intervention Trial. *Am J Epidemiol.* 144:537–547.
- [9] Geffken, D. F., M. Cushman, G. L. Burke, J. F. Polak, P. A. Sakkinen, and R. P. Tracy. 2001. Association between physical activity and markers of inflammation in a healthy elderly population. *Am J Epidemiol.* 153:242–250.
- [10] Rohde, L. E., C. H. Hennekens, and P. M. Ridker. 1999. Survey of C-reactive protein and cardiovascular risk factors in apparently healthy men. *Am J Cardiol.* 84:1018–1022.
- [11] Eckel, R. H., W. W. Barouch, and A. G. Ershow. 2002. Report of the National Heart, Lung, and Blood Institute–National Institute of Diabetes and Digestive and Kidney Diseases Working Group on the pathophysiology of obesity associated cardiovascular disease. *Circulation.* 105:2923–2928.
- [12] Church, T. S., C. E. Barlow, C. P. Earnest, J. B. Kampert, E. L. Priest, and S. N. Blair. 2002. Associations between cardiorespiratory fitness and C-reactive protein in men. *Arterioscler Thromb Vasc Biol.* 22:1869–1876.
- [13] Wannamethee, S. G., G. D. Lowe, P. H. Whincup, A. Rumley, M. Walker, and L. Lennon. 2002. Physical activity and

- hemostatic and inflammatory variables in elderly men. *Circulation*. 105:1785–1790.
- [14] Church, T. S., C. E. Finley, C. P. Earnest, J. B. Kampert, L. W. Gibbons, and S. N. Blair. 2002. Relative associations of fitness and fatness to fibrinogen, white blood cell count, uric acid and metabolic syndrome. *Int J Obes Relat Metab Disord*. 26:805–813.
- [15] Mora, S., I. M. Lee, J. E. Buring, and P. M. Ridker. 2006. Association of physical activity and body mass index with novel and traditional cardiovascular biomarkers in women. *JAMA*. 295:1412–1419.
- [16] Abramson, J. L. and V. Vaccarino. 2002. Relationship between physical activity and inflammation among apparently healthy middle-aged and older US adults. *Arch Intern Med*. 162:1286–1292.
- [17] Piros, S., S. Karlehagen, G. Lappas, and L. Wilhelmsen. 2000. Psychosocial risk factors for myocardial infarction among Swedish railway engine drivers. *J Cardiovasc Risk*. 7:389-394.
- [18] Martin, R., E. Matthias, P. Dominik, and M. Christoph. 2008. Cardiovascular mortality and exposure to extremely low frequency magnetic fields: a cohort study of Swiss railway workers. *Environmental Health*. 7:35.
- [19] Ridker, P. M. 1999. Evaluating novel cardiovascular risk factors: can we better predict heart attacks? *Ann Intern Med*. 130:933–937.
- [20] Ridker, P. M. 2001. High-sensitivity C-reactive protein. Potential adjunct for global risk assessment in the primary prevention of cardiovascular disease. *Circulation*. 103:1813–1818.
- [21] Ridker, P. M., C. H. Hennekens, J. E. Buring, and N. Rifai. 2000. C-reactive protein and other markers of inflammation in the prediction of cardiovascular disease in women. *N Engl J Med*. 342:836–843.
- [22] Yogendra, J., H. J. Yogendra, S. Ambardekar, R. D. Lele, S. Shetty, M. Dave, and N. Husein. 2004. Beneficial effects of yoga lifestyle on reversibility of ischaemic heart disease: caring heart project of International Board of Yoga. *J Assoc Physician India*. 52:283-299.
- [23] Manchanda, S. C., R. Narang, K. S. Reddy, U. Sachdeva, D. Prabhakaran, S. Dharmanand, M. Rajani, and R. Bijlani. 2000. Retardation of coronary atherosclerosis with yoga lifestyle intervention. *J Assoc Physician India*. 48:687-694.
- [24] Bijlani, R. L., R. P. Vempati, R. K. Yadav, R. B. Ray, V. Gupta, R. Sharma, N. Mehta, and S. C. Mahapatra. 2005. A Brief but comprehensive lifestyle education program based on yoga reduces risk factors for cardiovascular disease and diabetes mellitus. *J Altern Complement Med*. 11:267-274.
- [25] Zdrengea, D., L. Poanta, and D. Gaita. 2005. Cardiovascular risk factors and risk behaviors in railway workers. Professional stress and cardiovascular risk. *Rom J Intern Med*. 43(1-2):49-59.
- [26] Ridker, P. M., N. Rifai, L. Rose, et al. 2002. Comparison of C-reactive protein and low-density lipoprotein cholesterol levels in the prediction of first cardiovascular events. *N Engl J Med*. 347:1557– 1565.
- [27] Akosah, K. O., A. Schaper, C. Cogbill, et al. 2003. Preventing myocardial infarction in the young adult in the first place: how do the National Cholesterol Education Panel-III guidelines perform? *J Am Coll Cardiol*. 41:1475–1479.
- [28] Nasir, K., E. D. Michos, R. S. Blumenthal, and P. Raggi. 2005. Detection of high-risk asymptomatic adults with coronary calcium and National Cholesterol Education Panel-III guidelines. *J Am Coll Cardiol*. 46:1931–1936.
- [29] Ajani, U. A., E. S. Ford, and A. H. Mokdad. 2004. Prevalence of high C-reactive protein in persons with serum lipid concentrations within the recommended values. *Clin Chem*. 50:1618–1622.
- [30] Ford, E. S. 2003. The metabolic syndrome and C-reactive protein, fibrinogen, and leukocyte count: findings from the Third National Health and Nutrition Examination Survey. *Atherosclerosis*. 168:351–358.
- [31] Ridker, P. M., N. P. Paynter, N. Rifai, et al. 2008. C-reactive protein and parental history improve global cardiovascular risk prediction. The Reynolds risk score for men. *Circulation*. 118:2243–2251.
- [32] Heilbronn, L. K., M. Noakes, and P. M. Clifton. 2001. Energy restriction and weight loss on very-low fat diets reduce C-reactive protein concentrations in obese, healthy women. *Arterioscler Thromb Vasc Biol*. 21:881–883.
- [33] Kadoglou, N. P., F. Iliadis, N. Angelopoulou, et al. 2007. The anti-inflammatory effects of exercise training in patients with type 2 diabetes mellitus. *Eur J Cardiovasc Prev Rehabil*. 14:837–843.
- [34] Pullen, P. R., S. H. Nagamia, P. K. Mehta, W. R. Thompson, D. Benardot, R. Hammoud, J. M. Parrott, S. Sola, and B. V. Khan. 2008. Effects of yoga on inflammation and exercise capacity in patients with chronic heart failure. *J Card Fail*. 14:407–413.
- [35] Ornish, D., L. W. Scherwitz, J. H. Billings, S. E. Brown, K. L. Gould, T. A. Meritt, S. Sparler, W. T. Armstrong, T. A. Ports, R. L. Kirkeeide, C. Hogeboom, and R. J. Brand. 1998. Intensive lifestyle changes for reversal of coronary heart disease. *JAMA*. 280:2001-2007.
- [36] Damodaran, A., A. Malathi, N. Patil, N. Shah, S. Suryavanshi, S. Marathe. 2002. Therapeutic potential of yoga practices in modifying cardiovascular risk profile in middle aged men and women. *J Assoc Physicians India*. 50:633–640.
- [37] Mahapure, H. H., S. U. Shete, and T. K. Bera. 2008. Effect of yogic exercise on super oxide dismutase levels in diabetics. *International Journal of Yoga*. 1(1):21-26.