

Recent Research in Science and Technology 2013, 5(2): 04-06  
 ISSN: 2076-5061  
 Available Online: <http://recent-science.com/>



# Low cost spectro photometric determination of paraquat in environmental and biological sample

R.Khatoon<sup>1\*</sup>, M.K.Rai<sup>1\*</sup> and J.K.Rai<sup>2</sup>

<sup>1</sup>School of Studies in Chemistry, Pt.Ravi Shankar Shukla University Raipur-492010, India

<sup>2</sup>Chhattisgarh Council of Science & Technology, MIG 25, Indravati Colony, Raipur-492007, India.

## Abstract

An extractive, sensitive spectrophotometer method has been developed for the detection and determination of paraquat using glucose ( as easily available reducing agent ). Paraquat is reduced with glucose in alkaline medium to give a blue colored ion with an absorbance maxima at 610 nm. Beer's law is obeyed in the range 0.5-5.0  $\mu\text{g}$  of paraquat in 10ml of the final solution (ppm). The important analytical parameters and the optimum reaction conditions were evaluated. The method was applied successfully to the determination of paraquat in water, grain, plant material and biological sample.

**Keywords:** spectrophotometer, paraquat, biological samples.

## INTRODUCTION

Paraquat (1,1-Dimethyl-4,4-Bipyridinium Dichloride ion ), also known as Methyl Violegon, is a defoliant and desiccant agent used to control herbal growth in terrestrial and aquatic environments(1). Paraquat is a non-selective acutely toxic herbicide and one of the most commonly used herbicide in the world with a variety of agricultural uses, because of its rapid action, relatively low cost and broad spectrum of its activity(2). It is used to control weeds and grasses in many agricultural and non-agricultural areas. For instance, it used for pre-plant or pre emergence on vegetables grains, potatoes, and peanut areas; post emergence around fruit crops and soyabeans during the dormant season on clover and other legumes(3). Paraquat an oxidative stress inducing substance is a herbicide which is very toxic to all animals and to man as it is well characterized pneumotoxicant. It generates free radical and leads to multi-organ toxicity with necrotic damage to the lung, myocardial muscles, liver and kidney. Paraquat also induce extensive hemorrhagic incidents throughout the body and consequently leads to death(4).


Because of the toxicity and significance of paraquat, several methods, based on different analytical techniques have been reported for the detection and determination of paraquat i.e. Gas chromatography-ion trap mass spectrometry<sup>(6)</sup>, enzymatic-spectrophotometer determination<sup>(7)</sup>, spectrofluometric determination by manual and flow injection methods<sup>(8)</sup>, long-wavelength fluorescence detection method<sup>(9)</sup>, matrix assisted laser desorption/ionization mass spectrometry<sup>(10)</sup>, time - resolved fluoroimmunoassay<sup>(11)</sup>, differential pulse voltammetry<sup>(12)</sup>, High performance liquid chromatographic method<sup>(13)</sup>, etc. Many of the

earlier reported spectrophotometric methods are less sensitive and suffer from many drawbacks . A method for the determination of paraquat using sodium borohydride in an alkaline medium was reported.

## Property

Common Name <sup>(2)</sup>	: Paraquat
Molecular formula <sup>(2)</sup>	: C <sub>12</sub> H <sub>14</sub> Cl <sub>2</sub> N <sub>2</sub>
Chemical Name <sup>(2)</sup>	: 1,1-dimethyl-4,4-bipyridinium
Chemical group <sup>(2)</sup>	: Bipyridyl
Half life in Soil <sup>(5)</sup>	: 644 Days
Solubility <sup>(5)</sup>	: Soluble in water

Structural formula



## METHOD


In the present work, a new extractive, sensitive spectrophotometer method has been developed for the detection and determination of paraquat using glucose ( as easily available reducing agent ). Paraquat is reduced with glucose in alkaline medium to give a blue colored ion with an absorbance maxima at 610 nm . Beer's law is obeyed in the range .5 -5  $\mu\text{g}$  of paraquat in 10ml of the final solution (ppm). The important analytical parameters and the optimum reaction conditions were evaluated. The method was applied successfully to the determination of paraquat in water, grain, plant material and biological sample.

\*Corresponding Author


R.Khatoon  
 School of Studies in Chemistry, Pt.Ravi Shankar Shukla University  
 Raipur,492010, India

Tel: +91-9425520298;  
 Email: [mjkChem@gmail.com](mailto:mjkChem@gmail.com)

### Reaction Mechanism: Paraquat Formation of blue-coloured monocation radical




Scheme:1 Paraquat Reduction Oxidation




Effect of Temperature


## RESULT AND DISCUSSION



Absorbance curve of the Paraquat.



Calibration Curve for the determination of Paraquat.



Effect of pH

Table. Recovery of Paraquat in Water and Vegetables

Sample	Paraquat added μg b	Paraquat obtained in present method* c	% Recovery (c/b)×100
Water**	1	0.98	0.98 98
	2	1.99	0.99 99
Potato***	1	0.98	0.98 98
	2	1.95	0.97 97
Apple***	1	0.99	0.99 99
	2	1.95	0.97 97
Rice***	1	0.98	0.98 98
	2	1.97	0.98 98
Urine**	1	0.96	0.96 96
	2	1.95	0.97 97
Blood**	1	0.97	0.97 97
	2	1.95	0.97 97

\*Mean of three replicate analysis

\*\*Amount of sample 10ml

\*\*\*Amount of sample 5mg

## CONCLUSION

The present method is cheapest and more sensitive and extractive than the other spectrophotometer methods reported for the determination of paraquat. It can be successfully applied for the determination of paraquat in water, fruits, vegetables, cereals and biological sample.

## ACKNOWLEDGEMENTS

Authors are thankful to the Head School of Studies in Chemistry Pt. Ravi Shankar University and Director Genral Chattisgarh Council of Science & Technology for providing laboratory facilities and financial assistance.

## REFERENCES

- [1] Moure M.P., Couse A.B., Calvino D.F., Estvz M.A., Otero R.R., Gandara J.S., 2010. "Paraquat and Diquat Sorption on Iron Oxide Coated Quartz Particles and the Effects of Phosphates", *J. Chem. Eng. Data*, 55, 2668-2672.
- [2] Yamada, Dr. Y., 2002. "Paraquat (057)", *National Food Research Institute, Tsukuba, Japan*.
- [3] Morsheed, M.M., Omar.D., Mohamad, R., Wahed, S., Rahman,

- M.A., 2010. "Airborn Paraquat Measurement and its Exposure to Spray Operators in Treated Field Environment", *International Journal of Agriculture and Biology*, 5, 679-684.
- [4] Ayman, M., Gamal-el-Din, Abdulaziz, A., Yahya, A., Abdulhakeem, A., Majel, A., Abdullah, M., Bekairi, A., Mostofa, A.M., 2005. "Antioxidant Effect of Arabic Gum Against Paraquat Induced Lung Oxidative Stress in Mice", *J. Med. Sci.*, 5(2):55-100.
- [5] Oliveira, M.F., Lichtig, J., Masini, J.C., 2004. "Evaluation of a Nafion Coated Glassy Carbon Electrode For Determination of Paraquat by Differential Pulse Voltammetry", *J. Braz. Chem. Soc. Vol.15, No.5*.
- [6] Moreira, P.N., Pinho, P.G., Boltazar, M.T., Bastos, M.L., Carvalho, F., Dinis-oliveira, R.J., 2012. "Quantification of Paraquat in Postmortem Sample by Gas Chromatography-ion trap mass Spectrometry and Review of the Literature", *Biomedical chromatography*, vol. 26, issue 3, 338-349.
- [7] Yonamine, M., Almeida, R.M.de, 2010. "Enzymatic Spectrophotometer Determination of Paraquat in Urine Sample: A Method Based on its Toxic Mechanism", *Informa Health Care, Toxicology Mechanisms and Methods*, vol.20, No. 7, 424-427.
- [8] Perez-ruiz, T. Lozano, C.M., Tomas, V., Fenoll, J., 1998. "Spectrofluorimetric Determination of Paraquat by Manual and Flow Injection Methods", *Analyst*, Vol.123, 1577-1581.
- [9] Sendra, B., Panadero, S., Hens, A.G., 1999. "Selective Kinetic Determination of Paraquat Using Long-Wavelength Fluorescence Detection", *J. Agric. Food, Chem.* 47, 3733-3737.
- [10] Vermillion-Salsbury, R.L., Hercules, D.M., 1999. "Analysis of Diquat and Paraquat by Solid Phase Extraction Directly Coupled With Matrix Assisted Laser Desorption/Ionization Mass Spectrometry", *Intern. J. Environ. Anal. Chem.* Vol.73(4), 297-308.
- [11] Bacigalupo, M.A., Meroni, G., 2007. "Quantitative Determination of Diuron in Ground and Surface Water By Time-Resolved Fluoroimmunoassay: Seasonal Variations of Diuron, Carbofuran, and Paraquat in an Agricultural Area", *J. Agric. Food Chem.* 55, 3823-3828.
- [12] Bacigalupo, M.A., Meroni, G., 2007. "Quantitative Determination of Diuron in Ground and Surface Water By Time-Resolved Fluoroimmunoassay: Seasonal Variations of Diuron, Carbofuran, and Paraquat in an Agricultural Area", *J. Agric. Food Chem.* 55, 3823-3828.
- [13] Sellero, S.I., Cruz, A., Fernandez, P., Rivadulla, L.M., 2006. "Liquid Chromatographic Method for Simultaneous Determination of Paraquat and Diquat in Plasma, Urine, and Vitreous Humour", *Journal of Liquid Chromatography and Related technologies*, vol. 19, Issue 12, 2009-2024.
- [14] "Directorate of Agriculture", 2011. Indira Gandhi Agriculture University, Raipur (C.G.), Annual report