

Effect of nitrogen and phosphorus sources on amylase production in seed born fungi of maize

Anil U. Kulkarni ¹, Ashok M. Chavan ², Umesh T. Kesare ³ and Sanjay M. Pawar ⁴

¹Department of Botany, Lalbahadur Shashtri Sr. College, Partur (M.S.), India

²Seed Pathology and Fungal Biotechnology Laboratory, Department of Botany, Dr. Babasaheb Ambedkar Marathwada University, Aurangabad (M.S.), India

³Department of Botany, Adarsh College, Omerga, Dist. Osmanabad (M.S.), India

⁴Department of Botany, Shivaji College, Kannad, Dist. Aurangabad (M.S.), India

Abstract

Starch degrading amylolytic enzymes are of great significance in biotechnological application ranging from food, fermentation, and textile to paper industries. Amylase enzyme action of ten dominating fungi viz. *Alternaria alternata*, *Aspergillus flavus*, *A niger*, *A terreus*, *Curvularia lunata*, *Fusarium oxysporum*, *Helminthosporium tetramera*, *Penicillium notatum*, *Rhizoctonia solani* & *Trichoderma viride* isolated from different varieties of maize seeds were studied under the influences of nitrogen & phosphorus sources. The results are very helpful to minimize the bio-deterioration of maize seeds in different storage condition.

Keywords: Amylase, maize seed, seed borne fungi

INTRODUCTION

Production of extracellular hydrolytic enzyme by seeds born fungi has a role during the process of seed deterioration and has been considered helpful to their invasion and colonization. However, Vidhyasekharan et al (1966) [1] claimed that production of extracellular hydrolytic enzymes by *Fusarium moniliforme* and *Aspergillus flavus* was found to be responsible in the spoilage of paddy seeds. Fungi associated with seed in ill storage condition uses the seed content and release the extracellular hydrolytic enzymes. Such enzymatic activities cause bio-deterioration of seeds. Among inorganic nitrogen sources, potassium nitrate and sodium nitrate supported amylase production in case of *Aspergillus awamori* (Musaeva, 1966 [2] & Shilova 1967 [3]). *A tamari*, *Aspergillus flavus*, *A fumigatus*, *Percillum italicum* (Singh and Agrawal, 1981 [4]). Peptone and organic nitrogen sources, was found to be stimulatory in *Aspergillus awamori* (Musaeva 1966 [2]), in some seed born fungi of Bajra (Khairnar, 1967 [5]), Bhosale (1989) [6] reported that, gelatin and urea proved inhibitory amylase production in *Curvularia lunata* and *Fusarium oxysporum*.

The phosphorus sources used in the based medium, dipotassium hydrogen phosphate at very low concentration favoured amylase production in *Lentinus edodes* (El zalaki et al 1980) [7] and potassium dihydrogen phosphate in *Aspergillus niger* (Mahamood et al. 1978) [8].

Considering the fact attempt has been made to study the

impact of nutritional factors on amylase activity of seed borne fungi.

Materials and Methods

Monospore culture of *Alternaria alternate*, *Aspergillus flavus*, *A niger*, *A terreus*, *Curvularia lunata*, *Fusarium oxysporum*, *Helminthosporium tetramera*, *Penicillium notatum*, *Rhizoctonia solani* & *Trichoderma viride* isolated from different varieties of maize seeds and maintained on PDA slants.

Production of amylase was studied by growing the test fungi in liquid medium containing soluble Starch 1%, KNO₃ 0.25%, KH₂PO₄ 0.1% and MgSO₄.7H₂O 0.05%, PH of medium was adjusted at 5.5. Twenty five ml of the medium was poured in 100ml conical flasks autoclaved and inoculated separately with 0.1ml spore suspension of the test fungi which were grown for 7 days on PDA slants. The flasks were incubated for 6 days at 25±1°C with diurnal periodicity of light on 7th day, the flasks were harvested by filtering the contents through whatman filter no. 1, and the filtrates were collected in pre-sterilized bottles and termed as crude enzyme preparation.

Determination of amylase activity was done with the help of cup plate method which was adopted by Singh and Saxena (1982) [9], where 20ml of starch assay medium (Soluble starch - 10gm, Na₂HPO₄ - 2.84 gm, NaCl - 0.35 gm, Agaragar - 20 gm, distilled water 1000ml and PH 6.9) was poured in each Petriplate. On solidification of the medium a cavity (08 mm diameter) was made in the centre with the help of a corkborer (No. 4) and was filled with 1ml culture filtrates (crude enzyme preparation) of the test fungi. The plates were incubated at 28°C for 24 hours, than they were flooded with Lugol's iodine solution as an indicator. A clear non blue, circular zone obtained surrounding the central cavity; diameter was measured (mm) as the amylase activity zone similar procedure followed for the control. The KNO₃ served as control in the starch nitrate containing sources and in phosphorus sources potassium dihydrogen phosphate as control.

Received: July 12, 2012; Revised: Aug 22, 2012; Accepted: Sept 25, 2012.

*Corresponding Author

Anil U. Kulkarni
 Department of Botany, Lalbahadur Shashtri Sr. College, Partur (M.S.), India

Email: kulkarniau@rediffmail.com

RESULTS AND DISCUSSION

From table 1 and graph 1 reveals that calcium nitrate was stimulatory for amylase production in *Alternaria alternata*, *Aspergillus flavus*, *A niger*, *A terrus* and *Helminthosporium tetramera* where as sodium nitrate was found to be stimulatory for *Alternaria alternata*, *Aspergillus niger*, *A terrus*, *Curvularia lunata*, *Fusarium oxysporum*, *Helminthosporium tetramera*.

Sodium nitrate was stimulatory for amylase production in *Aspergillus niger*, *A terrus*. In ammonia form ammonium oxalate showed stimulatory effect for amylase production in *Aspergillus niger*, *A terrus* and *Curvularia lunata* all other forms of nitrogen sources showed minimum production of amylase in all ten tested fungi. Organic nitrogen sources are preferred for the production of α -amylase. A maximum α -amylase production was supported by yeast extract, peptone or beef extract (Hamilton et al., 1999 [10], Emanuilova and Toda, 1984 [11], Krishnan and Chandra, 1982 [12],

Hayashida et al., 1988 [13]). Urea & Casein strongly inhibited amylase production in all fungi. Gelatin stimulated the amylase production in *Alternaria alternata*, *Aspergillus niger*, where as peptone stimulated the amylase production in *Curvularia lunata*, *Helminthosporium tetramera*. About relationship between growth and amylase production, there was no any clear cut difference among them. However, *Aspergillus* genera viz *Aspergillus flavus*, *A niger*, *A terrus* showed maximum amylase production in the presence of calcium nitrate and sodium nitrate than any other nitrogen source.

It is clear from the data given in table 2 and graph 2 that in all fungi, potassium dihydrogen phosphate (control) was found to be better than the other phosphorus sources, on the contrary ammonium bi-phosphate for *Alternaria alternata*, *Trichoderma viride*, *Rhizoctonia solani* showed minimum production of amylase. Whereas *Aspergillus flavus*, *A niger* and *A terrus* showed maximum amylase production in potassium dihydrogen phosphate (control) followed by sodium dihydrogen phosphate.


Table 1. Effect of Nitrogen Sources on Amylase production in seed borne fungi

Nitrogen Sources (0.25% conc)	Fungi									
	Alal	Asfl	Asni	Aste	Culu	Fuox	Hete	Peno	Rhso	Trvi
A] Nitrates										
1] Cal nitrate	22	36	32	35	20	30	24	18	18	17
2] Sod nitrate	28	33	30	35	26	35	26	20	20	17
B] Nitrites										
Sodium nitrite	20	30	27	24	-	26	-	22	-	18
C] Ammonia forms										
1] Ammo oxalate	18	32	32	30	23	-	20	-	-	-
2] Amm chloride	-	-	-	-	-	-	-	-	-	-
3] Amm nitrate	-	24	23	20	-	-	-	-	-	-
4] Amm sulphate	-	23	24	21	-	-	-	-	-	-
D] Amids										
Urea	-	21	-	-	-	-	-	-	-	-
E] Organic forms										
1] Casein	12	23	12	20	20	10	21	10	16	16
2] Gelatin	25	20	30	23	-	30	-	-	-	-
3] Peptone	13	20	13	23	22	24	25	11	17	16
Control										
(Pot nitrate KNO ₃)	20	35	23	24	20	33	20	22	20	18


Table 2. Effect of Phosphorous sources on Amylase production in seed borne fungi

Phosphate Sources (0.1%)	Alal	Asfl	Asni	Aste	Culu	Fuox	Heyr	Peno	Rhso	Trvi
Phosphate Sources										
1] Amm biphos	14	22	23	20	19	20	19	18	16	15
2] Cal. Phos	15	24	24	21	17	20	20	19	17	15
3] Amm Phosphate	15	23	23	21	19	19	19	18	17	16
4] Plot diti.phos (C)	20	30	32	30	20	21	23	20	20	18
5] Sod di phos	20	28	26	27	20	20	20	19	18	18

Alal – <i>Alternaria alternata</i>	Fuox – <i>Fusarium oxysporum</i>
Asfl – <i>Aspergillus flavus</i>	Hetr – <i>Helminthosporium tetramere</i>
Asni – <i>Aspergillus niger</i>	Peno – <i>Penicillium notatum</i>
Aste – <i>Aspergillus terus</i>	Rhzo – <i>Rhizoctonia solani</i>
Culu – <i>Curvularia lunata</i>	Trvi – <i>Trichoderma viride</i>



Graph 1.



Graph 2.

ACKNOWLEDGMENT

We thank, Head Department of Botany Dr. Babasaheb Ambedkar Marathwada University, Aurangabad (M.S.) for providing laboratory facilities.

REFERENCES

[1] Vidhyasekaran, P.G., Murthaswamy and Subramanian, C.L., 1966. Role of seed borne microflora in paddy seeds. *Indan Phytopath.* 19: 333-341.

[2] Musaeva, T.L. 1966. Effect of inorganic nitrogen on the synthesis of amylolytic enzymes in submerged culture of *Aspergillus awamori*. *Uzbekskii Biolzh.* 9: 12-14.

[3] Shilova, A. A.1967. Effect of the composition of the medium on the synthesis of enzyme formed by submerged culture of fungus *Aspergillus awamori* Tr., Vses. Nauch-Issled inst ferment spirit from Ref. *Zh.Biol.*3:674.

[4] Singh, K.V. and Agrawal, S.C. 1981. Amylase production by some detmatophytes. *Act Bot. Indica.* 9: 32-34.

[5] Khairnar, D. N.1987. Studies on seed-born fungi of bajra, Ph.D. thesis Marathwada Uni. Aurangabad (M.S.).

[6] Bhosale, A. M.1989. Studies on fungal amylase and toxins in grain deterioration of sorghum. Ph.D. thesis, Dr. B.A.M.U. Aurangabad (M.S.).

[7] Zalaki, El. Esmat, M. and Hamza, M. A.1980. Propagation of lentinus endodes on modified malt sprout medium for amylase

production *Food chem.* 5: 131-138.

- [8] Mahmood, S.A. and Mashor, W.A. MEL Haddad and Reffaal, A.A.1978. Production α -amylase by *Aspergillus niger*, Ain Shan Uni. *Fac. Agric, Res. Bull.* 6: 1-16.
- [9] Singh, B. S. and Saxena, S. B.1982. Extra cellular amylolytic activity of penicillia. *Jou. Indian Bot. Soc.* 12:223.
- [10] Hamilton, L.M., Kelly, C.T. and Fogarty, W.M. 1999. Production and properties of the raw starch digesting α -amylase of *Bacillus* Sp. IMD 435. *Process. Biochem.* 35: 27-31.
- [11] Emanuilova, El. And Toda, K. 1984. α -amylase production in batch and continuous cultures by *Bacillus cladoycticus*. *Appl. Microbiol. Biotechnol.* 19 : 301-305.
- [12] Krishnan, T. Chandra, A.K.1982. Effect of oil seed cakes on α -amylase production by *Bacillus Licheniformis*, CUMC - 305 *Appl. Enoiron Microbiol.* 44: 270-274.
- [13] Hayashida, S., Teramotor, Inoue, T.1988. Production and characteristics of raw potato starch digesting α -amylase from *Bacillus subtilis*, *Appl. Environ. Micro Biol.* 54, 1561-1522.