

Influence of water application on photosynthesis, growth and biomass characteristics in *Jatropha curcas*

Krishan K. Verma¹, Shweta Vatsal¹, Ramwant K. Gupta², Sanjay Ranjan³, Chhedi Lal Verma⁴ and Munna Singh^{1,2}

¹Department of Botany, University of Lucknow, Lucknow- 226 007 (U.P.), India

²Department of Plant Physiology, College of Basic Science and Humanities (CBSH), G.B. Pant University of Agriculture and Technology, Pantnagar – 263145 (Uttarakhand), India

³Division of Plant Physiology, National Botanical Research Institute, Lucknow- 226001 (U.P.), India

⁴Central Soil Salinity Research Institute, Lucknow- 226 005 (U.P.), India

Abstract

The effect of CO₂ assimilation, stomatal conductance, transpiration rate, water use efficiency, growth and biomass productivity were studied in *Jatropha curcas* under different moisture levels of water (100, 75, 50 and 25% of field capacity). CO₂ assimilation, stomatal conductance, transpiration, growth and biomass were reduced in response to decreasing moisture content of water. The decreased CO₂ assimilation during irrigation stress was found largely dependent on stomatal closure, which reduced available internal CO₂ concentration and restricted water loss through transpiration based on leaf gas exchange hypothesis linked with stomatal limitation for photosynthesis to reduce carbon uptake followed by loss in leaf area expansion which declined total carbon uptake, growth and biomass in *Jatropha curcas* seedlings.

Keywords: Biomass, CO₂ assimilation, growth, stomatal conductance, transpiration rate, water use efficiency

INTRODUCTION

Fossil fuel reserves over the globe are decreasing and global prices soaring continuously putting tremendous pressure on developing economies. Therefore, alternate sources of fossil fuel are being searched by the researchers (1). *Jatropha curcas* has drawn much attention for producing biodiesel (2,3,4,5) being an energy plant, also shown drought- tolerance with good growth on poor quality of land (5,6). The plant parts have been used for various purposes such as animal feeding, medicine production and restoration of disturbed ecosystem (7,8,9). The seeds contain viscous oil used in soap industry, cosmetics and as a bio-diesel to substitute kerosene (8,9).

Water is essential for various metabolic activities. Its deficiency induces water stress on vegetation in combination with soil, plant and climate. All these factors interact to determine the water absorption and loss by the plants (10). In arid and semi-arid regions, plants are often exposed to water deficit stress, also known as moisture stress which negatively influence plant growth and biomass productivity (11,12,13). The plants can avoid moisture stress by maximizing water uptake (i.e., absorbing ground water by deep roots) or minimizing water loss through stomatal closure – small thick leaves, etc. (14). The occurrence of morphological and physiological responses may lead to some adaptation to moisture

stress may vary considerably among species (15). The apparent consequences of moisture stress is the progressive decline in photosynthetic capacity though photosynthetic system found to be resistant to irrigation levels linked with stomatal closure (16). The reduced leaf water reserve in relation to the limiting transpiration rate (10) eventually impairs biomass accumulation (17). The aim of this study was to be evaluate the responses of CO₂ assimilation, stomatal conductance, transpiration rate, water use efficiency, growth and biomass in *Jatropha curcas* subjected to different irrigation levels.

MATERIALS AND METHODS

Plant material and experimental design

One year old *Jatropha curcas* plants were chosen for obtaining their stem cuttings (18- 20 cm). These cuttings were shifted in polybags filled with 2 kg normal soil for 15 days to ensure sprouting (March, 2009). After two weeks, they were transplanted in earthen pots (30cm diameter and 30cm depth, filled with fertile soil and humus 3:1). The pots were watered daily to the field capacity to ensure that the plants did not experience drought. Fifty days old uniform plants were chosen for the study purpose and subjected to four moisture regimes (100, 75, 50 and 25% field capacity) in a glass house at Botany Department, Lucknow University, Lucknow. Microclimatic parameters such as temperature, relative humidity, rainfall and sunlight intensity were not controlled but recorded weekly (Fig. 1). The soil moisture content measured by calculating weight of 100gm of the soil from each pot under:

$$\text{Soil moisture content (\%)} = \frac{\text{Fresh weight} - \text{Dry weight}}{\text{Fresh weight}} \times 100$$

The average soil moisture content was retained ca. 40, 33, 21 and 10% in the pots during irrigation regimes viz., 100, 75, 50 and

Received: July 20, 2012; Revised: Aug 10, 2012; Accepted: Sept 25, 2012.

*Corresponding Author

Krishan K. Verma

Department of Botany, University of Lucknow, Lucknow- 226 007 (U.P.), India

Email: kverma22feb@yahoo.in

25%, throughout.

Measurement of CO₂ assimilation, stomatal conductance, transpiration and water use efficiency

CO₂ assimilation, stomatal conductance and transpiration were measured with an open system CIRAS-1 portable Infra Red Gas Analyzer photosynthesis system (IRGA, PP System, England). Measurements were made in natural sunlight (1500- 2000 $\mu\text{mol m}^{-2} \text{s}^{-1}$) between 10:00- 11:00am by using mature and fully expanded leaf (6- 8th) from control and moisture stress seedlings. Water use efficiency (WUE) was calculated (WUE=CO₂ assimilation/transpiration rate) as described by (18). All measurements were made in the forenoon to avoid high temperature and low humidity in the afternoon.

Growth and biomass characteristics

At the end of experiment, plants (n= 10) were harvested to analyse the pattern of growth and biomass i.e., height of plants, collar diameter, number of leaves, leaf area expansion, root length and their biomass. The height of plants were measured from the apex to the starting point of the basal leaf (19). The stem - collar diameter measured at ~15 cm above from the soil surface by using calliper. The leaf area expansion measured by using Leaf Area Meter CI- 202 (CID Inc., USA). The biomass measured by using different plants' parts of the each plant, separated and washed to

remove debris and then dried (68 ± 2 °C, 72 h) until constant weight achieved.

Statistical analysis

Data management and statistical analysis were performed using SPSS 16.0 software (SPSS Inc., Chicago. II USA).

RESULTS

The water stress as imposed by maintaining various irrigation levels compared to field capacity reduced CO₂ assimilation, stomatal conductance, transpiration and water use efficiency. The water stress (75, 50 and 25%) induced decline trends 15 days after the onset of water stress. The loss in CO₂ assimilation did occur in the range of 20 - 78%, incase *Jatropha curcas* subjected under water stress i.e., 75, 50 and 25% respectively (Fig. 2A).

The stomatal conductance, transpiration rate and water use efficiency were found down-regulated ca. 45- 63%, 30-51%, and 31- 53% respectively. *Jatropha curcas* seedlings after the application 60 days of various irrigation levels i.e., 75, 50 and 25% compared to 100% (upto field capacity) for stomatal conductance, transpiration and water use efficiency. The optimal reduction values of stomatal conductance, transpiration and water use efficiency was observed ca. 63, 51 and 53% incase irrigation level 25% of the field capacity ensured throughout (Fig. 2B, C and D).

Table 1. Growth and biomass characteristics of *Jatropha curcas* seedlings grown under different irrigation levels (75, 50 and 25 %). The control plants were irrigated to the level of field capacity (100%). Data are the mean (\pm) standard error of 8- 10 individuals.

Characteristics	% irrigation of water in relation to field capacity			
	100	75	50	25
Plant height (cm)	50.5 \pm 3.3	47.7 \pm 3.8	38.1 \pm 3.0	31.0 \pm 2.0
Loss (%)	-	5.54	24.55	38.61
Collar diameter (cm)	7.3 \pm 0.54	7.0 \pm 0.18	5.4 \pm 0.32	4.4 \pm 0.19
Loss (%)	-	4.11	26.03	43.84
Number of leaves	34 \pm 4	30 \pm 2	26 \pm 2	19 \pm 3
Loss (%)	-	11.76	23.53	44.12
Leaf area expansion (m ² plant ⁻¹)	10.0 \pm 1.3	8.9 \pm 0.31	6.1 \pm 0.5	4.53 \pm 0.23
Loss (%)	-	10.97	38.68	54.84
Specific leaf area (g m ⁻²)	1.97 \pm 0.02	1.8 \pm 0.02	1.32 \pm 0.01	1.19 \pm 0.02
Loss (%)	-	8.63	32.99	39.59
Biomass of leaf (g)	10.71 \pm 1.1	9.99 \pm 1	7.84 \pm 1.3	5.94 \pm 0.89
Loss (%)	-	6.72	26.8	44.54
Biomass of stem (g)	15.38 \pm 1.1	14.44 \pm 0.9	13.02 \pm 0.98	11.33 \pm 0.06
Loss (%)	-	8.06	15.34	26.33
Biomass of root (g)	4.19 \pm 0.3	4.02 \pm 0.42	3.13 \pm 0.04	2.82 \pm 0.03
Loss (%)	-	4.06	25.30	32.70
Total biomass (g)	30.28 \pm 1.4	28.15 \pm 1.1	23.99 \pm 0.2	20.09 \pm 1.9
Loss (%)	-	7.03	20.77	33.65
Length of root (cm)	37.5 \pm 2.1	35.8 \pm 1.7	25.8 \pm 1	21 \pm 1.8
Loss (%)	-	4.53	31.2	44.0
Leaf moisture content (%)	84.83 \pm 6	84.08 \pm 4.3	82.47 \pm 3.7	81.9 \pm 4.3
Loss (%)	-	0.88	2.78	3.40
Leaf weight ratio	0.57 \pm 0.006	0.55 \pm 0.008	0.47 \pm 0.003	0.41 \pm 0.002
Loss (%)	-	3.51	17.54	21.05

At the end (60 days) of the experiment, height of plants, collar diameter, number of leaves, leaf area expansion, specific leaf area, leaf water content, leaf weight ratio, root length and plant biomass monitored and found significantly reduced alongwith loss in the levels of applied irrigation levels compared to control which was experienced irrigation level to field capacity (100%) throughout. The height of plants and collar diameter was reduced 5- 39% and 4- 44%

(Table 2). The withdrawal of irrigation to the level of 25% severely impaired plant height, collar diameter ca. 39 and 40%. It has also declined number of leaves, leaf area expansion and specific leaf area ca. 12- 44%, 11- 55% and 8- 40%. Similarly, root length, leaf water content and leaf weight ratio also found affected to the level of 4 - 44%, 0.8 - 5% and 4 - 21% along with loss in biomass yield i.e., leaf, stem, root and total dry weights to the levels of 7- 45, 8- 26, 4-

33 and 7-34% (Table 1).




Fig 1. Environmental variables viz., average temperature (°C), relative humidity (%) and rainfall (mm) during the cultivation period (Feb. - June, 2009) of *Jatropha curcas*. (The seedlings- plants were grown in the range of PFDs ca. 1500 – 2000µmol m⁻² s⁻¹).




Fig 2. A - CO₂ assimilation rate (P_N), B - stomatal conductance (g_s), C - transpiration (E) and D - water use efficiency (WUE) in *Jatropha curcas* influenced by various irrigation levels (75, 50 and 25%) to establish moisture/ drought stress (75, 50 and 25%). The control were irrigated to the level of field capacity (100%), throughout. Vertical bars indicate SE (n = 6).

DISCUSSION

The effect of CO₂ assimilation, stomatal conductance, transpiration rate, water use efficiency, growth and biomass productivity were monitored in *Jatropha curcas* plants as influenced by various irrigation levels (100, 75, 50 and 25%). The reduction in

CO₂ assimilation (Fig. 1A) was largely found associated with stomatal closure, which decreased available internal CO₂, also restricted water loss through transpiration (15), resulted in a decline of photosynthetic capacity (20,21). Water stress induced a impaired stomatal conductance, transpiration with prevented water loss in *Jatropha curcas*. Consequently it decreased CO₂ assimilation (10)

because stomatal aperture affects both photosynthesis and transpiration (17). Similar results were previously obtained in legume species (22,23,24,25,26). The sustenance of the severe moisture stress also affects integrity of photosynthetic apparatus which may result in its irrepairable injuries linked with high temperature and low relative humidity (27). Hence, loss in the levels of water application which could cause rhizospheric. Moisture stress, negatively affected plant growth and biomass productivity due to availability of limited water which down-regulated leaf area expansion and carbon assimilation (21) alongwith impaired morphology – canopy i.e., plant height, collar diameter, number of leaves, leaf area expansion, specific leaf area (28,29,30). Our results are consistent with many previous studies (30,31,32,33,34). In conclusion, decreasing irrigation levels down-regulated biomass partitioning (10).

ACKNOWLEDGEMENT

The authors are grateful to the Head, Department of Botany, University of Lucknow, Lucknow, for providing necessary facilities during experimentation.

REFERENCES

- [1] Verma K K, Singh M, Verma C L. 2012. Developing a mathematical model for variation of physiological responses of *Jatropha curcas* leaves depending on leaf positions under soil flooding. *Acta Physiol Plant* Published online 28 Jan., 2012. Doi 10.1007/s11738-012-0941-y
- [2] Gubitz GM, Mittelbach M, Trabi M. 1999. Exploitation of the tropical oil seed plant *Jatropha curcas* L. *Bioresour Technol* 67:73–82
- [3] Achten WMJ, Mathijs E, Verchot L, Singh VP, Aerts R, Muys B. 2007. *Jatropha* biodiesel fueling sustainability? *Biofuels Bioprod Biorefining* 1:283–291
- [4] Fairless D. 2007. Biofuel: the little shrub that could: maybe. *Nature* 449:652–655
- [5] King AJ, He W, Cuevas JA, Freudenberger M, Ramiaramana D, Graham IA. 2009. Potential of *Jatropha curcas* as a source of renewable oil and animal feed. *J Exp Bot* 60:2897–2905
- [6] Wang WG, Li R, Liu B, Li L, Wang SH, Chen F. 2011. Effects of low nitrogen and drought stresses on proline synthesis of *Jatropha curcas* seedling. *Acta Physiol Plant* Published online 20 Jan., 2011. Doi 10.1007/s11738-010-0692-6
- [7] Heller J. 1996. Physic Nut. *Jatropha curcas* L. Promoting the conservation and use of underutilized and Neglected crops. 1. Institute of Plant Genetics and crop plant research, Gatersleben, International Plant Genetic Resources Institute, Rome, Italy
- [8] Openshaw K. 2000. A review of *Jatropha curcas*: an oil plant of unfulfilled promise. *Biomass Bioenergy* 19: 1-15
- [9] Liang Y, Chen H, Tang MJ, Yang PF, Shen S H. 2007. Responses of *Jatropha curcas* seedlings to cold stress: photosynthesis-related proteins and chlorophyll fluorescence characteristics. *Physiol Plant* 131: 508-517
- [10] Baquedano FJ, Castillo FJ. 2006. Comparative ecophysiological effects of drought on seedlings of the Mediterranean water-saver *Pinus halepensis* and water-spenders *Quercus coccifera* and *Quercus ilex*. *Trees* 20: 689-700. Doi 10.1007/s00468-006-0084-0
- [11] Wu FZ, Bao WK, Li FL, Wu N. 2008. Effects of drought stress and N supply on the growth, biomass partitioning and water-use efficiency of *Sophora davidii* seedlings. *Environ Exp Bot* 63:248–255
- [12] Efeoglu B, Ekmekci Y, Cicek N. 2009. Physiological response of three maize cultivars to drought stress and recovery. *S Afr J Bot* 75:34–42
- [13] Zhang M, Chen Q, Shen S. 2011. Physiological responses of two Jerusalem artichoke cultivars to drought stress induced by polyethylene glycol. *Acta Physiol Plant* 33: 313–318. Doi 10.1007/s11738-010-0549-z
- [14] Kozłowski, T.T., Pallardy, S.G., 2002. Acclimation and adaptive responses of woody plants to environmental stresses. *The Botanical Review* 68(2), 270-334
- [15] Souza RP, Machado EC, Silva JAB, Lagoa AMMA, Silveira JAG. 2004. Photosynthetic gas exchange, chlorophyll fluorescence and some associated metabolic changes in cowpea (*Vigna unguiculata*) during water stress and recovery. *Environ Exp Bot* 51: 45-56
- [16] Cornic G. 2000. Drought stress inhibits photosynthesis by decreasing stomatal aperture-not by affecting ATP synthesis. *Trends Plant Sci* 5: 187-188
- [17] Osipova S V, Permyakov A V, Permyakova M D, Pshenichnikova T A, Bořner A. 2011. Leaf dehydroascorbate reductase and catalase activity is associated with soil drought tolerance in bread wheat. *Acta Physiol Plant* Published online 13 April 2011 Doi 10.1007/s11738-011-0756-2
- [18] Mielke MS, De Azeiteiro AAF, Gomes FP, Aguiar MAG, Mangabeira PAO. 2003. Leaf gas exchange, chlorophyll fluorescence and growth responses of *Genipa americana* seedlings to soil flooding. *Environ Exp Bot* 50: 221-231
- [19] Wielgolaski FE. 1999. Starting dates and basic temperatures in phenological observations of plants. *Int J Biometeorol* 42: 158-168
- [20] Matos MC, Campos PS, Ramalho JC, Medeira MC, Maia MI, Semedo JM, Marques NM, Matos A. 2002. Photosynthetic activity and cellular integrity of the Andean legume *Pachyrhizus ahipa* (Wedd.) Parodi under heat and water stress. *Photosynthetica* 40(4): 493- 501
- [21] Singh B, Singh G. 2006. Effects of controlled irrigation on water potential, nitrogen uptake and biomass production in *Dalbergia sissoo* seedlings. *Environ Exp Bot* 55: 209-219
- [22] Ramalho JC, Chaves MM. 1992. Drought effects on plant water relations and carbon gain in two lines of *Lupinus albus* L. *Eur J Agron* 1: 271-280
- [23] Lauriano JA, Campos PS, Ramalho JC, Lidon FC, Guedes ME, Matos MdoC. 1997. Partial decline of *Arachis hypogaea* L. photosynthesis triggered by drought stress. *Photosynthetica* 33: 81-90
- [24] Lauriano JA, Lidon FC, Carvalho CA, Campos PS, Matos M do C. 2000. Drought effects on membrane lipids and photosynthetic

- activity in different peanut cultivars. *Photosynthetica* 38: 7-12
- [25] Matos MC, Lauriano JA, Campos PS, Ramalho JC, Semedo JM, Marques NM. 1998. Drought acclimation effects on gas exchanges and accumulation of some osmotic solutes in four *Arachis hypogaea* genotypes. – In: Garab G. 9ED.0: Photosynthesis: Mechanisms and effects. Vol. 4. Pp. 2557-2560. Kluwer Academic Publishers, Dordrecht- Boston-London
- [26] Campos PS, Ramalho JC, Lauriano JA, Silva MJ, Matos MdoC. 1999. Effects of drought on photosynthetic performance and water relations of four Vigna genotypes. *Photosynthetica* 36: 79-87
- [27] Xu ZZ, Zhou GS. 2005. Effects of water stress on photosynthesis and nitrogen metabolism in vegetative and reproductive shoots of *Leymus chinensis*. *Photosynthetica* 43(1): 29-35
- [28] Fernandez RJ, Wang MB, Reynolds JF. 2002. Do morphological changes mediate plant response to water stress? A steady-state experiment with two C₄ grasses. *New Phytol* 155: 79-88
- [29] Marron N, Dreyer E, Boudouresque E, Delay D, Petit JM, Delmotte FM, Brignolas F. 2003. Impact of successive drought and re-watering cycles on growth and specific leaf area of two *populus × canadensis* (Moench) clones, 'Dorskamp' and 'LuisaAvanzo'. *Tree Physiol* 23: 1225-1235
- [30] Zhang X, Zhang R, Li C. 2004. Population differences in physiological and morphological adaptations of *Populus davidiana* seedlings in response to progressive drought stress. *Plant Sci* 166: 791-797
- [31] Li C. 2000. Population differences in water-use efficiency of *Eucalyptus microtheca* seedlings under different watering regimes. *Physiol Plant* 108: 134-139
- [32] Amdt SK, Cliffork SC, Wanek W, Jones HG, Popp M. 2001. Physiological and morphological adaptations of the fruit tree *Ziziphus rotundifolia* in response to progressive drought stress. *Tree Physiol* 21: 705-715
- [33] Yin C, Duan B, Wang X, Li C. 2004. Morphological and physiological responses of two contrasting Poplar species to drought stress and exogenous abscisic acid application. *Plant Sci* 167: 1091-1097
- [34] Lei Y, Yin C, Li C. 2006. Differences in some morphological, physiological and biochemical responses to drought stress in two contrasting populations of *Populus przewalskii*. *Physiol Plant* 127: 182-191