

Research in Biotechnology, 4(1): 21-30, 2013

ISSN: 2229-791X

www.researchinbiotechnology.com

Regular Article

Impact of sodium and potassium ions in identification of offspring gender in high sugar rabbits

S. Chandraju^{1*}, Ashraf Beirami¹ and C. S. Chidan Kumar²

¹Department of Studies in Sugar Technology, Sir M.Vishweshwaraya Post-Graduate Center, University of Mysore, Tubinakere, Mandya, Karnataka, India

²Department of Engineering Chemistry, Alva's Institute of Engineering & Technology, Shobhavana Campus, Mijar, Moodbidri-574225, South Canara Dt. Karnataka, India

*Corresponding author E-mail:chandraju1@yahoo.com

Impact of sodium and potassium in diet to determine offspring gender in high-sugar rabbit has been investigated. Diabetes mellitus is a chronic, widely spread disease in living species. Sex determination has scientific basis for prevention of genetic diseases in addition to social backgrounds. 21 rabbits (obtained from veterinary college Bangalore), were divided in to three groups in the rabbits male to female 1:1. The first group was made diabetic with Na and K, the second group was non diabetic with Na and K, group third was control unit (Non-diabetic without Na and K). It was found that the delivered offsprings male to female ratio were 3.4:1.4:1 and 1:1 for the first, second and third groups respectively. Also, it was found that non diabetic rabbits fed with normal food yields maximum numbers of offspring (76), while non diabetic rabbits fed with Na and K yields 60 and diabetic rabbits with Na and K yield lowest numbers of offspring (40).

KeyWords: Sodium, Diabetes, Potassium, Sex ratio, Rabbit, Offspring.

Pregnancy is a state that allows a life form to develop with the support and protection from mother's body. The growth and development of the fetus in gestation is partially determined by the genome of the fetus, which produces its own growth factors as well as the majority of its hormones. However, this genetic influence is highly dependent upon interaction with environmental factors (Van Assche et al., 2001). One environmental factor vital in the growth and development of the fetus is nutrition. The fetus is solely dependent on the mother to supply its nutrients. It is also

dependent on the placenta, an essential organ in pregnancy, to transfer these nutrients from the maternal system to its own. Thus the fetal nutrition is a reflection of that of the mother's. This interaction exists in a sensitive equilibrium; if disturbed, there are fetal developmental consequences (Van Assche et al., 2001). Preselection of the gender of offspring is a subject that has held man's attention since the beginning of recorded history. Although scientific studies on genes have been conducted recently, sex selection and gender preference have been considered since ancient time. Anaxagoras, a greek

Received: 21.11.2012; Revised: 23.1.2013; Accepted: 12.2.2013

scientist was the first person who related the sex of fetus to testis (Mittwoch, 2005).

Gestational diabetes mellitus (GDM) is a frequent complication of pregnancy, affecting 3.5% of pregnancies in the United States (Engelgau et al., 1988). Because obesity and age are major risk factors for GDM (Feig et al., 2002), the prevalence of GDM is increasing (Solomon et al., 1997). Although GDM may represent a previously unrecognized state of continuous hyperglycemia (ie, diabetes), most women with GDM show glucose intolerance that does not persist after pregnancy. Women with GDM have been shown to be more insulin-resistant than normal pregnant women, and their insulin secretion is defective relative to the degree of insulin resistance (Xiang et al., 1999). There is a strong association between pregnancy in women with any form of diabetes and high infant mortality and morbidity in their offspring (Steninger et al., 1998).

Over the past 25 years, the rate of neonatal mortality among infants of diabetic mothers has declined from 250 per 1000 live births to approximately 20 per 1000 live births. About half of these deaths are due to congenital malformations that result from the diabetic intra-uterine environment (Weintrob et al., 1996). The frequency of congenital malformation, as well as morbidity associated with maternal diabetes, is directly related to the severity of the diabetes (Ornoy et al., 1998). Abnormalities in systems such as the cardiovascular system, musculoskeletal, and central nervous system occur 5 times more often in the offspring of diabetic mothers. Rare abnormalities like sacral agenesis and caudal regression syndrome occur between 200 and 400 times more frequently than the non-diabetic population (Weintrob et al., 1996). There are many Methods of sex selection such as: The consumption of particular foods, the use of various vaginal douches and the timing of

intercourse in relation to ovulation, Sperm sorting, Pre-implantation genetic diagnosis (PGD), Selective abortion, Infanticide, Periconceptual methods, postconceptual methods.

There are also methods which use different food combinations and especial diets to maximum the chance of having a baby with specific sex. The old believe is that eating salty, savory foods leads to delivering a male and sodium rich foods to a girl. Some believes that the ratios of the minerals sodium, potassium are important in determination of offspring gender. It was shown that pregnant female house mice maintained on a consistent low-food diet give birth to a lower proportion of males than do control females fed ad libitum (Meikle et al., 1995).

As a part of our ongoing research, we studied the Role of Sodium and Potassium ions in identification of offspring gender in High-sugar mammals (Chandarj et al., 2011).

Materials and Methods Medicine

Streptozotocin or Streptozocin or Izostazin or Zanosar (STZ) is a synthetic antineoplastic agent that is classically an anti-tumor antibiotic and chemically is related to other nitrosureas used in cancer chemotherapy. Streptozotocin sterile powders are provided and prepared as a chemotherapy agent. Each vial of sterilized Streptozotocin powder contains 1 g. of Streptozotocin active ingredient with the chemical name, 2-Deoxy-2- [(methyl-nitrosoamino) - carbonyl]amino]-D-glucopyranose and 200 K citric acid. Streptozotocin was supplied by Pharmacia Company. Streptozotocin is available for intravenous use as a dry-frozen, pale yellow, sterilized product. Pure Streptozotocin has alkaline pH. When it is dissolved inside the vial in distilled water as instructed, the pH in the solution inside the vial will be 3.5-4.5 because of the presence of citric acid. This material is

prepared in 1-g vials and kept in cold store and refrigerator temperature (2-8 °C) away from light. Control animals were given an equivalent volume of citrate buffer solution.

Experimental design

Polyphagia for at least one week were considered to be diabetes (STZ rats). 7 control rabbits [non diabetic (Na, K) and 7 neither diabetic nor (Na, K) were 21 Adult female rabbits weighting 1100-1300 g (9 months old), to Induction of diabetes, rabbits were fasted for 12 h before inducing diabetes, 7 adult rabbits were used for inducing diabetes. The rabbits were injected by a single intraperitoneal streptozotocin at the dose of 40 K/kg of the body weight. STZ was freshly dissolved in 0.05 M citrate buffer, pH 4.5. For the i.p. injection of STZ, the rabbits were held in one hand in dorsal position, the injection site was swabbed using povidone- iodine solution and the designated amount of STZ was injected within 10' after preparation in the caudal abdominal cavity using sterile 25g insulin needle. Streptozotocine induces diabetes within 3 days by destroying the beta cells (Karunanayake, 1975). Blood was collected for glucose determination using a glucometer (Accutrend Glucose, Roche Diagnostics, and Mannheim, Germany). Blood glucose levels were measured on the third day, STZ injected rabbits with blood glucose levels 15 m mol/l (270K/dl) as well as polydipsia, polyuria and injected with an equal volume of citrate buffer solution.

Diabetic rabbits and non-diabetic control group were kept in metabolic cages individually and separately and within 16-21 days, on the specified diets (Na, K and non Na, K) feeding and metabolism control. The first group diabetics (Na, K) and second group non diabetics (Na, K) was supplied with drinking water mixed with 1% sodium and potassium, the third group was chosen as a control group neither diabetics nor Na, K, pure drinking water was supplied. After

16-21 days, on the specified diets, the rabbits at the stage of the reproductive cycle were caged with male rabbit for mating and gestational 10 minutes. When rabbit is pregnant that it may start rejecting advances made the male rabbit. The number of litters and the gender of pups were recorded. Pups were sexed by means of the ano-genital distance, which is longer in males (Tarin et al., 1999), this was confirmed in later examinations during pre weaning development. The data were entered and analyses by SPSS software using t.test and the p-value less than 0.05 were considered as significant.

Results

In the first group diabetic mothers (Na, K), all of the 7 rabbits became pregnant which delivered 40 offspring. Their gender was 31 male 77.5% and 9 female 22.5%. In the second group, non diabetic (Na, K), all of the 7 rabbits became pregnant and delivered 60 offspring, their gender was 48 male 80% and 12 female 20% and in the third group, neither diabetic nor (Na, K) all 7 rabbits became pregnant and delivered 76 offspring that 38 male 50% and 38 female 50% (Table-2).

The sex ratio of female to male in the first group of diabetic mothers (Na, K) and in the second group, non diabetic (Na, K) were 3.4:1-4:1, While this ratio in the third group, neither diabetic nor (Na, K) was 1:1 respectively (Fig- 1,2). The percentage of the male offspring of diabetic mothers (Na, K) 77.5% was higher than the male offspring in control group 50% and also male offspring of non diabetic mothers (Na, K) 80% was higher than the male offspring in control group 50% (Fig-3).

The difference in the sex ratio between the first group diabetic mothers (Na, K) and the second group non diabetic mothers (Na, K) was not statistically significant, while the difference between the group of diabetic mothers (Na, K) with

control group and between group non diabetic mothers (Na, K) with control group were statistically significant (Table-1). The Total no of offspring in the first group

diabetic mothers (Na, K) 40 was lower than total no of offspring the second group, non diabetic (Na, K) 60, and also in the third group, neither diabetic nor Na, K 76 (Fig-4).

Table 1. Estimated Minerals Requirements

Mouse Minerals ** (g/Kg)	Amount diet (Kg)	Human *(mg-ug/day)
Calcium	5.0	1000
Chloride	0.5	750
Magnesium	0.5	2-5
Phosphorus	3.0	700
Sodium	0.5	500
Potassium	2.0	2000
Iron	35.0	8
Manganese	10.0	2-5
Zinc	150.0	10-12
Iodine	150.0	150-150
Molybdenum	150.0	75-250(ug)

**adapted from Nutrient Requirements of Nonhuman Primates.

* Adapted from Lanus Micronutrient information Center, Oregon State Unit.

Table 2. Sex ratio in different groups of rabbits.

Group	Total no of offspring	No. of male offspring	% age of male offspring	No. of female offspring	% age female offspring	Sex ratio
Diabetic (Na, K)	40	31	77.5	9	22.5	3.4
Non diabetic (Na, K)	60	48	80	12	20	4
Neither diabetic nor (Na, K)	76	38	50	38	50	1

Figure 1: Sex ratio of Male and female in different groups of rabbits




Figure 2: Compare of Male and female in different groups of rabbits




Figure 3: Percentage of Offspring sex in different groups of rabbits





Figure 4: Number of offspring in different groups of rabbits



Discussion

Streptozotocin has been widely used to induce type 1 diabetes in animal models especially rats and mice. It has been reported that STZ induce dose-dependent diabetes administered either intravenously or intraperitoneally (Tarin et al., 1999). Intraperitoneal injection of STZ led to physiologic

alterations consistent with reports of spontaneous and chemically induced diabetes in other animals (Hayashi, 2006, Holemans, 1997, Kaneko, 1964). Preselection of the gender of offspring is a subject that has held man's attention since the beginning of recorded history. While the natural sex ratio

at birth is usually 104–107 male to 100 female (Visaria, 1901; Bose, 1951; Allied, 1967).

The first medical technique that can be used to select for sex is that of pre-natal diagnosis [PND] and abortion. To utilize this method, the pregnant woman must undergo some sort of prenatal testing, such as an amniocentesis, chorionic villus sampling or an ultrasound, which will allow the doctor to determine the sex of the child, among other things. Once the woman has the information about the child's sex, she can obtain an abortion if the fetus is not of the desired sex. The use of PND and abortion in order to select for sex sounds extreme, and indeed, as Edgar Dahl points out, it is not common for Westerners to utilize such a technique. For example, a follow-up study of 578 patients having prenatal diagnosis at one Melbourne centre found that none of the women had a termination because of the sex of the fetus. Going through the traumatizing experience of an abortion is usually seen as too high a price to pay for a child of a particular sex (Visaria, 1967).

The second medical technique that can be used to select for sex is that of pre-implantation genetic diagnosis [PGD] with *in vitro* fertilization [IVF]. With this technique, the embryos are screened for sex prior to being implanted into the woman's uterus, thereby eliminating the need to later decide to terminate a pregnancy. PGD and IVF, however, are very invasive and potentially physically harmful, requiring the woman to go through at least one IVF cycle, which includes taking potent drugs to induce super-ovulation, extraction, fertilization and then testing and subsequent implantation of the embryos. Given the expense of IVF treatment cycles (according to IVF Canada in 2005 it cost \$5,500 for one cycle of IVF, not including drugs (Dahl, 2003) and, according to the same source, the drugs themselves can cost approximately \$3,000 for one cycle), it would be highly unlikely that it would be used as a

technique for sex selection alone. More likely, it could be used as a sex selection technique for those who are already undergoing IVF for other medical reasons.

The last medical technique that can be used to select for sex is sperm sorting. New technologies allow sperm to be sorted into those carrying X or Y chromosomes with varying degrees of accuracy. To date, the most successful way in which to sort sperm is flow cytometry, which has been branded as the MicroSort technique (Steinbock, 2003). Sex selection using flow cytometry results from distinguishing between the identifiable differences between the X and Y chromosomes, as the X chromosome is larger than the Y. The sorted sperm is then used to artificially inseminate the woman. Studies have shown that the Micro Sort technique is more effective in selecting for girls, a success rate of 91%, than for boys, with a success rate of only 76%.¹² Sperm sorting appears, then, to be the least invasive and least expensive (at about \$2,300US per cycle¹³) method of selecting for sex.

There are also methods which use different food combinations and especial diets to maximize the chance of having a baby with specific sex. The old belief is that eating salty, savory foods leads to delivering a male and sodium rich foods to a female. Some believe that the ratios of the minerals sodium, potassium are important in determination of baby gender. It was shown that pregnant female house mice maintained on a consistent low-food diet give birth to a lower proportion of males than do control females fed *ad libitum* (Meikle, 1995).

Today one of the good known methods on sex constitution is the preconception diet method. This method claims 80% accuracy and the theory is that by altering your diet to include and exclude certain food, the condition in the reproductive tract will be directly affected; increasing the odds of conceiving a particular sex it is also

recommended that both mother and father go on the diet. This is also consistent with the oriental philosophy that everything has a yin or yang quality and the foods supplied in the female diet, female and acid are all yin. The female diet is high in sodium but low in salt and potassium, containing acid forming foods. The diets nutritional content is questionable and contains multiple warnings. The diet may influence the conditions within the reproductive tract and the outer barrier surrounding the ovum. Enabling only one of the two types of sperm to penetrate the depending on which diet is adhered to. Langendon and Proctor first published 'the preconception Gender Diet 'based on results reported (Stolkowskowsky, 1982). The theory is that by altering your diet to include and exclude certain foods, the conditions in the reproductive tract will be directly affected, increasing the odds of conceiving a particular sex. This method under scrutiny claims of 80% accuracy based on one clinical trial of only 260 women, the results were published in the international journal of Gynecology and Obstetrics in 1980. The female diet is high in sodium but low in salt and potassium, containing acid forming foods. The diets nutritional content is questionable and contains multiple warnings. It is recommended to seek the advice of medical practitioner before going on such a restrictive diet, and stay on the diet for no longer than 3 months. The diet may influence the condition of the cervical mucus and within the reproductive tract and follicular fluid. Enabling only one of the two types of sperm to penetrate the egg depending on which diet is adhered to. The aim of this study was to elevate relationship between minerals and sex ratio in rabbits.

Acknowledgement

Authors are grateful to the Dean, Veterinary college of Bangalore, Bangalore for providing the facilities.

References:

- Abrevaya J. 2009. Are There Missing Girls in the United States? Evidence from Birth Data. *American Economic Journal: Applied Economics.*, 1(2): 1-34.
- American College of Obstetricians and Gynecologists, Press Release, ACOG Opposes Sex Selection for Family Planning Purposes. 2007. Feb: 1. http://www.acog.org/from_home/publications/press_releases/nr02-01-07-2.cfm.
- Bhaskar, V., Gupta B. 2007. India's Missing Girls: Biology, Customs and Economic Development, *Oxford Review of Economic Policy.*23: 221- 238.
- Bhat, M. 2006. Sex Ratio in India: *Correspondence, Lancet.* 367: 1725-1726.
- Bhat P.N. and Francis A. 2003. Fertility Decline and Gender Bias in Northern India, *Demography*, 40 (4): 637-657.
- Bolet ,G., Gueguen, L., Dando, P., Ollivoer ,L. 1982. Influence of mineral diet of the sow on the sex ratio of the new born. *Reported. Nutr. Dev.* 22:1073-81.
- Chandrabu S, Beirami A , Chidan Kumar C.S. 2011. Role of Sodium and Potassium ions in identification of baby gender in High-sugar mammal's. *International Journal of Pharmacy and Pharmaceutical Sciences.* 3(4): 303-306.
- Chandrabu S , Beirami A , Chidan Kumar C.S. 2011. Effect of calcium and magnesium ions in identification of baby in High-sugar mammals. *Research in Biotechnology.* 2(3):23-31.
- Chandrabu S, Beirami A, Chidan Kumar C.S. 2012. Effect of Sodium and Potassium ions in identification of baby in High-sugar hamsters. *Research in Journal of Pharmaceutical and Clinical Research.* 5(1):134-136.
- Chandrabu S, Beirami A, Chidan Kumar C.S. 2011. Impact of calcium and magnesium ions in identification of baby gender in High-sugar hamsters. *Research Journal of*

- Pharmaceutical Sciences and Research*. 3(12):1619-1624.
- Chandrabu S, Beirami A, Chidan Kumar C.S. 2012. Role of sodium and potassium ions in identification of offspring gender in rats. *Research in IOSR Journal of Pharmacy*. 2(6):54-59.
- Chandrabu S, Beirami A, C.S. Chidan Kumar.2012. Impact of calcium and magnesium ions in identification of offspring gender in high-Sugar rabbits. *Research in Journal of Chemical and Pharmaceutical Research*. 4(1):719-725
- Clark, S.2000. Son Preference and Sex Composition of Children: Evidence from India, *Demography*. 37(1): 95-108.
- Ebenstein, Avraham .2009. Estimating a Dynamic Model of Sex Selection in China, *Demography*, March, Forthcoming.
- Engelgau, MM.,Herman ,WH., Smith, PJ., German, RR., Aubert, RE.1988. The epidemiology of diabetes and pregnancy in the U.S .*Diabetes Care*. 18:1029-33.
- Feig, DS., Palda ,VA.2002. Type 2 diabetes in pregnancy: A growing concern. *Lancet*.359:1690-2.
- Gupta D, Bhat Mari,.1997. Fertility Decline and Increased Manifestation of Sex Bias in India, *Population Studies*. 51(3): 307-315.
- Jha Prabhat, Kumar R,Vasa P. Dhingra N,Thiruchelvam D, Moineddin R.2006. Low male-to-female [sic] sex ratio of children born in India: national survey of 1.1 million households, *Lancet*. 367: 211-218.
- Judith D.2005. ART and the Search for Perfectionism: On Selecting Gender, Genes, Gametes, 9 J. *GENDER RACE & JUST*. 241:265-66.
- Karunanayake EH, Hearse D, Mellows G. 1975. The metabolic fate and elimination of streptozocine. *Biochemical Society Transactions*.3: 410-14.
- Meikle ,DB.,Thornton, MW.1995. Premating and gestational effects of maternal nutrition on secondary sex ratio in house mice. *J. Reprod. Ferti.*, 105:193-196.
- Mittwoch ,U.2005. Sex in mythology and history. *Arq Bras Endocrinol Metab* .49:7-13.
- Ornoy A, Ratzon N. Greenbaum C,Peretz E,Soriano D, Dulitzky M .1998. Neurobehaviour of school age children born to diabetic mothers. Archives of Disease in Childhood. *Fetal and Neonatal Editio*.79: 94-99.
- ParkBin ,C.,Cho , N. 1995. Consequences of Son Preference in a Low-Fertility Society: Imbalance of the Sex Ratio at Birth in Korea, *Population and Development Review*.
- Retherford R , Roy D .2003. *Factors Affecting Sex-Selective Abortion in India and 17 Major States*, National Family Health Survey Subject Reports 21, International Institute for Population Sciences, Mumbai, India, January.
- Solomon CG, Willett WC, Carey VJ. 1997. A prospective study of pregravid determinants of gestational diabetes mellitus .*JAMA*.278:1078-83.
- Stenninger E, Flink R, Eriksson B,Sahlen C.1998. Long-term neurological dysfunction and neonatal hypoglycaemia after diabetic pregnancy. Archives of Disease in Childhood. *Fetal-Neonatal Edition*.79 (F):174-179.
- Stolkowski J. Lorrain J. 1982. Preconceptional selection of Fetal Sex. *int. J. Gynaecol. Obstet*. 18:440-3.
- Tarin JJ, Perez-Albala S,Aguilar A, Minarro J, Hermenegildo C. Cano A .1999. Long-Term Effects of Postovulatory Aging of Mouse Oocytes on Offspring: A Two-Generational Study. *Biol. Reprod*. 61: 1347-1355.
- Van Assche F, Holemans K,Aerts L.2001. Long-term consequences for offspring of diabetes during Pregnancy. *British Medical Bulletin*.60:173-182.
- Visaria P .1967. The sex ratio of the population of India and Pakistan and regional variations during 1901-61. In:

- Bose A, editor. Patterns of population change in India 1951-61. *Allied Publishers*. P: 334-71.
- Weintrob N, Karp M, Hod M. 1996. Short- and long-range complications in offspring of diabetic mothers. *Journal of Diabetes and Its Complications*. 10: 294-301.
- Xiang AH, Peters RK, Trigo E, Kjos SL, Lee WP, Buchanan TA. 1999. Multiple metabolic defects during late pregnancy in women at high risk for type 2 diabetes. *Diabetes*. 48:848-54.