

Diversity and dominance of flora around wetlands in Lakkavalli Range of Bhadra Wildlife Sanctuary, Karnataka

Vijaya Kumara*, Raghavendra Gowda, H.T., and Pramod, A.F.,

Department of Wildlife and Management, Bioscience Complex, Kuvempu University, Jnana Sahyadri, Shankaraghatta – 577 451, Karnataka, India

Abstract

The present communication is based upon a research work and the building a database for plants in the wetlands of Lakkavalli range of Bhadra Wildlife sanctuary. From these wetlands, about 133 species of plants have been identified of which herbs (27), shrubs (22) climbers (19) and trees (65). The objective of the work was to help wetland managers and ecologists by providing an account of the floral characteristics around wetland community.

Keywords: Bhadra Wildlife sanctuary, diversity, Lakkavalli range and Wetlands

INTRODUCTION

Forest constitutes a dynamic biological system with many organisms living together in perfect harmony with nature (Ellenberg, 1978). Biological diversity is a key issue of nature conservation and species diversity is one of important components of the biological diversity (Ito, 1997). Forest lands extend over a great number of ecosystems, harbouring a rich diversity of species and genes. Rio earth Summit held in 1992 emphasized the need to conserve the biodiversity of the globe in general and tropics in particular. The work of conservation must envisage wetlands since they cover about six percent of earth surface and are well known for high diversity in class, composition and four broad categories of functions viz., physical-hydrological, chemical, biological and socio-economic. Biodiversity assessment is often restricted on the red listing of threatened species and clarification of their habitat demands in forest practices and forest management plans. However, good data and appropriate indicators are necessary to assist policy making and monitoring to understand the causes of changes in biodiversity and to better implement protection strategies (Puumalainen *et al.*, 2003). Wetlands are well-known to nurture a good number of angiosperms providing conditions intermediate between true-aquatic and terrestrial ones.

India has over 1.08.276 species of bacteria, fungi, plants and animals already identified and described (Khoshoo, 1994). Out of these, over 84% species constitute fungi (21.2%), flowering plants (13.9) and insects (49.3%) etc.

Based on the data of the species already described, India is tenth among the plant-rich countries of the world, fourth among the Asian countries (Khoshoo, 1994). Moreover, India is also one among the 12-mega biodiversity centers of the world by having over 47.000 plant species. Its diversity is unmatched due to the presence of 16

different agro-climatic zones, 10 vegetative zones and 15 biotic provinces.

Flora lying at the base of food chain drive to energy flow in an aquatic system Their composition in the community has an enormous implication on local biodiversity. They strongly influence the water chemistry acting as both nutritional sinks through uptake and as nutrient pumps. They also have the property to improve the water quality by capturing heavy metals and ions.

Tropical forests are famous for being the most species rich ecosystems on the earth. These forests, which over only 7 per cent of the earth's land surface harbors more than half of the world species, are under constant threat of human interference resulting in the disappearance of plant species at an estimated rate of 0.8 to 2 per cent per year all over the world. This situation has raised global concern over the different constraints causing this damage to the tropical diversity. Human disturbance in terms of habitat destruction, introduction of exotic species, exploitation and pollution are considered as the major causes for the decline in species level both in India and worldwide (UNEP, 2001).

The most striking feature of the earth is the existence of life and the most striking feature of life is its diversity, topography, soil, climate and geographical location of a region influence the vegetation diversity of forest ecosystem.

Numerical quantification of biological diversity and / or its elements can be of great value because that kind of evaluation is objective and enables a comparison of current biodiversity status to be made between similar ecosystems. During the last century, a great number of different methods quantifying species diversity were developed (Ludwig and Reynolds, 1988 ; Patil and Taillie, 1982 ; Merganic and Smelko, 2004).

However disturbance sometime is the major factor in structuring communities by determining the forest dynamics, diversity and species richness (Shell, 1999). But the detailed description and discussion on the plant diversity along disturbance gradient from dry tropical forest is lacking (Sagar *et al.*, 2003). In this view, a study has been carried out to know the influence of disturbance on the structure, composition, dominance, species richness and diversity of plants in wetlands of Lakkavalli range of Bhadra Wildlife Sanctuary (BWLS), Karnataka to reveal the diversity and the similarity existing

Received: June 10, 2011; Revised August 21, 2011; Accepted August 21, 2011.

*Corresponding Author,

Vijaya Kumara
 Department of Wildlife and Management, Bioscience Complex, Kuvempu University, Jnana Sahyadri, Shankaraghatta – 577 451, Karnataka, India

Tel: +91-9448206428
 Email: vijay15675@gmail.com

among the plant species.

The objective of the present study is to describe the diversity, dominance, structure and composition of five wetlands in Lakkavalli range of Bhadra Wildlife Sanctuary (BWLS), Karnataka along a disturbance gradient and also to reveal the diversity and similarity existing among the plant species.

MATERIALS AND METHODS

Study Area

The study was carried out in Lakkavalli range of Bhadra Wildlife Sanctuary, consisting an area of 223.17 sq. kms (13°34' to 13°46'N latitude and 75°29' to 75°45'E longitude) in the Karnataka state of Southern India. The altitude varies from 650 mts-1875 mts above sea level with a general elevation of 1200-1500 mts. The sanctuary is located in the Malnad region of Karnataka about 50 kms to the east of western ghats. The temperature in the valley ranges from 9-35°C. The region receives an annual rainfall of 1600 to 2000 mm during the southwest monsoon between June and September. A distinct rainfall gradient result in a variation in vegetation type from semi evergreen forest and moist deciduous forest through dry deciduous forest shoals and grassland type forest was selected for this study during 2007-08.

Koramaguddakere lies in an altitude of 2302 feet (13°37'173" N and 75°39'95" S). Pickupkere lies in an altitude of 2413 feet (13°36'571" N and 75°39'307" S). Ramannanakere lies in (13°36'37" N and 75°37'56" E). Anegundikere lies in (13°34'846" N and 75°36'416" E). Mavinahalladakere (backwater) lies in (13°34'846" N and 75°36'416"E).

The study was conducted in five wetlands Lakkavalli range of Bhadra Wildlife sanctuary. Each forest type is named after the dominant tree species. The size and number of the samples were determined. Species composition for trees, shrubs and herbs was prepared after a thorough survey of different wetlands. Within each wetlands all the vegetation layers, i.e. trees, shrubs and herbs were analyzed for species richness and diversity. Herbs were analyzed during peak growing season, plants and flora were identified with the help of plant taxonomists.

Methodology

The best method to estimate the plant diversity in this type of habitat is Belt transects. Transects of the size 1000m X 5m, was laid in the adjacent forest cover of each wetlands. All plant species including trees, herbs and climbers which are present in the transect were identified to the species level and their number was counted

In all the five wetlands plant species present around were collected, identified and systematic enumeration was made with the available monographs relevant literatures and taxonomic revisions (Cambel, 1918; Mathew. 1983; Rath. 1999 and Dharamendra Singh, 1999).

Species similarity among the flora was computed using Sorenson's index (Wilson M.V. et al., 1984).

$$I = 2J/A+B.$$

Where I=similarity, J=Common species of both the series a and b. A=Total number of species in series a and B= Total number of species in series b.

RESULTS

A total of 133 species belongs to 56 families were identified during the study. Among the five study area Anegundikere wetland is more speciose (N=54) followed by Mavinahallakere and Ramananakere. Koramaguddakere and Pickupkere are less species compared to Anegundikere, however number of individuals is in case of Mavinahallakere (Back water) followed by Anegundikere and Ramannanakere (Table1). Among the pooled data a weed *Chromolina oderata* is most dominant followed by *Tectona grandis*, *Anogessus latifolia*, etc (Table2)

Shannon-Wiener diversity index have not shown significant difference among different samples, while Simpson's index showed that Koramaguddakere and Pickupkere are having less plant diversity compared to other three (Table3).

We computed Sorenson's index to compare the vegetation around the wetlands. It is resulted that Koramaguddakere and Pickupkere are significantly associated (>40%) with Ramannanakere. While all other associations are less than 40% (Table4).

The species with family of all the wetlands are been given in table 5.

Table 1: Mean species richness for all the vegetation layers in different wetlands

Sample	Mean Individuals	Variance	Standard Deviation	Standard Error	Total Individuals	Total Species
Mavinahallakere	3.314	34.749	5.895	0.498	464	51
Anegundikere	2.843	26.579	5.156	0.436	398	54
Ramannanakere	2.571	25.959	5.095	0.431	360	51
Koramaguddakere	1.607	15.147	3.892	0.329	225	31
Pickup kere	1.657	13.738	3.706	0.313	232	34




Table 2. Dominant plant species among pooled data

Species	Variance	Mean	Chi-sq
<i>Chromolina oderata</i>	101.4667	12.3333	41.1351
<i>Tectona grandis</i>	97.2	14	34.7143
<i>Anogessus latifolia</i>	84.7	5.5	77
<i>Randia dumetorum</i>	78.3	6.5	60.2308
<i>Cinnamomum malabathrum</i>	73.5	3.5	105
<i>Sida rhomboideae</i>	71.2	5	71.2
<i>Cyclea peltata</i>	70.6667	4.3333	81.5385
<i>Asparagus racemosus</i>	66.1667	5.1667	64.0323
<i>Allophylus cobbe</i>	64.1667	5.1667	62.0968
<i>Dalbergia latifolia</i>	63.6	10	31.8
<i>Mimosa pudica</i>	61.6	9	34.2222
<i>Cleome viscosa</i>	60.9667	4.8333	63.069
<i>Lagestroemia lanceolata</i>	57.0667	8.3333	34.24
<i>Macaranga peltata</i>	49.5	4.5	55
<i>Terminalia tomentosa</i>	49.5	9.5	26.0526
<i>Terminalia bellarica</i>	47.3667	5.8333	40.6
<i>Persea macrantha</i>	46.3	3.5	66.1429
<i>Cryptolepis buchanani</i>	45.7667	5.1667	44.2903
<i>Cassia fistula</i>	42.5667	7.1667	29.6977

Table 3. Diversity components among different wetlands in Lakkavalli range of Bhadra Wildlife sanctuary

Wetlands	Simpsons Diversity (1/D)	Shannon H' Natural Log
Mavinahallakere	36.388	3.669
Anegundikere	35.683	3.693
Ramannanakere	30.963	3.556
Koramaguddakere	22.48	3.177
Pickupkere	25.99	3.312

Table 4. Sorenson's index and correlation of species between different wetlands

Sorenson's index	Mavinahalla kere	Anegundi kere	Ramannana kere	Koramagudda kere	Pickup kere
Mavinahallakere	1				
Anegundikere	0.21	1			
Ramannanakere	0.22	0.36	1		
Koramaguddakere	0.27	0.31	0.44*	1	
Pickup kere	0.31	0.25	0.45*	0.37	1

* significant.

DISCUSSION

From the study it is clear that this forest ecosystem supports good plant species and the varied with flora of this region faces severe constraints on conservation and the sanctuary has a long history of protection hence it retained wonderful diversity plants in it.

The pace of development and the population pressure together with lack of knowledge on biodiversity are the major threats to the ecosystem conservation (Bentley, 1991). Diversity indices showed that Mavinahallakere and Anegundikere forests which are present in the core zone of the sanctuary are diverse in their vegetation than other three, which revealed that vegetation of the core are of the

sanctuary is more diverse than that of buffer area. Though major vegetation type of the study area is deciduous some of the evergreen species like *Persea macarantha* and *Hydnocarpus*

pentandra were appeared in Ramannanakere forest which is present in core area. There is no special pattern observed in the association of species in between the different forest type studied.




Fig. 2. Species richness in different wetlands

Table 5. Diversity of plant species among different wetlands in Lakkavalli range of Bhadra Wildlife Sanctuary.

Plant species	Family	Habit	Mavinahallakere	Ane Gundikere	Ramannanakere	Koramaguddadakere	Pickup kere
<i>Acacia concinna</i>	Mimoseae	C		7			
<i>Acacia instia</i>	Mimoseae	T	4				
<i>Ageratum conyzoides</i>	Asteraceae	H	3			2	
<i>Albizia lebbeck</i>	Mimoseae	T		5			
<i>Albizia procera</i>	Mimoseae	T	1				
<i>Allophylus cobbe</i>	Sapindaceae	S		16	15		
<i>Alocasia sp.</i>	Araceae	H					1
<i>Alseodaphne semicarpifolia</i>	Lauraceae	T			6		
<i>Alstonia scholaris</i>	Apocynaceae	T			1		
<i>Anogessus latifolia</i>	Combretaceae	T	22				
<i>Antidesma ghaesembla</i>	Euphorbiaceae	T		3			
<i>Ardesia solanaceae</i>	Myrsinaceae	S		5			
<i>Argeria nervosa</i>	Convolvulaceae	C					2
<i>Artocarpus lakoocha</i>	Moraceae	T			2		1
<i>Asparagus racemosus</i>	Liliaceae	C			21	5	
<i>Aspidopteris cordata</i>	Malpighiaceae	C	1				

<i>Bauhenia racemosa</i>	Papilionaceae	T	2				
<i>Bauhinia sp.</i>	Caesalpiniaceae	T		2			
<i>Bombax ceiba</i>	Malvaceae	T		2		2	
<i>Braynia retusa</i>	Euphorbiaceae	S			5		
<i>Butea monosperma</i>	Papilionaceae	T		7			
<i>Butea parviflora</i>	Papilionaceae	C		8			
<i>Caesalpinia mimosoides</i>	Caesalpiniaceae	S		12			
<i>Cansjera rheedii</i>	Opiliaceae	S		5			
<i>Carallia brachiata</i>	Rhizophoraceae	T			4		
<i>Careya arborea</i>	Myrtaceae	T	5	11			8
<i>Casaria tomentosa</i>	Flacourtiaceae	S			1		
<i>Cassia fistula</i>	Caesalpinaceae	T	17	7	13	1	2
<i>Cassia mimusoides</i>	Caesalpinaceae	H	7				
<i>Cassia sp.</i>	Caesalpinaceae	S	4				
<i>Cassia tora</i>	Caesalpinaceae	H	8				
<i>Centella asiatica</i>	Umbelliferae	H		2			
<i>Chromolina oderata</i>	Asteraceae	H		22	4	20	22
<i>Cinnamomum malabathrum</i>	Lauraceae	T			21		
<i>Cissus raphanda</i>	Vitaceae	C			1		
<i>Cleome viscosa</i>	Capparaceae	H				18	11
<i>Clerodendron viscosum</i>	Verbinaceae	S		9	1	3	
<i>Crotolaria sp.</i>	Fabaceae	H	3	2			
<i>Cryptolepis buchanani</i>	Asclapiadaceae	C	6		18		5
<i>Curculigo orchioides</i>	Palmae	T	9		2		
<i>Curcuma sp.</i>	Zingiberaceae	H	5				
<i>Cyclea peltata</i>	Menispermaceae	C			21	5	
<i>Cynodon dactylon</i>	Graminae	H	2				
<i>Dalbergia latifolia</i>	Papilionaceae	T	18	15	12		
<i>Dellinia pentagyna</i>	Dilliniaceae	T			15	7	9
<i>Dendrocalamus strictus</i>	Graminae	T	15				3
<i>Desmodium triquetrum</i>	Papilionaceae	H		5	14		6
<i>Dioscorea bulbifera</i>	Dioscoriaceae	C		8	1		
<i>Dioscorea pentaphylla</i>	Dioscoriaceae	C		5			
<i>Diospyros buxifolia</i>	Ebenaceae	T					4
<i>Diospyros montana</i>	Ebenaceae	T	9				
<i>Diploclisia glaucansis</i>	Menispermaceae	C		5			
<i>Drynaria quersifolia</i>		H		2			
<i>Elephantopus scaber</i>	Asteraceae	H	12				
<i>Embelia tsjerium -cottam</i>	Myrsinaceae	C	15				
<i>Emblica officinale</i>	Euphorbiaceae	T	12	7		8	
<i>Ficus callosa</i>	Moraceae	T			9		5
<i>Ficus hispida</i>	Moraceae	T		2			
<i>Ficus racemosa</i>	Moraceae	T		8	4	2	
<i>Ficus tsjehela</i>	Moraceae	T		1			
<i>Flacourtia indica</i>	Flacourtiaceae	T	1				
<i>Flemingia sp.</i>	Fabaceae	H	2				
<i>Glochidion sp.</i>	Euphorbiaceae	T					4
<i>Grewia tilifolia</i>	Teliaceae	T		11	1	8	13
<i>Gynura nitida</i>	Asteraceae	H					12
<i>Helicteres isora</i>	Sterculiaceae	S				5	5
<i>Hellotropium indica</i>	Boraginaceae	H	6			9	
<i>Hippocratea sp.</i>	Hippocrataceae	C		1			
<i>Holarrhina antidysentrica</i>	Apocynaceae	S	12				
<i>Homonioia riparia</i>	Euphorbiaceae	H		14			

<i>Hopea ponga</i>	Dipterocarpaceae	T			10	
<i>Hydnocarpus pentadra</i>	Flacourtiaceae	T		8		2
<i>Ichinocarpus fruitiscence</i>	Apocynaceae	C	3			
<i>Ipomea sp.</i>	Convolvulaceae	C	4			
<i>Ixora coccinea</i>	Rubiaceae	S			11	
<i>Jasminum sp.</i>	Oleaceae	C		3		
<i>Kydia calycina</i>	Malvaceae	T		2	2	2
<i>Lagestroemia flos-regini</i>	Lythraceae	T		7		
<i>Lagestroemia lanceolata</i>	Lythraceae	T	20		1	12
<i>Lannea coramandalica</i>	Anacardiaceae	T			12	
<i>Lantana camara</i>	Verbinaceae	S				8
<i>Leea talbotii</i>	Leeaceae	S	15	1		3
<i>Macaranga peltata</i>	Euphorbiaceae	T			19	12
<i>Mallotus philippinensis</i>	Euphorbiaceae	T	5	3	4	9
<i>Memecylon umbellatum</i>	Melastomaceae	T				7
<i>Meyna laxiflora</i>	Rubiaceae	T	8	12	2	2
<i>Michelia champaca</i>	Magnoliaceae	T			2	4
<i>Mimosa pudica</i>	Mimoseae	H	20			13
<i>Mitragyna parviflora</i>	Rubiaceae	T	4		14	
<i>Murraya koenigii</i>	Rutaceae	T			2	
<i>Musenda frondosa</i>	Rubiaceae	C			2	
<i>Narvelia zeylanica</i>	Ranunculaceae	C		1		
<i>Nothopegia racemosa</i>	Anacardiaceae	T		2		
<i>Ochlandra rheedi</i>	Graminae	S	2			
<i>Olea dioica</i>	Oleaceae	T			1	
<i>Pavetta tomentosa</i>	Rubiaceae	S		1		
<i>Persea macrantha</i>	Lauraceae	T			17	4
<i>Polygonum glabrum</i>	Polygonaceae	H		14		
<i>Polygonum sp.</i>	Polygonaceae	H				4
<i>Pongamia pinnata</i>	Papilionaceae	T		7		
<i>Pterocarpus marsupium</i>	Fabaceae	T		6		
<i>Randia dumetorum</i>	Rubiaceae	T	14	21	2	
<i>Rauvolfia serpentina</i>	Apocynaceae	H			1	2
<i>Rubus moluccanus</i>	Rosaceae	S	5			
<i>Santalum album</i>	Santalaceae	T				5
<i>Sapindus laurifolia</i>	Sapindaceae	T		1		
<i>Schlichera oleosa</i>	Araliaceae	T	8	7		5
<i>Securinuga leucopyrus</i>	Euphorbiaceae	S	2			
<i>Sida acuta</i>	Malvaceae	H	5			
<i>Sida rhomboideae</i>	Malvaceae	H	21			8
<i>Smilax zeylanica</i>	Smalacaeae	C				1
<i>Solanum sp</i>	Solanaceae	S	2			
<i>Stachytarpetta indica</i>	Verbinaceae	H	13			
<i>Sterculia guttata</i>	Sterculiaceae	T		6	8	
<i>Streptospermum personatum</i>	Bignoniaceae	T		4	7	7
<i>Syzygium cumuni</i>	Myrtaceae	T			7	8
<i>Taebermontana heyniiana</i>	Apocynaceae	T			5	
<i>Tamarindus indica</i>	Fabaceae	T	2			
<i>Tectona grandis</i>	Verbinaceae	T	21	28		16
<i>Terminalia bellarica</i>	Combretaceae	T		8	16	11
<i>Terminalia paniculata</i>	Combretaceae	T	21	18	15	13
<i>Terminalia tomentosa</i>	Combretaceae	T	18		2	12
<i>Thespesia lampas</i>	Malvaceae	T		2		
<i>Trema orientalis</i>	Urticaceae	T			5	2

<i>Trewia nudiflora</i>	Euphorbiaceae	T	5		
<i>Triumfetta rhomboidea</i>	Malvaceae	H	9		
<i>Urena lobata</i>	Malvaceae	S	9	4	
<i>Ventilago maderaspatensis</i>	Rhamnaceae	S	13		
<i>Volvulopsis nummularia</i>	Convolvulaceae	H	14		
<i>Xanthium stromarium</i>	Asteraceae	H	11		
<i>Zanthoxylum rhetsa</i>	Rutaceae	T		3	2
<i>Zizyphus oenoplia</i>	Rhamnaceae	S	12		2
<i>Zizyphus rugosa</i>	Rhamnaceae	S	15	4	

C = Climber, T = Tree, H= Herb, S = Shrub.

REFERENCES

- Bentley RW (1991) Deforestation: past present and future, TRI News 10(1) Spring.
- Camble TS (1918) Flora of Presidency of Madras London, Part 1 II III.
- Cannon CH, Peart DR and M Leighton (1998) Tree species diversity in commercially logged bornean rainforest, Science 282 : 1366-1368.
- Dharmendra Singh (1999) Proc. 86th Ses India Sci Cong Chennai, 33P.
- Ellenberg RW (1978) Vegetation Mitteleuropas mit den Alpen in Okologischer Sicht Ulmer, Stulgrat.
- Ito Y (1997) Diversity of forest tree species in Yanbaru, The northern part of Okinawa Island, Plant Ecology, 133 : 125-133.
- Khoshoo TN (1994) Curr. Sci. 67 (8) : 577- 582.
- Ludwig JA and JF Reynolds (2004) Statistical Ecology: A Primer on Methods and Computing, Wiley and Sons, New York.
- Mathew KM (1983) The flora of the Tamil Nadu, Carnatic Vol. I. II. III. Proc.
- Merganic J and S Smelko (2007) Quantification of tree species diversity in forest stands model Biodiverss, Eur. J. For. Res, 123 : 157-165.
- Merggalef R (1958) Information theory in ecology, General Systematic, 3 : 36-71.
- Pandeya SC, A Chandra and PS Pathak (2007) Genetic diversity in some perennial plant species within short distances, J. Environ. Biol, 28 : 83-86.
- Patil GP and C Taillie (1982) Diversity as a concept and its measurement, J. Am. Stat. Assoc. 77: 548-567.
- Puumalainen J, P Kennedy and S Folving (2003) Monitoring forest biodiversity : A European perspective with reference to temperate and Boreal forest zone, J. Environ. Manage, 67 : 5-14.
- Rath GC (1999) Proc 86th Ses, India Sci, Cong. Chennai, 28 PP.
- Sagar R, Raghubanshi AS and Singh JS (2003) Tree species composition, dispersion and diversity along a disturbance gradient in a tropical forest region of India, For. Eco. Manage, 186: 61-71.
- Shannon CE and Weaver W (1963) The Mathematical theory of communication, University of Illinois Press, Urbana.
- Sheil D (1999) Tropical forest diversity, environment change and species augmentation; after intermediate disturbance hypothesis, J. Veg. Sci, 10: 851-860.
- Simpson EH (1949) Measurement of diversity, Nature, 163 : 688.
- UNEP (2001) India State of the Environment-2001, United Nations Environment Programme, Nairobi.
- Whittaker RH (1972) Evolution and measurement of species diversity, Taxon, 21 : 213-251.
- Wilson MV and Shmida A (1984) Measuring beta diversity with presence absence data, J. Ecol, 72. 1055-64.
- Tsakaya H, Iokawa Y, Kondo M, Ohba H (2005) Large-scale general collection of wild-plant DNA in Mustang Nepal, J Plant Res, 118 : 57-60.