

# Toxic effects of chromium on the aquatic cyanobacterium *Oscillatoria* sp and removal of chromium by biosorption

S. Jayashree<sup>1</sup>, N. Thangaraju<sup>1</sup> and J. Joel Gnanadoss\*<sup>2</sup>

<sup>1</sup>CAS in Botany, University of Madras, Chennai – 600 025, India

<sup>2</sup>Dept., of Plant Biology and Biotechnology, Loyola College (Autonomous), Chennai – 600 034, India

## Abstract

“Chromium” is a non-essential metal for microorganisms and plants and a serious pollutant in diverse environment conditions. The hexavalent form of the metal Cr(VI) is considered to be more toxic than relatively harmless Cr(III) form. Certain microorganisms in the environment are able to tolerate high levels of Cr due to their resistance mechanism. In the present work we report the toxic effect of chromium on *Oscillatoria* sp and the removal of hexavalent using the cyanobacterium *Oscillatoria* sp., by biosorption method. *Oscillatoria* sp., was grown in BG-11 medium containing different concentration of Cr. Effect of Cr on cellular metabolism was studied by estimating the amount of chlorophyll a, carotenoids, c. phycocyanin, allophycocyanin, sugars, free amino acids and proteins at various metal concentration. An increase in metal concentration caused a decrease in the growth of the *Oscillatoria* sp., and also decreased the cellular contents like chlorophyll a, carotenoids, c.phycocyanin, allophycocyanin, sugars, free amino acids and proteins. Biosorption of chromium by *Oscillatoria* sp., was carried out using living cells, heat killed cells and pre-treated cells. Results showed that high amount of metal were adsorbed by heat killed and living cells of *Oscillatoria* sp. The experimental conditions used in the present work are simple and have low operational cost. The proposed biosorption method is economically feasible and eco-friendly in nature.

**Keywords:** Cyanobacteria, *Oscillatoria*, Biosorption, Chromium, Metal toxicity

## INTRODUCTION

Environmental pollution and contamination have become a key focus of concern during the recent times. Environmentally friendly processes need to be developed to clean up the environment without creating harmful waste products. Cyanobacteria are photosynthetic microorganisms and one of the largest and most important groups of prokaryotes on Earth (Saiqa Razi and Shahida Hasnain, 2006). Water contamination by heavy metals is often irreversible and in general it is tacit that the exposure to heavy metals leads to the establishment of a tolerant microbial population. Moreover, hexavalent chromium is considered to be the most hazardous form of chromium due to its solubility, mobility, and toxicity, as well as its carcinogenic and mutagenic properties. Bioavailability of Cr(VI) is largely a function of its ability to cross biological membranes, its powerful oxidizing capabilities and its interference with electron transport in respiration and photosynthesis. Heavy metals are widespread in the environment as they are non-degradable and thus persistent (Jasviri *et al.*, 2002). Microbial biomass has been used for removal of heavy metals. The use of microbial biomass is cost effective for industrial waste water treatment (Akthar *et al.*, 1995). The advantage of microbial biomass is that it can passively bind large amount of metals. This process is

called as biosorption, which is possible by both living and non-living biomass. In biosorption process, the biomass is more effectively used and exploited rather than in bioaccumulation (Rani Gupta *et al.*, 2000). Biosorption which is regarded as an emergent technology is now considered an ideal alternative method for decontamination of metal containing effluents. The advantage of biosorption technique is its low operating cost and that it will minimize the volume of chemical and biological sludge to be disposed.

Recent research indicates that many microorganisms can accumulate large concentration of metals (Ramteke *et al.*, 2000). Hence, the use of microbial cells for remediation of heavy metals offers a potential alternative to existing methods of decontamination of heavy metals. Algae, fungi and bacteria used for accumulation of heavy metals have been studied and its application in the biological processes in removal of heavy metals from wastewater are more effective (Kuyucak and Volesky 1989). The biomass is capable of absorbing and adsorbing metal ions from aqueous solution even when the cells have been killed (Garnham *et al.*, 1992).

Cyanobacteria are also useful in removing harmful metals from the environment. *Spirulina platensis*, a cyanobacterium has been shown to contain detectable levels of mercury and lead when grown under contaminated conditions (Slotton *et al.*, 1989), implying that this cyanobacterium can take up the toxic metals from the target environment. Studies confirmed that this cyanobacterium both adsorbs and takes up metal ions (Bender *et al.*, 1994). *Oscillatoria* sp., (filamentous) and *Synechocystis* sp., (unicellular), were also able to tolerate up to 200µg K<sub>2</sub>CrO<sub>4</sub> mL<sup>-1</sup> on BG11 medium under laboratory conditions (Hasnain *et al.*, 2005). A study also indicated that carboxyl groups on cyanobacterial biomass are responsible for binding with various ions (Gardea-Torresdey *et al.*, 1990). The biosorption of Hg<sup>2+</sup> under batch stirred reaction system by two cyanobacterial strains *Spirulina platensis* and *Aphanothece*

Received: June 12, 2012; Revised: July 24, 2012; Accepted: Aug 30, 2012.

\*Corresponding Author

J. Joel Gnanadoss

Dept., of Plant Biology and Biotechnology, Loyola College (Autonomous), Chennai – 600 034, India

Tel: +91-4428178200

Email: joelgna@gmail.com

*flocculosa*, showed that 98% of mercury was removed from the solution (Amber *et al.*, 2008). Cyanobacterial strains play a key role in the removal of chromium toxicity from leather industries and *Spirulina fusiformis* was capable of removing 93-98 % of the metal toxicity (Pandi *et al.*, 2009). The biosorption characteristics of arsenite and monomethyl arsonate from aqueous solution by *Acidithiobacillus ferrooxidans* BY-3 were studied and the results showed that both metals involved pseudo second order kinetics (Lei yan *et al.*, 2010). Recently chemically modified biomass (treating 0.1m/L HCL) of cyanobacteria *Microcystis* was used for the removal of anionic  $Sb(OH)_6^-$  by biosorption method (Fuhong sun *et al.*, 2011). The biological role of cyanobacterial strains in the removal of toxicity by heavy metal is rather scanty to completely understand the mechanism involved.

In the present study the biomass of the cyanobacterial strain *Oscillatoria sp.*, was used to remove the heavy metal chromium toxicity at various concentrations. The biochemical results showed that increase in metal concentration cause the decrease in the growth rate of *Oscillatoria sp.*, and also decreased the cellular contents.

## MATERIALS AND METHODS

### Organism and growth conditions

*Oscillatoria sp.*, a filamentous cyanobacterium, was obtained from the Culture Collection Centre, Centre for Advanced study in Botany, University of Madras, Guindy campus, Chennai. Cultures were grown at  $24 \pm 1^\circ\text{C}$  in a thermostatically controlled room illuminated with cool white fluorescent tubes (Philips 40 W) providing an irradiance of  $50 \mu\text{E}/\text{m}^2/\text{s}$  in a 12h: 12h light/dark regime.

### Heavy metal stock solutions

3.7g of  $K_2CrO_4$  was dissolved in 100 mL of glass distilled water. This stock solution was used as a heavy metal stock solution. From this stock different concentration of chromium (1, 2, 3, 4, 5,  $10 \mu\text{g}/\text{mL}$ ) were taken for the further analysis.

### Effect of chromium on growth of *Oscillatoria sp.*,

The cultures were maintained and subcultured using BG<sub>11</sub> medium (Rippka *et al.*, 1979) with different concentration of chromium. *Oscillatoria sp.*, was cultured and shaken with glass beads to get a uniform cell suspension and growth was measured by in terms of fresh weight.

### Estimation of pigments

*Oscillatoria sp.*, was grown in different concentrations of heavy metal solution (chromium). Medium without chromium served as control. At the end of growth period estimation of pigments like chlorophyll *a*, carotenoid was estimated using the method of Mackinney, 1941. Absorbance was measured using 10 mm width cuvette at 663 nm and 480 nm. Similarly c.phycocyanin (PC) and allophycocyanin (APC) were also estimated by Bennet and Bogoras, 1973.

### Estimation of sugars and free amino acids

For the estimation of sugars and free amino acids 100mg of

the sample was taken. The estimation of sugars and amino acids both are down by the modified method of Sadasivam and Manickam, 1996. The solution was cooled rapidly and the absorbance was read at 630 nm and 570 nm.

### Estimation of protein

Proteins were estimated following the method of Lowry *et al.*, 1951. *Oscillatoria sp.*, cells were centrifuged at 10000 rpm/min<sup>-10</sup>. The pellets were washed twice with Tris-HCL buffer (pH 7.5) and re-suspended in 5 mL of the buffer. Cells were disrupted in a motor and pestle for 10 minutes, then centrifuged and protein in the filtrate was precipitated with 10% Trichloroacetic acid. The precipitate was removed by centrifugation at 10000 rpm/min<sup>-10</sup> and dissolved in 1 mL of 0.1 N NaOH. After 30 minutes, 5mL of solution alkaline copper reagent was added to 1mL of the sample followed by 0.5 mL of Folin phenol reagent and optical density (O.D) was read at 650 nm in a spectrophotometer after 10 minutes. Amount of protein in the sample was calculated with a standard prepared using Bovine Serum Albumin (BSA).

### Biosorption of Chromium by *Oscillatoria sp.*,

Biosorption of chromium by *Oscillatoria sp* was done by three methods first one is living cells that means fresh biomass, uptake of heavy metal (chromium) in the living cells of *Oscillatoria sp.*, was monitored by amending the basal media with different concentrations (5 and  $10 \mu\text{g}/\text{mL}$ ) of the metals and the algal culture was incubated at an irradiance of  $50 \mu\text{E}/\text{m}^2/\text{s}$  and  $24 \pm 1^\circ\text{C}$  in a 12h: 12h light/dark regime. Heat killed cells were prepared by taking fresh biomass and incubated in hot water and maintained at  $100^\circ\text{C}$  for 5 minutes. For pre-treated cells, the fresh biomass was suspended in 0.1 N NaOH and incubated for 20 minutes. The biomass was later suspended in solutions containing different concentrations of chromium. The biomass was left for 1hr in the respective solutions. After an hour cells were harvested and centrifuged at 10000 rpm/min<sup>-10</sup>. The clear solution were collected and analyzed for metal using Atomic Absorption Spectrophotometer by the method described by Singh *et al.*, 1989.

## RESULTS AND DISCUSSION

### Effect of chromium on growth of *Oscillatoria sp.*,

It is well known that algal cells exposed to heavy metals may suffer serious morphological and biochemical alterations (Rocchetta *et al.*, 2006). The result of this investigation show that the growth of *Oscillatoria sp.*, decreased as the concentration of metal increased (1, 2, 3, 4, 5,  $10 \mu\text{g}/\text{mL}$ ). The metal concentration was low ( $1 \mu\text{g}/\text{mL}$  and  $2 \mu\text{g}/\text{mL}$ ) the growth of *Oscillatoria sp.*, is not affected. As the metal concentration increased ( $10 \mu\text{g}/\text{mL}$ ) the growth was found to be affected (Table 1). Davies *et al.*, (1978) stated that the metal content of organisms depends on the concentration of the ambient water as well as on bioavailability. Banerjee and Mishra, 2002 have also shown the mitigating effect of immobilization on cyanobacteria subjected to heavy metal stress. The growth of species, decreased as the concentration of metal increased. According to Daniel *et al.*, (1979) cyanobacteria of the genus *Oscillatoria* were the most tolerant to different metal concentrations.

Table 1. Effect of chromium on growth of *Oscillatoria sp.*,

Concentration of chromium (µg/mL)	Growth Measurement
C	+++
1	+++
2	+++
3	++
4	++
5	+
10	+

**Effect of Cr on photosynthetic pigments**

The result showed that chlorophyll *a*, carotenoids, phycocyanin and allophycocyanin concentration decreased in *Oscillatoria sp.*, as the metal concentration increased (1, 2, 3, 4, 5, 10 µg/mL). At increasing concentration, chlorophyll *a* gradually decreased from 1.765 to 0.517 mg/L. For carotenoids it decreased from 26.4 to 14.5 mg/L (Fig 1 and 2). Similarly for c.phycocyanin and allophycocyanin the percentage decrease over control was found to be high in 5µg/mL. At increasing concentration of metal both c.phycocyanin and allophycocyanin concentration gradually decreased (Table 1 and 2). The toxic effects of lethal concentration of Cr<sup>6+</sup> on photosynthetic pigments were more pronounced in *A.*

*nidulans* as compared to resistant strain Cr18. Pigment content has been identified as a valuable parameter for defining toxicity of heavy metals [Bolanos *et al.*, 1992, Rai *et al.*, 1981, Rai *et al.*, 1991, Raizada *et al.*, 1990, Sorentinom *et al.*, 1979]. It has been demonstrated that in the presence of heavy metal ions like Zn<sup>2+</sup>, Cu<sup>2+</sup>, Hg<sup>2+</sup>, and Pb<sup>2+</sup>, photosynthetic electron transport chain of cyanobacteria is inhibited [Mohanty *et al.*, 1985, Murthy *et al.*, 1999]. Decrease in photosynthetic pigments of *A. nidulans* in presence of Cr<sup>6+</sup>, thus, this indicates that biosynthesis of these pigments is affected by metal ions. On the other hand, biosynthesis of photosynthetic pigments of the resistant strain Cr18 is not much affected in sublethal concentration of Cr<sup>6+</sup>, leading to normal levels of photosynthetic pigments and growth in presence of Cr<sup>6+</sup> ions.




Fig1.Effect of Chromium on Chlorophyll *a* in *Oscillatoria sp.*,




Fig 2.Effect of chromium on Carotenoids in *Oscillatoria sp.*,  
Table 2. Effect of chromium on C. phycocyanin in *Oscillatoria sp.*,

Concentration of chromium (µg/ml)	C. phycocyanin(mg/mL)	% decrease over control
C	0.040	-
1	0.033	17.5
2	0.032	20
3	0.030	25
4	0.029	27.5
5	0.022	45
10	0.023	42.5

Table 3. Effect of chromium on allophycocyanin in *Oscillatoria sp.*,

Concentration of chromium (µg/mL)	Allophycocyanin (mg/mL)	% decrease over control
C	0.401	-
1	0.037	90.77
2	0.029	92.76
3	0.021	94.76
4	0.020	95.01
5	0.018	95.51
10	0.019	95.26

**Effect of Cr on sugars and free amino acids**

Many algae accumulate Cd, Cu, Pb, Zn and other metals which affect their general physiological and biochemical characteristics (Gupta and Singhal, 1995). The result showed that as the concentration of chromium (1-10µg/mL) increases the maximum amount of sugar and amino acids in 100 mg of cells decreased. The percentage over control was high when concentration of chromium was 10µg/mL. The percentage over control was low in 1µg/mL (Fig.3). The increased total carbohydrate accumulation following metals was accompanied by decrease Fathi *et al.* (2005) reported that the higher doses severely attenuate chlorophyll synthesis coupled with severe drop in protein resulting in increased carbohydrates. Torres *et al.* (1998) demonstrated that algae *Cylindrothica fusiformis* produce carbohydrate as a defense mechanism against copper toxicity in stationary phase when cells are exposed to 0.5 mg Cu L<sup>-1</sup>.

**Estimation of protein**

The result showed that as the concentration of chromium increases the amount of protein in the cells decreased (5.0 – 2.0). The percentage decrease over control was high in (62.96%) when concentration of chromium was 10µg/mL. the percentage decrease over control was low in (3.70%) when the concentration of chromium was 1µg/mL (Fig 3). The accumulation of protein at low heavy metal concentrations may be one of the ways through which the algae can abolish their toxic effects, or to increase respiration leading to the utilization of carbohydrate in favor of protein accumulation (Osman *et al.*, 2004). Whereas, the suppression of protein accumulation may be attributed to shortage of carbon skeleton results from low photosynthetic rate. Such results are in accordance with those of Fathi *et al.* (2000). However, some authors (Osman *et al.*, 2004; Tripathi and Gaur, 2006; Barbara and Michael, 1994) reported that the toxic action of heavy metals on the enzymatic reactions responsible for protein biosynthesis.




Fig 3.Effect of Chromium on Proteins, Amino acids and Sugars in *Oscillatoria sp.*,

### Biosorption of Chromium by *Oscillatoria sp.*,

The results were tabulated based on living cells, Heat killed cells, Pre-treated cells. The high amount of metal was adsorbed (1.90) in heat killed cells of *Oscillatoria sp.*, in 10µg/mL concentration and 5µg/mL the high amount of metal was adsorbed (0.86) in living cells (Table 4). Fisher et al., 1984 stated that heat killed cells accumulated as much metal as the live cells, further indicating that bioaccumulation of metal in the algae proceeded by non-metabolic adsorption also. Similarly Rabsch and Elarachker 1980 heatkilled

cells of *Coscinodiscus granii* accumulated 3 times more Cd and 4 times more Zn than did in live cells. Ahuja et al., 1999 also reported that, in a fresh water cyanobacterium, *Oscillatoria angustissima*, showed a very high capacity for the biosorption of zinc at 641 mg g<sup>-1</sup>. The uptake of the metals was higher in immobilized condition in comparison to free cells, due to the presence of abundant hydroxyl groups in the alginate, which bind the metal ions. Similar effects were observed by Brun et al., 1998; Gloaguen et al., 1996; Ye et al., 1997.

Table 4. Biosorption of chromium by *Oscillatoria sp.*,

Treatment	Initial concentration (µg/mL)	Final concentration (µg/mL)	Metal uptake by biomass	Metal removal (%)
Living cells	5	4.14	<b>0.86</b>	17.2
	10	9.06	0.94	9.4
Heat killed cells	5	4.19	0.81	16.2
	10	8.91	<b>1.09</b>	10.9
Pre-treated cells	5	4.72	0.28	5.6
	10	9.27	0.73	7.3

### CONCLUSIONS

The results of this study indicated that the biomass of *Oscillatoria sp.*, was suitable for the development of efficient biosorbent for the removal of heavy metal chromium from waste water. Sequentially, cyanobacteria grown in metal polluted environments had proven the ability to tolerate the high concentration of metals ions Cu, Cd and Zn. Our results has showed that high amount of metal was adsorbed in heat killed cells of *Oscillatoria sp.*, in 10µg/mL and 5µg/mL the high amount of metal was adsorbed in living cells. In fact, the pre-treated cells have shown less effect in the removal of chromium at various concentrations. An increase in metal concentration caused a decrease in the growth of the *Oscillatoria sp.*, and decreased the cellular contents like chlorophyll a, carotenoids, C.phycocyanin, allophycocyanin, sugars, free amino acids and proteins. The ability provided a preliminary data screening the potential of microbial biomass to be used as a biosorption for the removal of metal ions from contaminated waters. This will be useful to understand the absence/presence of chromium toxicity in waste water from industrial effluents, the bioabsorption tolerance of cyanobacteria could be a useful tool for characterizing chromium and other heavy metals contamination in the environment to monitor the bioremediation progressively.

### ACKNOWLEDGEMENTS

The authors S. Jayashree and J. Joel Gnanadoss express gratitude to the management, Loyola College for providing necessary facilities.

### REFERENCES

- [1] Saiqa Razi and Shahida Hasnain 2006. Gamma Irradiation: Impact on Chromate Resistant Cyanobacteria. *Science Asia* Vol .1, 107-111.
- [2] Jasvir. S., Khattar, Tangairalaa.Srama, Davinerp.singh, and Anuradha Sharma. 2002. Bioaccumulation of Chromium Ions by Immobilized Cells Of A Filamentous Cyanobacterium, *Anabaena Variabilis*. *J,microbial.biotechnol*, 137-141.
- [3] Akthar, N. M. D. and Mohan, P. M. 1995. Bioremediation of toxic metal ions from polluted lake waters and industrial effluents by fungal biosorbent. *Curr Sc*. 69 : 1028-1038.
- [4] Rani Gupta , Perna Ahuja, Seema Khan, Saxena, Harapriya mohapatra., 2000. microbial biosorbents: meeting challenges of heavy metal pollution in aqueous solution. *CurreInt Science*,vol 78.
- [5] Ramteke, P.W. 2000. Biosorption of nickel (II) by *Pseudomonas stutzeri*. *Environ. Biol.*, 21: 219-221.
- [6] Kuyucak, N and B. Volesky. 1989. Accumulation of Co by marine alga. *Biotechnol. Bioengineer.*, 33:809-814.
- [7] Garnham,G.W., G.A.Godd and G.M.Godd. 1992. Accumulation of Co, Zinc and Manganese by the estuarine green microalgae *Chlorella salina* immobilized in alginate beads. *Environ. Sci.Technol.*, 26:1764-1770.
- [8] Slotton, D.G., Goldman. C.R., and Frank. A., 1989. Commercially Grown *Spirulina* Found to Contain Low Levels of Mercury and Lead, *Nutrition Reports International*, vol. 2, 1165-1172.
- [9] Bender, J., Gould, J.P, Vatcharapijarn, Y., Young, J.S. and Phillips. P, 1994. Removal of Zinc and Manganese from Contaminated Water with *Cyanobacteria* Mats, *Water Environ.Research*, vol. 5 : 679-683.
- [10] Hasnain, Abdul Hameed., 2005. Cultural characteristics of chromium resistant unicellular cyanobacteria isolated from local environment in Pakistan. *Chinese Journal of Oceanology and Limnology* vol. 4 : 433-441.
- [11] Gardea-Torresdey, L., Becker-Hapak, M.K., Hosea, J.M., and Darnell, D.W. 1990. Effect of Chemical Modification of Algal Carboxyl Groups on Metal Ion Binding, *Environ. Sci. Technol.*, 19, pp. 1372-1379.



- [12] Amber Cain; Raveender Vannela; L. Keith Woo, Cyanobacteria as a biosorbent for mercuric ion. *Bioresource Technology*, Volume 99, Issue 14, September 2008, Pages 6578-6586.
- [13] Pandi. M., V; Shashirekha; Mahadeswara Swamy, Bioabsorption of chromium from retan chrome liquor by cyanobacteria, *Microbiological Research*, Volume 164, Issue 4, 2009, Pages 420-428.
- [14] Lei Yan., Huanhuan Yin; Shuang Zhang; Feifan Leng; Wenbin Nan; Hongyu Li, Biosorption of inorganic and organic arsenic from aqueous solution by *Acidithiobacillus ferrooxidans* BY-3, *Journal of Hazardous Materials*, Volume 178, Issues 1-3, 15 June 2010, Pages 209-217.
- [15] Fuhong Sun., Fengchang Wu., Haiqing Liao., Baoshan Xing., Biosorption of antimony(V) by freshwater cyanobacteria *Microcystis* biomass: Chemical modification and biosorption mechanisms, *Chemical Engineering Journal*, Volume 171, Issue 3, 15 July 2011, Pages 1082-1090.
- [16] Rippka R., Deruelles J., Waterbury J.B., Herdman M., Stanier R.Y., 1979. Generic assignments, strain histories and properties of pure culture of cyanobacteria. *J. Gen. Microbiol.*, 111: 1-61.
- [17] Mackinney, G., 1941. Absorption of light by chlorophyll solutions. *J. Biol. Chem.*, 140:315-322.
- [18] Bennet, A. and L. Bogoras, 1973. Complimentary chromatic adaptation in a filamentous blue-green alga. *J. cell Biol.* 48:419-433.
- [19] Sadasivam, S and Manickam, A, 1996. Biochemical Methods. New Age International (P) Limited, Publishers, New Delhi.
- [20] Lowry, O.H., Rosebrough, N.J., Farr, A.L. and Randall, R.J. 1951. Protein measurement with Folin-phenol reagent. *J. Biol. Chem.* 193: 265-275.
- [21] Singh, S.P., Verma, S.K.; Singh, R.K. and Pandey P.K. Copper uptake by free and immobilized cyanobacterium. *FEMS Microbiological Letters*, 1989, vol. 60, p. 193-196.
- [22] Rocchetta, I., V. Mazzuc and M.R. Carmen, 2006. Effect of chromium on the fatty acid composition of two strains of *Euglena gracili*. *Environ. Pollut.*, 141: 353-358. DOI: 10.1016/j.envpol.2005.08.035.
- [23] Davies, A.G. and Sleep, J.A. 1978. Photosynthesis in some British coastal waters may be inhibited by zinc pollution. *Nature, London* 227, 292-293.
- [24] Banerjee, M. and Mishra, S. Mitigating effect of immobilization on *Aulosira fertilissima* subjected to nickel and chromium stress. *Bulletin of Bioscience*, 2002, vol. 1, p. 43-46.
- [25] Daniel, S.F., Peters, T., Dean Adams, V., and J of Middlebrooks, E.: 1979, *Water Res.* 13, 305.
- [26] Bolanos, L., M. Garcia-Gonzalez, P. Mateo, and I. Bonilla. 1992. Differential toxicological response to cadmium in *Anabaena* strain PCC 7119 grown with NO<sub>3</sub> or NH<sub>4</sub> + as nitrogen sources. *J. Plant Physiol.* 140: 345-349.
- [27] Rai, L. C., J. P. Gaur, and H. D. Kumar. 1981. Protective effects of certain environmental factors on the toxicity of zinc, mercury and methyl mercury to *Chlorella vulgaris*. *Environ. Res.* 25: 250-259.
- [28] Rai, L. C., N. Mallick, J. B. Singh, and H. D. Kumar. 1991. Physiological and biochemical characteristics of a copper tolerant and wild type strain of *Anabaena doliolum* under copper stress. *J. Plant Physiol.* 38: 68-74.
- [29] Raizada. M., Rai, L. C., N. Mallick, Y. Husaini, A. K. Singh, and S. K. Dubey. 1990. Effect of four heavy metals on the biology of *Nostoc muscorum*. *Biol. Metals* 2: 229-234.
- [30] Sorentinom C. 1979. The effects of heavy metals on plankton: A review. *Phykos* 18: 149-161.
- [31] Mohanty, N. and P. Mohanty. 1985. Cation effects on primary processes of photosynthesis, pp. 1-18. In F. R. Singh and S. K. Sawhney (eds.), *Advances in Frontier Areas of Plant Biochemistry*. Prentice Hall of India, New Delhi.
- [32] Murthy, S. D. S. and P. Mohanty. 1999. Heavy metal ions induced inhibition in bioenergetic processes of photosynthesis, pp. 185-186. In G. S. Singhal and T. Ramasarma (eds.), *Trends in Bioenergetic and Biotechnological Processes*. Today and Tomorrow Publishers, New Delhi.
- [33] Gupta, A. and Singhal, G.S. 1995. Inhibition of PS II activity by copper and its effects on spectral properties of intact cells in *Anacystis nidulans*. *Environ. Exp. Bot.* 35 : 435-438.
- [34] Fathi, A.A., F.T. Zaki and H.A. Ibraheim, 2005. Response of tolerant and wild type strains of *Chlorella vulgaris* to Copper with special references to Copper uptake system. *Protistology*, 4: 73-78.
- [35] Torres, M., J. Goldberg and T.E. Jensen, 1998. Heavy metal uptake by polyphosphate bodies in living and killed cells of *Plectonema boryanum* (Cyanophyceae). *Microbios*, 96: 141-147.
- [36] Osman, M.E.H., A.H. El-Naggar, M.M. El-Sheekh and E. El-Mazally, 2004. Differential effects of Co<sup>2+</sup> and Ni<sup>2+</sup> on protein metabolism in *Scenedesmus obliquus* and *Nitzschia perminuta*. *Environ. Toxicol. Pharmacol.*, 16: 169-178. DOI: 10.1016/j.etap.2003.12.004
- [37] Fathi, A.A., F.T. Zaki and A.A. Fathy, 2000. Bioaccumulation of some heavy metals and their influence on the metabolism of *Scenedesmus bijuga* and *Anabaena spiroides*. *Egypt J. Biotechnol.*, 7: 293-307.
- [38] Tripathi, B.N.I. and J.P. Gaur, 2006. Physiological behavior of *Scenedesmus* sp. during exposure to elevated levels of Cu and Zn and after withdrawal of metal stress. *Protoplasma*, 229: 1-9. DOI: 10.1007/s00709-006-0196-9.
- [39] Barbara, W. and W. Michael, 1994. Comparative studies of the heavy metal uptake of whole cells and different types of cell walls from *Chlorella fusca*. *Biotechnol. Tech.*, 8: 227-232. DOI:10.1007/BF00155412
- [40] Fisher, N.S., Bohw, M. and Teysie, J.L. 1984. accumulation and toxicity of Cd, Zn, Hg, and Ag in the marine phytoplanktons. *Mar. Ecol. Prog. Ser* 18:201-204.
- [41] Rabsch, V. and Elbracher, R 1980 cadmium and zinc uptake growth and primary production in *Coscinodiscus granii* cultures containing low levels of cells and dissolved organic carbon. *Helgol. Wiss meeresuter* 33:79-88.
- [42] Ahuja P, Gupta R, saxena RK (1999) Zn<sup>2+</sup> biosorption by

- Oscillatoria angustissima*. Proc Biochem 34:77 -85.
- [43] Brun, L.A.; Maillet, J.; Richarte, J.; Herrmann, P. and Remy, J.C. Relationships between extractable copper, soil properties and copper uptake by wild plants in vineyard soils. *Environmental Pollution*, 1998, vol. 102, p. 151-161.
- [44] Gloaguen, V., Morvan, H. and Hoffmann, L. Metal accumulation by immobilized cyanobacterial mats from a thermal spring. *Journal of Environmental Science and Health*, 1996, vol. A31, no. 10, p. 2437-2451.
- [45] Ye, Z.H.; Baker, A.J.M.; Wong, M.H. and Willis, A.J. Zinc, lead and cadmium tolerance, uptake and accumulation in populations of *Phragmites australis* (Cav.) Trin. ex Steudel. *Annals of Botany*, 1997, vol.80, p. 363-370.