

Horticultural Biotechnology Research 2017, 3: 26-36
 doi: 10.25081/hbr.2017.v3.3450
<http://updatepublishing.com/journal/index.php/hbr/>



ISSN: 2455-460X

REGULAR ARTICLE

ORGANIC FERTILIZATION AND NATURAL SUBSTANCES TREATMENTS AFFECTS CHEMICAL CONSTITUENTS OF GUAR PLANTS

ABEER M. SHEHATA^{1,2}, M. A. H. ABDOU², A. A. EL-SAYED², F. A. ATTIA², R. A. TAHA²

¹Medicinal and Aromatic Plants Researches Department, Horticulture Research Institute, Agricultural Research Center, Egypt

²Horticulture Department, Faculty of Agriculture, Minia University, Egypt

ABSTRACT

This experiment was conducted at the Floriculture Nursery, Faculty of Agriculture, Minia University during the two successive seasons to study the effect of compost at rates (0, 25, 2.50 and 3.75 ton/fed.) and natural substances i.e. vitamins (ascorbic acid at 50 and 100 ppm, α -tocopherol at 10, 20 ppm and thiamine, at 25 and 50 ppm) and active yeast at 2.5 and 5 g/l. and their interaction on some chemical constituents of guar plants. The results indicated that, Addition of compost caused an increase in guaran %, Photosynthetic pigments, Total carbohydrates (%), N, P and K (%) and Protein (%). The maximum level in all previous traits was for the plants growing in the soil fertilized with the highest level of compost (3.75 t/fed.). Using the treatments of vit. C at 50 ppm followed by 5 g/l. active yeast, then vit. E at 10 ppm gave the highest values of all previous traits. The interaction between the two main factors (A×B) was significant for guaran %, photosynthetic pigments, total carbohydrates %, N and P %, The highest values were obtained from the interaction treatments of 3.75 ton/fed., compost in combination with active yeast at 5 g/l. and vit. C at 50 ppm. On the other hand, the interaction between compost, active yeast and vitamins treatments was not significant for protein content (%). We conclude that supply guar plants with compost at 3.75 ton/fed., and spraying plants with either active yeast at 5 g/l. or ascorbic acid (vit. C) at 50 ppm to improve the values of some chemical constituents under investigation condition.

Keywords: Guar, Compost, Vitamin C, Active yeast, Chemical constituents

INTRODUCTION

Guar, *Cyamopsis tetragonoloba*, L. Taub. (cluster bean vegetable), an annual summer adapted well to arid and semi-arid climatic zones [1]. It is also known as saline and drought resistant plant; it belongs to Family Fabaceae [2]. Guar is known and grown in Egypt as forage crop. Primarily, its green manure and seed production have considered as animal nutritive substances as its protein content 16 %. Recently, the interest has expanded to use a source of glectomannan gum. The endosperm contains guaran gum (20-30 %), a polymer of galactose (36 %) and manose (64 %), known as glectomannan [3]. The gum is used in food industry [4], it can also be used for treating diabetes mellitus, hyperglycemia, glycosuria and hyperlipoproteinemia and the seeds are used as a laxative [5]. Recently, the interest has extended to be use as a source of galactomannan gum.

Organic manures are important for medicinal plants to produce the best product in both quantity and quality and it is also very safe for human health and environment [6]. Composts are ecofriendly organic supplements which are

not harmful for even soil microbiota [7]. Recently, great attention has been focused on the possibility of using natural substances, i.e. vitamins (E or B₁ or C) and yeast in order to improve plant growth, flowering, fruit setting and yield.

Alpha-tocopherols (vitamin E), are secondary metabolites which protects plants from various abiotic stresses [8, 9]. Thiamine (vitamin B₁) is an essential for the synthesis of succinyl-Co-A and glycerin and for the reaction of them to form aminolevulinic acid [10], the main intermediate for formation of protoporphyrin in the precursor of chlorophyll [11] and a factor in the decarboxylation of pyruvate. In addition, various physiological processes depend more or less on the availability of vitamin B [12]. Moreover, [13] obtained pronounced increments in lemongrass yield, vegetative growth, as well as, essential oil percentage due to applying thiamine.

Ascorbic acid (vitamin C) an important antioxidant in plants which helps the plants to overcome stressful conditions [14]. In addition, yeast extract is a source of many natural plant growth substances (cytokinins), large

Received 21 November 2017; Accepted 30 December 2017

*Corresponding Author

Abeer M. Shehata

Medicinal and Aromatic Plants Researches Department, Horticulture Research Institute, Agricultural Research Center, Egypt

Email: abeershehata2006@yahoo.com

©This article is open access and licensed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>) which permits unrestricted, use, distribution and reproduction in any medium, or format for any purpose, even commercially provided the work is properly cited. Attribution – You must give appropriate credit, provide a link to the license, and indicate if changes were made.

amount of vitamin B and most of nutritional elements (P, K, S, Na, Ca and Mg), as well as, organic compound (high protein, carbohydrates, nucleic acids and lipids) [15].

Therefore, the present study was carried out to investigate the effect of organic manure (compost) and some natural substances (vitamin E, B₁ and C, as well as, active yeast) treatments on some chemical constituents of *Cyamopsis tetragonoloba*, L. plant.

MATERIALS AND METHODS

The present work was concluded at the Floriculture Nursery, Faculty of Agriculture, Minia University during the two successive seasons of 2010 and 2011 to study the effect of compost in combination with active yeast and some vitamins on growth, yield and guaran content of guar gum (*Cyamopsis tetragonoloba*, Taub.). Table (1) shows the physical and chemical properties of the used soil in both seasons.

The layout of this experiment during both seasons was split plot design with three replicates. The main-plots included four compost treatments (0, 1.25, 2.50 and 3.75 ton/feddani.), while the sub-plots were devoted to nine treatments; control, α -tocopherol (vit. E.) at 10 and 20 ppm, thiamine (vit. B₁) at 25 and 50 ppm, ascorbic acid (vit. C) at 50 and 100 ppm and active yeast (AY) at 2.5 and 5 g/l. Compost were added during soil preparation for cultivation in both seasons. The physical and chemical properties of the used compost are shown in table (2).

The plants were harvested of the fourth week of September in both seasons and the following data were recorded during the two seasons.

Guaran determinations

Concerning guaran determination of oven-dried seeds were determined according to [16].

Pigments determination

The contents of photosynthetic pigments namely; chlorophyll a, b and carotenoids (mg/g f. w.) of fresh leaves were determined according to [17] using the spectrophotometer at wave length of 656, 665 and 452.5 μ m, respectively.

Total carbohydrates percentage

Total carbohydrates including poly-saccharides in dry leaves of each experiment unit were colorimetrically determined with the anthrone sulphuric acid method [18]

N, P and K percentages

A sample weight of 0.2 g fine powder of the dry herb was digested using a mixture of hydrogen peroxide (H₂O₂) and concentrated sulphuric acid (H₂SO₄) (4:10). The clear digestion was quantitatively 100 ml volumetric flask. In this solution, the following elements were determined:

Nitrogen (%) was determined according to the modified Microkjeldahle method as described by [19]. Phosphorus (%) was determined colorimetrically by the spectrophotometer at wavelength of 650 μ m according to the method of [20]. Potassium (%) was determined using flame-photometry method according to [21].

Table 1: The physical and chemical properties of the used soil

Soil character	Values	Soil character	Values
Sand %	28.30	Available P %	15.12
Silt %	30.70	Exchangeable K ⁺ mg/100 g soil	2.11
Clay %	41.0	Exch. Ca ⁺⁺ mg/100 g soil	31.74
Soil type	Clay loam	Exch. Na ⁺ mg/100 g soil	2.41
Organic matter %	1.62		
CaCO ₃ %	2.09		Fe 8.54
pH 1:2.5	7.83	DTPA	Cu 2.06
E. C. m mhose/cm	1.04	Ext. ppm	Zn 2.75
Total N %	0.08		Mn 8.26

The experimental unit (plot) was 2 × 2.4 meters and contained 3 rows, 50 cm apart and seeds were sown in hills, 30 cm apart, on 30th March in both seasons. Seedlings were thinned to one plant/hill, after one month from sowing.

Table 2: The physical and chemical properties of the used compost

Properties	Value	Properties	Value
Dry weight of 1 m ³	450 kg	NaCl %	1.1-1.75
Fresh weight of 1 m ³	650-700 kg	Total P %	0.5-0.75
Moisture (%)	25-30	Total K %	0.8-1.0
pH (1:10)	7.5-8	Fe ppm	150-200
E. C. (m mhose/cm)	2-4	Mn ppm	25-56
Total N %	1-1.4	Cu ppm	75-150
Org. matter %	32-34	Zn ppm	150-225
Org. carbon %	18.5-19.7		
C/N ratio	18.5-14.1		

Each of vitamins and suspension of yeast were applied by hand sprayer 3 times. The first one was added after 5 w from planting date and two weeks thereafter. The plants were sprayed till run off. All agricultural practices were performed as usual, in the region for the production of guar plants.

Protein percentage

Protein percentage was estimated by multiplying nitrogen percent by 6.25. This was based on the assumption that the protein contains 16 % nitrogen, according to the method of [22].

Statistical analysis

The data of the two seasons were subjected to the statistical analysis of variance using [23]. L. SD test at 0.05 was used to compare the means of treatments.

RESULTS AND DISSECTION

Guaran percentage in the seeds

Regarding compost levels, data presented in table (3) revealed that guaran % in the dry seeds of *Cyamopsis tetragonoloba*, L. was significantly increased as a result of fertilizing plants with all used levels of compost treatments over the control in both seasons. The highest guaran % was obtained due to the treatment of compost at high level (3.75 ton/fed.) followed by the treatment of low level (1.25 ton/fed.) with significant differences detected among themselves in both growing seasons.

The enhancement of guaran % obtained in the present work due to using organic manure was also reported by [24] and [25] on guar plants.

Table (3) indicated that all examined eight treatments of vitamins and active yeast significantly increased guaran percentage in the dry seeds of *Cyamopsis tetragonoloba*,

L., except for the treatment of vit. C at 100 ppm in comparison with the control in the two growing seasons. The low concentrations of the three used vitamins treatments proved to be more effective than their high concentrations in this respect. However, active yeast at 5 g/l. was more effective than 2.5 g/l. in improving guaran %. The treatments of active yeast (5 g/l.) followed by vit. C (50 ppm) then B1 (25 ppm) resulted the highest guaran % with significant differences detected among themselves in both seasons. The least guaran % resulted from control treatment. The other treatments gave intermediate values. In this respect, [24] on guar plants found that the highest values of guaran % were obtained due to the application of farmyard manure accompanied by *Rhizobium* and yeast. The interaction between the two main factors (A×B) was significant for guaran % in the second season as illustrated in table (3). The superior interaction treatment over all was compost at high level (3.75 ton/fed.) in combination with active yeast at 5 g/l.

Photosynthetic pigments

The contents of the three studied photosynthetic pigments (chlorophyll a, b and carotenoids) were significantly enhanced by compost levels (1.25, 2.50 and 3.75 ton/fed.) in the two growing seasons as shown in Tables (4, 5 and 6). However, the maximum values were obtained by high level of compost (3.75 ton/fed.). In regard to the enhancement influence of organic fertilization, the previous results were in agreement with those of [26] on coriander; [27] on *Phaseolus vulgaris* and [28] on spider plant.

Table 3: Effect of compost, some vitamins and active yeast treatments on guaran % of *Cyamopsis tetragonoloba* seeds during 2010 and 2011 seasons

First season						
Vitamins and active yeast treatments (B)	Compost levels (ton/fed.) (A)					
	0.0	1.25	2.50	3.75	Mean (B)	
Control	11.56	11.95	12.31	12.68	12.13	
	12.67					
	11.98					
	12.78					
	12.04					
Vit. E at 10 ppm	12.67	12.78	13.10	13.61	13.04	
		12.40				
Vit. E at 20 ppm	11.98	12.40	12.75	13.15	12.57	
Vit. B ₁ at 25 ppm	12.78	12.89	13.21	13.73	13.15	
Vit. B ₁ at 50 ppm	12.04	12.45	12.79	13.19	12.62	
Vit. C at 50 ppm	13.17	13.30	13.73	14.32	13.63	
Vit. C at 100 ppm	11.59	12.00	12.36	12.72	12.17	
Active yeast (2.5 g/l)	12.25	12.66	12.99	13.40	12.83	
Active yeast (5.0 g/l)	13.50	13.82	14.25	14.65	14.06	
Mean (A)	12.39	12.69	13.05	13.49		
L. SD at 5 %	A:	0.16	B:	0.19	AB:	N. S.
Second season						
Vitamins and active yeast treatments (B)	Compost levels (ton/fed.) (A)					
	0.0	1.25	2.50	3.75	Mean (B)	
Control	11.95	12.45	12.90	13.27	12.64	
Vit. E at 10 ppm	13.05	13.29	13.73	14.23	13.58	
Vit. E at 20 ppm	12.39	12.91	13.36	13.76	13.10	
Vit. B ₁ at 25 ppm	13.19	13.40	13.82	14.35	13.69	
Vit. B ₁ at 50 ppm	12.46	12.97	13.38	13.80	13.15	
Vit. C at 50 ppm	13.59	13.81	14.36	14.95	14.18	
Vit. C at 100 ppm	11.97	12.50	12.95	13.30	12.68	
Active yeast (2.5 g/l)	12.67	13.19	13.60	14.00	13.37	
Active yeast (5.0 g/l)	13.93	14.03	14.69	15.31	14.49	
Mean (A)	12.80	13.17	13.64	14.11		
L. SD at 5 %	A:	0.11	B:	0.11	AB:	0.22

Vit. E. = α-tocopherol Vit. B₁. = Thiamine Vit. C. = Ascorbic acid

Table 4: Effect of compost, some vitamins and active yeast treatments on chlorophyll a content (mg/g f. w.) of *Cyamopsis tetragonoloba* during 2010 and 2011 seasons

First season						
Vitamins and active yeast treatments (B)	Compost levels (ton/fed.) (A)					
	0.0	1.25	2.50	3.75	Mean (B)	
Control	2.311	2.376	2.423	2.466	2.394	
	2.399					
Vit. E at 10 ppm	2.399	2.458	2.507	2.548	2.478	
		2.438				
Vit. E at 20 ppm	2.369	2.438	2.478	2.518	2.451	
Vit. B ₁ at 25 ppm	2.381	2.441	2.491	2.531	2.461	
Vit. B ₁ at 50 ppm	2.360	2.429	2.470	2.512	2.443	
Vit. C at 50 ppm	2.428	2.488	2.549	2.588	2.528	
Vit. C at 100 ppm	2.341	2.400	2.451	2.491	2.421	
Active yeast (2.5 g/l)	2.356	2.421	2.462	2.506	2.437	
Active yeast (5.0 g/l)	2.457	2.519	2.561	2.602	2.535	
Mean (A)	2.378	2.441	2.488	2.529		
L. SD at 5 %	A:	0.052	B:	0.011	AB:	0.220
Second season						
Vitamins and active yeast treatments (B)	Compost levels (ton/fed.) (A)					
	0.0	1.25	2.50	3.75	Mean (B)	
Control	2.351	2.411	2.462	2.500	2.431	
Vit. E at 10 ppm	2.401	2.461	2.518	2.555	2.484	
Vit. E at 20 ppm	2.375	2.439	2.481	2.521	2.454	
Vit. B ₁ at 25 ppm	2.380	2.445	2.490	2.528	2.461	
Vit. B ₁ at 50 ppm	2.365	2.431	2.476	2.516	2.447	
Vit. C at 50 ppm	2.448	2.508	2.568	2.571	2.523	
Vit. C at 100 ppm	2.352	2.418	2.477	2.520	2.442	
Active yeast (2.5 g/l)	2.359	2.421	2.486	2.529	2.449	
Active yeast (5.0 g/l)	2.468	2.520	2.573	2.548	2.527	
Mean (A)	2.389	2.450	2.503	2.536		
L. SD at 5 %	A:	0.041	B:	0.009	AB:	0.018

Vit. E. = α -tocopherol Vit. B₁. = Thiamine Vit. C. = Ascorbic acid**Table 5: Effect of compost, some vitamins and active yeast treatments on chlorophyll b content (mg/g f. w.) of *Cyamopsis tetragonoloba* during 2010 and 2011 seasons**

First season						
Vitamins and active yeast treatments (B)	Compost levels (ton/fed.) (A)					
	0.0	1.25	2.50	3.75	Mean (B)	
Control	0.771	0.794	0.807	0.822	0.799	
	0.801					
Vit. E at 10 ppm	0.801	0.820	0.853	0.967	0.860	
		0.815				
Vit. E at 20 ppm	0.790	0.815	0.841	0.851	0.824	
Vit. B ₁ at 25 ppm	0.795	0.813	0.846	0.859	0.828	
Vit. B ₁ at 50 ppm	0.787	0.812	0.835	0.841	0.819	
Vit. C at 50 ppm	0.809	0.829	0.856	0.980	0.869	
Vit. C at 100 ppm	0.781	0.800	0.817	0.830	0.807	
Active yeast (2.5 g/l)	0.785	0.807	0.821	0.836	0.812	
Active yeast (5.0 g/l)	0.819	0.839	0.866	0.985	0.877	
Mean (A)	0.793	0.814	0.838	0.886		
L. SD at 5 %	A:	0.018	B:	0.007	AB:	0.014
Second season						
Vitamins and active yeast treatments (B)	Compost levels (ton/fed.) (A)					
	0.0	1.25	2.50	3.75	Mean (B)	
Control	0.783	0.805	0.820	0.831	0.810	
Vit. E at 10 ppm	0.808	0.828	0.860	0.973	0.867	
Vit. E at 20 ppm	0.791	0.813	0.833	0.855	0.823	
Vit. B ₁ at 25 ppm	0.799	0.816	0.838	0.860	0.828	
Vit. B ₁ at 50 ppm	0.791	0.811	0.833	0.850	0.821	
Vit. C at 50 ppm	0.809	0.836	0.871	0.988	0.878	
Vit. C at 100 ppm	0.784	0.808	0.825	0.840	0.814	
Active yeast (2.5 g/l)	0.788	0.809	0.829	0.843	0.817	
Active yeast (5.0 g/l)	0.821	0.843	0.868	0.991	0.881	
Mean (A)	0.797	0.819	0.842	0.892		
L. SD at 5 %	A:	0.015	B:	0.003	AB:	0.006

Vit. E. = α -tocopherol Vit. B₁. = Thiamine Vit. C. = Ascorbic acid

Table 6: Effect of compost, some vitamins and active yeast treatments on carotenoids content (mg/g f. w.) of *Cyamopsis tetragonoloba* during 2010 and 2011 seasons

First season						
Vitamins and active yeast treatments (B)	Compost levels (ton/fed.) (A)					
	0.0	1.25	2.50	3.75	Mean (B)	
Control	0.793	0.811	0.819	0.834	0.814	
Vit. E at 10 ppm	0.822	0.850	0.867	0.888	0.857	
		0.851				
Vit. E at 20 ppm	0.809	0.839	0.851	0.876	0.844	
Vit. B1 at 25 ppm	0.815	0.843	0.861	0.881	0.850	
Vit. B1 at 50 ppm	0.805	0.831	0.841	0.866	0.836	
Vit. C at 50 ppm	0.829	0.859	0.880	0.899	0.867	
Vit. C at 100 ppm	0.796	0.818	0.828	0.846	0.822	
Active yeast (2.5 g/l)	0.799	0.826	0.836	0.858	0.830	
Active yeast (5.0 g/l)	0.838	0.866	0.910	0.930	0.886	
Mean (A)	0.812	0.838	0.855	0.875		
L. SD at 5 %	A:	0.019	B:	0.006	AB:	0.012
Second season						
Vitamins and active yeast treatments (B)	Compost levels (ton/fed.) (A)					
	0.0	1.25	2.50	3.75	Mean (B)	
Control	0.795	0.818	0.836	0.846	0.824	
Vit. E at 10 ppm	0.829	0.860	0.875	0.896	0.865	
Vit. E at 20 ppm	0.811	0.832	0.852	0.876	0.843	
Vit. B1 at 25 ppm	0.817	0.839	0.859	0.886	0.850	
Vit. B1 at 50 ppm	0.810	0.832	0.846	0.869	0.839	
Vit. C at 50 ppm	0.831	0.863	0.899	0.906	0.875	
Vit. C at 100 ppm	0.796	0.825	0.838	0.853	0.828	
Active yeast (2.5 g/l)	0.801	0.831	0.846	0.862	0.835	
Active yeast (5.0 g/l)	0.842	0.869	0.912	0.939	0.891	
Mean (A)	0.814	0.841	0.863	0.881		
L. SD at 5 %	A:	0.018	B:	0.011	AB:	0.022

Vit. E. = α -tocopherol Vit. B₁= Thiamine Vit. C.= Ascorbic acid

Table 7: Effect of compost, some vitamins and active yeast treatments on total carbohydrates (%) of *Cyamopsis tetragonoloba* during 2010 and 2011 seasons

First season						
Vitamins and active yeast treatments (B)	Compost levels (ton/fed.) (A)					
	0.0	1.25	2.50	3.75	Mean (B)	
Control	22.2	25.3	27.1	29.5	26.0	
Vit. E at 10 ppm	23.5	28.2	29.5	31.8	28.3	
		2.80				
Vit. E at 20 ppm	23.3	28.0	29.1	31.3	27.9	
Vit. B1 at 25 ppm	23.4	28.1	29.3	31.4	28.0	
Vit. B1 at 50 ppm	23.2	27.9	29.1	31.2	27.9	
Vit. C at 50 ppm	23.8	28.5	29.9	32.1	28.6	
Vit. C at 100 ppm	23.1	26.4	28.3	30.1	27.0	
Active yeast (2.5 g/l)	23.7	27.1	29.1	31.1	27.8	
Active yeast (5.0 g/l)	23.6	28.3	29.6	31.9	28.4	
Mean (A)	23.3	25.3	29.0	31.2		
L. SD at 5 %	A:	0.18	B:	0.4	AB:	0.8
Second season						
Vitamins and active yeast treatments (B)	Compost levels (ton/fed.) (A)					
	0.0	1.25	2.50	3.75	Mean (B)	
Control	23.5	26.7	27.9	29.4	26.9	
Vit. E at 10 ppm	23.6	28.6	29.4	31.8	28.4	
Vit. E at 20 ppm	23.4	28.1	29.2	31.6	28.1	
Vit. B1 at 25 ppm	23.4	28.2	29.2	31.5	28.0	
Vit. B1 at 50 ppm	23.3	28.0	29.1	31.5	28.0	
Vit. C at 50 ppm	23.7	29.5	31.1	31.3	28.9	
Vit. C at 100 ppm	23.2	26.5	28.3	30.3	27.2	
Active yeast (2.5 g/l)	23.9	27.2	29.2	31.2	27.9	
Active yeast (5.0 g/l)	23.6	28.7	29.6	31.9	28.5	
Mean (A)	23.5	27.9	29.4	31.3		
L. SD at 5 %	A:	2.5	B:	0.5	AB:	1.0

Vit. E. = α -tocopherol Vit. B₁= Thiamine Vit. C.= Ascorbic acid

Data presented in Tables (4, 5 and 6) showed that all spraying treatments significantly increased the chlorophyll a, b and carotenoids contents (mg/g. f. w.) over the check

treatment. The highest contents of the previous characters resulted from the plants which received active yeast at 5 g/l. followed by vit. C at 50 ppm in both experimental seasons.

The interaction between compost and spraying treatments was significant in both seasons for the three photosynthetic pigments. The highest contents of chlorophyll a, b and carotenoids in the fresh leaves of guar plants was obtained from the interaction treatments of 3.75 ton/fed., compost in combination with active yeast at 5 g/l. and vit. C at 50 ppm. Similar results were obtained by [29] on coriander; [30] on black cumin and anise and [31] on sage who found that the interaction between farmyard manure with yeast significantly increased chlorophyll a, b and carotenoids. Also, [32] on caraway; [33] on fennel and [34] on black cumin recorded that the highest contents of chl. a, b and carotenoids were obtained from the treatment of compost in combination with ascorbic acid.

Total carbohydrates percentage

Data presented in table (7) indicated that supplying guar plants with compost at 1.25, 2.50 and 3.75 ton/fed., resulted in a significant increase in the total carbohydrate percentage in the dry seeds of plant compared to control plants. The increase of carbohydrate % was gradually increased with the gradual increase of compost level in the two experimental seasons. The beneficial effect of organic fertilization on carbohydrate was obtained by [35] on *Sideritis monata*; [36] on oregano plants; [37] on *Lawsonia alba*; [38] on *Vicia faba*; [39] on kidney bean and [40] on sugar beet plant.

Concerning the influence of spraying treatments [(vit. C, E and B₁)] and active yeast, each at two concentrations on carbohydrate %, data presented in table (7) showed that all used eight treatments increased the percentage of carbohydrate compared to untreated plants in both seasons. Using the treatments of vit. C at 50 ppm followed by 5 g/l. active yeast, then vit. E at 10 ppm gave the highest values in this concern without significant differences detected among themselves.

Regarding ascorbic acid these results agree with those of [41] on *Vicia faba*. However, [39] on kidney bean; [42] on *Vicia faba* and [40] on sugar beet, concluded that active yeast increased total carbohydrates in plants. Moreover, [43] on *Syngonium podophyllum* found that total carbohydrates content was significantly increased when plants were treated with thiamine, ascorbic acid and kinetin.

Regarding interaction between the two main factors (A×B) it significantly increased carbohydrate % in both two seasons as shown in table (7). The highest values were obtained by the interaction treatments of compost (3.75 ton/fed.) in combination with vit. C (50 ppm), active yeast (5 g/l.) or vit. E (10 ppm) during the first season. Moreover, the interaction between high level of compost (3.75 ton/fed.) in combination with any tested spraying treatment, except for vit. C at 100 ppm gave the best interaction treatments.

Table 8: Effect of compost, some vitamins and active yeast treatments on N (%) of *Cyamopsis tetragonoloba* during 2010 and 2011 seasons

First season						
Vitamins and active yeast treatments (B)	Compost levels (ton/fed.) (A)					
	0.0	1.25	2.50	3.75	Mean (B)	
Control	1.76	2.16	2.45	2.68	2.263	
Vit. E at 10 ppm	2.13					
	2.13	2.57	2.91	3.13	2.685	
		2.45				
		2.50				
		2.40				
Vit. E at 20 ppm		2.63				
	2.01	2.45	2.78	3.00	2.560	
Vit. B ₁ at 25 ppm	2.06	2.50	2.84	3.06	2.615	
Vit. B ₁ at 50 ppm	1.97	2.40	2.72	2.97	2.515	
Vit. C at 50 ppm	2.19	2.63	2.98	3.19	2.748	
Vit. C at 100 ppm	1.88	2.29	2.59	2.83	2.398	
Active yeast (2.5 g/l)	1.93	2.35	2.66	2.90	2.460	
Active yeast (5.0 g/l)	2.24	2.68	2.03	3.24	2.798	
Mean (A)	2.019	2.448	2.773	3.000		
L. SD at 5 %	A:	0.026	B:	0.060	AB:	0.120
Second season						
Vitamins and active yeast treatments (B)	Compost levels (ton/fed.) (A)					
	0.0	1.25	2.50	3.75	Mean (B)	
Control	1.84	2.25	2.55	2.79	2.358	
Vit. E at 10 ppm	2.14	2.59	2.94	3.17	2.710	
Vit. E at 20 ppm	2.05	2.50	2.83	3.03	2.603	
Vit. B ₁ at 25 ppm	2.10	2.56	2.89	3.08	2.658	
Vit. B ₁ at 50 ppm	2.02	2.46	2.79	2.99	2.565	
Vit. C at 50 ppm	2.20	2.65	3.00	3.23	2.770	
Vit. C at 100 ppm	1.95	2.37	2.68	2.92	2.480	
Active yeast (2.5 g/l)	1.99	2.42	2.74	2.98	2.533	
Active yeast (5.0 g/l)	2.24	2.70	3.05	3.29	2.820	
Mean (A)	2.059	2.500	2.830	3.053		
L. SD at 5 %	A:	0.042	B:	0.055	AB:	0.110

Vit. E. = α-tocopherol Vit. B₁. = Thiamine Vit. C. = Ascorbic acid

Table 9: Effect of compost, some vitamins and active yeast treatments on P (%) of *Cyamopsis tetragonoloba* during 2010 and 2011 seasons

First season						
Vitamins and active yeast treatments (B)	Compost levels (ton/fed.) (A)					
	0.0	1.25	2.50	3.75	Mean (B)	
Control	0.32 0.36 0.35	0.33	0.34	0.35	0.335	
Vit. E at 10 ppm	0.36	0.38 0.37	0.39	0.39	0.380	
Vit. E at 20 ppm	0.35	0.37	0.37	0.37	0.365	
Vit. B1 at 25 ppm	0.36	0.38	0.38	0.39	0.378	
Vit. B1 at 50 ppm	0.34	0.36	0.36	0.37	0.358	
Vit. C at 50 ppm	0.37	0.39	0.39	0.39	0.385	
Vit. C at 100 ppm	0.33	0.34	0.35	0.36	0.345	
Active yeast (2.5 g/l)	0.33	0.35	0.36	0.37	0.353	
Active yeast (5.0 g/l)	0.37	0.39	0.40	0.40	0.390	
Mean (A)	0.348	0.366	0.371	0.377		
L. SD at 5 %	A:	0.015	B:	0.010	AB:	0.020
Second season						
Vitamins and active yeast treatments (B)	Compost levels (ton/fed.) (A)					
	0.0	1.25	2.50	3.75	Mean (B)	
Control	0.30	0.31	0.32	0.33	0.315	
Vit. E at 10 ppm	0.35	0.38	0.39	0.39	0.378	
Vit. E at 20 ppm	0.34	0.37	0.38	0.38	0.368	
Vit. B1 at 25 ppm	0.35	0.37	0.38	0.39	0.373	
Vit. B1 at 50 ppm	0.34	0.36	0.37	0.37	0.360	
Vit. C at 50 ppm	0.36	0.39	0.39	0.40	0.385	
Vit. C at 100 ppm	0.32	0.33	0.34	0.35	0.335	
Active yeast (2.5 g/l)	0.34	0.35	0.36	0.37	0.355	
Active yeast (5.0 g/l)	0.37	0.39	0.40	0.41	0.393	
Mean (A)	0.341	0.361	0.370	0.377		
L. SD at 5 %	A:	0.018	B:	0.016	AB:	0.032

Vit. E. = α -tocopherol Vit. B1.= Thiamine Vit. C.= Ascorbic acid

Table 10: Effect of compost, some vitamins and active yeast treatments on K (%) of *Cyamopsis tetragonoloba* during 2010 and 2011 seasons

First season						
Vitamins and active yeast treatments (B)	Compost levels (ton/fed.) (A)					
	0.0	1.25	2.50	3.75	Mean (B)	
Control	1.30 1.35	1.33	1.36	1.40	1.348	
Vit. E at 10 ppm	1.35	1.40 1.39 1.39 1.38 1.40	1.41	1.48	1.400	
Vit. E at 20 ppm	1.34	1.39	1.39	1.46	1.395	
Vit. B1 at 25 ppm	1.35	1.39	1.40	1.47	1.403	
Vit. B1 at 50 ppm	1.34	1.38	1.39	1.45	1.390	
Vit. C at 50 ppm	1.36	1.40	1.42	1.48	1.415	
Vit. C at 100 ppm	1.33	1.36	1.38	1.43	1.375	
Active yeast (2.5 g/l)	1.33	1.37	1.39	1.44	1.383	
Active yeast (5.0 g/l)	1.36	1.41	1.43	1.49	1.423	
Mean (A)	1.340	1.381	1.397	1.456		
L. SD at 5 %	A:	0.022	B:	0.024	AB:	N. S.
Second season						
Vitamins and active yeast treatments (B)	Compost levels (ton/fed.) (A)					
	0.0	1.25	2.50	3.75	Mean (B)	
Control	1.28	1.30	1.34	1.36	1.320	
Vit. E at 10 ppm	1.36	1.38	1.40	1.41	1.388	
Vit. E at 20 ppm	1.34	1.36	1.39	1.41	1.375	
Vit. B1 at 25 ppm	1.35	1.37	1.39	1.42	1.383	
Vit. B1 at 50 ppm	1.33	1.35	1.38	1.40	1.365	
Vit. C at 50 ppm	1.36	1.39	1.41	1.42	1.395	
Vit. C at 100 ppm	1.31	1.33	1.37	1.39	1.350	
Active yeast (2.5 g/l)	1.32	1.34	1.37	1.40	1.358	
Active yeast (5.0 g/l)	1.37	1.39	1.42	1.43	1.403	
Mean (A)	1.336	1.357	1.386	1.404		
L. SD at 5 %	A:	0.022	B:	0.017	AB:	N. S.

Vit. E. = α -tocopherol Vit. B1.= Thiamine Vit. C.= Ascorbic acid

Nitrogen, phosphorus and potassium %

Data presented in Tables (8, 9 and 10) showed that the increase in N, P and K % in the dry leaves of guar plants was parallel to the increase in compost level. The high level of compost (3.75 ton/fed.) followed by medium level (1.25 ton/fed.) significantly increased N, P and K % in both seasons in comparison with those of control treatment.

The promotive effect of organic manure was found by many authors, namely, [25] on guar; [44] on soybean; [45] on *Pisum sativum*; [36] on *Origanium syriacum*; [46] on rosemary; [47] on *Eclipta alba*; and [48] on roselle plants.

Concerning the treatments of vitamins and active yeast, data presented in Tables (8, 9 and 10) clearly revealed that all used treatments considerably increased N, P and K % in the dry levels of guar plants in comparison with control plants during both seasons. Active yeast at high level 5 g/l. followed by ascorbic acid at low concentration (50 ppm) then α -tocopherol at low concentration (10 ppm) gave the highest percentages of N, P and K with non-significant differences among themselves in case of P and K %. The treatment of vit. B₁ (25 ppm) gave the second order in this concern.

Many researchers came to similar results that active yeast augmented NPK % in the leaves of different plants such as coriander [49]; [26]. Lemongrass [50]. *Phaseolus vulgaris* [51] and [52]. *Vicia faba* [38]. The role of vit. C and vit. E in promoting NPK % was also, reported by [53] on sweet pepper and [54] on eggplant. Regarding the effect of vit. C,

[55] on black cumin; [56] on tomato and [57] on caraway plants concerning the effect of vit. E. Moreover, [58] on tomato; [43] on syngonium and [59] on snap bean found that spraying plants with vit. B₁ increased N, P and K total uptake by plant.

The interaction between the two main factors (A×B) was significant for N and P % in both seasons as shown in data presented in Tables (8, 9 and 10). The highest N and P % was obtained due to compost (3.75 ton/fed.) in combination with active yeast (5 g/l.) followed by vitamin C and E, each at low concentration.

Similar results were obtained by [60] on dill; [61] and [62] on faba bean and [63] on pea, concerning the effect of interaction treatment between organic manures and active. However, [64] on caraway and [65] on black cumin concluded that the highest contents of NPK were obtained in combination treatment of compost with ascorbic acid.

Protein (%)

Data presented in table (11) showed that the increase in protein % was due to the increase in compost level. All treatments of compost significantly increased protein % in the dry seeds of guar in both seasons in comparison with those of control. Moreover the best treatment which produced the highest values of protein content % was compost at 3.75 ton/fed. The promotive effect of organic manure was found by [24] and [25] on guar; [66] on pigeon pea; [67] on soybean and [48] on roselle plant.

Table 11: Effect of compost, some vitamins and active yeast treatments on protein (%) of *Cyamopsis tetragonoloba* seeds during 2010 and 2011 seasons

First season						
Vitamins and active yeast treatments (B)	Compost levels (ton/fed.) (A)					
	0.0	1.25	2.50	3.75	Mean (B)	
Control	15.1	15.6	15.9	16.1	15.68	
Vit. E at 10 ppm	16.1	16.6	16.9	17.1	16.68	
Vit. E at 20 ppm	15.4	16.0	16.2	16.7	16.08	
Vit. B ₁ at 25 ppm	15.6	16.2	16.3	16.9	16.25	
Vit. B ₁ at 50 ppm	15.3	15.9	15.9	16.5	15.90	
Vit. C at 50 ppm	15.9	16.5	16.6	16.7	16.43	
Vit. C at 100 ppm	15.2	15.7	15.9	16.3	15.78	
Active yeast (2.5 g/l)	15.2	15.8	15.9	16.4	15.83	
Active yeast (5.0 g/l)	16.4	16.9	17.4	17.6	17.08	
Mean (A)	15.58	16.13	16.33	16.70		
L. SD at 5 %	A:	0.35	B:	0.09	AB:	N. S.
Second season						
Vitamins and active yeast treatments (B)	Compost levels (ton/fed.) (A)					
	0.0	1.25	2.50	3.75	Mean (B)	
Control	15.2	15.5	15.8	16.2	15.68	
Vit. E at 10 ppm	16.3	16.8	16.9	17.5	16.88	
Vit. E at 20 ppm	15.4	16.2	16.3	16.7	16.15	
Vit. B ₁ at 25 ppm	15.6	16.2	16.4	16.8	16.25	
Vit. B ₁ at 50 ppm	15.4	16.0	16.2	16.6	16.05	
Vit. C at 50 ppm	15.9	16.5	16.6	17.2	16.55	
Vit. C at 100 ppm	15.2	15.8	15.9	16.4	15.83	
Active yeast (2.5 g/l)	15.1	16.0	16.1	16.4	15.90	
Active yeast (5.0 g/l)	16.7	17.3	17.5	17.9	17.35	
Mean (A)	15.64	16.26	16.41	16.86		
L. SD at 5 %	A:	0.40	B:	0.12	AB:	N. S.

Vit. E. = α -tocopherol Vit. B₁. = Thiamine Vit. C. = Ascorbic acid

Data presented in table (11) indicated that each of the eight treatments were positively and significantly effective on promoting protein % in the dry seeds of guar in the two seasons, over those of control. The best results were obtained due to the use of active yeast (5 g/l.) followed by vit. E at 10 ppm then vit. C at 50 ppm with significant differences between such three treatments in their capability in increasing the seeds percentage of protein. Similar results were obtained by [51] and [39] on *Phaseolus vulgaris*; [42] on *Vicia faba* and [40] on sugar beet, concerning the effect of active yeast. However, [68] on snap beans reported that vit. E had stimulative effect on protein content. Also, [41] on *Vicia faba*; [69] on *Saccharum* spp and [70] on *Pisum sativum* concluded that application of vit. C increased total protein in the seeds.

The interaction between compost, active yeast and vitamins treatments was not significant for protein % during both seasons as shown in table (11).

CONCLUSION

From the obtained results in this work, it could be recommended to supply *Cyamopsis tetragonoloba* Taub. plants with compost at 3.75 ton/fed., and spraying plants with either active yeast at 5 g/l. or ascorbic acid (vit. C) at 50 ppm to improve guaran production., as well as, enhancing the values of some chemical constituents of guar plant under investigation condition.

REFERENCES

1. Thakure, C. (1975): Scientific Crop Production. Metropolitan Book. Co-PVT. Limited I, Nalaji Subhash Marg, Delhi, India.
2. Vinizky, I. and Ray, D. T. (1988): Germination of guar seed under saline and temperature stress. *J. Am. Soc. Hort. Sci.*, 113:437-440.
3. Deraz, S. N. (1993): Biochemical studies on guar seeds (*Cyamopsis tetragonoloba* L.) and their flour incorporation in Balady bread making. *Mounofia J. Agric. Res.*, 18 (1-1): 75-88.
4. El-Etriby, H. M.; Al-Khamy, A. F. and Shahein, N. M. (1994): Evaluation of guar and Leucaena gum preparation in the manufacture of yoghurt. *Annals Agric. Sci.*, Moshtohor, 32 : 1635-1642.
5. Whistler, R. and Hymowitz, T. (1980): Guar: Production, Nutrition and Industrial Use. Purdue University Press, Lafayette, India 47907.
6. Dauda, S. N.; Ajayi, F. A. and Ndor, E. (2008): Growth and yield of watermelon (*Citrullus lanatus*) as affected by poultry manure application. *J. Agric. Soc. Sci.*, 4:121-211.
7. Zheljzkov, V. D. and Warman, P. R. (2004): Source-separated municipal solid waste compost application to Swiss chard and basil. *J. Environmental Quality*, 33:542-552.
8. Fryer, M. J. (1992): The antioxidative effects of thylakoid vitamin E (α -tocopherol). *Plant Cell and Environment*, 15:381-392.
9. Hess, J. L. (1993): Vitamin E, Alpha-tocopherol. In *Antioxidant in Higher Plants*. CRC Press Inc., Boca Raton, Florida, USA.
10. Strove, E. A. (1989): *Biochemistry*. First Published Eng. 1989. Revised from the 1986 Russian Ed. ISBN 5-03-000543-9, 348-355.
11. Hess, D. (1981): *Plant Physiology*. Narosa Publishing House, New Delhi, 1981, 11002, 2nd Ed., 78, 79, 160, 162, 205-208.
12. Samiullah, M. M.; Ansari, S. A. and Afridi, R. K. (1988): B-vitamins in relation to crop productivity. *Ind. Rew. Life Sci.*, 8:51-74.
13. Refaat, A. M. and Balbaa, L. K. (2001): Yield and quality of lemon grass plants (*Cymbopogon flexuosus* Stapf) in relation to foliar application of some vitamins and micro-elements. *Egypt Journal of Horticulture*, 28 : 41-57.
14. Blokhina, O.; Virdainen, E. and Fagersted, K. V. (2003): Antioxidant, oxidative damage and oxygen deprivations stress. *A Review Ann. Boi.*, 91:179-194.
15. Nagodawithana, W. T. (1991): *Yeast Technology*. Universal Foods Corporation Milwaukee, Wisconsin Published by Van Nostrand Reinhold, New York. 273 pp.
16. Anderson, E. (1949): Endosperm mucilage of legumes. Occurrence and composition. *Ind. Eng. Chem.*, 41:2887-2890.
17. Moran, R. (1982): Formula for determination of chlorophylls pigments extracted with N-N dimethyl-formamide. *Plant Physiol.*, 69:1376-1381.
18. Fales, F. W. (1951): The assimilation and degradation of carbohydrates by yeast cells. *J. Bio. Chem.*, 193:113-124.
19. Wilde, S. A.; Covey, R. P.; Lyer, J. C. and Voigt, G. K. (1985): *Soil and Plant Analysis for Tree Culture*. Oxford, IBH. Publishing Co., New Delhi, India.
20. Champan, H. D. and Pratt, P. F. (1975): *Methods of Analysis for Soil, Plant and Water Calif. Univ. Division of Agric. Sci.*, 172-174.
21. Cottenie, A.; Verloo, M.; Velghe, M. and Camerlynck, R. (1982): *Chemical Analysis of Plant and Soil*. Laboratory of Analytical and Agro Chemistry. State Univ., Ghent, Belgium.
22. Ranganna, S. (1978): *Manual of Analysis of Fruit and Vegetative Products*. Tata Mc Graw-Hill publishing company. Limited new Delhi, Second Reprinted, 634 pp.
23. MSTAT-C (1986): *A Microcomputer Program for the Design, Management and Analysis of Agronomic Research Experiments (Version 4.0)*, Michigan Stat Univ., U. S.
24. Gomaa, A. M. and Mohamed, M. H. (2007): Application of bio-organic agriculture and its effect on guar (*Cyamopsis tetragonoloba* L.) root nodules, forage, seed yield and yield quality. *World J. of Agric. Sci.*, 3 : 91-96.
25. Patel, C. S.; Patel, J. B.; Suthar, J. V. and Patel, P. M. (2010): Effect of integrated nutrient management on clusterbean [*Cyamopsis tetragonoloba* L. Taub.] seed production cv. Pusa navbahr. *Inter. J. of Agric. Sci.*, January to June, b : 206-208.
26. Abdalla, M. Y. A. (2009): Effect of organic, bio and mineral fertilization on growth, yield, oil productivity and chemical constituents of coriander plant. *J. Agric. Sci. Mansoura Univ.*, 34 : 5195-5208.
27. Fernandez, L. F.; Reyes, V. V.; Martinez, S. C.; Salomon, H. G.; Yanez, M. J.; Ceballos, R. J. and Dendooven, L. (2010): Effect of different nitrogen sources on plant characteristics and yield of common bean (*Phaseolus vulgaris* L.). *Bioresource Technology*, 101:396-403.
28. Ngetich, O. K.; Aguyoh, J. N. and Ogwenyo, J. O. (2012): Effects of composted farmyard manure on growth and yield of spider plant (*Cleome gynandra*). *Int. J. of Sci. and Nature*, 3 : 514-520.

29. Aly, M. K.; Mohamed, M. A. H.; Attia, F. A.; El-Sayed, A. A. and Abd El-Gawad, M. H. I. (2007): Effect of some organic and biofertilization treatments on coriander plants. 2-Chemical composition. Proc. 1st Inter. Conf. Desert. Cultivation, Problem and Solutions, Minia Univ., 27–29 March 2007.
30. Ahmed, E. F. A. (2007): Evaluation of certain fertilizing programs on anise and black cumin plants. Ph D. Thesis. Fac. Assiut Univ. Egypt.
31. Abdou, M. A. H.; Abdalla. M. Y.; Helaly, A. and Mosaad, S. (2013b): Physiological studies on sage plant. Minia J. of Agric. Res. and Develop, 32 : 161-186.
32. Abd-El-Naeem, L. M. A. (2008): Response of caraway plants to some organic and biofertilization treatments. M. Sc. Thesis, Fac. Agric., Minia Univ., Egypt.
33. Tanious, C. T. S. (2008): Effect of some organic fertilization treatments on fennel plants. M. Sc. Thesis, Fac. Agric., Minia Univ., Egypt.
34. Abdou, M. A. H.; Aly, M. K.; Attia, F. A.; Ahmed, E. T. and Al-Shareif, A. M. O. (2013a): Physiological studies on black cumin plant. The 1st Under Press Inter of Hort. Agric. Assuit, February 2013.
35. El-Sherbeny, S. E.; Khalil, M. Y. and Naguib, N. Y. (2005): Influence of compost levels and suitable spacings on the productivity of *Sideritis monata* L. plants recently cultivated under Egyptian conditions. Bulletin of Fac. Agric., Cairo Univ., 56 : 373-392.
36. El-Leithy, A. S.; El-Hanafy, S. H.; Ahmed, G. F.; Etman, M. A. and Ali, M. Y. (2007): Effect of compost, bio-fertilizer and active dry yeast on *Origanum syriacum* L. plants under Sinai conditions. J. Product and Dev., 12 : 153-171.
37. Abdel-Kader, A. A.; El-Keltawi, N.; Tawfik, A. A. and Khater, M. R. (2008): Growth, yield and quality of henna (*Lawsonia alba* Lam.) plant as affected by soil type and application of active dry yeast. Assiut J. of Agric. Sci., 39 : 143-165
38. Mady, M. A. (2009b): Effect of foliar application with yeast extract and zinc on fruit setting and yield of faba bean (*Vicia faba* L.). J. Biol. Chem. Environ. Sci., 4 (20):109-127.
39. Nasser, R. M.; Ahmed, Y. M. and Nassar, D. M. (2011): Effect of foliar spray with active yeast extract on morphological, anatomical and yield characteristics of kidney bean (*Phaseolus vulgaris* L.) Australian J. of Basic and Applied Sci., 5: 1071-1079.
40. Agamy, R.; Hashem, M. and Al-Amri, S. (2013): Effect of soil amendment with yeast as biofertilizers on the growth and productivity of sugar beet. African J. of Agric. Res., 8 : 46-56.
41. El-Bassiouny, H. M. S.; Gobarah, M. E and Ramadan (2005): Effect of antioxidants on growth, yield, savism causative agents in seeds of *Vicia faba* L. plants grown under reclaimed sandy soils. J. Agric. Pakistan.
42. Abou El-Yazied, A. and Mady, M. A. (2012): Effect of boron and yeast extract foliar application on growth, pod setting and both green pod and seed yield of broad bean (*Vicia faba* L.). J. of American Sci., 8 : 517-533
43. Abd El-Aziz, N. G.; El-Quesni, F. E. and Farahat, M. M. (2007b): Response of vegetative growth and some chemical constituents of *Syngonium podophyllum* L. to foliar application of thiamine, ascorbic acid and kinetin at Nubaria. World J. of Agric. Sci., 3 : 301-305.
44. Ngoc Son, T.; Van Tnu, V.; Hong Man, L. and Hiraoka, H. (2001): Effect of organic and biofertilizer on quality, grain yield and soil properties of soybean under rice based cropping system. Japan Inter. Center for Agric. Sci., 9:55-61.
45. Solieman, T. H.; El-Khatib, H. A. and El-Araby, S. M. (2003): Effects of organic manure, mineral nitrogen and bio-fertilizer application on vegetative growth and chemical composition of pea (*Pisum sativum* L.). Zagazig J. Agric. Res., 30 : 751-767.
46. Abd El-Aziz, M.; Pokluda, R. and Abdel Wahab, M. (2007a): Influence of compost, microorganisms and NPK fertilizer upon growth, chemical composition and essential oil production of *Rosmarinus officinalis* L. Not. Bot. Hort. Agrobot. Cluj., 35 : 86-90.
47. Priya, S. and Elakkiya, R. (2012): Effect of organic and biofertilizers on growth and yield of *Eclipta alba* L. Int. J. of Pharm Tech Research, 4: 1703-1705.
48. Gendy, A. S. H.; Said-Al-Ahl and Mahmoud, A. (2012): Growth, productivity and constituents of roselle (*Hibiscus sabdariffa* L.) plants as influenced by cattle manure and biofertilizers treatments. Australian Journal of Basic and Applied Science, 6 : 1-12.
49. El-Sayed, A. A.; Ali, M. K. and Abd El-Gawad (2002): Response of coriander plants to some phosphorus, zink and active yeast treatments Proc. 2nd Inter. Conf. Hort. Sci., Kafr El-Sheikh, Tanta Univ., Egypt, Sept. 10-12:434-446.
50. Naguib, N. Y. (2002): Yield and quality of lemongrass plants (*Cymbopogon flexuous* Stapf) as influenced by farmyard manure and foliar application of bread yeast. Annals of Agric., Sci., Ain Shams Univ., Cairo. 47 : 859–873.
51. El-Tohamy, W. A. and El-Greadly, N. H. M. (2007): Physiological response, growth, yield and quality of snap beans in response to foliar application of yeast, vitamin E and zinc under sandy soil conditions. Australian Journal of Basic and Applied Sci., 1 294-299.
52. Abdel-Hakim, W. M.; Moustafa, Y. M. M. and Gheeth, R. H. (2012): Foliar application of some chemical treatments and planting date affecting snap bean (*Phaseolus vulgaris* L.) plants grown in Egypt. J. of Hort. Sci. and Ornamental Plants, 4 : 307-317.
53. Talaat, N. B. (2003): Physiological studies on the effect of salinity, ascorbic acid and putrescine on sweet pepper plant. Ph D. Thesis, Fac, Agric., Cairo Univ., Egypt
54. El-Tohamy, W. A.; El-Abagy, H. M. and El-Greadly, N. H. M. (2008): Studies on the effect of putrescine, yeast and vitamin C on growth, yield and physiological responses of eggplant (*Solanum melongena* L.) under sandy soil conditions. Australian J. of Basic and Applied Sciences, 2 : 296-300.
55. Ismail, S. I. I. (2008): Anatomical and physiological studies on *Nigella sativa*, L. plant. Ph. D. Dess. Fac. Agric Mansoura Univ., Egypt
56. Mady, M. A. (2009a): Effect of foliar application with salicylic acid and vitamin E on growth and productivity of tomato (*Lycopersicon esculentum* Mill) plant. J. Agric. Sci. Mansoura Univ., 34 : 6735-6746.
57. Botros, W. S. E. (2013): Physiological studies on caraway plants. M. Sc. Thesis, Fac. Agric., Minia Univ. Egypt.
58. El-Mansi, A. A. Bardisi, H. and El-Robae, M. M. (2004): Comparative study on the effect of yeast, some vitamins, IBA and boron on growth and yield of tomato under sandy soil conditions. Zagazig J. Agric. Res., 31 (4A): 1409-1424.
59. Mohammad, O.; Bardisi, A. and El-Bassiouny, R. (2009): Effect of foliar spray with Ca, Mg and vitamin

- B1 on the productivity and storability of snap bean grown in sandy soil. Zagazig J. Agric. Res., 36 : 41-62.
60. Menesi, F. A.; El-Mahrouk, E. M.; Sidky, M. A. and Rashed, N. M. (2002): Effect of bio-organic and chemical fertilization on dill, coriander and parsley, J. Agric. Sci., Mansoura Univ., 27 : 8337-8345.
61. Mohamed, H. A. and Gomaa, A. M. (2005): Faba bean growth and green yield and its quality as influenced by the application of bio-organic farming system. Journal of Applied Sciences Research, 1 : 350-385.
62. Gomaa, A. M.; Afifi, M. H. M.; Mohamed, M. F. and El-Dewiny, C. Y. (2010): Nodulation, growth characters and yield quality of faba bean cultivated in newly reclaimed sandy soil under bio-organic agriculture system. Inter. J. Academic Research, 2: 134-139.
63. Mahmoud, R.; El-Desuki, M.; Abdel-Mouty, M. and Ali, A. (2013): Effect of compost levels and yeast extract application on the pea plant growth, pod yield and quality. Journal of Applied Sciences Research, 9 : 149-155.
64. Abdou, M. A. H.; Attia, F. A.; Ahmed, E. T. and Abd El-Naeem, L. (2009): Response of caraway plants to some organic, antioxidants and biofertilization treatments. Proc. 4th Inter. Environ. Conf., Mansoura Univ. on Environmental and Healthy Safety, 28-29 October, 2009, p: 109-119.
65. Al-Shareif, A. M. O. (2012): Physiological studies on black cumin plant. Ph. D. Thesis. Fac. of Agric. Minia Univ. Egypt.
66. Shafeek, M. R.; El-Zeiny, O. A. and Ahmed, M. E. (2004): Pigeon pea (*Cajanus cajan*) growth and yield as affected by the interaction between bio and organic fertilizers. Annals of Agricultural Science (Cairo), 49 : 619-630.
67. Maheshbabu, H. M.; Hunje, R.; Patil, N. K. B. and Babalad, H. B. (2008): Effect of organic manures on plant growth, seed yield and quality of soybean. Karnataka J. Agric. Sci., 21 : 219-221.
68. Nour, K. A. M.; Mansour, N. T. S. and Eisa, G. S. A. (2012): Effect of some antioxidants on some physiological and anatomical characters of snap bean plants under sandy soil conditions. New York Sci. J., 5 : 1-9
69. Batool, E.; Ahmed, S. and Faheem, A. (2012): Effect of exogenous application of ascorbic acid on antioxidant enzyme activities, proline contents and growth characters of *Saccharum* spp. hybrid cv. HSF-240 under salt stress. Turk J. Biol., 36 (2012): 630-640.
70. Gheeth, R. H. M.; Moustafa, Y. M. M and Abdel-Hakeem, W. H. (2013): Enhancing growth and increasing yield of peas (*Pisum sativum*, L.) by foliar application of ascorbic acid and cobalt chloride. J. Novel Applied Sciences, 2 : 106-115.