

## SHORT COMMUNICATION

# Effectiveness of some integrated disease management factors (IDM) on *Fusarium* wilt infected tomato

Hussein Ali Salim<sup>1\*</sup>, Sobita Simon<sup>1</sup>, Abhilasha A. Lal<sup>1</sup>, Abbas Lateef Abdulrahman<sup>2,3</sup>

<sup>1</sup>Department of Plant Pathology, Sam Higginbottom Institute of Agriculture, Technology and Sciences, (SHIATS), Allahabad, U.P. India.

<sup>2</sup>Directorate of Agricultural Extension and Training, Ministry of Agriculture, Iraq.

<sup>3</sup>Faculty of Agriculture, University Putra Malaysia, 43400 UPM, Serdang, Selangor, Malaysia.

## Abstract

The study was conducted to investigate efficacy of Integrated disease management (IDM) viz., solarized soil, *Trichoderma harzianum*, spent mushroom compost and carbendazim on growth characteristics of infected tomato by Fusarial wilt (*Fusarium oxysporum* f.sp. lycopersici) under pots conditions in the Bio-net house during 2013-2014. Seven treatments and five replicates were taken up in completed randomized design. Maximum shoot length, fresh and dry shoot weight were noticed in T4 (solarized soil+ Spent mushroom compost+ *T. harzianum*) 118.10 cm, 67.25 and 42.20 g respectively as compared to *T. harzianum* or spent mushroom compost that alone treated plants. Maximum root length, fresh and dry root weight were also recorded in T4 (solarized soil+ spent mushroom compost+ *T. harzianum*) 28.35 cm, 4.65 and 2.55 g respectively. The yield of tomato plants was significantly increased in T6 (tomato plant alone without *Fusarium oxysporum*) 170.15 g/plant followed by T4 137.52 g/plant as compared with T0- *Fusarium oxysporum* alone and T1- Solarized soil + *Fusarium oxysporum* 0.0 g.

*Key words:* *Fusarium oxysporum*, *Trichoderma harzianum*, spent mushroom, carbendazim

## Introduction

Tomato (*Lycopersicon esculentum* Mill.) is one among the most used vegetable in family solanaceae and is cultivated in many areas throughout the world (Villareal, 1979). There are many diseases for this plant, and *Fusarium* wilt is major among them causing 10-50 % yield loss (Larkin and Fravel, 1998; Borrero et al., 2004). It is of worldwide importance where at least 32 countries had reported the disease, which is particularly severe in countries with warm climate. Mui-Yun Wong, (2003). It is a devastating disease causing considerable

economic losses ranging from 10-80% yield loss in many tomatoes producing area of the country (Kesavan and Chaudhary, 1977). Integrated disease management (IDM), is an important strategy to control this deadly disease (El Khoury and Makkouk, 2010).

The genus, *Trichoderma* (Deuteromycota, Dematiaceae), is a common saprophytic fungus in the rhizosphere soil. This has some mycoparasite ability against plant pathogens (Papavizas, 1985; Elad et al., 1993; Elad, 2000; Freeman et al., 2004; Dubey et al., 2007) and

Received 29 July 2017; Accepted 24 August 2017; Published 21 September 2017

\*Corresponding Authors

Hussein Ali Salim

Department of Plant Pathology, Sam Higginbottom Institute of Agriculture, Technology and Sciences, (SHIATS), Allahabad, U.P. India

Email: [H\\_salim11111@yahoo.com](mailto:H_salim11111@yahoo.com)

©This article is open access and licensed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>) which permits unrestricted, use, distribution and reproduction in any medium, or format for any purpose, even commercially provided the work is properly cited. Attribution – You must give appropriate credit, provide a link to the license, and indicate if changes were made.

nematodes (Windham et al., 1989; Sharon et al., 2001). This fungus can be used as a biocontrol measure for many diseases (Sivan and Chet, 1986; Calvet et al., 1990).

Similarly, mushroom compost (spent mushroom substrate (SMS), mushroom soil) also has some antimicrobial effects and can be used against plant diseases caused by fungi (Davis et al. 2005). This study was conducted to evaluate some IDM elements: Solarized soil, *Trichoderma harzianum*, spent mushroom compost and carbendazim on some growth parameters of infected tomato by *Fusarium wilt*.

## Materials and Methods

### Study area

The present study was carried out under pots conditions in the Bio-net house, Department of Plant Pathology, Sam Higginbottom Institute of Agriculture, Technology & Sciences, Naini, Allahabad during 2013-2014 with seven treatments *Fusarium oxysporum* alone, Solarized soil + *F. oxysporum*, Solarized soil + spent mushroom compost + *F. oxysporum*, Solarized soil + *Trichoderma harzianum* + *F. oxysporum*, Solarized soil+ spent mushroom compost + *Trichoderma harzianum* + *F. oxysporum*, Carbendazim + *F. oxysporum* and Tomato plant alone (without *F. oxysporum*) with five replicates. Tomato cultivar seeds CO - 3 were obtained from IIVR (Indian Institute of Vegetable Research), Varanasi, Uttar Pradesh, India. *Trichoderma harzianum* were obtained as biocont from Yash Trichoguard, DBT Referral Lab, SHIATS, Allahabad, Uttar Pradesh, India. Spent mushroom compost acquired from Department of Plant Pathology, Sam Higginbottom Institute of Agriculture, Technology & Sciences, Uttar Pradesh, India. *Fusarium oxysporum* f.sp. *lycopersici* was isolated from infected roots of tomato by tissue segment method. The infected roots of tomato were cut separately into small pieces into the wash glass with the help of sterile blade and surface sterilized with mercuric chloride (0.1%) for 20-30 seconds followed by three times rinsing with sterilized distilled water. The diseased pieces were then placed on pre-sterilized blotting paper to remove excess moisture. The surface sterilized diseased pieces were then aseptically, transferred on PDA medium and incubated at 28±2°C for five days.

After incubation, colonies were observed and identified on the basis of morphological and reproductive characters.

Pathogenicity of *Fusarium oxysporum* f.sp. *lycopersici* was tested on healthy tomato seedling at the three-true-leaf stage according to Koch's postulate, their roots were cut and dipped into conidial suspension (10<sup>6</sup> spores/ml) of tested isolate for 30 minute, and then the seedlings were transplanted into sterilized soils in pots (10 cm in diameter) and kept in Bio-net. After symptoms appeared the pathogen was re-isolated from infected portion of tomato seedling and again identified through microscopic method in the laboratory.

Farm yard manure (FYM) @ 100 g / pot was mixed with solarized and unsolarized soil and was filled in thirty-five pots with capacity 10 kg soil / pot. The bioagent *Trichoderma harzianum* was applied @ 2 g / pot, four days before sowing the seeds of tomato variety (CO-3), carbendazim 50 % W.P was applied @ 2 kg a.i / ha, whereas spent mushroom compost was unsterile and mixed with soil in the pots @ 20 g / kg of soil. Seven days after germination, four seedlings per pot were maintained in each treatment. Observations were recorded on shoot length (cm), fresh and dry shoot weight (g), root length (cm), fresh and dry root weight (g) and yield / plant (g). In the experiment Complete Randomized Design (CRD) was adopted. Soil solarization was conducted for 2 months from 15th April to 15th June 2013 at research field of SHIATS, Allahabad, The soil was tilled, leveled and Irrigation was given prior to laying of the polythene sheet which was transparent and thick.

The analysis of variance (ANOVA) technique was applied for drawing conclusion from data. The calculated values were compared the tabulated values at 5% level of probability (Fisher and Yates, 1968).

## Results and discussion

### Shoot length (cm)

The results presented in Table 1 revealed that significant increase of shoot length was found in T<sub>6</sub> (Tomato plant alone) and T<sub>4</sub> (solarized soil + Spent mushroom compost + *Trichoderma harzianum*+ *F.o*) (124.81 and 118.10 cm respectively) as compared with other treatments, However, (T<sub>6</sub>, T<sub>4</sub>), (T<sub>3</sub>, T<sub>4</sub>), (T<sub>3</sub>, T<sub>5</sub>) and (T<sub>1</sub>, T<sub>0</sub>), were found non-significant from each other.

Table 1. Effect of solarized soil with spent mushroom compost, *Trichoderma harzianum* and carbendazim on plant growth parameters of tomato.

Treatment	Shoot length (cm)	Fresh Shoot Weight (g)	Dry Shoot Weight (g)	Root Length (cm.)	Fresh Root Weight (g)	Dry Root Weight (g)	Yield / plant (g)
To	50.65	7.50	5.05	10.35	0.60	0.39	0.00
T1	55.10	8.65	5.60	11.55	0.79	0.52	0.00
T2	81.25	40.70	21.95	23.85	4.35	1.90	128.25
T3	103.45	53.45	32.80	28.25	4.50	1.60	107.12
T4	118.10	67.25	42.20	28.35	4.65	2.55	137.52
T5	100.55	50.40	30.45	20.00	4.40	1.85	115.77
T6	124.81	91.35	61.60	28.65	6.20	3.30	170.15
C. D. 0.05	16.105	23.326	15.308	5.724	1.422	0.865	17.546

T<sub>0</sub>- *Fusarium oxysporum* alone

T<sub>1</sub>- Solarized soil + *F. oxysporum*

T<sub>2</sub> - Solarized soil + spent mushroom compost + *F. oxysporum*

T<sub>3</sub>- Solarized soil + *Trichoderma harzianum* + *F. oxysporum*

T<sub>4</sub>-Solarized soil+ spent mushroom compost + *Trichoderma harzianum* + *F. oxysporum*

T<sub>5</sub>- Carbendazim + *F. oxysporum*

T<sub>6</sub>- Tomato plant alone (without *F. oxysporum*)

### Root length (cm)

The results presented in Table 1 revealed that T<sub>6</sub> (Tomato plant alone, 28.65 cm), T<sub>4</sub> (solarized soil + Spent mushroom compost + *Trichoderma harzianum*+ *F.o*, 28.35 cm), T<sub>3</sub> (solarized soil + *Trichoderma harzianum*+ *F.o* ,28.25 cm) and T<sub>2</sub> (solarized soil + Spent mushroom compost+ *F.o*, 23.85 g) significantly increased in root length from T<sub>5</sub> (carbendazim+ *F.o*, 20.00 cm), T<sub>1</sub> (solarized soil + *F.o* ,11.55 cm) and T<sub>0</sub> (*F.o* alone,10.35 cm). Among the treatments (T<sub>0</sub>, T<sub>1</sub>), (T<sub>6</sub>, T<sub>4</sub>, T<sub>3</sub>, T<sub>2</sub>), (T<sub>5</sub>, T<sub>2</sub>) were not significant from each other.

### Fresh and dry shoot weight (g)

The results presented in Table 1 revealed that treatments T<sub>6</sub>, T<sub>4</sub>, T<sub>3</sub>, T<sub>5</sub> and T<sub>2</sub> significantly increased in fresh and dry shoot weight from (T<sub>1</sub>, T<sub>0</sub>), However (T<sub>4</sub>, T<sub>3</sub>, T<sub>5</sub>), (T<sub>4</sub>, T<sub>5</sub>, T<sub>2</sub>) and (T<sub>1</sub>, T<sub>0</sub>), were found non-significant from each other.

### Root fresh weight (g)

The results presented in Table 1 revealed that significantly increased the fresh root weight in treatments T<sub>6</sub> (6.20 g), T<sub>4</sub> (4.65 g), T<sub>3</sub>

Najat and Kakhkashan (2012) reported that under pots culture 117.5 % increase in plant height can be achieved for tomato plants treated with *T. harzianum*. Similar results were reported by Sundaramoorthy and Balabaskar (2013). Rasool et al. (2011) reported that *Trichoderma* species can be used as biocontrol agents against phytopathogenic fungi and increase fresh shoot and root weight

(4.50 g), T<sub>5</sub> (4.40 g) and T<sub>2</sub> (4.35 g) from T<sub>1</sub> (0.79 g) and T<sub>0</sub> (0.60 g).

### Root dry weight (g)

The results presented in Table 1 revealed that significantly increased the dry root weight in the treatments T<sub>6</sub>, T<sub>4</sub>, T<sub>2</sub>, T<sub>5</sub> and T<sub>3</sub> as compared with T<sub>1</sub> and T<sub>0</sub>. However, the treatments (T<sub>6</sub>, T<sub>4</sub>), (T<sub>4</sub>, T<sub>2</sub>, T<sub>5</sub>), (T<sub>2</sub>, T<sub>5</sub>, T<sub>3</sub>) and (T<sub>1</sub>, T<sub>0</sub>) were found non-significant from among themselves.

### Yield / plant (g)

Fruit yield (g) per plant of tomato were taken after 150 days of germination in the Table 1 revealed that significantly increased the yield (g) of tomato plants was found in T<sub>6</sub> (170.15 g, plant without fusarium and treatments), T<sub>4</sub> (137.52 g ,S.S + S.m.c + *T.h* +*F.o*) , T<sub>2</sub>( 128.25 g, S.S + S.m.c + *F.o*), T<sub>5</sub> (115.70 g ,C +*F.o*) and T<sub>3</sub>( 107.12 g, S.S + *T.h* +*F.o*), as compared with the treatments T<sub>1</sub>(0.0 g, S.S + *F.o* ) and T<sub>0</sub> (0.0 g, *F.o* alone), however the treatments (T<sub>4</sub>, T<sub>2</sub>) and (T<sub>2</sub>,T<sub>3</sub>,T<sub>5</sub>) and (T<sub>1</sub>,T<sub>0</sub> ) were found non-significant among themselves.

in tomato seedlings. Increases in dry root weight in tomato seedlings were observed in treated soil with *T. harzianum* T969 as compared with the control, dry root weight also increased in seed that inoculated by *Trichoderma* isolates. Ebtsam et al. (2009) observed that control treatment was recorded low values of dry weight of plants in comparison with other treatments. Ebtsam et

al. (2009); Niknejad et al. (2000) and Zaghoul et al. (2007) showed the treatment of *B. subtilis*, *T. harzianum* enhanced the number, weight and yield of fruits of tomato. The probable reasons for such findings may be that *Trichoderma harzianum* have many positive effects on plant and systemic resistance to plant diseases (Harman et al., 2004; Harman, 2006). The growth enhancement might be due to secretion of auxins, gibberellins and cytokinins (Sundaramoorthy and Balabaskar, 2013).

### Conclusion

From the findings, it's concluded that effectiveness of some integrated disease management factors (IDM) on some characteristics of growth of infected tomato by fusarium wilt using *T. harzianum*, spent mushroom compost, chemical method (Carbendazim) and physical method (Soil solarization). Plants treated with a combination of *T. harzianum* and spent mushroom compost with solarized soil showed significant increasing in shoot length (cm), fresh and dry shoot weight (g), root length (cm), fresh and dry root weight (g) and yield / plant (g).

### Acknowledgements

Authors wish to express their sincere gratitude and appreciation to the Department of Plant Pathology, Sam Higginbottom Institute of Agriculture, Technology & Sciences for awarding the opportunity to accomplish this work

### Authors' contributions

SS and AAL planned and coordinated the research, HAS participated in, set up and carried out the field experiment, collected data, analysis and data interpretation, and wrote the article. ALA reviewed and submitted article to Journal of Scientific Agriculture. All the authors revised manuscript.

### References

Borrero C, Trillas MI, Ordales J, Tello JC, Aviles M. (2004). Predictive factors for the suppression of Fusarium wilt of tomato in plant growth media. *Phytopathology*. vol 94, pp 1094 – 1101.

Calvet C, Pera J, Bera JM (1990). Interaction of *Trichoderma* spp with *Glomus mossaeae* and two wilt pathogenic fungi. *Agric. Ecosyst. Environ.* vol 9, pp 59-65.

Davis D. Donald, Larry J. Kuhns, and Tracey L. Harpster. (2005). Use of mushroom compost to suppress artillery fungi. *J. Environ. Hort.* vol 23, (4), pp 212–215.

Dubey RC, Kumar H, Pandey RR (2009). Fungitoxic effect of Neem extracts on growth and sclerotial survival of *Macrophomina phaseolina in vitro*. *J. Am. Sci.* vol 5, pp 17-24.

Ebtsam M. Morsy; K.A. Abdel-Kawi and M.N.A. Khalil, (2009) Efficiency of *Trichoderma viride* and *Bacillus subtilis* as Biocontrol Agents gainst *Fusarium solani* on Tomato Plants, Egypt. *J. Phytopathol.*, Vol. 37, No. 1, pp. 47-57 (2009)

El Khoury W. and K. Makkouk, (2010). Integrated plant disease management in developing countries, *Journal of Plant Pathology*. vol 92, (4), pp 35- 42.

Elad Y (2000). Biological control of foliar pathogens by means of *Trichoderma harzianum* and potential modes of action. *Crop Prot.* vol 19, pp709-714.

Elad Y, Zimmand G, Zags Y, Zuriel S, Chet I (1993). Use of *Trichoderma harzianum* in combination or alternation with fungicides to control Cucumber grey mold (*botrytis cinerea*) under commercial greenhouse condition. *Plant Pathology*. vol 42, pp 324-356.

Fisher, R. A. and Yates, (1968). Statistical method for research workers. Oliver and boyd ltd. Edinburgh and London, 10.

Freeman Gajalakshmi S, Abbasi SA (2004). Neem leaves as a source of fertilizer-cum-pesticide vermicompost. *Bioresource Technol.* vol 92, pp 291-296.

Harman GE (2006). Overview of mechanisms and uses of *Trichoderma* spp. *Phytopathology*, vol 96, pp 190-194.

Harman GE, Lorito M, Lynch JM (2004). Uses of *Trichoderma* spp. to alleviate or remediate soil and water pollution. *Adv. Appl. Microbiol.* vol. 56, pp313-330.

Kesavan, V. and B. Chaudhary. (1977). Screening for resistance to *Fusarium* wilt of tomato. *SABRO J.*, vol 9, pp 51-65.

Larkin RP, Fravel DR. (1998). Efficacy of various fungal and bacterial biocontrol organisms for the control of Fusarium wilt of tomatoes. *Plant Disease*. vol 82, pp 1022 – 1028.

Mui-Yun Wong. (2003). Soilborne Plant Pathogen Class Project, pp 728.

- Najat A. Bokhari and Kahkashan Perveen. (2012) Antagonistic action of *Trichoderma harzianum* and *Trichoderma viride* against *Fusarium solani* causing root rot of tomato, African Journal of Microbiology Research Vol. 6(44), pp. 7193-7197.
- Niknejad, M.; Sharfi-Tehani, A. and Okhovat, M. (2000). Effect of antagonistic fungi *Trichoderma spp.* on the control of *Fusarium* wilt of tomato caused *Fusarium oxysporum f. sp. lycopersici* under greenhouse conditions. Iranian Agric. Sci., 1: 31 - 37.
- Papavizas GC (1985). *Trichoderma* and *Gliocladium* biology, ecology and the potential for biocontrol. Annual Review Phytopathology. vol 23, pp 23-77.
- Rasool Azarmi, Behzad Hajieghrari and Abolfazl Giglou, (2011). Effect of *Trichoderma* isolates on tomato seedling growth response and nutrient uptake African Journal of Biotechnology Vol. 10(31), pp. 5850-5855, 29 June, 2011
- Sharon E, Bar-Elad M, Chet I, Herrera-Estrella A, Kleifeld O, Spiegel Y (2001). Biological control of root knot nematode *Meloidogyne javanica* by *Trichoderma harzianum*. Phytopathology, vol 91, (7), pp 687-693.
- Sivan A, Chet I (1986). Biological control of *Fusarium* spp in cotton, Wheat and muskmelon by *Trichoderma harzianum*. J. Phytopathol. vol 116, pp 39-47.
- Sundaramoorthy S. and P. Balabaskar. (2013) Biocontrol efficacy of *Trichoderma spp.* against wilt of tomato caused by *Fusarium oxysporum f. sp. Lycopersici*. Journal of Applied Biology & Biotechnology Vol. 1 (03), pp. 036-040.
- Villareal, R. L. (1979). Tomato is the tropical. Westview Press, Boulder, Colorado. pp 174.
- Windham GL, Windham MT, Williams WP (1989). Effects of *Trichoderma spp* on Maize growth and *Meloidogyne arenaria* reproduction. Plant Disease. vol 73, pp 493-494.
- Zaghloul, R.A.; Hanafy, Ehsan A.; Neweigy, N. A. and Khalifa, Neamat A. (2007). Application of biofertilization and biological control for tomato production. 12<sup>th</sup> Conference of Microbiology; Cairo, Egypt, (18-22), 198-212.