

Journal of Spices and Aromatic Crops
Vol. 16 (2) : 118–121 (2007)


Indian Society for Spices

Seed germination studies in *Garcinia* spp.

Ancy Joseph¹, K N Satheeshan & T G Jomy

Department of Horticulture
College of Agriculture, Kerala Agricultural University
Padannakad-671 328, Kerala, India.
E mail: ancy24@rediffmail.com

Received 24 January 2007; Revised 13 August 2007; Accepted 10 September 2007

Abstract

Germination studies conducted at Padannakad (Kerala) on *Garcinia gummi-gutta* and *Garcinia cowa* indicated that soaking seeds in hydrogen peroxide (30%) for 30 min and overnight soaking in cow milk and kinetin (500 ppm) were effective in breaking the dormancy of seeds and resulted in 60.05%, 39.96% and 34.41% germination, respectively in *G. gummi-gutta* and 65.33%, 69.33% and 54.67% germination in *G. cowa* at 7 months after sowing. In *Garcinia tinctoria*, soaking in hydrogen peroxide (30%) for 30 min and overnight soaking in kinetin (500 ppm) and gibberellic acid (500 ppm) induced early germination resulting in 78.6%, 93.3%, and 64.0% germination respectively, at 2 months after sowing.

Key words: *Garcinia* species, seed dormancy, seed germination.

The genus *Garcinia* (Clusiaceae) contains several important economic species such as *G. mangostana* L. (mangosteen), *G. gummi-gutta* (L.) Rob. (camboge) and *G. indica* Choisy (kokum). Thirty one species of *Garcinia* occur in India (Maheshwari 1964), but most of these are under exploited. Most of these tropical fruit species show dormancy, *G. mangostana* being an exception (Normah *et al.* 1997). Seed dormancy either induced or innate, imparts seeds with capacity to germinate at favourable environmental conditions (Vazquez-Yanes & Orozco-Segovia 1993). But prolonged dormancy in many *Garcinia* spp. is a limiting factor for large scale production of planting material. Though *Garcinia* spp. can be propagated by seeds, androdioeciousness (George *et al.* 1992) and long pre-bearing age necessitates vegetative

method of propagation. Softwood grafting technique has been standardized to overcome the dioecious nature of *Garcinia* spp. (KAU 2002). For grafting, either its own seedlings or that of closely related species like *G. cowa* Roxb. or *G. tinctoria* Wight can be used as root stocks. Prolonged seed dormancy of *Garcinia* spp. causes delay in rootstock production and timely grafting. In case of *G. gummi-gutta*, seeds have a dormancy period of 8 to 9 months and on removal of seed coat, germination occurs within 2 months of sowing (KAU 2001). But it is laborious and time consuming when done on a commercial scale. Hence, a study was undertaken to find out simple techniques for breaking the seed dormancy of *Garcinia* spp.

The experiment was conducted at College of Agriculture, Padannakad (Kerala), during

¹ Present address: Aromatic and Medicinal Plants Research Station, Asamanoor-683 549, Ernakulam, Kerala, India.

2000–02 in a completely randomized design with three replications. Seeds of *G. gummi-gutta*, *G. cowa* and *G. tinctoria* were collected from 20 year old trees from District Agricultural Farm, Taliparamba (Kannur District, Kerala). The treatments (Table 1) were determined after conducting preliminary studies. For *G. tinctoria*, all the treatments in Table 1 except T₉ were given. In each treatment, 40 seeds were used per replication. Observations on germination were recorded at weekly interval in all the three species.

Table 1. Treatments imposed for germination studies in *G. cambogia* and *G. cowa* seeds

Treatment	Treatment	Duration
T ₁	Control (soaking in water)	18 h
T ₂	Sulphuric acid (H ₂ SO ₄) 20%	15 min
T ₃	Hot water (55°C)	15 min
T ₄	Potassium nitrate (KNO ₃) 0.2%	18 h
T ₅	Thiourea 2%	18 h
T ₆	Hydrogen peroxide (H ₂ O ₂) 30%	30 min
T ₇	Gibberellic acid (GA) (500 ppm)	18 h
T ₈	Kinetin (500 ppm)	18 h
T ₉	Cow milk (fresh)	18 h

The data revealed that at 4 months after sowing (MAS), seed treatment with H₂O₂ (T₆) was highly effective in inducing early germination in *G. gummi-gutta* (Table 2). No germination was noticed in rest of the treatments, within this period. At 5 MAS, treatments with kinetin (T₈), cow milk (T₉) and H₂O₂ (T₆) were significantly superior to control and other treatments and T₆ was the most effective among all treatments. At 6 MAS, treatments with H₂O₂ (T₆), GA (T₇), kinetin (T₈) and cow milk (T₉) were highly effective in enhancing seed germination. Hydrogen peroxide treatment was significantly superior and at 8 MAS 71.04% germination was recorded, whereas germination just started in control at 8 MAS (2.22%).

In *G. cowa*, germination was earliest in H₂O₂ (T₆) and cow milk treated (T₉) seeds at 3 MAS

(Table 3) and their superiority was retained in subsequent stages also. Seed treatment with kinetin (T₈) was also highly effective. Gibberellic acid did not give any desirable effect in *G. cowa* even though it was effective in *G. gummi-gutta*. At 10 MAS, H₂O₂ and cow milk treatments gave 76% germination compared to 36% in control. After 10 months, no significant difference was observed between the treatments and control.

In *G. tinctoria*, soaking in KNO₃ (T₄) and H₂O₂ (T₆) resulted in early germination (17% germination at 1 MAS). At 1½ MAS, KNO₃ (T₄), kinetin (T₈) and H₂O₂ (T₆) gave higher germination compared to other treatments (Table 4). Thiourea was ineffective in breaking dormancy at all stages. At 2 MAS, H₂O₂, gibberellic acid and kinetin gave more than 60% germination against 1.3% in control. But at 4 MAS, H₂SO₄ (T₂), H₂O₂ (T₆), GA (T₇) and kinetin (T₈) were on par for their effect on germination. Acid treatment and hot water treatment were also superior to control though their beneficial effect became apparent only 3 MAS. In *G. tinctoria*, polyembryony was observed in all the replications (4%–6% of germinated seeds).

Mathew & George (1995) reported that *G. gummi-gutta* seeds take more than 1 year for germination if they are sown soon after harvest and as the storage time increases, the time taken for germination decreases. In their studies, GA 250 ppm gave the highest germination percentage. In the present study, treatments like soaking in H₂O₂ 30%, kinetin 500 ppm and cow milk were found to be highly effective in inducing early germination. Similar results were also observed in *G. cowa*. Overnight soaking in cow milk was equally effective in breaking the dormancy and favoured early germination in *G. gummi-gutta* and *G. cowa* which otherwise required one year or more for germination. However, in *G. tinctoria*, along with H₂O₂, KNO₃ was also effective in inducing early germination. Both H₂O₂ and KNO₃ are reported to remove inhibitors from seed coat and enhance germination (Sadhu 1996).

Table 2. Evaluation of various treatments on germination of *Garcinia gummi-gutta* seeds

Treatment	4 MAS	5 MAS	6 MAS	7 MAS	8 MAS	9 MAS	10 MAS	11 MAS
Control	0.00 (0.09)	0.00 (0.09)	0.00 (0.09)	0.00 (0.09)	2.22 (0.15)	2.22 (0.12)	3.33 (0.18)	8.88 (0.18)
H ₂ SO ₄ 20% - 15 min	0.00 (0.09)	0.00 (0.09)	6.66 (0.24)	11.10 (0.33)	13.32 (0.37)	17.76 (0.43)	21.09 (0.48)	28.86 (0.48)
Hot water (55°C) - 15 min	0.00 (0.09)	0.00 (0.09)	0.00 (0.09)	1.11 (0.09)	2.22 (0.15)	7.77 (0.28)	9.99 (0.32)	14.43 (0.32)
KNO ₃ 0.2% - 18 h	0.00 (0.09)	2.22 (0.15)	4.44 (0.21)	7.77 (0.28)	11.10 (0.34)	18.87 (0.45)	26.64 (0.54)	34.41 (0.54)
Thiourea 2% - 18 h	0.00 (0.09)	0.00 (0.09)	2.22 (0.15)	4.44 (0.21)	5.55 (0.24)	8.88 (0.29)	15.54 (0.40)	21.09 (0.40)
H ₂ O ₂ 30% - 30 min	13.32 (0.37)	39.96 (0.68)	43.29 (0.72)	60.05 (0.90)	71.04 (1.01)	73.26 (1.03)	77.7 (1.08)	77.70 (1.08)
GA 500 ppm - 18 h	0.00 (0.09)	4.44 (0.21)	15.54 (0.40)	32.19 (0.60)	35.52 (0.64)	36.63 (0.65)	42.18 (0.71)	47.73 (0.71)
Kinetin 500 ppm - 18 h	0.00 (0.09)	8.88 (0.29)	23.31 (0.50)	34.41 (0.62)	44.40 (0.73)	52.17 (0.81)	55.5 (0.84)	60.00 (0.84)
Cow milk (fresh) - 18 h	0.00 (0.09)	21.09 (0.47)	28.86 (0.56)	39.96 (0.68)	48.84 (0.77)	53.28 (0.82)	63.27 (0.92)	64.44 (0.92)
CD (P=0.05)	0.06	0.12	0.12	0.15	0.18	0.18	0.18	0.18

Values indicate % germination; Values in parenthesis indicate transformed values ; MAS=Months after sowing; H₂SO₄=Sulphuric acid; KNO₃=Potassium nitrate; H₂O₂=Hydrogen peroxide; GA - Gibberellic acid

Table 3. Evaluation of various treatments on germination of *Garcinia cowa* seeds

No.	Treatment	3 MAS	4 MAS	5 MAS	6 MAS	7 MAS	8 MAS	9 MAS	10 MAS
T ₁	Control	0.00 (0.09)	0.00 (0.09)	0.00(0.09)	2.67 (0.15)	24.00 (0.45)	34.67 (0.57)	36.00 (0.58)	36.00 (0.58)
T ₂	H ₂ SO ₄ 20% - 15 min	0.00 (0.09)	2.67 (0.15)	20.00(0.41)	32.00 (0.54)	34.67 (0.56)	42.67 (0.64)	42.67 (0.64)	44.00 (0.65)
T ₃	Hot water (55°C) - 15 min	0.00 (0.09)	0.00 (0.09)	4.00(0.18)	13.33 (0.33)	21.33 (0.43)	36.00 (0.58)	38.67 (0.60)	41.34 (0.63)
T ₄	KNO ₃ 0.2% - 18 h	0.00 (0.09)	9.33 (0.26)	14.67(0.32)	28.00 (0.49)	30.67 (0.52)	38.67 (0.60)	40.00 (0.61)	44.00 (0.65)
T ₅	Thiourea 2% - 18 h	0.00 (0.09)	2.67 (0.15)	16.00(0.36)	21.33 (0.43)	25.33 (0.47)	32.00 (0.54)	37.33 (0.59)	37.33 (0.59)
T ₆	H ₂ O ₂ 30% - 30 min	12.00 (0.32)	32.00 (0.54)	49.33(0.70)	64.00 (0.82)	65.33 (0.83)	73.33 (0.90)	73.33 (0.90)	76.00 (0.92)
T ₇	GA 500 ppm - 18 h	0.00 (0.09)	0.00 (0.09)	5.30(0.20)	16.00 (0.37)	25.33 (0.48)	26.67 (0.49)	29.33 (0.52)	29.33 (0.52)
T ₈	Kinetin 500 ppm - 18 h	0.00 (0.09)	18.67 (0.40)	26.67(0.49)	42.67 (0.64)	54.67 (0.74)	57.33 (0.76)	57.33 (0.76)	62.67 (0.81)
T ₉	Cow milk (fresh) - 18 h	8.00 (0.26)	20.00 (0.42)	46.67(0.67)	60.00 (0.79)	69.33 (0.86)	73.33 (0.90)	73.33 (0.90)	73.33 (0.90)
	CD (P=0.05)	0.06	0.09	0.12	0.12	0.12	0.09	0.09	0.09

Values indicate % germination; Values in parenthesis indicate transformed values; MAS=Months after sowing; H₂SO₄=Sulphuric acid; KNO₃=Potassium nitrate; H₂O₂=Hydrogen peroxide; GA=Gibberellic acid

Table 4. Evaluation of various treatments on germination of *Garcinia tinctoria* seeds

No.	Treatment	1 MAS	1½ MAS	2 MAS	2½ MAS	3 MAS
T ₁	Control	0.00 (0.09)	0.00 (0.09)	1.30 (0.12)	41.30 (0.62)	44.00 (0.65)
T ₂	H ₂ SO ₄ 20% - 15 min	0.00 (0.09)	0.00 (0.09)	22.70 (0.45)	76.00 (0.92)	80.00 (0.96)
T ₃	Hot water (55°C) - 15 min	0.00 (0.09)	2.70 (0.15)	28.00 (0.50)	40.00 (0.62)	48.00 (0.69)
T ₄	KNO ₃ 0.2% - 18 h	17.30 (0.39)	32.00 (0.54)	41.30 (0.62)	68.00 (0.85)	81.30 (0.97)
T ₅	Thiourea 2% - 18 h	0.00 (0.09)	4.00 (0.17)	5.30 (0.20)	50.70 (0.71)	56.00 (0.75)
T ₆	H ₂ O ₂ 30% - 30 min	17.30 (0.38)	62.70 (0.81)	78.60 (0.98)	88.00 (1.03)	89.30 (1.04)
T ₇	GA 500 ppm - 18 h	6.70 (0.22)	9.30 (0.27)	64.00 (0.84)	80.00 (0.96)	85.30 (1.01)
T ₈	Kinetin 500 ppm - 18 h	4.00 (0.17)	48.00 (0.68)	93.30 (1.08)	96.00 (1.11)	96.00 (1.19)
	CD (P=0.05)	0.09	0.18	0.21	0.12	0.12

Values indicate % germination; Values in parenthesis indicate transformed values; MAS=Months after sowing; H₂SO₄=Sulphuric acid; KNO₃=Potassium nitrate; H₂O₂=Hydrogen peroxide; GA = Gibberellic acid

Prolonged seed dormancy found in *Garcinia* spp. increases the time and expenses required for raising rootstocks for grafting. The study established that seed germination can be significantly improved in *G. gummi-gutta* and *G. cowa* by pre-treatment with H₂O₂, kinetin or cow milk instead of practicing the laborious and time consuming process of seed coat removal. In *G. tinctoria* also, a satisfactory level of germination can be attained within 2 months by pre-treating seeds with H₂O₂, gibberellic acid or kinetin as against 4 months in control. Adoption of appropriate pre-treatment in each *Garcinia* species will reduce the cost of graft production by reducing the dormancy period and increasing percentage germination.

References

- George S T, Baby Latha A K, Mathew K L & Geetha C K 1992 Pattern of flowering and flower development in *Kodampuli* (*Garcinia cambogia* Desr). Indian Cocoa Arecanut Spices J. 16: 68-70.
- KAU (Kerala Agricultural University) 2001 Three Decades of Spices Research at KAU. Kerala Agricultural University, Thrissur.
- KAU (Kerala Agricultural University) 2002 Camboge. In: Package of Practices Recommendations: Crops. 12th Edition (pp. 11-120). Kerala Agricultural University, Thrissur.
- Maheshwari J K 1964 Taxonomic studies on Indian Guttiferae. III. The genus *Garcinia* L. Bull. Bot. Survey India 6: 107-135.
- Mathew K L & George S T 1995 Dormancy and storage of seeds in *Garcinia cambogia* Desr. (*Kodampuli*). J. Trop. Agric. 33: 77-79.
- Normah M N, Ramiya S D & Gintangga M 1997 Desiccation sensitivity of recalcitrant seeds - a study on tropical fruit species. Seed Sci. Res. 7: 179-183.
- Sadhu M K 1996 Plant propagation. New Age International (P) Ltd., New Delhi.
- Vazquez-Yanes C & Orozco-Segovia A 1993 Patterns of seed longevity and germination in the tropical rain forest. Ann. Rev. Ecol. Sys. 24: 69-87.