

Advances in glioblastoma management

B. Iliescu¹, I. Poeta²

¹“Prof. Dr. N. Oblu” Clinical Emergency Hospital Iasi

²“Gr.T. Popa” University of Medicine and Pharmacy Iasi

Introduction

Resection of the maximum amount of tumoral tissue while preserving the function of the brain within or neighboring the tumor region established itself as a fundamental goal of modern oncologic neurosurgery. Maximal safe removal of the tumoral tissue has been documented extensively as a major prognostic factor for patients with glioblastoma (8). As such it became imperative for the oncologic neurosurgeon to be able to identify reliably the functional structures, cortical, subcortical, and vascular both preoperatively and in the operating room (5).

Mapping the eloquent areas of the brain has become a well-established technique using the intraoperative electrophysiology, with reliable methods and results (9,10,11). As well, preoperative mapping of the same areas using functional MRI and/or magnetoencephalography (MEG) has given clinically useful results in experienced hand (2, 3). However these methods do not provide any information about the white matter tracts that may be affected by invasive, intrinsic brain tumors. The advances in diffusion-tensor (DT) imaging techniques have been used to map white matter tracts in the normal brain (1, 4).

Methodology

The aim of our paper is to demonstrate

the role of DT imaging in preoperative mapping of white matter tracts in relation to cerebral neoplasms and the way it can influence the operative decision in our practice. We discuss as well the methods used intraoperatively to identify and recognize functional areas and neighboring white matter tracts in the cases with glioblastoma (6, 7, 8). Ten patients suffering from this brain malignancy within or neighboring eloquent cortical areas underwent DT imaging examinations prior to tumor excision. Anatomical information about white matter tract location, orientation, and projections was obtained in every patient. Depending on the tumor type and location, evidence of white matter tract edema (one patient), infiltration (one patient), displacement (five patients), and disruption (three patients) could be assessed with the aid of DT imaging in each case. Diffusion-tensor imaging allowed for visualization of white matter tracts and was found to be beneficial in the surgical planning for patients with intrinsic brain tumors.

Conclusion

Our report is both a general and personal. In the first part we analyze the decision making process and protocol in the era of complex preoperative planning, microsurgery, ultrasound aspiration, neuronavigation, awake craniotomy, and

white matter tracts stimulation techniques in the surgical treatment of these cases. We present our experience using these technologies in combination with tractography in the treatment of 10 glioblastoma cases, affecting eloquent areas of the brain. The data strongly suggests that using both pre and intraoperative functional techniques improves significantly the results in terms of postoperative neurological status allowing for a safe maximal tumor removal.

References

1. Handler T, et. Al. Delineating gray and white matter involvement in brain lesions: three-dimensional alignment of functional magnetic resonance and diffusion-tensor imaging. *J Neurosurg* 99:1018–1027, 2003.
2. Kim P., Singh M.S. Functional magnetic resonance imaging for brain mapping in neurosurgery. *Neurosurg Focus*, 15(1):1-7, 2003.
3. Kober H, Nimsky C, Möller M, Hastreiter P, Fahlbusch R, Ganslandt O. Correlation of sensorimotor activation with functional magnetic resonance imaging and magnetoencephalography in presurgical functional imaging: a spatial analysis. *Neuroimage*, 14(5):1214–28, 2001.
4. Krings T, Reinges MHT, Thiex R, Gilsbach JM, Thron M. Functional and diffusion-weighted magnetic resonance images of space-occupying lesions affecting the motor system: imaging the motor cortex and pyramidal tracts. *J Neurosurg* 95:816–824, 2001.
5. Lehericy S, Duffau H, Cornu P, et al: Correspondence between functional magnetic resonance imaging somatotopy and individual brain anatomy of the central region: comparison with intraoperative stimulation in patients with brain tumors. *J Neurosurg* 92:589–598, 2000
6. Martino, J., Taillandier, L., Moritz-Gasser, S., Gatignol, P., Duffau, H. Re-operation is a safe and effective therapeutic strategy in recurrent WHO grade II gliomas within eloquent areas. *Acta Neurochir.* 151(5):427-36; discussion 436, 2009.
7. McDonald JD, Chong BW, Lewine JD, et al: Integration of preoperative and intraoperative functional brain mapping in a frameless stereotactic environment for lesions near eloquent cortex. Technical note. *J Neurosurg* 90:591–598, 1999
8. McGirt M.J., Chaichana K.L., Gathinji M., Attenell F.J., Than K., Olivi A., Weingart J.D., Brem H., Quiñones-Hinojosa A. Independent association of extent of resection with survival in patients with malignant brain astrocytoma. *J Neurosurg* 110:156–162, 2009
9. Meyer, B. Introduction: Surgery of limbic and paralimbic gliomas. *Neurosurg Focus* 27 (2):E1, 2009.
10. Plaza, M., Gatignol, P., Leroy, M., Duffau, H. Speaking without Broca's area after tumor resection. *Neurocase*, Mar 9:1-17, 2009.
11. Von Lehe, M., Schramm, J. Gliomas of the cingulate gyrus: surgical management and functional outcome. *Neurosurg Focus* 27 (2):E9, 2009.