

## Demographic pattern, clinical profile and outcome of traumatic spinal cord injuries at a tertiary care hospital

K. Aswani Kumar<sup>1</sup>, B.V. Subrahmanyam<sup>2</sup>, S.V. Phanindra<sup>2</sup>, S. Satish Kumar<sup>3</sup>, P.N. Harish<sup>4</sup>, P. Ramamohan<sup>5</sup>, Amit Agrawal<sup>6</sup>

Narayana Medical College Hospital, Chinthareddypalem, Nellore, Andhra Pradesh (India)

<sup>1</sup>Resident Forensic Medicine, Department of Forensic Medicine

<sup>2</sup>Professor of Forensic Medicine, Department of Forensic Medicine

<sup>3</sup>Associate Professor of Emergency Medicine, Department of Emergency Medicine

<sup>4</sup>Associate Professor of Neurosurgery, Department of Neurosurgery

<sup>5</sup>Associate Professor of Pharmacology, Department of Forensic Pharmacology

<sup>6</sup>Professor of Neurosurgery, Department of Neurosurgery

**Abstract:** *Background:* Traumatic spinal cord injury (SCI) is recognized as a serious public health problem resulting in significant morbidity, mortality and permanent disability. The present study is aimed to describe the epidemiological characteristics and outcome of patients with traumatic spinal cord injury in rural tertiary referral care center form South India. *Material and methods:* The present study was conducted at Narayana Medical College and Hospital, Nellore. All patients admitted and managed for traumatic spinal cord injury were retrieved and data collected in a pre-designed proforma. Patient characteristics, details of etiology, mechanism of injury, level of injury, extent of neurological deficits, details of investigations, details of management and immediate outcome were recorded. *Results:* A total 152 patients were included in the present study. The mean age was 38.45 years and majority the patients were young adult males. The mean hospital stay was 19.12 days. 71.7% percent patients were non-agriculture workers (mainly involved in construction work) and 28.3% patients were farmers. 61.2% of the patients sustained injuries due to fall from height and 34.2% patients sustained injuries due road traffic accidents. Cervical spine injuries were most common (44.1%), followed by thoraco-lumbar region (36.8%) and dorsal spinal region (19.1%). 9 patients expired in post-injury during hospital stay and all of them had complete cervical spinal cord injury. All patients received aggressive rehabilitation care. *Conclusion:* In accordance with the literature our results reflect that traumatic spinal cord injuries affect young population and can leave these persons with significant functional and physical morbidity. The major limitation of the study is that it is a single institution based and may not reflect the true spectrum of traumatic spinal cord injuries in the population.

**Key words:** Spinal cord injury, spinal injuries, spinal trauma

### Introduction

Traumatic spinal cord injury (SCI) is recognized as a serious public health problem that can result in significant morbidity, mortality and permanent disability (leading to emotional and financial hardships). (1-3) Only few studies report the epidemiological characteristics of traumatic SCI in developing

countries and describe their etiology, demographic and clinical characterizes and management option. (4, 5) The present study is aimed to describe the epidemiological characteristics and outcome of patients with traumatic spinal cord injury in rural tertiary referral care center form South India.

### Material and methods

The present study was conducted at Narayana Medical College and Hospital, Nellore. After approval from the Institutional Ethical Committee, the medical records of all patients admitted and managed for traumatic spinal cord injury were retrieved and data collected in a pre-designed proforma. Patient characteristics including age, gender, occupation were noted. Details of etiology, mechanism of injury, level of injury, extent of neurological deficits, details of investigations, details of management and immediate outcome were recorded. The details of motor and sensory functions were recorded by using the American Spinal Injury Association (ASIA) scale and the severity was classified as complete or incomplete injury. (6) The data were analyzed on Epiinfo ® version 7. The descriptive statistics were used to calculate mean and frequencies of data.

### Results

A total of 152 patients were included in the present study. The mean age was 38.45 years (range 6-72 years  $SD\pm 14.566$  years). In present study majority of the patients were young adult males (Figure 1). The mean

hospital stay was 19.12 days (range 4-59 days  $SD\pm 10.699$  days). 71.7% percent patients were non-agriculture workers (mainly involved in construction work) and 28.3% patients were farmers. 61.2% of the patients sustained injuries due to fall from height and 34.2% patients sustained injuries due road traffic accidents. Cervical spine injuries were most common (44.1%), followed by thoracolumbar region (36.8%) and dorsal spinal region (19.1%). In our series majority of the patients had incomplete spinal cord injury. 18 patients had associated traumatic brain injury and had low Glasgow Coma Score at the time of presentation. CT scan was performed for all these patients and two patients had lesions (one patient extradural hematoma and one patient acute subdural hematoma) those required neurosurgical intervention. Twenty two patients were managed conservatively and 130 patients underwent spinal fusion and fixation. Elective mechanical ventilation was required in 19 cases (all cases were of cervical spine injuries). 9 patients expired post-injury during hospital stay and all of them had complete cervical spinal cord injury. All patients received aggressive rehabilitation care (chest physiotherapy, bowel, bladder and back care).




Figure 1 - Age and gender distribution (n=152)




Figure 2 - Distribution of the patients according to the month of the year

### Discussion

As we observed and many studies have reported in the past, traumatic spinal cord

injury young adult males in their productive years. (7-9) (1, 9-13) As reported in many, fall from height was the leading cause of spinal cord injuries in our series (mainly in non-agricultural workers followed by agricultural workers). (14-19) In contrary to this many other studies have reported where the authors found agricultural workers (9, 20) and road traffic injuries (21-23) as the commonest causes of traumatic spinal cord injuries respectively. We found incomplete injuries more than the complete injuries and cervical spine injuries were being the most common (44.1%), followed by thoraco-lumbar injuries (36.8%) and dorsal spinal injuries (19.1%). In literature the distribution of incomplete and complete injuries varies as per the anatomical involvement, location and mechanism of the injuries. (1, 9, 20, 24-26)

Most of the traumatic spinal injuries are managed by neurosurgeons, spinal surgeons and by orthopedic surgeons (followed by intensive rehabilitation) and the management of spinal cord injury patients is determined by their neurological status, instability of the spine and ability to tolerate neurosurgical intervention. (4, 27, 28) As described in the literature, in majority of our patients surgical interventions were performed for different indication to stabilize the spine and/or to decompress the spinal cord. (29-32) Depending on the severity of level of the spinal cord injury, the duration of the hospital stay varies. In present study we noted that the mean hospital stay was 19.12 days (range 4-59 days SD±10.699 days). Lan et al reported that the average hospital stay for paraparetic was 62 days, for paraplegics 118 days, for tetraparetics 102 days and for tetraplegics 132 days. (24) In another study the authors reported that the mean hospital stay for paraparetic was 44 days, for paraplegics, for tetraparetics 90 days and for tetraplegics patients it was 100 days. (1) In present study we found the mortality rate of 5.9%. Chen et al (1) reported a mortality rate of 6.0% and Lan

et al (24) reported a mortality rate of 10.1%. respiratory failure (particularly in tetraplegic) was the leading cause of mortality. (1, 24) Other causes those can influence the outcome include sepsis (from pressure ulcers), renal failure and other associated systemic injuries (abdominal/chest injuries, head injury). (1, 24)

In present study we did not collect data regarding financial implications of spinal cord injuries. However because of the difficult to cure and their potential to give rise to much human stress traumatic spinal cord injury can cause great loss not only to the individuals but also to the society. (33) The lifetime cost of the traumatic spinal cord injuries has been shown to be directly attributed to the age at the time of injury and the severity of injury. (30) It has been estimated that the lifetime costs for an incomplete motor lesion at age 50 years at any level range from \$300,000 to \$1.7 million for a person who sustain higher injuries (tetraplegia) at the age of 25 years. (34) Type of transportation depends on locations of treating facilities and availabilities of pre-hospital care. In major cities there is greater availability of ambulance networks equipped with trained and semi-trained personnel's, in contrary to this in rural areas these patients are transferred by untrained personnel's with different means. (35, 36)

## Conclusion

In spite of much advancement for the management of spinal cord injury patients, there is no definitive treatment modality which can help to recover the neurological functions. Hence the preventive programs have been promoted to prevent spinal cord injuries. (17, 20) Preventive strategies and policies needs to be focused on subjects with high risk activities: sports activities (diving and motorcycles accidents) for young people; road traffic accidents for adults; injuries due to fall in elderly population. (37) The major limitation of the study is that it is a single institution based and may not reflect the true

spectrum of traumatic spinal cord injuries in the population. However in accordance with the literature our results reflect that traumatic spinal cord injuries affect young population and can leave these persons with significant functional and physical morbidity.

### Correspondence

Dr. Amit Agrawal  
Professor of Neurosurgery  
Department of Neurosurgery  
Narayana Medical College Hospital  
Chinthareddyapalem  
Nellore-524003  
Andhra Pradesh (India)  
Email- dramitagrawal@gmail.com  
dramit\_in@yahoo.com  
Mobile- +91-8096410032

### References

- Chen CF, Lien IN. Spinal cord injuries in Taipei, Taiwan, 1978-1981. *Paraplegia*. 1985;23(6):364-70.
- McDonald JW, Sadowsky C. Spinal-cord injury. *Lancet*. 2002;359(9304):417-25.
- Tascu A, Exergian F, Daniel S, Iliescu A, Pascal C, Popescu G, et al. Traumatic spine injuries in children. Experience on 434 cases and therapeutic perspectives. *Romanian Neurosurgery*. 2015;XXIX(2):135-40.
- Rathore MFA, Hanif S, Farooq F, Ahmad N, Mansoor SN. Traumatic spinal cord injuries at a tertiary care rehabilitation institute in Pakistan. *JPM*. 2008;58(2):53-7.
- Rahimi-Movaghar V, Sayyah MK, Akbari H, Khorramirouz R, Rasouli MR, Moradi-Lakeh M, et al. Epidemiology of traumatic spinal cord injury in developing countries: a systematic review. *Neuroepidemiology*. 2013;41(2):65-85.
- American Spinal Injury Association. International standards for neurological classification of spinal cord injury revised Chicago ASIA. 2002.
- Girard R, Boisson D, Depassio J, Boucand M, Eyssette M. The female paraplegic: a statistical survey. *Spinal Cord*. 1983;21(3):149-53.
- Girard R, Minaire P, Castanier M, Berard E, Perrineriche B. Spinal cord injury by falls: comparison between suicidal and accidental cases. *Paraplegia*. 1980;18(6):381.
- Minaire P, Castanier M, Girard R, Berard E, Deidier C, Bourret J. Epidemiology of spinal cord injury in the Rhône-Alpes Region, France, 1970-75. *Paraplegia*. 1978;16(1):76-87.
- Burke DC, Toscano J. Incidence and distribution of spinal cord injury. Menzies Foundation Technical Report. 1987;1 13-47.
- Gjone R, Nordlie L. Incidence of traumatic paraplegia and tetraplegia in Norway: a statistical survey of the years 1974 and 1975. *Paraplegia*. 1978;16(1):88-93.
- Stover SL, Fine PR. The epidemiology and economics of spinal cord injury. *Paraplegia*. 1987;25(3):225-8.
- Chacko V, Joseph B, Mohanty SP, Jacob T. Management of spinal cord injury in a general hospital in rural India. *Paraplegia*. 1986;24(5):330-5.
- Dryden DM, Saunders LD, Rowe BH, May LA, Yiannakoulis N, Svenson LW, et al. The epidemiology of traumatic spinal cord injury in Alberta, Canada. *The Canadian journal of neurological sciences*. 2003;30(2):113-21.
- Pickett GE, Campos-Benitez M, Keller JL, Duggal N. Epidemiology of traumatic spinal cord injury in Canada. *Spine*. 2006;31(7):799-805.
- Pickett W, Simpson K, Walker J, Brison RJ. Traumatic spinal cord injury in Ontario, Canada. *The Journal of trauma*. 2003;55(6):1070-6.
- Shingu H, Ohama M, Ikata T, Katoh S, Akatsu T. A nationwide epidemiological survey of spinal cord injuries in Japan from January 1990 to December 1992. *Paraplegia*. 1995;33(4):183-8.
- Kannus P, Palvanen M, Niemi S, Parkkari J. Alarming rise in the number and incidence of fall-induced cervical spine injuries among older adults. *The journals of gerontology Series A, Biological sciences and medical sciences*. 2007;62(2):180-3.
- Nobunaga AI, Go BK, Karunas RB. Recent demographic and injury trends in people served by the Model Spinal Cord Injury Care Systems. *Archives of physical medicine and rehabilitation*. 1999;80(11):1372-82.
- Dincer F, Oflazer A, Beyazova M, Celiker R, Basgöze O, Altıoklar K. Traumatic spinal cord injuries in Turkey. *Paraplegia*. 1992;30(9):641-6.
- Hung CC, Chiu WT, Tsai JC, Laporte RE, Shih CJ. [An epidemiological study of head injury in Hualien County, Taiwan]. *Journal of the Formosan Medical Association*. 1991;90(12):1227-33.
- Thurman DJ, Burnett CL, Beaudoin DE, Jeppson L, Sniezek JE. Risk factors and mechanisms of occurrence in motor vehicle-related spinal cord injuries: Utah. *Accident Analysis & Prevention*. 1995;27(3):411-5.
- Thurman DJ, Burnett CL, Jeppson L, Beaudoin DE, Sniezek JE. Surveillance of spinal cord injuries in Utah, USA. *Paraplegia*. 1994;32(10):665-9.
- Lan C, Lai JS, Chang KH, Jean YC, Lien IN. Traumatic spinal cord injuries in the rural region of Taiwan: an epidemiological study in Hualien county, 1986-1990. *Paraplegia*. 1993 Jun;31(6):398-403. PubMed PMID: 8337004. Epub 1993/06/01. eng.

25. Richardson RR, Meyer PR. Prevalence and incidence of pressure sores in acute spinal cord injuries. *Paraplegia*. 1981;19(4):235-47.
26. Tator CH, Fehlings MG. Review of the secondary injury theory of acute spinal cord trauma with emphasis on vascular mechanisms. *Journal of neurosurgery*. 1991;75(1):15-26.
27. Scivoletto G, Morganti B, Molinari M. Early versus delayed inpatient spinal cord injury rehabilitation: an Italian study. *Archives of physical medicine and rehabilitation*. 2005;86(3):512-6.
28. Guttman L. New hope for spinal sufferers: Ludwig Guttman. Reproduced from *Medical Times*, November 1945. *Paraplegia*. 1979;17(1):6-15.
29. Esce, Haines. Acute Treatment of Spinal Cord Injury. Current treatment options in neurology. 2000;2(6):517-24.
30. Becker D, Sadowsky CL, McDonald JW. Restoring function after spinal cord injury. *The neurologist*. 2003;9(1):1-15.
31. Chen TY, Dickman CA, Eleraky M, Sonntag VK. The role of decompression for acute incomplete cervical spinal cord injury in cervical spondylosis. *Spine*. 1998;23(22):2398-403.
32. Mirza SK, Krengel WF, Chapman JR, Anderson PA, Bailey JC, Grady MS, et al. Early versus delayed surgery for acute cervical spinal cord injury. *Clinical orthopaedics and related research*. 1999 (359):104-14.
33. Shingu H, Ikata T, Katoh S, Akatsu T. Spinal cord injuries in Japan: a nationwide epidemiological survey in 1990. *Paraplegia*. 1994;32(1):3-8.
34. Westergren H, Farooque M, Olsson Y, Holtz A. Spinal cord blood flow changes following systemic hypothermia and spinal cord compression injury: an experimental study in the rat using Laser-Doppler flowmetry. *Spinal cord*. 2001;39(2):74-84.
35. Srinivasan U. acute Spinal Cord Injury: managing at the Site of Impact and addressing Reality Gap. *JAPI* 2012;60:7-9.
36. Singh R, Sharma SC, Mittal R, Sharma A. Traumatic spinal cord injuries in Haryana: An epidemiological study. *Indian J Community Med*. 2003;28(4):184-86.
37. Kiwerski J. The causes, sequelae and attempts at prevention of cervical spine injuries in Poland. *Spinal Cord*. 1993;31(8):527-33.