



“The Silk Road” via subarachnoid cisterns. Cerebrospinal fluid dissemination of meningiomas

A.I. Cucu¹, Mihaela Dana Turliuc^{1,2}, Claudia Florida Costea^{1,2}, B. Costăchescu^{1,2}, Cristina Mihaela Ghiciuc², B. Dobrovăţ^{1,2}, Camelia Margareta Bogdănici², Daniela Maria Tănase², Gabriela Florenţa Dumitrescu¹, Anca Sava^{1,2}, I. Poeată^{1,2}

¹ “Prof. Dr. Nicolae Oblu” Emergency Clinical Hospital Iaşi, ROMANIA

² “Grigore T. Popa” University of Medicine and Pharmacy Iaşi, ROMANIA

ABSTRACT

Meningiomas are generally slow-growing extra-axial benign tumours and in rare cases they can metastasize both neural and extra-neural. Intracranial meningiomas with leptomeningeal dissemination are extremely rare and the exact pathogenesis still remains unknown. The aim of this review is to analyse the pathways of intracranial and spinal metastatic spread of intracranial meningiomas and to discuss their particular clinical and pathological features. We highlight the fact that there is a possibility of leptomeningeal dissemination, even if cerebrospinal fluid cytology is negative, in patients with a medical history of a resected meningioma. We identified three possible ways of dissemination: haematogenous, through the CSF, or during surgery. From a histopathological point of view, the more malignant the meningioma, the more likely its leptomeningeal dissemination.

INTRODUCTION

Meningiomas are generally benign intracranial tumours and represent approximately 30% of all primary central nervous system tumours, with an incidence that has increased in recent years (1, 2, 3, 4, 5). Usually occurring on the surface of the brain as they originate in the arachnoid cells, meningiomas are generally slow-growing extra-axial benign tumours. In very rare cases, especially when the tumours become malignant, meningiomas can metastasize both in neural and extra-neural sites (6).

Only 0.1% of meningiomas are thought to metastasize (7), and usually these cases were atypical and anaplastic meningiomas (8, 9, 10), also known for their tendency to relapse after surgery (2, 11, 12, 13). Ather Enam et al. report an overall metastatic risk of 5% for atypical meningiomas and of 30% for anaplastic meningiomas (8), although

Keywords
meningioma,
metastatic spread,
CSF disseminated
meningioma



Corresponding author:
Mihaela Dana Turliuc

“Grigore T. Popa” University of
Medicine and Pharmacy Iaşi,
Romania

turliuc_dana@yahoo.com

Copyright and usage. This is an Open Access article, distributed under the terms of the Creative Commons Attribution Non-Commercial No Derivatives License (<https://creativecommons.org/licenses/by-nc-nd/4.0/>) which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is unaltered and is properly cited.

The written permission of the Romanian Society of Neurosurgery must be obtained for commercial re-use or in order to create a derivative work.

ISSN online 2344-4959
© Romanian Society of
Neurosurgery




First published
September 2019 by
London Academic Publishing
www.lapub.co.uk

other authors consider these percentages to be overestimated (11, 14, 15).

Meningiomas can metastasize in various organs (lung, liver, spleen, adrenal gland, thyroid, parotid gland) or bones, skin and deep soft tissue (16), but also in the intracranial or spinal space through cerebrospinal fluid (CSF) pathways. According to the existing literature, metastatic dissemination of meningiomas through CSF is a rare presentation (17, 18, 19, 20, 21) and, since 1950, only 45 cases have been reported (22). Nevertheless, different other authors consider that the CSF dissemination risk of meningiomas is an uncommon event, as it may occur

in 4% of metastasizing meningiomas (17, 23).

However, the first historical description of leptomeningeal metastasis was made by Oliver in 1837, followed by Eberth in 1870 (24, 25). Also, the first description of carcinoma cells in the CSF was realized by Dufour (26) and, in 1912, Berman was the first who used the term "meningeal carcinomatosis" (27). However, Kalm, in 1950, was the first who published a case of a malignant tentorium tumour, i.e. an anaplastic meningioma that metastasized in the medulla oblongata and in the leptomeningeal space (28).



PATHWAYS OF METASTATIC SPREAD IN MENINGIOMAS

Meningioma cells gain access to the subarachnoid space in several ways: haematogenous pathway, through the CSF, or during surgery. More frequently,

the spread is haematogenous via the venous system and tumour cells gain access to the CSF through the dural sinuses or epidural plexus, especially in meningiomas that invade dural sinuses (29, 30). A

second route of dissemination is by CSF with the tumour spreading throughout the neuroaxis (Figure 1).

In a review, Rawat *et al.* highlighted the fact that approximately 75% of patients with meningiomatous metastasis regardless of the route they followed have had previous operation for the primary tumour (31). Also, it was hypothesized that tumour cells gain access to the vascular channels and meningeal surface by seeding during surgery (32). Although this theory has earlier been rejected (33, 34), now it is accepted that surgical manipulation may release tumour cells into the CSF (30, 31).

Metastatic spread during or after surgery exists theoretically, even though this theory is unlikely. But the incidence of leptomeningeal dissemination after meningiomas surgery is low and there are also cases with leptomeningeal dissemination through CSF without surgical intervention (35). Regarding malignant transformation of meningiomas after surgical resection, there is also the theoretical one (22). Koenig *et al.* reported that at the site of surgical injury, the growth factors can promote malignant changes inside the meningeal tumour (36) and this event was observed by Morantz and Shain in an experimental rat model (37).

INTRACRANIAL AND SPINAL CORD DISSEMINATION THROUGH CSF PATHWAYS

There are variable sites of meningioma metastasis through CSF pathways. The metastasis occurred due to leptomeningeal seeding from the neighboring meningioma, and the spinal canal is the second most common site (17) (Table 1).

In literature, spinal metastasizing meningiomas are rarely reported (17, 38, 39, 40, 41). In spinal intradural dissemination, the tumour cells have a tendency to accumulate more frequently at dorsal nerve root level, especially in the medullary cone and cauda equine, probably due to directional active flowing of the CSF throughout the neuroaxis, and also to the effect of gravity (42, 43). In a large study on 200 consecutive meningiomas, of WHO grade I, which were followed prospectively during a median time of 8.5 years, Chamberlain and Glantz reported that 4 patients (2%) were diagnosed with spinal metastasis (21). Vries *et al.* also reported CSF or drop metastasis in 5% of non-benign meningiomas (44).

Meningioma metastasis may be simultaneous, both in the intracranial space and in the spinal cord

(17, 45). In 1992, Akimura *et al.* reported a malignant meningioma metastasizing through the CSF pathways, both in the cerebellopontine angle cistern and in the thoracic spinal cord. The primary tumour was a parasagittal malignant meningioma two-times operated, the second time for recurrence. At the first surgery, the frontal horn of the lateral ventricle was opened because the meningioma infiltrated into the deep frontal brain. The authors concluded that this artificial communication between the meningioma cavity and the CSF pathways enhanced the probability for tumour metastasis into cerebellopontine angle cistern and thoracic spinal cord (17).

NEOPLASTIC MENINGITIS

It is well known that neoplastic meningitis is more common with solid carcinoma such as lung, breast and gastrointestinal cancer (46), but literature also reported few malignant meningioma cases with CSF dissemination into the brain ventricles (47). There were only eight cases (47, 48) with intracranial or intraspinal malignant meningioma arising from low-grade meningiomas, which disseminated throughout the CSF (47). In neoplastic meningitis, CSF dissemination of malignant meningioma cells may cause a variety of neurological symptoms such as disturbances of multiple cranial nerves, hydrocephalus, cerebellar dysfunction and multiple spinal nerve roots or cauda equine symptoms (49).

Brainstem damage by CSF dissemination of malignant cell is rare, literature reporting a few cases with central hyperventilation, Wallenberg syndrome and diplopia, facial nerve palsy and unsteadiness of gait secondary of CSF dissemination of malignant tumour cells (20, 47, 50).

It is important to mention, in terms of diagnosis, that only 54% of all cases with leptomeningeal meningiomatous dissemination revealed malignant cells in the CSF on initial lumbar puncture, and only 8% of these cases remained negative, even after repeated examinations (51). The reason for this low specificity of the lumbar puncture remains unclear, although Fujimaki *et al.* speculate the fact that malignant cells adhere rather than float freely in the CSF (47). Although identification of malignant cells by CSF cytology has been considered the diagnostic gold standard, these paradigms have changed nowadays due to limited sensitivity of cytology.

HISTOPATHOLOGY OF CSF-DISSEMINATED MENINGIOMAS

Even though the pathogenesis of CSF - disseminated meningiomas is not completely understood, over time researchers have issued various theories. In this regard, Engelhard proposed three different pathways of dissemination: (1) tumour cells could be "shed" away directly into the CSF due to direct contact between an anaplastic meningioma and CSF pathways; (2) tumour cells might invade the leptomeningeal space during its progression; 3) the tumour cells might be inoculated within the CSF at the time of the surgery (52).

Other authors, such as Russel and Rubinstein, considered that tumour friability may play an important role in meningioma dissemination within CSF (42). However, considering that meningiomas arise from arachnoid cells and are naturally exposed to CSF during their growth, it is difficult to explain the scarcity of meningioma dissemination through CSF (22) as Chamberlain and Glantz reported that only 4% of the meningiomas could present leptomeningeal dissemination and positive CSF cytology at the time of the diagnosis (21).

More theoretically, the risk of intraventricular meningiomas to metastasize through CSF pathways should be high, but literature reports only nine cases of intraventricular meningiomas (22). As an explanation, Miller and Ramsden considered that the dynamics of the CSF pathways might prevent fragment formation and deposition of tumour cells (22, 53). However, in a review of 45 cases of meningioma with leptomeningeal dissemination through CSF, Park et al. noticed that the period of time needed for leptomeningeal dissemination in cases with intraventricular meningiomas is the shortest when compared to other intracranial and spinal meningiomas (22).

All histological subtypes of meningiomas can metastasize (Table 1), even benign meningiomas (45, 54, 55, 56, 57, 58), but meningeal tumours with clearly malignant features have a higher metastatic rate (59). Metastatic meningiomas are associated more frequently with aggressive meningiomas (WHO grade 2, and WHO grade 3), with a range of occurrence of 10-25% (21), i.e. the more malignant the meningioma, the more likely its leptomeningeal dissemination (9, 17, 41, 60).

According to literature, from a histological point of view, several factors are predictive of meningioma metastasis, including high cellularity, nuclear

pleomorphism (Figure 2), high mitosis rate, tumour necrosis, and invasion of blood vessels (7, 8). On the other hand, other authors consider that the metastasizing behaviour of these tumours is not correlated with their histological features (30, 31).




FIGURE 2. Microphotographs of atypical meningioma, WHO grade II. A. A tumour made up of meningotheelial cells arranged in a sheet-like pattern, with increased cellularity, pleomorphism, and areas with small cells having high Nucleus/Cytoplasm ratio (HE staining, x20); B. Ki67 labelling index has high values and indicates a malignant neoplasia (Anti-Ki67 antibody immunohistochemical staining, X40).

CONCLUSION

Intracranial meningiomas with CSF dissemination are extremely rare and, despite reports from literature, their pathogenesis remains unknown. In the case of a patient with a resected meningioma, the possibility of CSF dissemination of tumour cells should be borne in mind, even if CSF cytology is negative. Also, a spinal MRI should be performed, especially when spinal signs and symptoms are present.

TABLE 1. Literature review of meningioma cases with CSF- dissemination

Year of publication, Author	Primary site of meningioma	WHO Grade	Intracranial metastasis	Spinal metastasis	Time to CSF disseminated disease
2013, Tsuda et al. (43)	T10-11 intradural	I →II	-	+	12 years
2011, Wu et al. (61)	Convexity	III	+	-	1.7 years
2011, Kim et al. (62)	Posterior fossa	III	+	-	0
2011, Peng et al. (63)	Medial temporal	III	+	-	2 years
2009, Kuroda et al. (64)	Skull base	I→II→III I	+	+	6.3 years
2009, Eom et al. (6)	Lateral ventricle	II→III	-	+	1.3 years
2008, Erkutlu et al. (65)	Posterior fossa	III	-	+	2.7 years
2008, Santhosh et al. (65)	Convexity	III	+	+	9 months
2007, Shintaku et al. (67)	Lateral ventricle	I→III	+	-	4.3 years
2006, Chuang et al. (68)	Convexity	III	-	+	3 months
2005, Cramer et al. (60)	C1-C3 intradural	II	-	+	1.4 years
2005, Al-Habib et al. (69)	Not mentioned	III	+	+	2 months
2005, Chamberlain and Glantz (21)	Not mentioned, 8 cases	I	+ (8 cases)	+ (6 cases)	-
2005, Wakabayashi et al. (70)	Frontal convexity	III	+	-	13 years
2005, Koenig et al. (36)	Temporal lobe	III	-	+	1 month
2005, Darwish et al. (71)	Lateral ventricle	II→III	+	+	7 months
2002, Ramakrishnamurthy et al. (35)	Lateral ventricle	I	+	-	4 years
2001, Conrad et al. (72)	Convexity	I→II→III I	+	+	6.4 years
2000, Meinsma-VdTuin et al. (73)	C2-C4 intradural	III	+	+	6 months
2000, Lee et al. (74)	Convexity	III	+	+	9 years
1998, Lee and Landy (41)	Convexity	III	+	+	1.5 years
	Skull base	I	+	+	3.8 years
	Convexity	III	-	+	8.5 years

1995, Peh and Fan (39)	Lateral ventricle	III	+	+	5 years
1993, Greenberg et al. (75)	Lateral ventricle	III	+	+	2 months
1992, Satoh et al. (18)	Skull base	I	+	+	0
1992, Akimura et al. (17)	Convexity	III	+	+	1.8 years
1989, Kamiya et al. (76)	Lateral ventricle	III	-	+	6 months
1987, Strenger et al. (77)	Third ventricle	III	+	-	1.5 months
1985, Kleinschmidt-DeMasters and Avakian (20)	Lateral ventricle	III	+	+	1.7 years
1975, Ludwin and Conley (78)	Convexity	III	+	+	10 months
1972, Miller and Ramsden (53)	Convexity	III	+	-	-
1971, Riley et al. (22, 78)	Convexity	I→III	-	+	-
1970, Shuanghoti et al. (22, 78)	Convexity	III	-	+	-
1963, Russell et al. (22, 78)	Skull base	I	+	-	-
1960, Hoffman et al. (22, 78)	Convexity	I→III	+	+	-
1954, Winkelman (22, 78)	Skull base	I	+	-	-
1950, Kalm (22, 78)	Posterior fossa	III	+	-	-

REFERENCES

- Louis DN, Scheitauer BW, Budka H, von Deimling A, Kepes JJ. Meningiomas. In: Kleihues P, Cavenee WK (Eds). Pathology and Genetics of Tumors of the Nervous System. Lyon: IARC Press, 2000.
- Cucu AI, Costea CF, Poeata I, et al. Prognostic factors in atypical meningioma. Rom Neurosurg. 2017; 31(2):165-171.
- Cucu AI, Costea CF, Poeata I, et al. Anatomical localization of atypical meningiomas: our experience on 81 patients. Rev Med Chir Soc Med Nat Iasi. 2018; 122:744-752
- Cucu AI, Costea CF, Carauleanu A, et al. Meningiomas related to the Chernobyl irradiation disaster in North-Eastern Romania between 1990 and 2015. Revista de Chimie Bucharest. 2018; 69:1562-1565.
- Cucu AI, Turliuc MD, Carauleanu A, et al. Chemical aspects of peritumoral cerebral edema in atypical meningiomas. Rev Chim (Bucharest). 2018; 69:2804-2807.
- Eom KS, Kim DW, Kim TY. Diffuse craniospinal metastases of intraventricular rhabdoid papillary meningioma with glial fibrillary acidic protein expression: a case report. Clin Neurol Neurosurg. 2009; 111(7):619-623.
- Figueroa BE, Quint DJ, McKeever PE, Chandler WF. Extracranial met-astatic meningioma. Br J Radiol. 1999; 72:513-516.
- Ather Enam S, Abdulrauf S, Mehta B, et al. Metastasis in meningioma. Acta Neurochir. 1996; 138: 1172-1178, discussion 1177-1178.
- Gezen F, Kahraman S, Canakci Z, Bedük A. Review of 36 cases of spinal cord meningioma. Spine (Phila Pa 1976). 2000; 25: 727-731.
- Scognamiglio G, D'Antonio A, Rossi G, et al. CD90 expression in atypical meningiomas and meningioma metastasis. Am J Clin Pathol. 2014; 141:841-849.
- Forest F, Berremila SA, Gyenes C, et al. Metastatic meningiomas: an unusual clinical and pathological diagnosis with highly variable outcome. J Neurooncol. 2014; 120(2): 411-421.
- Cucu AI, Costea CF, Turliuc MD, et al. Mirror, mirror on the wall, who's the fairest of them all? Atypical meningioma associated with multiple meningiomas. Romanian Neurosurgery. 2018; 32:563-572.
- Cucu AI, Turliuc MD, Costea CF, et al. Atypical meningiomas of cerebellopontine angle. A five case series. Rev Med Chir Soc Med Nat Iasi. 2018; 122: 533-545.

14. Gordon A, Maloney A. A case of metastasizing meningioma. *J Neurol Neurosurg Psychiatry*. 1965; 28:159-162.
15. Kabus D, Sidhu GS, Wiczorek RL, Choi HS. Metastatic meningioma. Hemangiopericytoma or angioblastic meningioma? *Am J Surg Pathol*. 1993; 17:1144-1150.
16. Cucu AI, Turliuc DM, Costea CF, et al. Pathways of metastatic spread in meningiomas. *Romanian Neurosurgery*. 2019; 33(1):12-16.
17. Akimura T, Orita T, Hayashida O, et al. Malignant meningioma metastasizing through the cerebrospinal pathway. *Acta Neurol Scand*. 1992; 85(5):368-371.
18. Satoh T, Kageyama T, Yoshimoto Y, et al. Intra-thecal dissemination of meningiomas; a case report. *No Shinkei Geka*. 1992; 20:805-808.
19. Noterman J, Depierreux M, Raftopoulos C, Brotchi J. Metastases of meningioma. Apropos of 2 cases. *Neurochirurgie*. 1987; 33:184-189.
20. Kleinschmidt-DeMasters BK, Avakian JJ. Wallenberg syndrome caused by CSF metastasis from malignant intraventricular meningioma. *Clin Neuropathol*. 1985; 4:214-219.
21. Chamberlain MC, Glantz MJ. Cerebrospinal fluid-disseminated meningioma. *Cancer*. 2005; 103(7):1427-1430.
22. Park KS, Kim KH, Park SH, et al. Intracranial meningioma with leptomeningeal dissemination: retrospective study with review of the literature. *J Korean Neurosurg Soc*. 2015; 57(4):258-265.
23. Ludwin SK, Rubinstein LJ, Russell DS. Papillary meningioma: a malignant variant of meningioma. *Cancer*. 1975; 36(4):1363-1373.
24. Cantillo R, Jain J, Singhakowinta A, Vaitkevicius V. Blindness as initial manifestation of meningeal carcinomatosis in breast cancer. *Cancer*. 1979; 44:755-757.
25. Eberth C. Zur entwickl pung des ephheliomas (cholesteatomas) der pia and der lung. *Virchow S Arch*. 1870; 49:51-63.
26. Dufour H. Meningite sarcomateuse diffuse avec envahissement de las melle et das racines: cytologie positive et speciale du liquide cephalorachidien. *Rev Neurol*. 1904; 12:104-106.
27. Berman WF. Meningeal carcinomatosis. *JAMA*. 1912; 58:1437-1439.
28. Kalm H. Ein malignes Tentoriummeningeom mit Metastasierung in die Medulla oblongata und in die subarachnoidalen Liquorräume. *Deutsche Zeitschrift für Nervenheilkunde*. 1950, 163:131-140.
29. DeAngelis LM. Current diagnosis and treatment of leptomeningeal metastasis. *J Neurooncol*. 1998; 38(2-3):245-252.
30. Fulkerson DH, Horner TG, Hattab EM. Histologically benign intraventricular meningioma with concurrent pulmonary metastasis: case report and review of the literature. *Clin Neurol Neurosurg*. 2008; 110(4):416-419.
31. Rawat B, Franchetto AA, Elavathil J. Extracranial metastases of meningioma. *Neuroradiology*. 1995; 37:38-41.
32. Gyepes MT, DAngio GJ. Extracranial metastases from central nervous system tumors in children and adolescents. *Radiology*. 1966; 87:55-63.
33. Christensen E, Kiaer W, Winblad S. Meningeal turnouts with extracerebral metastases. *Br J Cancer*. 1949; 3:485-493.
34. Wilkenman NW Jr., Cassell C, Schlesinger B. Intracranial tumors with extracranial metastases. *J Neuropathol Exp Neurol*. 1952; 11:149-168.
35. Ramakrishnamurthy TV, Murty AV, Purohit AK, Sundaram C. Benign meningioma metastasizing through CSF pathways: a case report and review of literature. *Neurol India*. 2002; 50:326-329.
36. Koenig MA, Geocadin RG, Kulesza P, et al. Rhabdoid meningioma occurring in an unrelated resection cavity with leptomeningeal carcinomatosis. Case report. *J Neurosurg*. 2005; 102(2):371-375.
37. Morantz RA, Shain W. Trauma and brain tumors: an experimental study. *Neurosurgery*. 1978; 3(2):181-186.
38. Wippold II FJ, Smirnoitopoulos JG, Pilgram TK. Lesions of the cauda equina: a clinical pathology review from the Armed Forces Institute of Pathology. *Clin Neurol Neurosurg*. 1997; 9:229-234.
39. Peh WC, Fan YW. Case report: intraventricular meningioma with cerebellopontine angle and drop metastases. *Br J Radiol*. 1995; 68(808): 428-430.
40. Solero CL, Fornari M, Giombini S, et al. Spinal meningiomas: review of 174 operated cases. *Neurosurgery*. 1989; 125:153-160.
41. Lee TT, Landy HJ. Spinal metastases of malignant intracranial meningioma. *Surg Neurol*. 1998; 50(5): 437-441.
42. Russel DS, Rubinstein LJ. Pathology of tumors of the nervous system, 3rd. London: Edward Arnold, 1971.
43. Tsuda K, Akutsu H, Yamamoto T, et al. Benign spinal meningioma without dural attachment presenting delayed CSF dissemination and malignant transformation. *Brain Tumor Pathol*. 2013; 30(3):185-191.
44. de Vries J, Wakhloo AK. Repeated multifocal recurrence of grade I, grade II, and grade III meningiomas: regional multicentricity (primary new growth) or metastases? *Surg Neurol*. 1994; 41(4):299-305.
45. Miller DC, Ojemann RG, Proppe KH, et al. Benign metastasizing meningioma. *J Neurosurg*. 1985; 62:763-766.
46. Clarke JL, Perez HR, Jacks LM, et al. Leptomeningeal metastases in the MRI era. *Neurology*. 2010; 74:1449-1454.
47. Fujimaki M, Takanashi M, Kobayashi M, et al. Cerebrospinal fluid dissemination of anaplastic intraventricular meningioma: report of a case presenting with progressive

- brainstem dysfunction and multiple cranial nerve palsies. *BMC Neurol.* 2016; 16:82.
48. Tao CY, Wang JJ, Li H, You C. Malignant intraventricular meningioma with craniospinal dissemination and concurrent pulmonary metastasis. *World J Surg Oncol.* 2014; 12:238.
 49. Maria GP, Teodoro S, Jenny F, et al. Leptomeningeal metastasis from solid tumors: a diagnostic and therapeutic challenge. *Neurol Sci.* 2015; 36:117-123.
 50. Toyooka T, Miyazawa T, Fukui S, et al. Central neurogenic hyperventilation in a conscious man with CSF dissemination from a pineal glioblastoma. *J Clin Neurosci.* 2005; 12:834-837.
 51. William RW, J Peter G, Jerome BP. Diagnosis and Treatment of Leptomeningeal Metastases from Solid Tumors: Experience with 90 Patients. *Cancer.* 1982; 49:759-772.
 52. Engelhard HH, Corsten LA. Leptomeningeal metastasis of primary Central Nervous System (CNS) Neoplasm. In: Abrey LE, Chamberlain MC, Engelhard HH (Eds.). *Leptomeningeal metastases. Cancer Treatment and Research.* Vol 125. Boston: Springer, 2005.
 53. Miller AA, Ramsden F. Malignant meningioma with extracranial metastases and seeding of the subarachnoid space and the ventricles. *Pathol Eur.* 1972; 7:167-175.
 54. Erman T, Hanta I, Hacıyakupolu S, et al. Huge bilateral pulmonary and pleural metastasis from intracranial meningioma: a case report and review of the literature. *J Neurooncol.* 2005; 74:179-181.
 55. Aumann J, Vandenbosch J, Elbers J, Wagenaar J. Metastatic meningioma of the lung. *Thorax.* 1986; 41:487-488.
 56. Kovoor J, Jayakumar PN, Srikanth SG, et al. Solitary pulmonary metastasis from intracranial meningioma. *Australas Radiol.* 2002; 46:65-68.
 57. Som PM, Sacher M, Strenger SW, et al. "Benign" metastasizing meningiomas. *AJNR.* 1987; 8:127-130.
 58. Cerda-Nicolas M, Lopez-Givens C, Perez-Bacete M, et al. Histologically benign metastatic meningioma: morphological and cytogenetic study. *J Neurosurg.* 2003; 98:194-198.
 59. Kaminski J, Movsas B, King E, et al. Metastatic meningioma to the lung with multiple pleural metastases. *Am J Clin Oncol.* 2001; 24:579-582.
 60. Cramer P, Thomale UW, Okuducu AF, et al. An atypical spinal meningioma with CSF metastasis: Fatal progression despite aggressive treatment. Case report. *J Neurosurg Spine.* 2005; 3:153-158.
 61. Wu YT, Ho JT, Lin YJ, Lin JW. Rhabdoid papillary meningioma: a clinicopathologic case series study. *Neuropathology.* 2011; 31:599-605.
 62. Kim JP, Park BJ, Lim YJ. Papillary meningioma with leptomeningeal seeding. *J Korean Neurosurg Soc.* 2011; 49:124-127.
 63. Peng J, Liang ZG, Li KC. Intracranial malignant meningioma with cerebrospinal fluid dissemination: a case report. *Chin Med J (Engl).* 2011; 124:1597-1599.
 64. Kuroda H, Kashimura H, Ogasawara K, et al. Malignant intracranial meningioma with spinal metastasis—case report. *Neurol Med Chir (Tokyo).* 2009; 49:258-261.
 65. Erkutlu I, Buyukhatipoglu H, Alptekin M, et al. Spinal drop metastases from a papillary meningioma : a case report and review of the literature : utility of CSF sampling. *Med Oncol.* 2009; 26:242-246.
 66. Santhosh K, Kesavadas C, Radhakrishnan VV, et al. Rhabdoid and papillary meningioma with lepto-meningeal dissemination. *J Neuroradiol.* 2008; 35:236-239.
 67. Shintaku M, Hashimoto K, Okamoto S. Intraventricular meningioma with anaplastic transformation and metastasis via the cerebrospinal fluid. *Neuropathology.* 2007; 27:448-452.
 68. Chuang HC, Lee HC, Cho DY. Intracranial malignant meningioma with multiple spinal metastases—a case report and literature review: case report. *Spine (Phila Pa 1976).* 2006; 31:E1006-E1010.
 69. Al-Habib A, Lach B, Al Khani A. Intracerebral rhabdoid and papillary meningioma with leptomeningeal spread and rapid clinical progression. *Clin Neuropathol.* 2005; 24:1-7.
 70. Wakabayashi K, Suzuki N, Mori F, et al. Rhabdoid cystic papillary meningioma with diffuse subarachnoid dissemination. *Acta Neuropathol.* 2005; 110:196-198.
 71. Darwish B, Munro I, Boet R, et al. Intraventricular meningioma with drop metastases and subgaleal metastatic nodule. *J Clin Neurosci.* 2004; 11:787-791.
 72. Conrad MD, Schonauer C, Pelissou-Guyotat I, et al. Recurrent lumbosacral metastases from intracranial meningioma. Report of a case and review of the literature. *Acta Neurochir (Wien).* 2001; 143: 935-937.
 73. Meinsma-vdTuin M, Molenaar WM, Mooij JJ. Spinal papillary meningioma: a case report and review of the literature. *Acta Neurochir (Wien).* 2000; 142:703-708.
 74. Lee WH, Chen A, Chao DG, et al. Malignant meningioma with rhabdoid transformation. *Zhonghua Yi Xue Za Zhi (Taipei).* 2000; 63:492-497.
 75. Greenberg SB, Schneck MJ, Faerber EN, Kanev PM. Malignant meningioma in a child: CT and MR findings. *AJR Am J Roentgenol.* 1993; 160:1111-1112.
 76. Kamiya K, Inagawa T, Nagasako R. Malignant intraventricular meningioma with spinal metastasis through the cerebrospinal fluid. *Surg Neurol.* 1989; 32:213-218.
 77. Strenger SW, Huang YP, Sachdev VP. Malignant meningioma within the third ventricle: a case report. *Neurosurgery.* 1987; 20:465-468.
 78. Ludwin SK, Conley FK. Malignant meningioma metastasizing through the cerebrospinal pathways. *J Neurol Neurosurg Psychiatry.* 1975; 38:136-142.