

Showa Univ J Med Sci 25(3), 193~201, September 2013

Original

Effect of Hydrogen Peroxide and High Glucose on the Glucose Metabolism of Lymphoma-derived U937 Cells

Yasunori TAKEDA¹⁾, Yuya NAKAMURA^{1, 2)}, Masahiro INAGAKI³⁾,
Mayumi TSUJI¹⁾, Hitomi HASEGAWA¹⁾, Masanobu WATANABE¹⁾,
Hiromichi GOTOH²⁾, Yoshikazu GOTO²⁾ and Katsuji OGUCHI¹⁾

Abstract: Our study aimed to clarify specific oxidative stress and glucose metabolic disorders in hemodialysis patients, by examining hydrogen peroxide (H₂O₂)- and high glucose-induced oxidative stress, glucose transport and the failure of glycolysis. As an *in vitro* blood cell model of end-stage renal disease (ESRD) in patients with diabetes, human monocytic U937 cells of malignant lymphoma origin were exposed to high glucose (28.9 mM) for 6 days, with 5 mM H₂O₂ added on the last day. The generation of intracellular reactive oxygen species (ROS), glucose levels, lactate levels, AMP-activated protein kinase (AMPK) activity and Glut4 levels were examined. Exposure of U937 cells to H₂O₂ resulted in a significant increase in intracellular ROS generation and glucose levels. Under high glucose conditions, treatment with H₂O₂ significantly promoted these actions. In H₂O₂-induced U937 cells, AMPK activity and Glut4 levels were significantly increased, but lactate and pyruvate levels were significantly decreased. Thus, exposure of U937 cells to H₂O₂ and a high glucose load promoted an increase in intracellular ROS, and exposure to H₂O₂ induced increased glucose transport and high intracellular glucose due to reduced glycolytic metabolism. This suggests that reduced glycolytic metabolism might be induced in states of high oxidative stress in hemodialysis patients with diabetes.

Key words: glucose, hydrogen peroxide, lactate, oxidative stress, U937 cells

Introduction

Patients with diabetes mellitus (DM) undergoing hemodialysis are exposed to significant oxidative stress¹⁻⁶⁾. We previously reported increased Cu/Zn-superoxide dismutase (SOD) in the blood and upregulated Cu/Zn-SOD mRNA in the leukocytes of hemodialysis patients, as well as a correlation between leukocyte membrane lipid peroxidation and expression of Cu/Zn-SOD mRNA in hemodialysis patients; this causes oxidative stress which may also occur due to blood cell contact with the dialysis membrane^{7, 8)}. Dialysis-associated oxidative stress is associated with complications such as arteriosclerosis⁹⁻¹¹⁾, however few studies have investigated the effect of oxidative stress on glucose metabolism in dialysis patients. Sugar transport increases

¹⁾ Department of Pharmacology, Showa University School of Medicine, 1-5-8 Hatanodai, Shinagawa-ku, Tokyo 142-8555, Japan.

²⁾ Saiyu Soka Hospital.

³⁾ Department of Chemistry, College of Arts and Sciences, Showa University.

in response to depletion of the intracellular ATP supply, shifting to other intracellular glucose metabolic pathways, such as the polyol pathway from the glycolytic system, hexosamine pathway, pentose phosphate pathway, and protein kinase C pathway^{12, 13}), however many aspects of this shift are not well defined. Strict control of blood glucose is necessary in dialysis patients to improve prognosis and prevent myocardial infarction, cerebral infarction, and lower extremity arterial disease¹⁴⁻¹⁷). Therefore, given the potential changes in glycolytic metabolism due to the influence of dialysis on blood glucose, it is important to investigate the effect of oxidative stress on glycolytic metabolism.

In this *in vitro* study, we cultured malignant lymphoma-derived U937 cells in a high glucose medium to model end-stage renal disease (ESRD) associated with DM. We then induced oxidative stress by adding hydrogen peroxide (H_2O_2) to these cells and analyzed the effects on reactive oxygen species (ROS), glucose transport, and glucose metabolism.

Methods

Study materials and cell culture

Human monocytes (U937 cells, EC85011440) were purchased from the European Collection of Animal Cell Culture (ECACC, UK). The following reagents were also used in the study: H_2O_2 (Wako Co., Osaka, Japan); fetal bovine serum (FBS; Sigma Aldrich, St. Louis, MO); penicillin G sodium (Invitrogen, Carlsbad, CA); streptomycin sulfate (Invitrogen); and 5-(and-6)-chloromethyl-2', 7'-dichlorodihydrofluorescein diacetate (CM-H2DCF-DA; Molecular Probes, Eugene, OR). In order to treat the U937 cells with a high glucose load, cells were cultured in Roswell Park Memorial Institute medium (RPMI-1640) containing 28.9 mM glucose for 5 days (high glucose group), while the normal glucose group consisted of U937 cells cultured in standard RPMI-1640 medium containing 11.1 mM D-glucose. Previous experimental studies showed no dead cells due to apoptosis when the mannitol osmotic pressure was adjusted to equal the osmotic pressure of the glucose (30 mM)¹⁸). U937 cells were cultured by routine methods with 10% FBS, 100 units/mL of penicillin G sodium, and 100 μ g/mL of streptomycin sulfate, at 37°C and 5% CO_2 .

Drug treatment

U937 cells in the normal and high glucose groups (two groups each) were cultured for 5 days, and each medium was replaced with or without H_2O_2 (5 mM), and then cells were incubated for 24 h, after that all the parameters were measured. The cultural experiment was carried out under aseptic conditions.

Evaluation of ROS

CM-H2DCFDA was used to evaluate intracellular ROS formation in U937 cells treated with H_2O_2 (5 mM). After cellular uptake, CM-H2DCFDA is first modified to 2',7'-dichlorofluorescein by intracellular esterases, and then by intracellular ROS to create oxidized fluorescent 2', 7'-dichlorofluorescein¹⁹). After dissolving CM-H2DCFDA with dimethyl sulfoxide, 7 μ M

CM-H2DCFDA was added to the cells (3×10^6 cells / mL) and incubated for 15 min at 37°C, 5% CO₂. After treatment, the cells were washed with heated phosphate-buffered saline. Fluorescence intensity was measured on a plate reader (Berthold Technologies, Bad Wildbad, Germany) at an excitation wavelength of 488 nm and a measuring wavelength of 525 nm.

Glucose uptake and glycolysis

1. Intracellular glucose

Intracellular glucose concentrations in U937 cells after 5 mM H₂O₂ treatment were measured by high performance liquid chromatography with electrochemical detection (HPLC-ECD), using a separation column ACT-gelPAC ($\phi 4.6 \times 150$ mm, Act-Science, Tokyo, Japan) and a glucose oxidase-stabilized enzyme-immobilized column GO-ENZ ($\phi 4.0 \times 5$ mm, Eicom Co., Kyoto, Japan). Endogenous H₂O₂ and glucose-derived H₂O₂ were separated with 100 mM phosphate buffer solution (pH 6.5) in the mobile phase.

2. AMP-activated protein kinase (AMPK)

AMPK was measured with an AMPK Kinase Assay Kit (CycLex, Nagano, Japan). Extracted samples were added to a microplate coated with AMPK substrate consisting of mouse Insulin Receptor substrate-1 (IRS-1) serine 789-peptide, which induced the phosphoric acid reaction. After detection using phosphorylated IRS-1 mouse monoclonal antibody and horseradish peroxidase-labeled anti-mouse IgG, the sample was stained using tetra-methylbenzidine and measured at 450 nm on a plate reader (Berthold Technologies). Fluorescence intensity was measured for each protein.

3. Glut4

Glut4 was measured using an SLC2A4 (Human) ELISA kit (Abnova, Walnut, CA). Extracted samples and Glut4 were treated with specific biotin-conjugated polyclonal antibody and avidin-conjugated horseradish peroxidase, and then placed in a microplate. After incubation, tetra-methylbenzidine substrate was added, and the sample was measured at 450 nm.

4. Intracellular pyruvate

Intracellular pyruvate concentrations were measured with a Pyruvate Assay Kit (BioVision Research Products, Milpitas, CA). Extracted samples were mixed with 50 μ L Reaction Mix (Pyruvate Assay Buffer 46 μ L, Pyruvate Probe 2 μ L, Enzyme Mix 2 μ L), incubated in the dark for 30 min, and then measured at 570 nm on a plate reader (Berthold Technologies). The fluorescence intensity was measured for each protein.

5. Intracellular lactic acid

Intracellular lactic acid concentrations were evaluated in a series using an LT-GEL separation column ($\phi 4.6 \times 50$ mm; Eicom) and a lactic acid oxidase-immobilized enzyme-immobilized column LT-ENZ ($\phi 4.0 \times 5$ mm; Eicom) and measured by HPLC-ECD. Endogenous H₂O₂ and lactic acid-derived H₂O₂ were separated using 100 mM phosphate buffer solution containing 200 mg tetra-n-hexylammonium bromide / L (pH 7.0).




Fig. 1. Effect of hydrogen peroxide (H₂O₂) and high glucose on intracellular reactive oxygen species (ROS) production. U937 cells were cultured for 6 days in RPMI-1640 medium containing either 11.1 mM D-glucose (Control) or 28.9 mM glucose (High glucose). For the last 24 h, 5 mM H₂O₂ was added to the control group (5 mM H₂O₂) and the high glucose group (High glucose + 5 mM H₂O₂). **P* < 0.05 vs. Control; #*P* < 0.05 vs. 5 mM H₂O₂; ▼*P* < 0.05 vs. High glucose.

Statistical Analysis

Results are presented as means + standard error. Dunnett's tests were used, and statistical significance was set at *P* < 0.05.

Results

Effect of H₂O₂ and high glucose on intracellular ROS production

Compared with untreated cells, ROS production was significantly higher (2-fold) in the normal and high glucose groups treated with 5 mM H₂O₂. Compared with the normal glucose group, ROS production in the high glucose group was significantly higher (1.4-fold), confirming that H₂O₂ increases ROS production (Fig. 1).

Intracellular glucose uptake (glucose transport mechanism) and glycolysis

1. Effect of H₂O₂ and high glucose on intracellular glucose

Compared with untreated cells, intracellular glucose was significantly higher (8-fold) in H₂O₂-treated cells. Intracellular glucose was 2-fold higher in the high glucose group than in the normal group (*P* < 0.05; Fig. 2a), thus validating the H₂O₂-associated increase in intracellular glucose.

2. Effect of H₂O₂ and high glucose on AMPK production

AMPK production was significantly higher (1.3-fold) in H₂O₂-treated cells than in untreated cells, however there were no significant differences between the normal and high glucose groups (Fig. 2b).

3. Effect of H₂O₂ and high glucose on Glut4 production

Compared with untreated cells, Glut4 production was significantly higher (1.5-fold) in cells from the normal glucose group treated with 5 mM H₂O₂. Glut4 production tended to be higher




Fig. 2.

- (a) Effect of hydrogen peroxide (H₂O₂) and high glucose on intracellular glucose. U937 cells were cultured for 6 days in RPMI-1640 medium containing either 11.1 mM D-glucose (Control) or 28.9 mM glucose (High glucose). For the last 24 h, 5 mM H₂O₂ was added to the control group (5 mM H₂O₂) and the high glucose group (High glucose + 5 mM H₂O₂). **P* < 0.05 vs. Control; #*P* < 0.05 vs. 5 mM H₂O₂; ▼*P* < 0.05 vs. High glucose.
- (b) Effect of hydrogen peroxide (H₂O₂) and high glucose on AMP-activated protein kinase (AMPK) production. U937 cells were cultured for 6 days in RPMI-1640 medium containing either 11.1 mM D-glucose (Control) or 28.9 mM glucose (High glucose). For the last 24 h, 5 mM H₂O₂ was added to the control group (5 mM H₂O₂) and the high glucose group (High glucose + 5 mM H₂O₂). **P* < 0.05 vs. Control; ▼*P* < 0.05 vs. High glucose.
- (c) Effect of hydrogen peroxide (H₂O₂) and high glucose on Glut4 production. U937 cells were cultured for 6 days in RPMI-1640 medium containing either 11.1 mM D-glucose (Control) or 28.9 mM glucose (High glucose). For the last 24 h, 5 mM H₂O₂ was added to the control group (5 mM H₂O₂) and the high glucose group (High glucose + 5 mM H₂O₂). **P* < 0.05 vs. Control.
- (d) Effect of hydrogen peroxide (H₂O₂) and high glucose on intracellular pyruvic acid concentration. U937 cells were cultured for 6 days in RPMI-1640 medium containing either 11.1 mM D-glucose (Control) or 28.9 mM glucose (High glucose). For the last 24 h, 5 mM H₂O₂ was added to the control group (5 mM H₂O₂) and the high glucose group (High glucose + 5 mM H₂O₂). **P* < 0.05 vs. Control; ▼*P* < 0.05 vs. High glucose.
- (e) Effect of hydrogen peroxide (H₂O₂) and high glucose on intracellular lactic acid concentration. U937 cells were cultured for 6 days in RPMI-1640 medium containing either 11.1 mM D-glucose (Control) or 28.9 mM glucose (High glucose). For the last 24 h, 5 mM H₂O₂ was added to the control group (5 mM H₂O₂) and the high glucose group (High glucose + 5 mM H₂O₂). **P* < 0.05 vs. Control; ▼*P* < 0.05 vs. High glucose.

in cells treated with 5 mM H₂O₂, but the change was not significant. Comparison of the normal and high glucose groups revealed no significant changes (Fig. 2c).

4. Effect of H₂O₂ and high glucose on intracellular pyruvic acid

Treatment with 5 mM H₂O₂ significantly reduced intracellular pyruvic acid concentrations in cells from both the normal and high glucose groups, compared to untreated cells. There was no significant difference between groups (Fig. 2d).

5. Effect of H₂O₂ and high glucose on intracellular lactic acid

Compared with untreated cells, the amount of intracellular lactic acid was significantly reduced in cells of the normal and high glucose groups treated with 5 mM H₂O₂. Intracellular lactic acid

was 1.5-fold higher than the control in the high glucose group ($P < 0.05$; Fig. 2e).

Discussion

This *in vitro* study using U937 cells revealed that intracellular ROS production was 2-fold higher after H₂O₂ treatment, whereas high glucose treatment caused a 1.4-fold increase in intracellular ROS, compared to cells treated with normal glucose levels. Previously, Sakai and colleagues²⁰⁾ cultured human and mouse pancreatic β cells in a high glucose medium (30 mM glucose) for 4 days, and found that intracellular ROS production was 4 times higher than in cells cultured with 5 mM glucose. In this study, increased ROS production due to H₂O₂ was augmented by a high glucose load. Malignant lymphoma-derived U937 cells provide an *in vitro* model of blood cells for DM-associated hemodialysis, raising the possibility that high oxidative stress during treatment is worsened in patients with high blood glucose. This *in vitro* study also proves the importance of strict blood glucose control in order to reduce oxidative stress in hemodialysis patients with various complications such as arteriosclerosis⁹⁻¹¹⁾.

AMPK is activated when intracellular ATP levels decrease; the enzyme phosphorylates members of the acetyl-CoA carboxylase metabolic pathway²¹⁾ and mammalian target of rapamycin²²⁾. Adjustment of these enzyme activities and the activity of transcription factors²³⁾ and cofactors²⁴⁾ modulates intracellular energy metabolism. Glut4 is a glucose transporter belonging to class I of the Glut family and mediates activation of AMPK and insulin. Glut4 promotes transcription and translocation to the plasma membrane, and is also involved in promoting glucose uptake. Horie *et al* treated rat cardiomyocytes with H₂O₂ and measured the time- and concentration-dependent effects. Their study showed that a long duration and high concentration of H₂O₂ increased the time required for Glut4 to travel to the cell membrane. Delayed Glut4 transport to the cell membrane in AMPK-deficient cells treated with 100 μ M H₂O₂ (25 vs. 15 min for control) demonstrated the importance of AMPK in Glut4 translocation to the cell membrane after H₂O₂ treatment²⁵⁾. In our study, AMPK was also significantly higher (1.3-fold) in cells treated with 5 mM H₂O₂ for 24 h, while Glut4 was approximately 1.5-fold higher. Thus, reduction of either enzyme of glycolytic metabolism reduces intracellular ATP. Furthermore, inhibition of glycolytic metabolism is suspected to inhibit activation of glyceraldehyde-3-phosphate dehydrogenase (GAPDH) due to H₂O₂ load^{20, 26)}. We also demonstrated that production of lactic acid was suppressed by approximately 70%, whereas intracellular glucose increased 8-fold after H₂O₂ exposure. Decreased glycolytic metabolism due to H₂O₂ was studied by Sakai *et al*²⁰⁾, who treated mouse pancreatic β cells with 50 μ M H₂O₂ for 30 min and found an approximate 50% reduction in pyruvic acid, whereas the amount of lactic acid remained unchanged. In contrast, Colussi *et al*²⁶⁾ treated U937 cells with 1 mM H₂O₂ for 1 h and found that intracellular glucose consumption was 97%, whereas lactic acid production was suppressed by 96%. Our study also observed these reductions, unlike the study by Sakai *et al*²⁰⁾ which found no such changes. This discrepancy may be due to differences in cell type, H₂O₂ concentration, or duration of administration. Our study and that of Colussi *et al*²⁶⁾ used U937 cells, but in well-differentiated cancer cells such as these, even under oxygenated conditions,

lactic acid is generated by translocation of lactate dehydrogenase from pyruvic acid produced by glycolysis, in what is known as the Warburg effect²⁷⁻²⁹). However, if glycolytic metabolism is strongly suppressed by H₂O₂, inhibition of lactic acid production is stronger in cancer cells, such as U937 cells with their initially high production of lactic acid, compared with other cells.

Mitochondria-derived ROS production increases under high glucose conditions, and then, resulting in inhibition of the glycolytic metabolic enzyme, GADPH. Additionally, ROS cause DNA damage, leading to activation of poly (ADP-ribose) polymerase¹²). After culturing mouse pancreatic β cells in 25 mM glucose for 2 days, Sakai and colleagues found that GADPH activity was reduced by 36% compared with cells incubated in 5 mM glucose²⁰). In addition, Du *et al*¹³) cultured bovine endothelial cells in 30 mM glucose for 2 days and found that GADPH activity was reduced by 66% compared with cells treated with 5 mM glucose. In these studies, GADPH activation was reduced by a high glucose load, but reduced activation of GADPH usually leads to a reduction in intracellular ATP. However, in our study, despite the increase in intracellular ROS production due to high glucose, the amounts of pyruvic acid and lactic acid were not reduced, suggesting no ATP reduction-dependent increase in AMPK or Glut4. Glycolysis downstream of pyruvic acid is maintained, as is intracellular ATP. The type of ROS produced might be different under conditions of high glucose load and hydrogen peroxide load, or it might be the unique property of cancer cells. Advanced glycation end products (AGE) can serve as an indicator for changes in GADPH activation due to high glucose load. High glucose induces various intracellular glucose pathways, including the glycolytic system by reducing GADPH activity via mitochondria-derived ROS production, activation of the hexosamine pathway, activation of protein kinase C, increased production of AGE, and activation of the polyol pathway¹²). We previously reported an increase of AGE associated with a high glucose load³⁰), indicating the possibility of reduced GADPH activation.

As an *in vitro* blood cell model of DM-associated ESRD, this study has a number of limitations. Firstly, our use of malignant lymphoma-derived cancer cells as the study material, and secondly, replacing the mechanical stress of contact between blood cells and the dialysis membrane with an H₂O₂ load. A third limitation of the study is the inability to consider changes in glucose metabolism and glucose transport mechanisms due to the change in H₂O₂ concentration. Because blood cells come into contact with the hemodialysis membrane during dialysis treatment, the use of monocytic line-derived cells is a valid *in vitro* model of blood cells during hemodialysis, but there is the possibility of metabolic differences, despite the absence of the Warburg effect²⁷⁻²⁹). These results do not mean that mechanical stress influences ROS and glycolytic metabolism. However, despite these limitations, this model can serve as a useful *in vitro* screening tool for DM-associated ESRD.

In conclusion, this study demonstrated an increase of intracellular ROS due to H₂O₂ and high glucose load, and indicated an increase in glucose transport with reduced glycolytic metabolism due to H₂O₂. Reduced glycolytic metabolism might be induced in states of high oxidative stress in hemodialysis patients with DM.

Conflict of interest

The authors have declared no conflict of interest.

References

- 1) Pawlak K, Pawlak D, Mysliwiec M. Cu / Zn superoxide dismutase plasma levels as a new useful clinical biomarker of oxidative stress in patients with end-stage renal disease. *Clin Biochem.* 2005;**38**:700–705.
- 2) Nakayama H, Akiyama S, Shiotani S, *et al.* Evaluation of superoxide dismutase activity in dialyzed patients by electron spin resonance spectroscopy. *Am J Nephrol.* 2002;**22**:6–10.
- 3) Bonomini M, Stuard S, Carreno MP, *et al.* Neutrophil reactive oxygen species production during hemodialysis: role of activated platelet adhesion to neutrophils through P-selectin. *Nephron.* 1997;**75**:402–411.
- 4) Descamps-Latscha B, Goldfarb B, Nguyen AT, *et al.* Establishing the relationship between complement activation and stimulation of phagocyte oxidative metabolism in hemodialyzed patients: a randomized prospective study. *Nephron.* 1991;**59**:279–285.
- 5) Maher ER, Wickens DG, Griffin JF, *et al.* Increased free-radical activity during haemodialysis? *Nephrol Dial Transplant.* 1987;**2**:169–171.
- 6) Nguyen AT, Lethias C, Zingraff J, *et al.* Hemodialysis membrane-induced activation of phagocyte oxidative metabolism detected in vivo and in vitro within microamounts of whole blood. *Kidney Int.* 1985;**28**:158–167.
- 7) Akiyama S, Inagaki M, Tsuji M, *et al.* mRNA study on Cu / Zn superoxide dismutase induction by hemodialysis treatment. *Nephron Clin Pract.* 2005;**99**:c107–c114.
- 8) Washio K, Inagaki M, Tsuji M, *et al.* Correlation between leukocyte membrane lipid peroxidation and expression of Cu / Zn-superoxide dismutase mRNA in hemodialysis patients. *Blood Purif.* 2012;**33**:59–65.
- 9) Taki K, Takayama F, Tsuruta Y, *et al.* Oxidative stress, advanced glycation end product, and coronary artery calcification in hemodialysis patients. *Kidney Int.* 2006;**70**:218–224.
- 10) Locatelli F, Canaud B, Eckardt KU, *et al.* Oxidative stress in end-stage renal disease: an emerging threat to patient outcome. *Nephrol Dial Transplant.* 2003;**18**:1272–1280.
- 11) Canaud B, Cristol J, Morena M, *et al.* Imbalance of oxidants and antioxidants in haemodialysis patients. *Blood Purif.* 1999;**17**:99–106.
- 12) Brownlee M. The pathobiology of diabetic complications: a unifying mechanism. *Diabetes.* 2005;**54**:1615–1625.
- 13) Du XL, Edelstein D, Rossetti L, *et al.* Hyperglycemia-induced mitochondrial superoxide overproduction activates the hexosamine pathway and induces plasminogen activator inhibitor-1 expression by increasing Sp1 glycosylation. *Proc Natl Acad Sci U S A.* 2000;**97**:12222–12226.
- 14) Hayashino Y, Fukuhara S, Akiba T, *et al.* Diabetes, glycaemic control and mortality risk in patients on haemodialysis: the Japan Dialysis Outcomes and Practice Pattern Study. *Diabetologia.* 2007;**50**:1170–1177.
- 15) Oomichi T, Emoto M, Tabata T, *et al.* Impact of glycemic control on survival of diabetic patients on chronic regular hemodialysis: a 7-year observational study. *Diabetes Care.* 2006;**29**:1496–1500.
- 16) Morioka T, Emoto M, Tabata T, *et al.* Glycemic control is a predictor of survival for diabetic patients on hemodialysis. *Diabetes Care.* 2001;**24**:909–913.
- 17) Wu MS, Yu CC, Yang CW, *et al.* Poor pre-dialysis glycaemic control is a predictor of mortality in type II diabetic patients on maintenance haemodialysis. *Nephrol Dial Transplant.* 1997;**12**:2105–2110.
- 18) Leal EC, Aveleira CA, Castilho AF, *et al.* High glucose and oxidative / nitrosative stress conditions induce apoptosis in retinal endothelial cells by a caspase-independent pathway. *Exp Eye Res.* 2009;**88**:983–991.
- 19) Afzal M, Matsugo S, Sasai M, *et al.* Method to overcome photoreaction, a serious drawback to the use of dichlorofluorescein in evaluation of reactive oxygen species. *Biochem Biophys Res Commun.* 2003;**304**:619–624.
- 20) Sakai K, Matsumoto K, Nishikawa T, *et al.* Mitochondrial reactive oxygen species reduce insulin secretion by

- pancreatic beta-cells. *Biochem Biophys Res Commun.* 2003;**300**:216–222.
- 21) Sim AT, Hardie DG. The low activity of acetyl-CoA carboxylase in basal and glucagon-stimulated hepatocytes is due to phosphorylation by the AMP-activated protein kinase and not cyclic AMP-dependent protein kinase. *FEBS Lett.* 1988;**233**:294–298.
 - 22) Bolster DR, Crozier SJ, Kimball SR, *et al.* AMP-activated protein kinase suppresses protein synthesis in rat skeletal muscle through down-regulated mammalian target of rapamycin (mTOR) signaling. *J Biol Chem.* 2002;**277**:23977–23980.
 - 23) Kawaguchi T, Osatomi K, Yamashita H, *et al.* Mechanism for fatty acid “sparing” effect on glucose-induced transcription: regulation of carbohydrate-responsive element-binding protein by AMP-activated protein kinase. *J Biol Chem.* 2001;**277**:3829–3835.
 - 24) Yang W, Hong YH, Shen XQ, *et al.* Regulation of transcription by AMP-activated protein kinase: phosphorylation of p300 blocks its interaction with nuclear receptors. *J Biol Chem.* 2001;**276**:38341–38344.
 - 25) Horie T, Ono K, Nagao K, *et al.* Oxidative stress induces GLUT4 translocation by activation of PI3-K / Akt and dual AMPK kinase in cardiac myocytes. *J Cell Physiol.* 2008;**215**:733–742.
 - 26) Colussi C, Albertini MC, Coppola S, *et al.* H₂O₂-induced block of glycolysis as an active ADP-ribosylation reaction protecting cells from apoptosis. *FASEB J.* 2000;**14**:2266–2276.
 - 27) Gogvadze V, Zhivotovsky B, Orrenius S. The Warburg effect and mitochondrial stability in cancer cells. *Mol Aspects Med.* 2010;**31**:60–74.
 - 28) Samudio I, Fiegl M, Andreeff M. Mitochondrial uncoupling and the Warburg effect: molecular basis for the reprogramming of cancer cell metabolism. *Cancer Res.* 2009;**69**:2163–2166.
 - 29) Vander Heiden MG, Cantley LC, Thompson CB. Understanding the Warburg effect: the metabolic requirements of cell proliferation. *Science.* 2009;**324**:1029–1033.
 - 30) Oishi S, Tsuji M, Hasegawa H, *et al.* High glucose promotes oxidative stress-induced apoptosis via a caspase-independent pathway in human monocytic U937 cells. *J Showa Med Assoc.* 2010;**70**:315–325. (in Japanese).

[Received January 25, 2013 : Accepted February 19, 2013]