



PHYSICAL FITNESS LEVELS AMONG 200, 500 AND 1000 METER CANOEING PLAYERS: A COMPARATIVE STUDY

**Baljinder Singh Bal¹,
Bhupinder Singh²,
Gurpinder Singh³,
Lovepreet Singh⁴**

¹Dr., Department of Physical Education (T),
Guru Nanak Dev University, Amritsar,
Punjab, India

^{2, 3 & 4}Department of Physical Education (T),
Guru Nanak Dev University, Amritsar,
Punjab, India

Abstract:

The purpose of this study was to compare Physical Fitness levels among 200, 500 and 1000 Meter Canoeing Players. To obtain data, the investigators had selected Twenty one (N = 21) male Canoeing Player of 19 to 25 years (Mean \pm SD: Age: 19.761 \pm 2.488 years; Body Mass: 1.749 \pm 7.046 kilograms; Body Height: 65.047 \pm 9.286 meters) of age to act as subjects. Components of Physical fitness (i.e., Agility, Balance, Coordination, Power, Reaction Time and Speed) were taken up for the present study. Statistical analysis was performed using the Statistical Package for the Social Sciences for Windows version 16.0 software (SPSS Inc., Chicago, IL). Data is expressed as the mean \pm SD. One way Analysis of Variance (ANOVA) was employed to find out the intra-group differences. To test the hypothesis, the level of significance was set at 0.05. No significant differences were found in Agility, Balance, Coordination, Power and Speed. However, significant differences were found in Reaction Time.

Keywords: canoeing, agility, balance, coordination, power, reaction time, speed

1. Introduction

Canoeing is a sport which involves endurance and strength [1]. Work intensity is varied, mostly high, sub-maximal and maximal [2]. Research conducted on canoeists [3] and kayakers [4, 5] contributed to the scientific definition of the required somatic model.

The previous studies showed that canoeists are characterized by very strong skeletal build, tallness, large body mass, long upper limbs, muscularity of the chest and

upper limbs and athletic build [6,7], having at the same time narrow hips and slim lower limbs [8]. Considering the potential role of physical characteristics, when examining paddlers using the ergometry system, all subjects overcome the same resistance in order to perform work, irrespective of body mass [9, 10].

Within particular sports, there exist various disciplines or playing positions with specific demands that require different approaches in training and are associated with different physical and morphological characteristics [11, 12]. Traditionally, the determination of a physical profile in a given sport involves the use of predictive testing as a measure of power and strength [13], speed [14], aerobic fitness [15] or flexibility [16].

2. Material and Methods

2.1 Selection of Subjects

To obtain data, the investigators had selected Twenty one (N = 21) male Canoeing Player of 19 to 25 years (Mean \pm SD: Age: 19.761 \pm 2.488 years; Body Mass: 1.749 \pm 7.046 kilograms; Body Height: 65.047 \pm 9.286 meters) of age to act as subjects. The data were collected from All India Open Canoe Sprint Championship organized by Sant Seechewal water sports centre Sultanpur Lodhi, Kapurthala, Punjab from 16-18 June, 2017.

All the subjects were informed about the objective and protocol of the study. Distribution and demographics of subjects are brought forth in Table 1.

Table 1: Distribution and demographics of subjects

Variables	Sample Size (N=21)
	Mean \pm SD
Age	19.761 \pm 2.488
Body Mass	1.749 \pm 7.046
Body Height	65.047 \pm 9.286




Figure 1: Distribution and demographics of subjects

2.2 Selection of Variables

The following components of Physical fitness were taken up for the present study:

- i. Agility
- ii. Balance
- iii. Coordination
- iv. Power
- v. Reaction Time
- vi. Speed

3. Statistical Analysis

Statistical analysis was performed using the Statistical Package for the Social Sciences for Windows version 16.0 software (SPSS Inc., Chicago, IL). Data is expressed as the mean \pm SD. One way Analysis of Variance (ANOVA) was employed to find out the intra-group differences. To test the hypothesis, the level of significance was set at 0.05.

4. Results

For each of the chosen variable, the result pertaining to components of Physical fitness of Canoeing Player are brought forth in the following tables:

Table 2: Analysis of variance (ANOVA) results among 200 Meter, 500 Meter and 1000 Meter men Canoeing players with regard to the sub-parameter Agility

Source of Variation	Sum of Squares	d.f.	Variance	F-value	p-value
Between Groups	1.2457	2	0.6229	F = 1.47408	.255352
Within Groups	7.6057	18	0.4225		
Total	8.8514	20			

The f-ratio value is 1.47408. The p-value is .255352. The result is not significant at $p < .05$.

It is concluded that the null hypothesis H_0 is not rejected. Therefore, there is not enough evidence to claim that not all 3 population means are equal, at the $\alpha=.05$ significance level.

Table 3: Analysis of variance (ANOVA) results among 200 Meter, 500 Meter and 1000 Meter men Canoeing players with regard to the sub-parameter Balance

Source of Variation	Sum of Squares	d.f.	Variance	F-value	p-value
Between-treatments	9.5238	2	4.7619	F = 0.12014	.887498
Within-treatments	713.4286	18	39.6349		
Total	722.9524	20			

The f-ratio value is 0.12014. The p-value is .887498. The result is not significant at $p < .05$.

It is concluded that the null hypothesis H_0 is not rejected. Therefore, there is not enough evidence to claim that not all 3 population means are equal, at the $\alpha=.05$ significance level.

Table 4: Analysis of variance (ANOVA) results among 200 Meter, 500 Meter and 1000 Meter men Canoeing players with regard to the sub-parameter Coordination

Source of Variation	Sum of Squares	d.f.	Variance	F-value	p-value
Between-treatments	82.5714	2	41.2857	F = 1.0342	.375699
Within-treatments	718.5714	18	39.9206		
Total	801.1429	20			

The f-ratio value is 1.0342. The p-value is .375699. The result is not significant at $p < .05$.

It is concluded that the null hypothesis H_0 is not rejected. Therefore, there is not enough evidence to claim that not all 3 population means are equal, at the $\alpha=.05$ significance level.

Table 5: Analysis of variance (ANOVA) results among 200 Meter, 500 Meter and 1000 Meter men Canoeing players with regard to the sub-parameter Power

Source of Variation	Sum of Squares	d.f.	Variance	F-value	p-value
Between-treatments	15.5238	2	7.7619	F = 1.04043	.373606
Within-treatments	134.2857	18	7.4603		
Total	149.8095	20			

The f-ratio value is 1.04043. The p-value is .373606. The result is not significant at $p < .05$.

It is concluded that the null hypothesis H_0 is not rejected. Therefore, there is not enough evidence to claim that not all 3 population means are equal, at the $\alpha=.05$ significance level.

Table 6: Analysis of variance (ANOVA) results among 200 Meter, 500 Meter and 1000 Meter men Canoeing players with regard to the sub-parameter Reaction time

Source of Variation	Sum of Squares	d.f.	Variance	F-value	p-value
Between-treatments	0.0069	2	0.0034	F = 4.70599	.022698
Within-treatments	0.0132	18	0.0007		
Total	0.0201	20			

The f-ratio value is 4.70599. The p-value is .022698. The result is significant at $p < .05$.

It is concluded that the null hypothesis H_0 is rejected. Therefore, there is enough evidence to claim that not all 3 population means are equal, at the $\alpha=.05$ significance level.

Table 7: Analysis of least significant difference (LSD) post-hoc test among 200 Meter, 500 Meter and 1000 Meter men Canoeing players with regard to the sub-parameter Reaction time

Analysis Multiple Comparisons

Group (A)	Group (B)	Mean Difference	Sig.
200 M (0.1631)	500 M	-.00814	.580
	1000 M	-.04186*	.010
500 M (0.1713)	200 M	.00814	.580
	1000 M	-.03371*	.032
1000 M (0.205)	200 M	.04186*	.010
	500 M	.03371*	.032

*. The mean difference is significant at the 0.05 level.




Figure 2: Graphical representation of mean scores among 200 Meter, 500 Meter and 1000 Meter men Canoeing players with regard to the sub-parameter Reaction time

Table 8: Analysis of variance (ANOVA) results among 200 Meter, 500 Meter and 1000 Meter men Canoeing players with regard to the sub-parameter Speed

Source of Variation	Sum of Squares	d.f.	Variance	F-value	p-value
Between-treatments	0.0543	2	0.0271	F = 2.06024	.156423
Within-treatments	0.2371	18	0.0132		
Total	0.2914	20			

The f-ratio value is 2.06024. The p-value is .156423. The result is not significant at $p < .05$.

It is concluded that the null hypothesis H_0 is not rejected. Therefore, there is not enough evidence to claim that not all 3 population means are equal, at the $\alpha=.05$ significance level.

5. Conclusions

In summary, the current study exhibits an intervening attempt to determine the difference of Physical Fitness levels among 200, 500 and 1000 Meter Canoeing Players. No significant differences were found in Agility, Balance, Coordination, Power and Speed. However, significant differences were found in Reaction Time.

Acknowledgements

A special acknowledgement of appreciation for this work in preparing the original manuscript is due to assistance from male Canoeing Player those who have participated in All India Open Canoe Sprint Championship organized by Sant Seechewal water sports centre Sultanpur Lodhi, Kapurthala, Punjab from 16-18 June, 2017.

I am indebted to Padmashri S. Baba Balbir Singh Seechewal for their inspiration and propositions rendered during the study.

References

1. Pendergast D. R., Bushnell D., Wilson D. W., Cerretelli P. (1989). Energetics of kayaking. *Eur J Appl Physiol Occup Physiol.* 59 (5): 342–50.
2. Michael J. S., Rooney KB, Smith R. (2008). The metabolic demands of kayaking: a review. *J Sports Sci Med.* 7 (1): 1–7.
3. Telford R., Tumulty D., Damm G. (1984). Skinfold measurements in well performed Australian athletes. *Sports Science and Medicine Quarterly.* 1: 13–6.
4. Ackland T., Kerr D., Hume P., Ridge B., Clark S., Broad E., Ross W. (2001). A Sports Medicine Odyssey: Challenges, Controversies & Change, ACT, Australia, Sports Medicine Australia. *Anthropometric normative data for Olympic rowers and paddlers.*
5. Ackland T. R., Ong K. B., Kerr D. A., Ridge B. (2003). Morphological characteristics of Olympic sprint canoe and kayak paddlers. *J Sci Med Sport.* 6 (3): 285–94.

6. Cressie N. A. C., Withers R. T., Craig N. P. (1986). The statistical analysis of somatotype data. *Am J Phys Anthropol.* 29:197–208.
7. Ziemilska A. (1973). Budowa somatyczna zawodników wysoko kwalifikowanych. *Wyniki badań AWF.* 17:1–88. In Polish.
8. Carter J. E. L., Ross W. D., Aubry S. P., Hebbelinck M., Borms J. (1986). Anthropometry of Montreal athletes. In: Carter JEL, editor. *Medicine and sport: physical structure of Olympic athletes.* Basel: Karger. pp. 25–52.
9. Bishop, D. (2000). Physiological predictors of flat-water kayak performance in women. *European Journal of Applied Physiology.* 82, 91-97.
10. Van Someren, K.A., Phillips, G.R.W. and Palmer, G.S. (1999). Comparison of physiological responses to open water kayaking and kayak ergometry. *International Journal of Sports Medicine.* 21, 200-204.
11. Gabbett T., Georgieff B. (2007). Physiological and anthropometric characteristics of Australian junior national, state, and novice volleyball players. *J Strength Cond Res.* 21: 902-908.
12. Mielgo-Ayuso J., Calleja-Gonzalez J., Clemente-Suarez V. J., Zourdos M. C. (2015). Influence of anthropometric profile on physical performance in elite female volleyballers in relation to playing position. *Nutricion Hospitalaria.* 31: 849-857.
13. Cronin J. B., Hansen K. T. (2005). Strength and power predictors of sports speed. *J Strength Cond Res.* 19: 349-357.
14. Gabbett T., Georgieff B. (2007). Physiological and anthropometric characteristics of Australian junior national, state, and novice volleyball players. *J Strength Cond Res.* 21: 902-908.
15. Leone M., Lariviere G., Comtois A. S. (2002). Discriminant analysis of anthropometric and biomotor variables among elite adolescent female athletes in four sports. *J Sports Sci.* 20: 443-449.
16. Simoneau G. G. (1998). The impact of various anthropometric and flexibility measurements on the sit-and-reach test. *J Strength Cond Res.* 12: 232-237.

Creative Commons licensing terms

Authors will retain the copyright of their published articles agreeing that a Creative Commons Attribution 4.0 International License (CC BY 4.0) terms will be applied to their work. Under the terms of this license, no permission is required from the author(s) or publisher for members of the community to copy, distribute, transmit or adapt the article content, providing a proper, prominent and unambiguous attribution to the authors in a manner that makes clear that the materials are being reused under permission of a Creative Commons License. Views, opinions and conclusions expressed in this research article are views, opinions and conclusions of the author(s). Open Access Publishing Group and European Journal of Physical Education and Sport Science shall not be responsible or answerable for any loss, damage or liability caused in relation to/arising out of conflict of interests, copyright violations and inappropriate or inaccurate use of any kind content related or integrated on the research work. All the published works are meeting the Open Access Publishing requirements and can be freely accessed, shared, modified, distributed and used in educational, commercial and non-commercial purposes under a [Creative Commons attribution 4.0 International License \(CC BY 4.0\)](https://creativecommons.org/licenses/by/4.0/).