

Enantioselective Formal [3+2] Cycloaddition of Epoxides with Imines under Bronsted Base Catalysis: Synthesis of 1,3 Oxazolidines with Quaternary Stereogenic Center

著者	Azusa Kondoh, Shiori Akahira, Masafumi Oishi, Masahiro Terada
journal or publication title	Angewandte Chemie
volume	57
number	21
page range	6299-6303
year	2018-04-12
URL	http://hdl.handle.net/10097/00125839

doi: 10.1002/anie.201802468


Enantioselective Formal [3+2] Cycloaddition of Epoxides with Imines under Brønsted Base Catalysis: Synthesis of 1,3-Oxazolidines with Quaternary Stereogenic Center

Azusa Kondoh,^[b] Shiori Akahira,^[a] Masafumi Oishi,^[a] and Masahiro Terada^{*[a]}

Abstract: The formal [3+2] cycloaddition of epoxides with unsaturated compounds is a powerful methodology for the synthesis of densely functionalized five-membered heterocyclic compounds containing oxygen. We have developed a novel enantioselective formal [3+2] cycloaddition of epoxides under Brønsted base catalysis. The bis(guanidino)iminophosphorane as a chiral organosuperbase catalyst enabled the enantioselective reaction of β,γ -epoxysulfones with imines, owing to its strong basicity and high stereocontrolling ability, to provide enantioenriched 1,3-oxazolidines having two stereogenic centers including a quaternary one in a highly diastereo- and enantioselective manner.

The ring expansion of strained cyclic compounds has attracted considerable attention as a useful strategy for constructing polysubstituted cyclic frameworks.^[1] Among a variety of reactions based on this strategy, the formal [3+2] cycloaddition of epoxides with unsaturated compounds is a particularly powerful methodology for the synthesis of densely functionalized five-membered heterocyclic compounds containing oxygen. These reactions are generally catalyzed by transition metal complexes,^[2] Lewis acids^[3] or the combination of Lewis acids with halides.^[4] Epoxides formally serve as the synthetic equivalent of a 1,3-dipole under the influence of these catalysts, which is the key to the reaction proceeding with a variety of unsaturated compounds. Recently, development of asymmetric variants has also been advanced by utilizing transition metal catalysts or Lewis acid catalysts with chiral ligands.^[5] However, catalytic systems that can construct multiple stereogenic centers in a highly stereoselective manner are still rather limited. Therefore, the expansion of the repertoire for this methodology through the establishment of new catalytic systems is highly anticipated. We recently established a conceptually different catalytic system for the formal [3+2] cycloaddition of epoxides under Brønsted base catalysis, which is complementary to the conventional catalytic systems. We have developed a formal [3+2] cycloaddition of β,γ -epoxyesters with imines providing 2,4,5-trisubstituted 1,3-oxazolidines in a highly diastereoselective manner.^[6] As the next stage of our research, we envisioned the development of an asymmetric variant of this catalytic system by utilizing a chiral Brønsted base catalyst.

Specifically, we designed an enantioselective formal [3+2] cycloaddition of β,γ -epoxysulfones **2** having a substituent on the β -carbon with imines **3**, which involves the construction of two stereogenic centers, including a quaternary one, through an enantioconvergent process to provide enantioenriched 1,3-oxazolidines **4** (Scheme 1).



Scheme 1. Designed Reaction System.

Treatment of racemic β,γ -epoxysulfone **2** with a chiral Brønsted base would result in deprotonation at the position α to the sulfonyl group, followed by epoxide opening to provide alkoxide **B** possessing an alkenyl sulfone moiety. The driving force for this would be the release of ring strain. At this stage, the chiral information of starting **2** has disappeared. This intermediate would then formally serve as the synthetic equivalent of a 1,3-dipole, and the cycloaddition with imine **3** would proceed in a stepwise fashion, i.e., addition of alkoxide **B** to imine **3** followed by intramolecular aza-Michael addition of intermediate **C**, to afford 1,3-oxazolidine **4**. The main challenge of the intended reaction is the stereocontrol of the two stereogenic centers. To this end, we expected that a suitable choice of chiral Brønsted base would enable the enantioselective addition of alkoxide **B** to imine **3** although such precedents are rather limited.^[7] In addition, the diastereocontrol of the subsequent aza-Michael addition of **C** would be achieved by substrate control and/or catalyst control, thus providing stereocontrolled 1,3-oxazolidines **4**. Enantioenriched 1,3-oxazolidines can be utilized as synthetically versatile intermediates, chiral auxiliaries,^[8] and ligands in transition metal catalysis.^[9] They are also an important structural motif found in many biologically active compounds.^[10] Therefore, the intended reaction would provide new efficient access to enantioenriched 1,3-oxazolidines that are difficult to synthesize by other methods.^[11] Based on this idea, we describe herein an enantioselective formal [3+2] cycloaddition of β,γ -epoxysulfones with imines under Brønsted base catalysis. A chiral bis(guanidino)iminophosphorane (*M*)-**1** (Figure 1),^[12] as a chiral organosuperbase catalyst, enabled the efficient synthesis of

[a] S. Akahira, M. Oishi, Prof. Dr. M. Terada
Department of Chemistry, Graduate School of Science
Tohoku University
Aramaki, Aoba-ku, Sendai 980-8578 (Japan)
E-mail: mterada@m.tohoku.ac.jp

[b] Dr. A. Kondoh
Research and Analytical Center for Giant Molecules
Graduate School of Science, Tohoku University
Aramaki, Aoba-ku, Sendai 980-8578 (Japan)

Supporting information for this article is given via a link at the end of the document.

enantioenriched 1,3-oxazolidines having two stereogenic centers including a quaternary one in a highly diastereo- and enantioselective manner.





Figure 1. Chiral Bis(guanidino)iminophosphoranes (*M*)-1.


We began our investigation by evaluating the viability of our designed reaction by using achiral organobases having different basicities (Table 1, entries 1-4). β,γ -Epoxy sulfone **2a** having a phenyl group at the β position was chosen as the primary substrate and treated with *N*-Boc imine **3a** in the presence of 10 mol% organobase in toluene at -20 °C. As a result, the use of DBU ($pK_{\text{BH}^+} = 24.3$ in CH_3CN),^[13] MTBD ($pK_{\text{BH}^+} = 25.4$), and TBD ($pK_{\text{BH}^+} = 26.0$) resulted in the formation of *N,O*-acetal **5aa** in almost quantitative yield with a high *Z/E* ratio, and only a trace amount of the desired 1,3-oxazolidine **4aa** was observed in the crude ^1H NMR spectra (entries 1-3).^[14] In contrast, P2-*t*Bu having much stronger basicity ($pK_{\text{BH}^+} = 33.5$) provided **4aa** in good yield with good diastereoselectivity (entry 4). These results clearly suggest that the use of a catalyst having strong basicity, which facilitates the intramolecular aza-Michael addition of the anion of *N,O*-acetal **5aa**, is essential for completing the tandem catalytic process due to the low electrophilicity of the β,β -disubstituted sulfone moiety of the intermediate. These preliminary results prompted us to start the investigation of the enantioselective reaction of **2a** with **3a** by using chiral bis(guanidino)iminophosphoranes (*M*)-1, which were developed by our group,^[12] possessing comparably high basicity to P2-*t*Bu.^[15] The initial experiment was conducted with the catalyst generated in situ by treating 11 mol% (*M*)-1a·HBr with 10 mol% $\text{KN}(\text{SiMe}_3)_2$ prior to use. However, **4aa** was not formed, and instead, *N,O*-acetal **5aa** was obtained quantitatively with 82% ee for the major *Z* isomer (entry 5). The use of other inorganic bases, such as $\text{NaN}(\text{SiMe}_3)_2$ and NaOtBu , for generation of the catalyst provided results similar to that with $\text{KN}(\text{SiMe}_3)_2$ (entries 6 and 7). We assumed that the failure of the aza-Michael addition was attributed to the detrimental effect of alkali metal cations, such as potassium cation and sodium cation, which would reduce the requisite nucleophilicity of the anion of the *N,O*-acetal **5aa**. Based on this hypothesis, we next attempted the reaction by using $\text{KN}(\text{SiMe}_3)_2$ with 30 mol% 18-crown-6 as an additive. As a result, the formal [3+2] cycloaddition proceeded to afford **4aa** in good yield as a single diastereomer with 75% ee (entry 8), which is a similar level of enantioselectivity to that of *N,O*-acetal **5aa** without 18-crown-6 (entries 5 vs. 8). The extension of the reaction time further improved the yield of **4aa** (entry 9). In order to increase the enantioselectivity, other precatalysts (*M*)-1·HX having different substituents were examined (entries 10-12). The reaction with precatalyst (*M*)-1b·HCl possessing mesityl groups on the 7,7-

Table 1. Initial Screening of Reaction Conditions^[a]



entry	base catalyst	yield [%] ^[b]		ee [%] ^[c]	
		4aa (dr)	5aa (<i>Z/E</i>)	4aa	5aa
1	DBU	<5 (-)	95 (98/2)	-	-
2	MTBD	2 (-)	98 (99/1)	-	-
3	TBD	1 (-)	99 (98/2)	-	-
4	P2- <i>t</i> Bu	85 (87/13)	3 (<1/99)	-	-
5	(<i>M</i>)-1a·HBr/ $\text{KN}(\text{SiMe}_3)_2$	<1 (-)	>99 (96/4)	-	82 ^[i]
6	(<i>M</i>)-1a·HBr/ $\text{NaN}(\text{SiMe}_3)_2$	1 (-)	98 (98/2)	-	77
7	(<i>M</i>)-1a·HBr/ NaOtBu	<1 (-)	95 (97/3)	-	82
8 ^[d]	(<i>M</i>)-1a·HBr/ $\text{KN}(\text{SiMe}_3)_2$	84 (>99/1)	12 (86/14)	75	-
9 ^[d,e]	(<i>M</i>)-1a·HBr/ $\text{KN}(\text{SiMe}_3)_2$	98 (>99/1)	<1 (-)	75	-
10 ^[d,e]	(<i>M</i>)-1b·HCl/ $\text{KN}(\text{SiMe}_3)_2$	97 (>99/1)	<3 (-)	-3	-
11 ^[d,e]	(<i>M</i>)-1c·HCl/ $\text{KN}(\text{SiMe}_3)_2$	40 (>99/1)	60 (93/7)	91	97
12 ^[d,e]	(<i>M</i>)-1d·HCl/ $\text{KN}(\text{SiMe}_3)_2$	95 ^[f] (>99/1)	3 (<1/99)	93	-
13 ^[d,e,g]	(<i>M</i>)-1d·HCl/ $\text{KN}(\text{SiMe}_3)_2$	97 ^[h] (>99/1)	3 (<1/99)	93	-

[a] Reaction conditions: **2a** (0.10 mmol), **3a** (0.12 mmol), organobase (0.010 mmol) or (*M*)-1·HX (0.011 mmol) with inorganic base (0.010 mmol), toluene (4.0 mL), -20 °C. [b] NMR yields unless otherwise noted. Diastereomeric ratio of **4aa** and *Z/E* ratio of **5aa** were determined by ^1H NMR analysis. [c] Enantiomeric excess of the major isomer of **4aa** or (*Z*)-**5aa** was determined by chiral stationary phase HPLC analysis. [d] 0.030 mmol of 18-crown-6 (30 mol%) was used. [e] The reaction was conducted for 14 h. [f] 88% isolated yield. [g] The reaction was performed in 0.50 mmol scale. [h] Isolated yield. [i] Enantiomeric excess of (*E*)-**5aa** was 46%.




membered spirocyclic rings proceeded smoothly, however, nearly racemic **4aa** was obtained (entry 10). In contrast, (*M*)-1c·HCl with 1-naphthyl groups substantially increased the

enantioselectivity albeit with moderate yield (entry 11). Finally, (*M*)-**1d**·HCl, which had 1-naphthyl groups on the spirocyclic rings and *t*Bu groups on the nitrogen of the guanidine moieties, was found to be the best precatalyst to provide **4aa** in 88% isolated yield with 93% ee (entry 12). The reaction in larger scale proceeded without any problem (entry 13). The absolute configuration of **4aa** was unambiguously determined to be (*2R,4R*) by single-crystal X-ray diffraction analysis of racemic **4aa** and enantioenriched amino alcohol **7** derived from enantioenriched **4aa** (vide infra).^[16]

With the optimum reaction conditions in hand, the scope of substrates was investigated (Table 2). First, the substituent at the β -position of the epoxysulfones **2** was screened (entries 1-8). The reaction of substrates **2b-2d** having a halogen moiety at the *para* position of the phenyl ring provided the corresponding products **4ba-4ea** in high yields with high enantioselectivities

Table 2. Substrate Scope^[a]




entry	2 or 6	R ¹	R ²	4	yield [%] ^[b]	ee [%] ^[c]
1	2b	4-F-C ₆ H ₄	Ph	4ba	89	92
2	2c	4-Cl-C ₆ H ₄	Ph	4ca	90	88
3	2d	4-Br-C ₆ H ₄	Ph	4da	92	87
4	2e	4-CF ₃ -C ₆ H ₄	Ph	4ea	87	72
5	6f	4-Me-C ₆ H ₄	Ph	4fa	99	87
6	6g	3-Me-C ₆ H ₄	Ph	4ga	90	84
7	6h	3-MeO-C ₆ H ₄	Ph	4ha	90	88
8	6i	2-naphthyl	Ph	4ia	99	92
9	2a	Ph	4-Cl-C ₆ H ₄	4ab	88	93
10	2a	Ph	4-MeO-C ₆ H ₄	4ac	86	93
11	2a	Ph	2-Me-C ₆ H ₄	4ad	92	41
12	2a	Ph	2-naphthyl	4ae	92	89
13	2a	Ph	2-thienyl	4af	99	90
14	2a	Ph	2-furyl	4ag	97	76

[a] Reaction conditions: **2** or **6** (0.10 mmol), **3** (0.12 mmol), (*M*)-**1d**·HCl (0.011 mmol), KN(SiMe₃)₂ (0.010 mmol), 18-crown-6 (0.030 mmol), toluene (4.0 mL), -20 °C. [b] Isolated yields. [c] Enantiomeric excess was determined by chiral stationary phase HPLC analysis.

(entries 1-3). 4-Trifluoromethylphenyl-substituted **2e** also underwent the reaction smoothly albeit with moderate ee (entry 4). Some β,γ -epoxysulfones were difficult to prepare in pure form. In these cases, the β,γ -epoxysulfones **2** were converted to the corresponding allylic alcohols **6** by treatment with a catalytic amount of TBD in THF, and the pure isolated (*Z*)-**6**, which were formed as the major isomer (*Z/E* > 95/5 in each case), were used as substrates in the reaction with imine **3a**.^[17] With this alternative protocol, each substrate possessing *para*-tolyl, *meta*-tolyl, 3-methoxyphenyl, or 2-naphthyl groups, afforded the corresponding oxazolidines in high yields with high enantioselectivities (entries 5-8). An alkyl substituent, such as a methyl group, was also examined. However, the corresponding 1,3-oxazolidine was not formed and an unidentified mixture of products was obtained.^[18] Next, the scope of *N*-Boc imines was examined by using **2a** as a substrate (entries 9-14). Both aryl imines having an electron-withdrawing chloro group and an electron-donating methoxy group at the *para* position provided **4ab** and **4ac**, respectively, in high yields with high enantioselectivities (entries 9-10). In contrast, the reaction with **3d** having an *ortho*-tolyl group provided **4ad** in good yield with only modest ee value (entry 11). 2-Naphthyl-substituted **3e** and heteroaryl imines, such as 2-thienyl- and 2-furyl-substituted imines, underwent the reaction without any problem, and the corresponding products **4ae-4ag** were obtained in high yields with good to high enantioselectivities.


It is worth noting that all of the reactions conducted with (*M*)-**1** provided 1,3-oxazolidines **4** as a single diastereomer. In order to gain some insight into the origin of the diastereoselectivity, some control experiments were carried out (Scheme 2). Specifically, both *Z* and *E* isomers of racemic *N,O*-acetal (\pm)-**5aa** were treated with P2-*t*Bu or (*M*)-**1a**·HBr/KN(SiMe₃)₂/18-crown-6. In the case of P2-*t*Bu as a catalyst, (\pm)-(*Z*)-**5aa** provided (\pm)-**4aa** as a mixture of diastereomers in a 86/14 ratio, which was almost identical to that obtained in the reaction of **2a** with **3a** (Table 1, entry 4), while (\pm)-(*E*)-**5aa** provided (\pm)-**4aa** in only low yield in a 55/45 ratio (Scheme 2a). In contrast, (*M*)-**1a** provided (\pm)-**4aa** in 81% yield as a single diastereomer from (\pm)-(*Z*)-**5aa** (Scheme 2b). These results suggest that the Brønsted base catalysts are partially responsible for the diastereocontrol in the intramolecular aza-Michael addition of **5aa** although substrate control is mainly operative. Therefore, in this tandem catalytic process, the key roles of (*M*)-**1** are as follows: 1) facilitating the reaction with its strong basicity, 2) controlling the enantioselectivity in the addition of the alkoxide to the imine, and 3) assisting the diastereocontrol of the aza-Michael addition. Furthermore, the control experiment revealed that (\pm)-(*E*)-**5aa** was far less reactive than (\pm)-(*Z*)-**5aa**. In the case of (*M*)-**1a** as a catalyst, the cyclization of (\pm)-(*E*)-**5aa** did not proceed, and 93% of starting (\pm)-(*E*)-**5aa** was recovered (Scheme 2b). In addition, the preliminary experiment revealed that the enantioselectivity in the addition of *Z* isomer of the alkoxide is much higher than that in the addition of *E* isomer (Table 1, entry 5), suggesting that (*M*)-**1** can effectively control the enantioselectivity with *Z* isomer compared with *E* isomer.^[19] Therefore, the selective formation of the (*Z*) configuration of the alkoxide intermediate through the ring opening of β,γ -epoxysulfone **2**, which would be independent

of the choice of Brønsted base catalyst, was critical for achieving both high yield and high stereoselectivity of **4**.



Scheme 2. Control Experiments.

Finally, derivatization of 1,3-oxazolidine **4aa** was conducted (Scheme 3). **4aa** was easily convertible to the corresponding amino alcohol **7** in almost quantitative yield by treatment with aqueous HBr (Scheme 3a). Further transformation of **7** was attempted to utilize the sulfone moiety as a handle for manipulation (Scheme 3b). Thus, treatment of **7** with 1,1'-carbonyldiimidazole (CDI) provided cyclic carbamate **8** in good yield. The Julia-Lithgove olefination was then conducted, and the desired **10** was obtained in good overall yield with perfect *E* selectivity by using Sml_2 as a reductant.^[20] The direct desulfonation of **8** was also operable by using Na(Hg) to provide **11** in good yield (Scheme 3c).



Scheme 3. Derivatization of **4aa**.

In conclusion, we have developed a novel enantioselective formal [3+2] cycloaddition of epoxides under Brønsted base catalysis. A bis(guanidino)iminophosphorane as a chiral organosuperbase efficiently catalyzed the enantioselective reaction of β,γ -epoxysulfones with imines to provide enantioenriched 1,3-oxazolidines having two stereogenic centers including a quaternary one in a highly diastereo- and enantioselective manner. This reaction involves: 1) the generation of the key alkoxide intermediate through epoxide opening, which is a formal synthetic equivalent of a 1,3-dipole, 2) the enantioselective addition of the intermediate to the imine, and 3) a diastereoselective intramolecular aza-Michael addition. Both strong basicity and high stereocontrolling ability were the required properties of the catalysts for achieving this tandem catalytic process, which emphasized the usability of a bis(guanidino)iminophosphorane as a chiral organosuperbase catalyst. Further studies, including a mechanistic study on the stereocontrol of the reaction, are in progress.

Experimental Section

The reaction of **2a** with **3a** is representative (Table 1, entry 12). To a solution of **2a** (27 mg, 0.10 mmol) and **3a** (24 μL , 0.12 mmol) in toluene (2.0 mL) was added a toluene solution (2.0 mL) containing (*M*)-**1d**-HCl (12 mg, 0.011 mmol), $\text{KN}(\text{SiMe}_3)_2$ (0.50 M in toluene, 20 μL , 0.010 mmol), and 18-crown-6 (1.0 M in toluene, 30 μL , 0.030 mmol) dropwise in 15 seconds at -20°C . After stirred for 14 h at that temperature, the reaction was quenched with sat. aq. NH_4Cl . The product was extracted with AcOEt for three times. The combined organic layer was washed with brine, dried over Na_2SO_4 and evaporated. The crude mixture was purified by silica-gel column chromatography (hexane/AcOEt = 3/1) to afford **4aa** (46 mg, 0.095 mmol, 95%, >99/1 dr, 93% ee) as a white solid.


Acknowledgements

This research was supported by a Grant-in-Aid for Scientific Research (S) (JP16H06354) and (C) (JP16K05680) from the JSPS.

Keywords: Brønsted base • cycloaddition • organocatalyst • asymmetric catalysis • quaternary stereogenic center


- [1] For reviews, see: a) C.-Y. Huang, A. G. Doyle, *Chem. Rev.* **2014**, *114*, 8153-8198; b) D. J. Mack, J. T. Njardarson, *ACS Catal.* **2013**, *3*, 272-286; c) M. A. Cavitt, L. H. Phun, S. France, *Chem. Soc. Rev.* **2014**, *43*, 804-818; d) E. A. Ilardi, J. T. Njardarson, *J. Org. Chem.* **2013**, *78*, 9533-9540; e) J. T. Njardarson, *Synlett* **2013**, *24*, 787-803; f) D. Garayalde, C. Nevado, *Beilstein J. Org. Chem.* **2011**, *7*, 767-780; g) T. Hudlicky, J. W. Reed, *Angew. Chem.* **2010**, *122*, 4982-4994; *Angew. Chem. Int. Ed.* **2010**, *49*, 4864-4876.
- [2] a) B. M. Trost, S. R. Angle, *J. Am. Chem. Soc.* **1985**, *107*, 6123-6124; b) T. Fujinami, T. Suzuki, M. Kamiya, S. Fukuzawa, S. Sakai, *Chem. Lett.* **1985**, *14*, 199-200; c) B. M. Trost, A. R. Sudhakar, *J. Am. Chem. Soc.* **1987**, *109*, 3792-3794; d) B. M. Trost, A. R. Sudhakar, *J. Am. Chem. Soc.* **1988**, *110*, 7933-7935; e) J.-G. Shim, Y. Yamamoto, *J. Org. Chem.* **1998**, *63*, 3067-3071.
- [3] For a review, see: a) V. D'Elia, J. D. A. Pelletier, J.-M. Basset, *ChemCatChem* **2015**, *7*, 1906-1917. For selected recent examples, see: b) H. Jiang, H. Jie, J. Li, *RSC Adv.* **2016**, *6*, 100307-100311; c) J.

- Zhang, Y. Xiao, J. Zhang, *Adv. Synth. Catal.* **2013**, *355*, 2793-2797; d) Z. Chen, Y. Xiao, J. Zhang, *Eur. J. Org. Chem.* **2013**, 4748-4751; e) Z. Chen, Z. Tian, J. Zhang, J. Ma, J. Zhang, *Chem. Eur. J.* **2012**, *18*, 8591-8595.
- [4] For a review, see: a) C. Martín, G. Fiorani, A. W. Kleij, *ACS Catal.* **2015**, *5*, 1353-1370. For selected recent examples, see: b) J. A. Castro-Osma, A. Earlam, A. Lara-Sánchez, A. Otero, M. North, *ChemCatChem* **2016**, *8*, 2100-2108; c) P. Wang, J. Qin, D. Yuan, Y. Wang, Y. Yao, *ChemCatChem* **2015**, *7*, 1145-1151; d) C. Beattie, M. North, *Chem. Eur. J.* **2014**, *20*, 8182-8188, and references cited therein.
- [5] a) J.-J. Suo, J. Du, Q.-R. Liu, D. Chen, C.-H. Ding, Q. Peng, X.-L. Hou, *Org. Lett.* **2017**, *19*, 6658-6661; b) Q. Cheng, H.-J. Zhang, W.-J. Yue, S.-L. You, *Chem* **2017**, *3*, 428-436; c) C. Ma, Y. Huang, Y. Zhao, *ACS Catal.* **2016**, *6*, 6408-6412; d) X. Yuan, L. Lin, W. Chen, W. Wu, X. Liu, X. Feng, *J. Org. Chem.* **2016**, *81*, 1237-1243; e) W. Chen, Y. Xia, L. Lin, X. Yuan, S. Guo, X. Liu, X. Feng, *Chem. Eur. J.* **2015**, *21*, 15104-15107; f) W. Chen, X. Fu, L. Lin, X. Yuan, W. Luo, J. Feng, X. Liu, X. Feng, *Chem. Commun.* **2014**, *50*, 11480-11483; g) W. Chen, L. Lin, Y. Cai, Y. Xia, W. Cao, X. Liu, X. Feng, *Chem. Commun.* **2014**, *50*, 2161-2163; h) Z. Liu, X. Feng, H. Du, *Org. Lett.* **2012**, *14*, 3154-3157; i) W.-Q. Wu, C.-H. Ding, X.-L. Hou, *Synlett* **2012**, *23*, 1035-1038; j) M. B. Shaghafi, R. E. Grote, E. R. Jarvo, *Org. Lett.* **2011**, *13*, 5188-5191; k) C. Larksarp, H. Alper, *J. Org. Chem.* **1998**, *63*, 6229-6233; l) C. Larksarp, H. Alper, *J. Am. Chem. Soc.* **1997**, *119*, 3709-3715.
- [6] A. Kondoh, K. Odaira, M. Terada, *Angew. Chem.* **2015**, *127*, 11392-11396; *Angew. Chem. Int. Ed.* **2015**, *54*, 11240-11244.
- [7] a) D. Zhou, X. Yu, J. Zhang, W. Wang, H. Xie, *Org. Biomol. Chem.* **2016**, *14*, 6193-6196; b) J. Liu, F.-M. Zhu, Y.-B. Chu, L.-H. Huang, Y.-F. Zhou, *Tetrahedron: Asymmetry* **2015**, *26*, 1130-1137; c) T.-Z. Li, X.-B. Wang, F. Sha, X.-Y. Wu, *Tetrahedron* **2013**, *69*, 7314-7319.
- [8] For a review, see: C. Agami, F. Couty, *Eur. J. Org. Chem.* **2004**, 677-685.
- [9] a) H. Nakano, Y. Okuyama, E. Kwon, *Heterocycles* **2014**, *89*, 1-26; b) C. Wolf, H. Xu, *Chem. Commun.* **2011**, *47*, 3339-3350; c) T. Arai, Y. Ogino, T. Sato, *Chem. Commun.* **2013**, *49*, 7776-7778.
- [10] For selected examples, see: a) J. D. Scott, R. M. Williams, *Chem. Rev.* **2002**, *102*, 1669-1730; b) L. Yu, W. Zhou, Z. Wang, *Bio. Med. Chem. Lett.* **2011**, *21*, 1541-1544; c) K. Ii, S. Ichikawa, B. Al-Dabbagh, A. Bouhss, A. Matsuda, *J. Med. Chem.* **2010**, *53*, 3793-3813; d) J. D. Scott, R. M. Williams, *J. Am. Chem. Soc.* **2002**, *124*, 2951-2956; e) G. P. Moloney, D. J. Craik, M. N. Iskander, T. L. Nero, *J. Chem. Soc., Perkin Trans. 2.* **1998**, 199-206.
- [11] For selected recent examples on catalytic enantioselective synthesis of 1,3-oxazolidines, see: a) S. Tanaka, R. Gunasekar, T. Tanaka, Y. Iyoda, Y. Suzuki, M. Kitamura, *J. Org. Chem.* **2017**, *82*, 9160-9170; b) X. Wu, W. Zhou, H.-H. Wu, J. Zhang, *Chem. Commun.* **2017**, *53*, 5661-5664; c) Y. Liao, X. Liu, Y. Zhang, Y. Xu, Y. Xia, L. Lin, X. Feng, *Chem. Sci.* **2016**, *7*, 3775-3779; d) L. Yang, A. Khan, R. Zheng, L. Y. Jin, Y. J. Zhang, *Org. Lett.* **2015**, *17*, 6230-6233; e) S. K. Nimmagadda, Z. Zhang, J. C. Antilla, *Org. Lett.* **2014**, *16*, 4098-4101; f) D. M. Rubush, T. Rovis, *Synlett* **2014**, *25*, 713-717; g) Y. Fukata, K. Asano, S. Matsubara, *Chem. Lett.* **2013**, *42*, 355-357; h) K. S. Williamson, T. P. Yoon, *J. Am. Chem. Soc.* **2012**, *134*, 12370-12373; i) Z. Jin, X. Wang, H. Huang, X. Liang, J. Ye, *Org. Lett.* **2011**, *13*, 564-567; j) D. J. Michaelis, K. S. Williamson, T. P. Yoon, *Tetrahedron* **2009**, *65*, 5118-5124.
- [12] a) T. Takeda, M. Terada, *J. Am. Chem. Soc.* **2013**, *135*, 15306-15309; b) T. Takeda, M. Terada, *Aust. J. Chem.* **2014**, *67*, 1124-1128; c) A. Kondoh, M. Oishi, T. Takeda, M. Terada, *Angew. Chem.* **2015**, *127*, 16062-16065; *Angew. Chem. Int. Ed.* **2015**, *54*, 15836-15839; d) T. Takeda, A. Kondoh, M. Terada, *Angew. Chem.* **2016**, *128*, 4812-4815; *Angew. Chem. Int. Ed.* **2016**, *55*, 4734-4737.
- [13] *Superbases for Organic Synthesis* (Ed.: T. Ishikawa), John Wiley & Sons, Chippingham, **2009**.
- [14] (Z) configuration of major isomer of **5aa** was unambiguously determined by X-ray crystallographic analysis. CCDC No.1822256. See the Supporting Information for details.
- [15] The pK_{BH^+} of achiral bis(guanidino)iminophosphorane **12** possessing a backbone similar to **1** was reported to be 26.8 in THF, see: A. A. Kolomeitsev, I. A. Koppel, T. Rodima, J. Barten, E. Lork, G.-V. Röschenthaler, I. Kaljurand, A. Kütt, I. Koppel, V. Mäemets, I. Leito, *J. Am. Chem. Soc.* **2005**, *127*, 17656-17666.



- [16] CCDC No.1822255 for racemic **4aa** and CCDC No.1822257 for (R)-**7**. See the Supporting Information for details.
- [17] See the Supporting Information for details.
- [18] The reaction of β,γ -epoxysulfone having an ethoxycarbonyl group at the R¹ position was also examined under the optimized reaction conditions. However, the conversion of the substrate was very low (ca. 30%), and the corresponding oxazolidine was not obtained. The introduction of such electron-withdrawing groups seems to have a detrimental effect on the reaction system.
- [19] The reaction of (E)-**6a** with **3a** under the optimized reaction conditions resulted in the formation of (E)-**5aa** in 95% yield with <1% ee.
- [20] E. O. Volz, G. W. O'Neil, *J. Org. Chem.* **2011**, *76*, 8428-8432.

COMMUNICATION



Azusa Kondoh, Shiori Akahira,
Masafumi Oishi, Masahiro Terada*

Page No. – Page No.

**Enantioselective Formal [3+2]
Cycloaddition of Epoxides with
Imines under Brønsted Base
Catalysis: Synthesis of 1,3-
Oxazolidines with Quaternary
Stereogenic Center**

A novel enantioselective formal [3+2] cycloaddition of epoxides was developed under Brønsted base catalysis. The bis(guanidino)iminophosphorane as a chiral organosuperbase catalyst enabled the enantioselective reaction of β,γ -epoxysulfones with imines, owing to its strong basicity and high stereocontrolling ability, to provide enantioenriched 1,3-oxazolidines having two stereogenic centers including a quaternary one in a highly diastereo- and enantioselective manner.