

STARS

University of Central Florida
STARS

UCF Patents

Technology Transfer

1-29-2013

Mesenchymal stem cells with increased developmental potency by expressing Nanog

Kiminobu Sugaya

Angel Alvarez

University of Central Florida

Find similar works at: <https://stars.library.ucf.edu/patents>

University of Central Florida Libraries <http://library.ucf.edu>

This Patent is brought to you for free and open access by the Technology Transfer at STARS. It has been accepted for inclusion in UCF Patents by an authorized administrator of STARS. For more information, please contact STARS@ucf.edu.

Recommended Citation

Sugaya, Kiminobu and Alvarez, Angel, "Mesenchymal stem cells with increased developmental potency by expressing Nanog" (2013). *UCF Patents*. 755.

<https://stars.library.ucf.edu/patents/755>





(12) **United States Patent**
Sugaya et al.

(10) **Patent No.:** **US 8,361,791 B2**
(45) **Date of Patent:** **Jan. 29, 2013**

(54) **MESENCHYMAL STEM CELLS WITH INCREASED DEVELOPMENTAL POTENCY BY EXPRESSING NANOG**

(75) Inventors: **Kiminobu Sugaya**, Winter Park, FL (US); **Angel Alvarez**, Orlando, FL (US)

(73) Assignee: **University of Central Florida Research Foundation, Inc.**, Orlando, FL (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **13/483,807**

(22) Filed: **May 30, 2012**

(65) **Prior Publication Data**

US 2012/0238015 A1 Sep. 20, 2012

Related U.S. Application Data

(63) Continuation of application No. 11/258,401, filed on Oct. 24, 2005, now Pat. No. 8,192,988.

(60) Provisional application No. 60/621,901, filed on Oct. 22, 2004, provisional application No. 60/650,438, filed on Feb. 4, 2005.

(51) **Int. Cl.**
C12N 5/00 (2006.01)
C12N 5/02 (2006.01)

(52) **U.S. Cl.** **435/325; 435/366**

(58) **Field of Classification Search** None
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

6,833,269	B2	12/2004	Carpenter
7,618,621	B2	11/2009	Sugaya et al.
7,635,467	B2	12/2009	Sugaya et al.
2006/0110440	A1	5/2006	Sugaya et al.
2006/0134789	A1	6/2006	Sugaya et al.
2006/0188489	A1	8/2006	Sugaya et al.

FOREIGN PATENT DOCUMENTS

WO	WO 03/064463	A2	8/2003
WO	WO 2004/072226	A2	8/2004

OTHER PUBLICATIONS

Hatano, Shin-ya et al., "Pluripotential competence of cells associated with Nanog activity", *Mechanisms of Development*, 2005, vol. 122, pp. 67-79.

Silva, Jose et al., "Nanog promotes transfer of pluripotency after cell fusion", *Nature*, Jun. 22, 2006, vol. 44, pp. 997-1001, Nature Publishing Group.

Hart, Adam H. et al., "Identification, Cloning and Expression Analysis of the Pluripotency Promoting Nanog Genes in Mouse and Human", *Developmental Dynamics*, 2004, vol. 230, pp. 187-198.

Yu, Junying et al., "Induced Pluripotent Stem Cell Lines Derived from Human Somatic Cells", *Science*, 2007; 318, 1917, doi:10.1126/science.1151526.

Takahashi, Kazutoshi et al., "Induction of Pluripotent Stem Cells from Mouse Embryonic and Adult Fibroblast Cultures by Defined Factors", *Cell*, Aug. 25, 2006, vol. 126, pp. 663-676.

Liu, et al, (Jul. 1004), *Xi Bau Yu Fen Zi Mian Yi Xue Za Zhi*, 20(4): 495-498 (Abstract only).

Sugaya et al. (2006) *Panminerva Med.* 48:87-96.

"The Human Embryonic Stem Cell and the Human Embryonic Germ Cell" *Stem Cells: Scientific Progress and Future Research Directions*. Department of Health and Human Services. Jun. 2001., http://stemcells.nih.gov/info/scireport/2001_report, downloaded Feb. 8, 2008.

Mattson (2001) *Expert Rev. Neurotherapeutics* 1:267-273.

Isacson (2003) *Lancet Neurol* 2:417-424.

Lindvall et al. (2004) *Nature Med* 10 Suppl:S42-S50.

Snyder et al. (2004) *J Neurosci* 24:157-168.

Richards (1997) *Cell Mol Life Sci* 53:790-802.

Weissman (200) *Science* 287:1442-1446.

Sumer et al. (2011) "NANOG is a Key Factor for Induction of Pluripotency in Bovine Adult Fibroblasts", *Journal of Animal Science*, 441:997-1001 (Abstract only).

Zhao et al. (2010) "Rapid and Efficient Reprogramming of Human Amnion-Derived Cells into Pluripotency by Three Factors OCT4/SOX2/ANANOG", *Differentiation*, 80(2-3): 123-29 (Abstract only).

Pierantozzi et al. (2011) "Pluripotency Regulators in Human Mesenchymal Stem Cells: Expression of NANOG but not of OCT-4 and SOX-2", *Stem Cells and Development* 20(5): 915-23 (Abstract only).

Ma et al. (2008) "G9a and Jhd2a Regulate Embryonic Stem Cell Fusion-Induced Reprogramming of Adult Neural Stem Cells", *Stem Cells*, 26:2131-41.

Wang et al. (Apr. 2008) "Requirement of Nanog Dimerization for Stem Cell Self-Renewal and Pluripotency", *Proceedings of the National Academy of Science USA*, 105(17): 6326-31.

Theunissen et al. (Jan. 2011) "Nanog Overcomes Reprogramming Barriers and Induces Pluripotency in Minimal Conditions", *Current Biology*, 21(1): 65-71.

Silva et al. (2009) "Nanog is the Gateway to the Pluripotent Ground State", *Cell*, 138:722-37.

Go et al. (2008) "Forced Expression of Sox2 or Nanog in Human Bone Marrow Derived Mesenchymal Stem Cells Maintains Their Expansion and Differentiation Capabilities", *Experimental Cell Research*, 314(5): 1147-54 (Abstract only).

Liu et al. (2009) "Effects of Ectopic Nanog and Oct4 Overexpression on Mesenchymal Stem Cells", *Stem Cells and Development*, 18(7): 1013-22 (Abstract only).

Okano et al (2009) "Strategies Toward CNS-Regeneration Using Induced Pluripotent Stem Cells", *Genome Information*, 23(1): 217-20 (Abstract only).


Patel et al. (2010) "Advances in Reprogramming Somatic Stem Cells to Induced Pluripotent Stem Cells", *Stem Cell Reviews*, 6(3): 367-80.

Primary Examiner — Robert M Kelly

(74) *Attorney, Agent, or Firm* — Timothy H. Van Dyke; Beusse, Wolter, Sanks, Mora & Maire, P.A.

(57) **ABSTRACT**

Disclosed herein are methods and materials for producing a more developmentally potent cell from a less developmentally potent cell. Specifically exemplified herein are methods that comprise introducing an expressible dedifferentiating polynucleotide sequence into a less developmentally potent cell, wherein the transfected less developmentally potent cell becomes a more developmentally potent cell capable of differentiating to a less developmentally potent cell of its lineage or origin or a different lineage.



PRIOR ART

FIG. 1.




FIG. 2




FIG. 3




FIG. 4






FIG. 5




FIG. 6





FIG. 7



Green: β III-tubulin, Red: GFAP, Blue: DAPI


FIG. 8

FIG. 9: NanogP8 Sequence

```

tactaac
atgagtghggatccagattgtcccaaaagccttgcccttgccttgaagaatccgactgt.aaa
M S V D F A C P Q S L P C F E E S D C K
gaatcttcaacctatgcctgtgatttgtgggcoctgaagaaaaactatccatccttgc aaatg
E S S P M P V I C G P E E N Y P S L Q M
tcttctgctgagatgcctcaacacagagactgtctctcctccttccctccatggatctg
S S A E M F H S E S Y S P L P S S M D L
cttattcaggacagccctgattcttcaccagtcaccaaggcaaacaccactctctgaa
L I Q D S P D S S S P K G K Q P S S A
gagaatagtgtcgcaaaaaaggaagacaaggtcccggtcaagaaacagaagaccagaact
E N S V A K K E D K V P V K
gtgttctcttccaccccaagctgtgtgtactcaatgatagatttcaaaagacagaastacctc
agcctccagcagatccaaagaactctccaacatcctgaaacctcaactacaaaccaggtgaag
acctggttccagaaccagagaastgaaatctcaagaggtcccagaaaaacaactggccgaag
aatagcaatggtgtgacgcagaaggcoctcagcaactacctaccccagccctctactctcc
N S N G V S Q K A S A P S Y P S L Y S S
taccaccagggatgectggtgaaccocgacchgggaaccttccaatgtgagaaaccagarc
Y H Q G C L V N P S G N L P M S N Q S
tggaaacaattcaacctggagcaaccagaccagaaacatccagtcctggagcaaccactcc
S N N S S S N Q S Q N I Q S S N H S
tggaaactcagacctggtgcaccaatcctggaacaatcaggcoctggaacagtcccttc
S N S Q S C S Q S S N N Q A S N S P F
tataactgtggagaggaatctctgcagtcctgcacgacttccagccaaattctcctgcc
Y N C G E E S L Q S C M H F Q P N S P A
agtgacttggaggctgccttggaaagctgctggggaaaggcocttaatgtaatacagcagacc
S D L E A A L E A A G E G L N V I Q Q S
actaggtatcttagtactccacaaaccatggatttatctctaaactactccatgaacatg
S R Y F S S P Q S M D L F L N Y S M N M
caacctgaagacgtgtga
Q P E D V -
agatgagtgaacctgatattactcaatttcagtcctggacactggctgaactccttctc
cctcctcccatccctcataggatcttctcttggtttggaaaccacgtgcttcgggttccat
gatgcctatccagtcaatctcatggagggtggagtatggttggagcctaactcagcgaggt
ttcttttttttttttctattggatcttctggaaaaaatcttttttttttttttttgg
aaagggatcttgcctgcggccaggtggagtgcagtggggggtcttggctcactgc
aagetccgctccgggttccagccatctcctgectcagcctcccgagcagctggact
acagggcccccacctccgccggctaatattttgatttttagtagaacaggggttca
ctgtgttagccaggtggtctcgatctcctgaccttctgatccgcccgcctgggctccc
taaacagctggattccagggctgagccacgggcctgcctagaaaagacattttaataa
ccttggctgctaaaggacaacattgatagaagccgtctctggctatagatagtagatcta
atactagtttggatattcttaggggtttagaatctaacctcaagaatagaastacaagta
cgaattggctgatggaggtggtat

```



Key:
 Lane 3: 100bp ladder
 Lane 4: 10μM 5-aza-C
 Lane 5: 3μM 5-aza-C
 Lane 6: 1μM 5-aza-C

Electrophoresis 050101

Key:
 Lane 7: Negative control
 Lane 8: Positive control 1x
 Lane 9: Positive control 2x
 Lane 10: 100bp ladder

Treatment with 10, 3 or 1μM of 5azaC for 21 days, 5 days coculture.


FIG. 10

Electrophoresis 050115
Troponin I and hANP



FIG. 11

GATA-4 and MLC-2v



3 weeks of treatment (3uM BrdU or 5-azaC) and coculture for 7 days

Lane 2: ladder

Lane 3: Troponin I or GATA-4 low RNA of BrdU treatment

Lane 4: Troponin I or GATA-4 high RNA of BrdU treatment

Lane 5: Troponin I or GATA-4 low RNA of 5azaC treatment

Lane 6: Troponin I or GATA-4 high RNA of 5azaC treatment

Lane 7: ladder

Lane 8: hANP or MLC-2v low RNA of BrdU treatment

Lane 9: hANP or MLC-2v high RNA of BrdU treatment

Lane 10: hANP or MLC-2v low RNA of 5azaC treatment

Lane 11: hANP or MLC-2v high RNA of 5azaC treatment





FIG. 12



- Lane 1: 100bp ladder
- Lane 2: 3uM combined treatment 3 weeks
- Lane 3: 1uM combined treatment 3 weeks
- Lane 4: 3uM 5azaC 3 weeks
- Lane 5: 3uM BrdU 3 weeks
- Lane 6: 3uM Control (nt 12/27)
- Lane 7: 3uM combined treatment 3 weeks
- Lane 8: 1uM combined treatment 3 weeks
- Lane 9: 3uM 5azaC 3 weeks
- Lane 10: 3uM BrdU 3 weeks
- Lane 11: 3uM Control (nt 12/27)
- Lane 12: 100bp ladder

January 24, 2005

GATA-4 and MLC-2v



Troponin I and hANP




FIG. 13

- Lane 1: DNA ladder
- Lane 2: 10uM 5azaC treatment
- Lane 3: 3uM 5azaC treatment
- Lane 4: 1uM 5azaC treatment
- Lane 5: 3uM 5azaC treatment
- Lane 6: 3uM BrdU treatment
- Lane 7: 10uM 5azaC treatment
- Lane 8: 3uM 5azaC treatment
- Lane 9: 1uM 5azaC treatment
- Lane 10: 3uM 5azaC treatment
- Lane 11: 3uM BrdU treatment
- Lane 12: DNA ladder




FIG. 14




FIG. 15




FIG. 15

Nanog vector sequence analysis




FIG. 17

1

MESENCHYMAL STEM CELLS WITH INCREASED DEVELOPMENTAL POTENCY BY EXPRESSING NANOG

CROSS-REFERENCE TO RELATED APPLICATIONS

This application is a continuation of U.S. patent application Ser. No. 11/258,401 filed Oct. 24, 2005 (Now U.S. Pat. No. 8,192,988), and claims priority to U.S. Provisional Application No. 60/621,901 filed Oct. 22, 2004 and 60/650,438 filed Feb. 4, 2005, both of which are incorporated herein in their entirety.

BACKGROUND OF THE INVENTION

The use of stem cells for the treatment of neurodegenerative conditions offers the hope of curing diseases like Alzheimer's and Parkinson's by means of transplantation [1]. However, major obstacles regarding cell procurement, directing cell fate and avoiding immune response hinder clinical development [2-4] Research has focused on both adult and embryonic stem cells and attempted to balance limitations in regulating their development and preventing immune response. Increased potency of stem cells can be achieved by epigenetic modifications through nucleotide derivatives [5] and their lineage can be directed by gene transfection [6,7].

Patients currently suffering from neurodegenerative conditions have limited treatment options. Conventional drug therapy helps delay or reduce the symptoms of disease but is unable to restore complete functionality of the brain or repair damaged tissue. Through stem cell-based therapies, scientists aim to transplant cells in order to regenerate damaged tissue and restore proper function. However, the best source of stem cells for transplantation remains an unresolved issue; with debate focusing around embryonic or adult derived stem cells. Embryonic stem cells can be readily differentiated to multiple neuronal fates but pose the risk of tumor formation or immune response; whereas adult stem cell technology is easily accessible, but provides limited capacity for transdifferentiation. An optimal approach may be to increase cellular plasticity of adult stem cells for use in autologous transplantation.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 shows the vector system for cloning nanog according to the teachings in Example 1.

FIG. 2 shows images of cells before and after transfection with nanog: A: shows mesenchymal stem cells pre-transfection; B: shows nanog-transfected mesenchymal stem cells 12 days post-transfection; C: shows nanog-transfected mesenchymal stem cells 4 weeks post-transfection.

FIG. 3 shows images of transfected mesenchymal stem cells 9 days (A and B) and 2 months (C and D) post transfection.

FIG. 4 shows images a co-culture system in accord with one embodiment of the subject invention.

FIG. 5 shows images of Co-culturing experiments which demonstrated that embryoid body-like clusters began differentiation within 48 hours (A). Control cells with our empty vector treatments failed to show any signs of neural differentiation (B). Embryoid-like bodies adhered to membrane and differentiation occurred as neural cells migrated radially outward (C).

2

FIG. 6-7 show images of the clustering of nanog transfected cells.

FIG. 8 shows images of MeSC-derived neurons and astrocytes.

FIG. 9 shows sequence of Nanog encoding polynucleotide (SEQ ID NO: 1) and corresponding polypeptide sequence (SEQ ID NO: 2).

FIGS. 10-13 shows gel images of gene expression in cells subjected to various treatments demonstrating an ability to increase potency of mesenchymal stem cells and differentiation into cardiac cells.

FIGS. 14-16 show photograph images of cells subjected to various treatments.

FIG. 17 shows a schematic representation of the nanog sequence cloned inside a CMV mammalian promoter vector.

DETAILED DESCRIPTION

In reviewing the detailed disclosure which follows, and the specification more generally, it should be borne in mind that all patents, patent applications, patent publications, technical publications, scientific publications, and other references referenced herein are hereby incorporated by reference in this application, in their entirety to the extent not inconsistent with the teachings herein.

Reference to particular buffers, media, reagents, cells, culture conditions and the like, or to some subclass of same, is not intended to be limiting, but should be read to include all such related materials that one of ordinary skill in the art would recognize as being of interest or value in the particular context in which that discussion is presented. For example, it is often possible to substitute one buffer system or culture medium for another, such that a different but known way is used to achieve the same goals as those to which the use of a suggested method, material or composition is directed.

It is important to an understanding of the present invention to note that all technical and scientific terms used herein, unless defined herein, are intended to have the same meaning as commonly understood by one of ordinary skill in the art. The techniques employed herein are also those that are known to one of ordinary skill in the art, unless stated otherwise. For purposes of more clearly facilitating an understanding of the invention as disclosed and claimed herein, the following definitions are provided.

The differentiation of stem cells along multiple lineages has been intensely studied given their great therapeutic potential. However, the mechanisms that underlie proliferation, self-renewal and differentiation in cells with the capacity for further development remains poorly understood. A recently discovered gene, nanog, is required to sustain pluripotency in embryonic stem cells and acts concomitantly with embryonic transcription factor Oct-4, yet utilizes a STAT-3 independent mechanism. The subject invention is based on the inventor's discovery that gene transfection of adult stem cells with nanog, an embryonic stem cell gene maintaining pluripotency [8,9], can allow for the production of neurons and astrocytes from bone marrow cells via a two-step process. First, mesenchymal stem cells are modified by nanog transfection, and the cells form embryoid-like bodies. Then cells are committed to neuronal lineage in a co-culture system with differentiated neural stem cells separated by a semi-permeable membrane. This technology may be a means of generating effective autologous stem cell transplants to improve neuroreplacement strategies. The inventors have discovered that adult stem cells can be dedifferentiated through introduction and expression of the nanog gene or other dedifferentiating genes.

Thus, in one embodiment, the invention provides methods for making a more developmentally potent cell from a less developmentally potent cell. In a typical embodiment, the method comprises the step of introducing an expressible dedifferentiating polynucleotide sequence into a less developmentally potent cell, wherein the transfected less developmentally potent cell becomes a more developmentally potent cell capable of differentiating to a less developmentally potent cell of its lineage of origin or a different lineage. In certain embodiments, the inventive methods further comprise the step of co-culturing the transfected less developmentally potent with neural-lineage cells or media conditioned with neural-lineage cells, wherein the transfected cells become a more developmentally potent cells capable of differentiating to a less developmentally potent cells of its lineage of origin or a different lineage.

In the practice of one embodiment of the invention, the phenotype of the less developmentally potent cell is changed when it becomes a more developmentally potent cell. Thus, the invention provides methods for changing a first phenotype of a less developmentally potent cell into a second phenotype of more developmentally potent cell. The change from a certain potency to a higher level of potency is considered "dedifferentiation" in accord with the teachings herein. In preferred embodiments, the less developmentally potent cell is a stem cell, more preferably a hematopoietic stem cell, a neural stem cell, an epithelial stem cell, an epidermal stem cell, a retinal stem cell, an adipose stem cell and a mesenchymal stem cell.

In yet further aspects of the invention are provided pharmaceutical compositions comprising said more developmentally potent cells prepared according to the methods of the invention and a pharmaceutically-acceptable carrier or excipient. The invention provides such pharmaceutical compositions comprising said more developmentally potent cells that are tissue stem cells for use in cell or tissue regeneration or for correcting a disease or disorder in a tissue or animal in need thereof.

Thus, the invention also provides methods for using the pharmaceutical compositions provided herein to treat an animal in need thereof by administering the more developmentally potent cells thereto. In certain preferred embodiments, the more developmentally potent cells comprise a cluster of two or more of the more developmentally potent cells. Preferably, the animal has a corporal or neurological deficit that can be treated or ameliorated by administration of said more developmentally potent cells, such as a deficit caused by a neurodegenerative disease, a traumatic injury, a neurotoxic injury, ischemia, a developmental disorder, a disorder affecting vision, an injury or disease of the spinal cord, a demyelinating disease, an autoimmune disease, an infection, an inflammatory disease, or corporal disease, disorder, injury, trauma, malfunction, degeneration or loss. In preferred embodiments, the one or plurality of more developmentally potent cells are capable of migrating to an area of tissue damage, differentiating in a tissue-specific manner and functioning in a manner that reduces the neurological or corporal deficit. As provided by the methods of the invention herein, the cells are administered by injecting one or a plurality of more developmentally potent cells with a syringe, inserting the more developmentally potent cells with a catheter or surgically implanting the more developmentally potent cells. In certain embodiments, the more developmentally potent cells are injected with a syringe into a body cavity that is fluidly-connected to the area of neurological or corporal deficit. In certain preferred embodiments, the body cavity is a brain ventricle. In other embodiments, the more developmen-

tally potent cells are inserted with a catheter into a body cavity that is fluidly-connected to the area of neurological or corporal deficit. In certain preferred embodiments, the body cavity is a brain ventricle. In still further additional embodiments, the more developmentally potent cells are surgically implanted into a body cavity that is fluidly-connected to the area of neurological or corporal deficit. In certain preferred embodiments, the body cavity is a brain ventricle. The more developmentally potent cells can also alternatively be inserted using a syringe or catheter or surgically implanted directly at the site of the neurological or corporal deficit or systemically (e.g., intravenously).

Also, the more developmentally potent cells may be further genetically modified through introduction of polynucleotide sequences the bias against differentiation to certain cell types or bias toward differentiation to certain cell types. Provisional Application Nos. 60/621,483 and 60/621,902 naming Dr. Kiminobu Sugaya as an inventor and entitled "Methods and Products For Biasing Development" are incorporated by reference to provide examples of negative and positive biasing that could be subjected to the more developmentally potent cells produced by the methods taught herein.

Administration of the one or a plurality of more developmentally potent cells into an animal results in said cells differentiating into a terminally-differentiated cell. Thus, the invention provides methods for making a terminally-differentiated cell, comprising the step of administering the more developmentally potent cells of the invention into an animal in need thereof. As provided by the methods of the invention herein, the cells are administered by injecting the more developmentally potent cells with a syringe, inserting the more developmentally potent cells with a catheter or surgically implanting the more developmentally potent cells. In certain embodiments, the more developmentally potent cells are injected with a syringe into a body cavity that is fluidly-connected to the area of neurological or corporal deficit. In certain preferred embodiments, the body cavity is a brain ventricle. In other embodiments, the more developmentally potent cells are inserted with a catheter into a body cavity that is fluidly-connected to the area of neurological or corporal deficit. In certain preferred embodiments, the body cavity is a brain ventricle. In still further additional embodiments, the more developmentally potent cells are surgically implanted into a body cavity that is fluidly-connected to the area of neurological or corporal deficit. In certain preferred embodiments, the body cavity is a brain ventricle. The more developmentally potent cells can also alternatively be inserted using a syringe or catheter or surgically implanted directly at the site of the neurological or corporal deficit or systemically (e.g., intravenously).

In yet another embodiment, the invention relates to treating a stem cell, excluding those of neural origin, such that it is converted into a more developmentally potent cell, which enables it to differentiate into the various cell types found in eye tissue, inter alia, choroid, iris and retinal pigment epithelium cells, rod and cone photoreceptor cells, horizontal cells, bipolar neurons, amacrine, ganglion and optic nerve cells. These non-limiting, exemplary cell types found in eye tissue are collectively referred to as retinal cells. The methods comprising the step of contacting more developmentally potent cells of the invention with an effective amount of one or a combination of growth factor selected from the group consisting of TGF- β 3, IGF-1 and CNTF for an effective period such that the growth factor-contacted cells can differentiate into retinal cells.

As used herein, the terms "multipotent neural stem cells (MNSCs)," "neural stem cells (NSCs)" and "neural progeni-

tor cells (NPCs)" refer to undifferentiated, multipotent cells of the CNS. Such terms are commonly used in the scientific literature. MNSCs can differentiate into tissue-specific cell types, for example astrocytes, oligodendrocytes, and neurons when transplanted in the brain. MNSCs of the invention are distinguished from natural MNSCs by their adaptation for proliferation, migration and differentiation in mammalian host tissue when introduced thereto.

As used herein, a "less developmentally potent cell" is a cell that is capable of limited multi-lineage differentiation or capable of single-lineage, tissue-specific differentiation, for example, an untreated mesenchymal stem cell can differentiate into, inter alia, osteocytes and chondrocytes, i.e., cells of mesenchymal lineage, but has only limited ability to differentiate into cells of other lineages (e.g., neural lineage).

As used herein, a "more developmentally potent cell" is a cell that is readily capable of differentiating into a greater variety of cell types than its corresponding less developmentally potent cell. For example, a mesenchymal stem cell can readily differentiate into osteocytes and chondrocytes but has only limited ability to differentiate into neural or retinal lineage cells (i.e., it is a less developmentally potent cell in this context). Mesenchymal stem cells treated according to the methods of the invention become more developmentally potent because they can readily differentiate into, for example, mesenchymal-lineage and neural-lineage cell types; the plasticity of the cells is increased when treated according to the methods of the invention.

The invention provides methods of delivery and transplantation of the more developmentally potent cells of the invention to ameliorate the effects of age, physical and biological trauma and degenerative disease on the brain or central nervous system of an animal, as well as other tissues such as, for example, retinal tissue. It is well recognized in the art that transplantation of tissue into the CNS offers the potential for treatment of neurodegenerative disorders and CNS damage due to injury. Transplantation of new cells into the damaged CNS has the potential to repair damaged circuitries and provide neurotransmitters thereby restoring neurological function. It is also recognized in the art that transplantation into other tissue, such as eye tissue, offers the potential for treatment of degenerative disorders and tissue damage due to injury. As disclosed herein, the invention provides methods for generating more developmentally potent cells adapted for proliferation, migration and differentiation in mammalian tissue when introduced thereto. The use of more developmentally potent cells in the treatment of neurological disorders and CNS damage, as well as the use of more developmentally potent cells in the treatment of other tissue damage or degeneration, can be demonstrated by the use of established animal models known in the art.

In one embodiment dedifferentiated cells or more developmentally potent cells of the invention can be administered to an animal with abnormal or degenerative symptoms obtained in any manner, including those obtained as a result of age, physical or biological trauma, or neurodegenerative disease and the like, or animal models created by man using recombinant genetic techniques, such as transgenic and "gene knockout" animals.

Recipients of the more developmentally potent cells of the invention can be immunosuppressed, either through the use of immunosuppressive drugs such as cyclosporin, or through local immunosuppression strategies employing locally applied immunosuppressants, but such immunosuppression need not necessarily be a prerequisite in certain immunoprivileged tissues such as, for example, brain and eye tissues. In certain embodiments, the delivery method of the invention

can cause less localized tissue damage to the site of cell damage or malfunction than existing methods of delivery.

More developmentally potent cells of the invention can be prepared from the recipient's own tissue. In such instances, the progeny of the more developmentally potent cells can be generated from dissociated or isolated tissue and proliferated in vitro using the methods described herein. In the case of mesenchymal stem cells (MeSCs), progeny can be generated from MeSCs isolated from, for example, bone marrow. Upon suitable expansion of cell numbers, the stem cells of the invention can be harvested and readied for administration into the recipient's affected tissue.

There are significant differences in the method of delivery to the brain of the more developmentally potent cells compared to the prior art. One exemplary difference is as follows: the more developmentally potent cells of the invention are transplanted intraventricularly. Further, while the transplantation of one or more separate more developmentally potent cells is efficacious, the more developmentally potent cells of the invention are preferably transplanted in the form of clusters of two or more cells via a surgical procedure or injection using a syringe large enough to leave the clusters substantially intact. The results disclosed in the Examples below indicate that ventricular delivery of more developmentally potent cells of the invention in the form of a cluster of two or more cells can result in migration to the area of damage in the brain and proper neuronal differentiation. Another benefit of intraventricular injection is less tissue destruction, resulting in less localized recruitment of immune cells by the host. This is evidenced by the lack of ventricular distortion, tumor formation, and increased host astrocyte staining without any immunosuppression.

The method of delivery of the more developmentally potent cells of the invention to the brain can be essentially duplicated for other immunoprivileged tissue such as, for example, the eye. Delivery of one or more separate or two or more of the more developmentally potent cells in the form of a cluster via injection using a syringe large enough to leave the any cluster of two or more cells that is present substantially intact can result in migration to the area of damage in the eye and proper tissue-specific differentiation.

In the context of the present application, a polynucleotide sequence is "homologous" with the sequence according to the invention if at least 70%, preferably at least 80%, most preferably at least 90% of its base composition and base sequence corresponds to the sequence according to the invention. According to the invention, a "homologous protein" is to be understood to comprise proteins which contain an amino acid sequence at least 70% of which, preferably at least 80% of which, most preferably at least 90% of which, corresponds to the amino acid sequence shown in FIG. 9; wherein corresponds is to be understood to mean that the corresponding amino acids are either identical or are mutually homologous amino acids. The expression "homologous amino acids" denotes those which have corresponding properties, particularly with regard to their charge, hydrophobic character, steric properties, etc. Thus, in one embodiment the protein may be from 70% up to less than 100% homologous to nanog.

Homology, sequence similarity or sequence identity of nucleotide or amino acid sequences may be determined conventionally by using known software or computer programs such as the BestFit or Gap pairwise comparison programs (GCG Wisconsin Package, Genetics Computer Group, 575 Science Drive, Madison, Wis. 53711). BestFit uses the local homology algorithm of Smith and Waterman, *Advances in Applied Mathematics* 2: 482-489 (1981), to find the best segment of identity or similarity between two sequences. Gap

performs global alignments: all of one sequence with all of another similar sequence using the method of Needleman and Wunsch, *J. Mol. Biol.* 48:443-453 (1970). When using a sequence alignment program such as BestFit, to determine the degree of sequence homology, similarity or identity, the default setting may be used, or an appropriate scoring matrix may be selected to optimize identity, similarity or homology scores. Similarly, when using a program such as BestFit to determine sequence identity, similarity or homology between two different amino acid sequences, the default settings may be used, or an appropriate scoring matrix, such as *blosum45* or *blosum80*, may be selected to optimize identity, similarity or homology scores.

The term "isolated" means separated from its natural environment.

The term "polynucleotide" refers in general to polyribonucleotides and polydeoxyribonucleotides, and can denote an unmodified RNA or DNA or a modified RNA or DNA.

The term "polypeptides" is to be understood to mean peptides or proteins which contain two or more amino acids which are bound via peptide bonds.

The polypeptides for use in accord with the teachings herein include polypeptides corresponding to *nanog*, and also includes those, at least 70% of which, preferably at least 80% of which, are homologous with the polypeptide corresponding to *nanog*, and most preferably those which exhibit a homology of least 90% to 95% with the polypeptide corresponding to *nanog* and which have dedifferentiating influence. See polypeptide sequence provided in FIG. 9. Thus, the polypeptides may have a homology of from 70% to up to 100% with respect to *nanog*.

As used herein, a "polypeptide sequence exhibiting dedifferentiating influence" is a polypeptide whose presence in the cell causes an increase in potency, or transformation from a less developmentally potent cell to a more developmentally potent cell. Examples of such polypeptide sequences include the expression products of the *nanog* gene, and polynucleotide sequences that hybridize to the complement of the sequence in FIG. 9, as well as expression products of the polynucleotide sequences listed in Table 1 below in Example 3.

The terms "stringent conditions" or "stringent hybridization conditions" includes reference to conditions under which a polynucleotide will hybridize to its target sequence, to a detectably greater degree than other sequences (e.g., at least 2-fold over background). Stringent conditions are sequence-dependent and will be different in different circumstances. By controlling the stringency of the hybridization and/or washing conditions, target sequences can be identified which are 100% complementary to the probe (homologous probing). Alternatively, stringency conditions can be adjusted to allow some mismatching in sequences so that lower degrees of similarity are detected (heterologous probing).

Typically, stringent conditions will be those in which the salt concentration is less than about 1.5 M Na ion, typically about 0.01 to 1.0 M Na ion concentration (or other salts) at pH 7.0 to 8.3 and the temperature is at least about 30° C. for short probes (e.g., 10 to 50 nucleotides) and at least about 60° C. for long probes (e.g., greater than 50 nucleotides). Stringent conditions may also be achieved with the addition of destabilizing agents such as formamide. Exemplary low stringency conditions include hybridization with a buffer solution of 30 to 35% formamide, 1 M NaCl, 1% SDS (sodium dodecyl sulphate) at 37° C., and a wash in 1× to 2×SSC (20×SSC=3.0 M NaCl/0.3 M trisodium citrate) at 50 to 55° C. Exemplary moderate stringency conditions include hybridization in 40 to 45% formamide, 1 M NaCl, 1% SDS at 37° C., and a wash in

0.5× to 1×SSC at 55 to 60° C. Exemplary high stringency conditions include hybridization in 50% formamide, 1 M NaCl, 1% SDS at 37° C., and a wash in 0.1×SSC at 60 to 65° C.

Specificity is typically the function of post-hybridization washes, the critical factors being the ionic strength and temperature of the final wash solution. For DNA—DNA hybrids, the T_m can be approximated from the equation of Meinkoth and Wahl, *Anal. Biochem.*, 138:267-284 (1984): $T_m = 81.5^\circ C. + 16.6 (\log M) + 0.41 (\% GC) - 0.61 (\% \text{ form}) - 500/L$; where M is the molarity of monovalent cations, % GC is the percentage of guanosine and cytosine nucleotides in the DNA, % form is the percentage of formamide in the hybridization solution, and L is the length of the hybrid in base pairs. The T_m is the temperature (under defined ionic strength and pH) at which 50% of a complementary target sequence hybridizes to a perfectly matched probe. T_m is reduced by about 1° C. for each 1% of mismatching; thus, T_m , hybridization and/or wash conditions can be adjusted to hybridize to sequences of the desired identity. For example, if sequences with approximately 90% identity are sought, the T_m can be decreased 10° C. Generally, stringent conditions are selected to be about 5° C. lower than the thermal melting point (T_m) for the specific sequence and its complement at a defined ionic strength and pH. However, severely stringent conditions can utilize a hybridization and/or wash at 1, 2, 3, or 4° C. lower than the thermal melting point (T_m); moderately stringent conditions can utilize a hybridization and/or wash at 6, 7, 8, 9, or 10° C. lower than the thermal melting point (T_m); low stringency conditions can utilize a hybridization and/or wash at 11, 12, 13, 14, 15, or 20° C. lower than the thermal melting point (T_m). Using the equation, hybridization and wash compositions, and desired T_m , those of ordinary skill will understand that variations in the stringency of hybridization and/or wash solutions are inherently described. If the desired degree of mismatching results in a T_m of less than 45° C. (aqueous solution) or 32° C. (formamide solution) it is preferred to increase the SSC concentration so that a higher temperature can be used. An extensive guide to the hybridization of nucleic acids is found in *Current Protocols in Molecular Biology*, Chapter 2, Ausubel, et al., Eds., Greene Publishing and Wiley-Interscience, New York (2000).

Accordingly, polynucleotide sequences that hybridize to the complement of the sequence in FIG. 9 are contemplated for use in dedifferentiating cells as taught herein.

US. Patent Application Nos. 2003/0219898, 2003/0148513, and 2003/0139410 are incorporated by reference to the extent they are not inconsistent with the teachings herein. These first two of these patent applications describe multiple uses of increased potency cells obtained from the taught methods, and in particular, the implantation of stem cells for different therapeutic treatments of neurological trauma and degenerative conditions. The third patent application is directed to the use of certain compounds to stimulate proliferation and migration of stem cells. Those skilled in the art will readily appreciate that the dedifferentiated cells of the subject invention could be substituted in place of the potent cells taught in the aforementioned patent applications, without undue experimentation. Also, the methods of the third patent may be combined with the present invention without undue experimentation.

According to another embodiment, the subject invention comprises a method of influencing transcription of an endogenous polynucleotide sequence comprising contacting a non-embryonic cell or cellular component comprising an endogenous polynucleotide sequence with a *nanog* protein or protein encoded by a polynucleotide sequence that hybridizes

to a complement of the sequence shown in FIG. 9 under stringent conditions (i.e. nanog-like protein). Such influence may further include, but is not limited to, demethylation of DNA and reversal histone acetylation. The nanog protein or nanog-like protein may be one expressed by a polynucleotide sequence introduced in the cell or cellular component, or protein delivered into the cell or cellular component, or protein expressed by an endogenous polynucleotide sequence that has been activated. Nanog expression may be activated by the provision of Oct 4 and/or Sox2, which typically form a dimer. In a specific embodiment, the cellular component is a nucleus, liposome, or mitochondria. Such endogenous polynucleotide sequence or cellular component contacted by nanog or nanog may be removed from a cell or cellular component and introduced into another cell or cellular component.

In another specific embodiment, the invention pertains to increasing the efficacy of nuclear transfer comprising transfecting a nucleus with a polynucleotide encoding nanog or nanog-like protein to obtain a treated nucleus and introducing the treated nucleus into a cell. The cell may be any suitable cell but would typically be an ovum with its nucleus removed.

EXAMPLE 1

Dedifferentiation of Mesenchymal Stem Cells

Introduction

Embryonic stem cells are derived during the blastocyst stage from the inner cell mass of prenatal mammalia; and possess the intrinsic properties of rapid self-renewal and pluripotency. Under the influence of endogenous and extracellular signals, these cells migrate and differentiate during the developmental process. Extracellular signals regulating self-renewal or differentiation have been demonstrated in vitro by differentiating embryonic stem cells into cell types comprising all three germ layers. These varieties include neuronal, pancreatic, cardiac and hematopoietic tissue using well-established culturing protocols. Embryonic stem cells form embryoid bodies, non-adherent proliferating clusters, in the presence of leukemia-inhibitory factor (LIF) and a feeder layer of typically fibroblast cells. Upon removal of LIF or transfer to non-feeder cell cultures, embryoid bodies undergo spontaneous differentiation. Early differentiation is characterized by loss of stem cell-specific surface antigens (SSEA-1) and alkaline phosphatase activity. Additionally, endogenous signals, including regulatory intracellular proteins, continually change throughout development. Numerous gene expression studies show distinct variations among different embryonic and adult stem cells, pointing toward underlying mechanisms responsible for the continual loss of potency corresponding with differentiation. Several key genes, namely Oct-3/4, LIF, DNMT3B and Nanog, are repeatedly shown to be almost exclusively expressed in embryonic stem cells that regulate pluripotency [10-14]. The immediate down-regulation of these genes may explain irreversible loss of potency, making embryonic stem cells an attractive source for clinical therapies. However, serious questions remain concerning the production of these cells in sufficient quantities for therapies, bioethical potential immune response and tumor formation [15].

The inventors believe that adult stem cells offer a practicable alternative to the use of embryonic tissue as they are easily harvested and potentially taken from autologous sources to preclude immune response. Stem cell populations have been found in several adult tissues including adipose [16], muscle [17], pancreas [18] and liver [19] and primarily

bone marrow [20-23]; all potential sources for cellular transplants [24]. Previous in vitro studies with adult bone marrow-derived stem cells have demonstrated the ability to differentiate into brain [21], liver [23] and cardiac cells [22]. In vivo studies have shown evidence that adult stem cells can migrate and differentiate into various tissues, albeit at extremely low frequencies [20]. However, challenges have been raised over the plasticity of these cells given both the low frequencies of detected cells and new found evidence of cell fusion, in conjunction with false positives [25-27]. An ideal therapeutic alternative may exist if adult cells can be dedifferentiated to an embryonic-like state and recommitted to differentiate to a desired cell fate.

Nanog, also referred to as early embryo specific NK (ENK) [28], is a recently discovered gene responsible for maintaining pluripotency in embryonic stem cells [8,9,28, 29]. This unique gene and its cousin, Nanog2, are genetically distinct members of the ANTP class of homeodomain proteins [30] and have at least twelve identified pseudogenes [31]. Structurally, Nanog contains three alpha helices encoded within the homeodomain portion and can be divided into three regions with respect to the central homeodomain sequence [30]. The N-terminal region is rich in serine and threonine residues indicating phosphate-regulated transactivation, possibly through SMAD4 interactions, while the C-terminal domain is seven times as active with an unusual motif of equally spaced tryptophans separated by four amino acids, each flanked with serine or threonine residues. Gene expression studies have shown nanog to be active in embryonic stem cells, tumors and some adult tissue. Nanog expression precipitously decreases with differentiation and maintains self-renewal in embryonic stem cells by gene transfection. In culture, nanog guards against differentiation and acts concomitantly with Oct-4, Wnt and BMP-4, yet utilizes a STAT-3 independent mechanism to maintain an undifferentiated state. Inventors believe that the role of nanog in regulating pluripotency makes this gene a potential candidate for increasing the potency of adult stem cells.

Previous studies regulating gene expression in stem cell lines have provided valuable insight into underlying mechanisms of proliferation, self-renewal and differentiation. Gene manipulation experiments can either prevent or enhance differentiation. In particular, differentiation can be prevented in embryonic stem cell lines by over expression of Pem or nanog, genes that regulate pluripotency. Conversely, over-expression of lineage specific gene Nurr1 promotes the differentiation of neural stem cell lines to produce dopamine-secreting cells. Taken together, gene vectors can maintain cells in a specific state or allow for lineage committed cells to develop into a specific subpopulation. In one embodiment, the subject invention pertains to a method of dedifferentiating adult stem cells by expressing genes regulating pluripotency to enhance transdifferentiation. This technology allows for adult cells to be used for autologous transplantation and thereby provide a greater understanding of stem cell biology.

FIG. 17 shows a schematic representation of the nanog sequence cloned inside a CMV mammalian promoter vector. The 5'UTR contains an Oct-4 and Sox2 binding region. The nanog protein coding sequence can be divided into an N-terminal, homeodomain and C-terminal region. The C-terminal region can be further subdivided into a C1, Cw and C2 domains. The 3' UTR contains an Alu sequence element.

Methods and Results

Human mesenchymal stem cells (hMeSC) are initially plated in 6-well plates, adhere to the surface and allowed to divide to varying degrees of confluence. They are cultured in serum-DMEM (Dulbecco's modified Eagle's medium) con-

- nocompetent rat: immune response following grafting of expanded neural precursor cells. *Neuroscience*. 2001; 106(1):201-16.
3. Trojanowski J Q, Kleppner S R, Hartley R S, Miyazono M, Fraser N W, Kesari S, Lee V M. Transfectable and transplantable postmitotic human neurons: a potential "platform" for gene therapy of nervous system diseases. *Exp Neurol*. 1997; 144(1):92-7.
 4. Zlokovic B V, Apuzzo M L. Cellular and molecular neurosurgery: pathways from concept to reality—part I: target disorders and concept approaches to gene therapy of the central nervous system. *Neurosurgery*. 1997 April; 40(4): 789-804
 5. Qu T Y, Dong X J, Sugaya I, Vaghani A, Pulido J, Sugaya K. Bromodeoxyuridine increases multipotency of human bone marrow-derived stem cells. (in press).
 6. Chang G, Sugaya K. Inner ear hair cell generation from human neural stem cells through Math1 transfection. Program No. 303.1. 2004 Abstract Viewer/Itinerary Planner. Washington, D.C.: Society for Neuroscience, 2004.
 7. Wagner J, Akerl J, Castro D S, Holm P C, Canals J M, Snyder E Y, Perlmann T, Arenas E. Induction of a midbrain dopaminergic phenotype in Nurr1-overexpressing neural stem cells by type 1 astrocytes. *Nat Biotechnol*. 1999 July; 17(7):653-9.
 8. Chambers I, Colby D, Robertson M, Nichols J, Lee S, Tweedie S, Smith A. Functional expression cloning of Nanog, a pluripotency sustaining factor in embryonic stem cells. *Cell*. 2003; 113(5):643-55.
 9. Mitsui K, Tokuzawa Y, Itoh H, Segawa K, Murakami M, Takahashi K, Maruyama M, Maeda M, Yamanaka S. The homeoprotein Nanog is required for maintenance of pluripotency in mouse epiblast and ES cells. *Cell*. 2003; 113(5):631-42.
 10. Bhattacharya B, Miura T, Brandenberger R, Majido J, Luo Y, Yang A X, Joshi B H, Ginis I, Thies R S, Amit M, Lyons I, Condie B G, Itskovitz-Eldor J, Rao M S, Puri R K. Gene expression in human embryonic stem cell lines: Unique molecular signature. *Blood*. 2004; 103(8): 2956-2964.
 11. Calhoun J D, Rao R R, Warrenfeltz S, Rekaya R, Dalton S, McDonald J, Stice S L. Transcriptional profiling of initial differentiation events in human embryonic stem cells. *Biochem Biophys Res Commun*. 2004 Oct. 15; 323(2):453-64.
 12. Clark A T, Rodriguez R T, Bodnar M S, Abeyta M J, Cedars M I, Turek P J, Firpo M T, Reijo Pera R A. Human STELLAR, NANOG, and GDF3 genes are expressed in pluripotent cells and map to chromosome 12p13, a hotspot for teratocarcinoma. *Stem Cells*. 2004; 22(2): 169-179.
 13. Spencer J M, Chen X, Draper J S, Antosiewicz J E, Chon C H, Jones S B, Brooks J D, Andrews P W, Brown P O, Thomson J A. Gene expression patterns in human embryonic stem cells and human pluripotent germ cell tumors. *Proc Nat Acad Sci USA*. 2003; 100(23): 13350-13355.
 14. Tanaka T S, Kunath T, Kimber W L, Jaradat S A, Stagg C A, Usada M, Yokota T, Niwa H, Rossant J, Ko M. Gene expression profiling of embryo-derived stem cells reveals candidate genes associated with pluripotency and lineage specificity. *Genome Res*. 2002; 12: 1921-1928.
 15. Erdo F, Buhrle C, Blunk J, Hoehn M, Xia Y, Fleischmann B, Focking M, Kustermann E, Kolossov E, Hescheler J, Hossmann K A, Trapp T. Host-dependent tumorigenesis of embryonic stem cell transplantation in experimental stroke. *J Cereb Blood Flow Metab*. 2003 July; 23(7):780-5.
 16. Gimble J, Guilak F. Adipose-derived adult stem cells: isolation, characterization, and differentiation potential. *Cytotherapy*. 2003; 5(5):362-9.

17. Cao B, Huard J. Muscle-derived stem cells. *Cell Cycle*. 2004 February; 3(2):104-7.
18. Hess D, Li L, Martin M, Sakano S, Hill D, Strutt B, Thyssen S, Gray D A, Bhatia M. Bone marrow-derived stem cells initiate pancreatic regeneration. *Nat Biotechnol*. 2003 July; 21(7):763-70.
19. Muller-Borer B J, Cascio W E, Anderson P A, Snowwaert J N, Frye J R, Desai N, Esch G L, Brackham J A, Bagnell C R, Coleman W B, Grisham J W, Malouf N N. Adult-derived liver stem cells acquire a cardiomyocyte structural and functional phenotype ex vivo. *Am J Pathol*. 2004 July; 165(1):135-45.
20. Jiang Y, Jahagirdar B N, Reinhardt R L, Schwartz R E, Keene C D, Ortiz-Gonzalez X R, Reyes M, Lenvik T, Lund T, Blackstad M, Du J, Aldrich S, Lisberg A, Low W C, Largaespada D A, Verfaillie C M. Pluripotency of mesenchymal stem cells derived from adult marrow. *Nature*. 2002 Jul. 4; 418(6893):41-9.
21. Brazelton T R, Rossi F M, Keshet G I, Blau H M. From marrow to brain: expression of neuronal phenotypes in adult mice. *Science*. 2000 Dec. 1; 290(5497):1775-9.
22. Fukuda K. Use of adult marrow mesenchymal stem cells for regeneration of cardiomyocytes. *Bone Marrow Transplant*. 2003 August; 32 Suppl 1:S25-7.
23. Theise N D, Nimmakayalu M, Gardner R, Illei P B, Morgan G, Teperman L, Henegariu O, Krause D S. Liver from bone marrow in humans. *Hepatology*. 2000 July; 32(1):11-6.
24. Hedrick M H, Daniels E J. The use of adult stem cells in regenerative medicine. *Clin Plast Surg*. 2003 October; 30(4):499-505.
25. Alvarez-Dolado M, Pardal R, Garcia-Verdugo J M, Fike J R, Lee H O, Pfeffer K, Lois C, Morrison S J, Alvarez-Buylla A. Fusion of bone-marrow-derived cells with Purkinje neurons, cardiomyocytes and hepatocytes. *Nature*. 2003 Oct. 30; 425(6961):968-73
26. Terada N, Hamazaki T, Oka M, Hoki M, Mastalerz D M, Nakano Y, Meyer E M, Morel L, Petersen B E, Scott E W. Bone marrow cells adopt the phenotype of other cells by spontaneous cell fusion. *Nature*. 2002 Apr. 4; 416(6880): 542-5.
27. Wu T, Cieply K, Nalesnik M A, Randhawa P S, Sonzogni A, Bellamy C, Abu-Elmagd K, Michalopolous G K, Jaffe R, Kormos R L, Gridelli B, Fung J J, Demetris A J. Minimal evidence of transdifferentiation from recipient bone marrow to parenchymal cells in regenerating and long-surviving human allografts. *Am J Transplant*. 2003 September; 3(9):1173-81.
28. Wang S, Tsai M, Chiang M, Li H. A novel NK-type homeobox gene, ENK (early embryo specific NK), preferentially expressed in embryonic stem cells. *Gene Exp Patterns* 2003; 3: 99-103.
29. Hart A H, Hartley L, Ibrahim M, Robb L. Identification, cloning and expression analysis of the pluripotency promoting Nanog genes in mouse and human. *Dev Dyn*. 2004; 230(1): 187-198.
30. Pan G J, Pei D Q. Identification of two distinct transactivation domains in the pluripotency sustaining factor nanog. *Cell Res*. 13(6): 499-502.
31. Booth H A, Holland P W. Eleven daughters of NANOG. *Genomics*. 2004; 84(2): 229-238.

EXAMPLE 4

Cardiac Differentiation of Human Mesenchymal Cells

Cardiac differentiation of human mesenchymal stem cells (MeSCs) is achieved through treatment with nucleotide

15

derivatives BrdU and 5-azaC and/or forced expression of embryonic stem cell gene nanog. Following treatment, cells were placed in a co-culture with cardiac cells (cardiomyocyte cell line H9c2). Human MeSCs plated in 6-well culture plates and expanded in serum-DMEM (Delbecco's Modified Eagle's Medium) containing 10% MeSC formulated fetal bovine serum (FBS, Stem Cell, Inc) containing antibiotics/antimycotics. MeSCs were treated with varying concentrations (1-10 uM) of BrdU and/or 5-azaC for 3 weeks or transfected with mammalian expression vector containing a nanog encoding sequence. Cell media was changed every three days prior to co-culture with cardiomyocytes. Cardiomyocytes were expanded and grown to near confluence in serum-DMEM with 10% non-conditioned FBS and antibiotics/antimycotics. To differentiate cardiac cells, serum media was replaced with plain DMEM containing only antibiotics and allowed to differentiate. Co-cultures were created by combining MeSCs with cardiac cells and culturing in cardiac media. Following co-culture, cells were treated with TRIzol and gene expression was assessed using RT-PCR and cardiac specific primers. Gel electrophoresis of samples revealed expression of cardiac specific genes following treatment.

The first (Electrophoresis 050101, FIG. 10) shows screens for each primer tested. The key shows which sample is in a given lane. Negative control represents untreated mesenchymal stem cells and Positive controls 1x and 2x are nanog transfected MeSCs (2x indicates that well received twice the number of rat cardiac cells, not twice the amount of nanog transfected cells). Each sample was co-cultured with rat car-

16

diomyocytes and primers are human specific and represent markers of cardiomyocyte related gene expression. The second data set (Electrophoresis 050115, FIG. 11) shows gene expression for each primer following 3 uM treatment of either BrdU or 5azaC. The high and low RNA is because we had low cell numbers and tested one well (low RNA) against combining two wells of equal treatment (high RNA). The third attachment (Electrophoresis 050116, FIG. 12) shows the effects of three weeks treatment of combined (3 uM or 1 uM of both 5azaC and BrdU), 3 uM of either 5azaC or BrdU, or nanog transfected cells (marked "control"). The poor quality is the result of low cell numbers and the use of a different RT-PCR kit (BioRad instead of the usual Invitrogen). See also FIG. 13. FIGS. 14-16 pertain to photograph images of MeSCs treated with 3 uM of 5azaC (A 050123 3 uM 5azaC 3weeks MSC.jpg, FIG. 14), of 3 uM of BrdU (A 050123 3 uM BrdU 3 weeks MSC.jpg, FIG. 15) and of nanog transfected cells combined with rat cardiac cells in co-culture (A 41227 of ntMSC 1213 2.jpg, FIG. 16). The cell differentiation is due to environmental signals and cell to cell contacts.

The inventors demonstrate that treatment with nucleotide derivatives and/or nanog transfection provides for cardiac differentiation of mesenchymal stem cells. Accordingly, an embodiment of the invention pertains to a method of increasing the potency of a cell comprising introducing a gene comprising nanog activity, optionally in conjunction with treatment of such cell with a compound, such as a nucleotide derivative, known to exert a dedifferentiating influence on cells.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 9

<210> SEQ ID NO 1

<211> LENGTH: 1628

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: CDS

<222> LOCATION: (28) .. (945)

<400> SEQUENCE: 1

```

tttttcctcc tcttctctta tactaac atg agt gtg gat cca gct tgt ccc caa      54
                               Met Ser Val Asp Pro Ala Cys Pro Gln
                               1                               5

agc ttg cct tgc ttt gaa gaa tcc gac tgt aaa gaa tct tca cct atg      102
Ser Leu Pro Cys Phe Glu Glu Ser Asp Cys Lys Glu Ser Ser Pro Met
10                               15                               20                               25

cct gtg att tgt ggg cct gaa gaa aac tat cca tcc ttg caa atg tct      150
Pro Val Ile Cys Gly Pro Glu Glu Asn Tyr Pro Ser Leu Gln Met Ser
30                               35                               40

tct gct gag atg cct cac aca gag act gtc tct cct ctt cct tcc tcc      198
Ser Ala Glu Met Pro His Thr Glu Thr Val Ser Pro Leu Pro Ser Ser
45                               50                               55

atg gat ctg ctt att cag gac agc cct gat tct tcc acc agt ccc aaa      246
Met Asp Leu Leu Ile Gln Asp Ser Pro Asp Ser Ser Thr Ser Pro Lys
60                               65                               70

ggc aaa caa ccc act tct gca gag aat agt gtc gca aaa aag gaa gac      294
Gly Lys Gln Pro Thr Ser Ala Glu Asn Ser Val Ala Lys Lys Glu Asp
75                               80                               85

aag gtc ccg gtc aag aaa cag aag acc aga act gtg ttc tct tcc acc      342
Lys Val Pro Val Lys Lys Gln Lys Thr Arg Thr Val Phe Ser Ser Thr
90                               95                               100                               105

```

-continued

cag ctg tgt gta ctc aat gat aga ttt cag aga cag aaa tac ctc agc Gln Leu Cys Val Leu Asn Asp Arg Phe Gln Arg Gln Lys Tyr Leu Ser 110 115 120	390
ctc cag cag atg caa gaa ctc tcc aac atc ctg aac ctc agc tac aaa Leu Gln Gln Met Gln Glu Leu Ser Asn Ile Leu Asn Leu Ser Tyr Lys 125 130 135	438
cag gtg aag acc tgg ttc cag aac cag aga atg aaa tct aag agg tgg Gln Val Lys Thr Trp Phe Gln Asn Gln Arg Met Lys Ser Lys Arg Trp 140 145 150	486
cag aaa aac aac tgg ccg aag aat agc aat ggt gtg acg cag aag gcc Gln Lys Asn Asn Trp Pro Lys Asn Ser Asn Gly Val Thr Gln Lys Ala 155 160 165	534
tca gca cct acc tac ccc agc ctc tac tct tcc tac cac cag gga tgc Ser Ala Pro Thr Tyr Pro Ser Leu Tyr Ser Ser Tyr His Gln Gly Cys 170 175 180 185	582
ctg gtg aac ccg act ggg aac ctt cca atg tgg agc aac cag acc tgg Leu Val Asn Pro Thr Gly Asn Leu Pro Met Trp Ser Asn Gln Thr Trp 190 195 200	630
aac aat tca acc tgg agc aac cag acc cag aac atc cag tcc tgg agc Asn Asn Ser Thr Trp Ser Asn Gln Thr Gln Asn Ile Gln Ser Trp Ser 205 210 215	678
aac cac tcc tgg aac act cag acc tgg tgc acc caa tcc tgg aac aat Asn His Ser Trp Asn Thr Gln Thr Trp Cys Thr Gln Ser Trp Asn Asn 220 225 230	726
cag gcc tgg aac agt ccc ttc tat aac tgt gga gag gaa tct ctg cag Gln Ala Trp Asn Ser Pro Phe Tyr Asn Cys Gly Glu Glu Ser Leu Gln 235 240 245	774
tcc tgc atg cac ttc cag cca aat tct cct gcc agt gac ttg gag gct Ser Cys Met His Phe Gln Pro Asn Ser Pro Ala Ser Asp Leu Glu Ala 250 255 260 265	822
gcc ttg gaa gct gct ggg gaa ggc ctt aat gta ata cag cag acc act Ala Leu Glu Ala Ala Gly Glu Gly Leu Asn Val Ile Gln Gln Thr Thr 270 275 280	870
agg tat ttt agt act cca caa acc atg gat tta ttc cta aac tac tcc Arg Tyr Phe Ser Thr Pro Gln Thr Met Asp Leu Phe Leu Asn Tyr Ser 285 290 295	918
atg aac atg caa cct gaa gac gtg tga agatgagtga aactgatatt Met Asn Met Gln Pro Glu Asp Val 300 305	965
actcaatttc agtctggaca ctggctgaat ccttcctctc ccctcctccc atccctcata	1025
ggatttttct tgtttgaaa ccacgtgttc tggtttccat gatgcctate cagtcaatct	1085
catggagggt ggagtatggt tggagcctaa tcagcgaggt ttcttttttt ttttttecta	1145
ttggatcttc ctggagaaaa tacttttttt tttttttttg agacggagtc ttgctctgtc	1205
gcccaggctg gactgcagtg gcgcgggtctt ggctcactgc aagctccgcc tcccgggttc	1265
acgccattct cctgcctcag cctcccagc agctgggact acaggcgccc gccacctcgc	1325
ccggctaata ttttgatatt tttagtagaga cagggtttca ctgtgttagc caggatggtc	1385
tcgatctcct gaccttgta tccgccgcc tcggcctccc taacagctgg gattacaggc	1445
gtgagccacc gcgccctgcc tagaaaagac attttaataa ccttggtctgc taaggacaac	1505
attgatagaa gccgtctctg gctatagata agtagatcta atactagttt ggatatcttt	1565
agggtttaga atctaacctc aagaataaga aatacaagta cgaattgggtg atgaagatgt	1625
att	1628

<210> SEQ ID NO 2

<211> LENGTH: 305

<212> TYPE: PRT

-continued

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 2

```

Met Ser Val Asp Pro Ala Cys Pro Gln Ser Leu Pro Cys Phe Glu Glu
 1           5           10           15
Ser Asp Cys Lys Glu Ser Ser Pro Met Pro Val Ile Cys Gly Pro Glu
 20           25           30
Glu Asn Tyr Pro Ser Leu Gln Met Ser Ser Ala Glu Met Pro His Thr
 35           40           45
Glu Thr Val Ser Pro Leu Pro Ser Ser Met Asp Leu Leu Ile Gln Asp
 50           55           60
Ser Pro Asp Ser Ser Thr Ser Pro Lys Gly Lys Gln Pro Thr Ser Ala
 65           70           75           80
Glu Asn Ser Val Ala Lys Lys Glu Asp Lys Val Pro Val Lys Lys Gln
 85           90           95
Lys Thr Arg Thr Val Phe Ser Ser Thr Gln Leu Cys Val Leu Asn Asp
 100          105          110
Arg Phe Gln Arg Gln Lys Tyr Leu Ser Leu Gln Gln Met Gln Glu Leu
 115          120          125
Ser Asn Ile Leu Asn Leu Ser Tyr Lys Gln Val Lys Thr Trp Phe Gln
 130          135          140
Asn Gln Arg Met Lys Ser Lys Arg Trp Gln Lys Asn Asn Trp Pro Lys
 145          150          155          160
Asn Ser Asn Gly Val Thr Gln Lys Ala Ser Ala Pro Thr Tyr Pro Ser
 165          170          175
Leu Tyr Ser Ser Tyr His Gln Gly Cys Leu Val Asn Pro Thr Gly Asn
 180          185          190
Leu Pro Met Trp Ser Asn Gln Thr Trp Asn Asn Ser Thr Trp Ser Asn
 195          200          205
Gln Thr Gln Asn Ile Gln Ser Trp Ser Asn His Ser Trp Asn Thr Gln
 210          215          220
Thr Trp Cys Thr Gln Ser Trp Asn Asn Gln Ala Trp Asn Ser Pro Phe
 225          230          235          240
Tyr Asn Cys Gly Glu Glu Ser Leu Gln Ser Cys Met His Phe Gln Pro
 245          250          255
Asn Ser Pro Ala Ser Asp Leu Glu Ala Ala Leu Glu Ala Ala Gly Glu
 260          265          270
Gly Leu Asn Val Ile Gln Gln Thr Thr Arg Tyr Phe Ser Thr Pro Gln
 275          280          285
Thr Met Asp Leu Phe Leu Asn Tyr Ser Met Asn Met Gln Pro Glu Asp
 290          295          300

```

Val
305

<210> SEQ ID NO 3

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 3

tttttctctcc tcttctctcta

20

<210> SEQ ID NO 4

<211> LENGTH: 20

-continued

```

<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        primer

<400> SEQUENCE: 4

attggtgatg aagatgtatt                                20

<210> SEQ ID NO 5
<211> LENGTH: 6
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        6 His Tag

<400> SEQUENCE: 5

His His His His His His
  1             5

<210> SEQ ID NO 6
<211> LENGTH: 1417
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 4

tcccttcgca agccctcatt tcaccaggcc cccggcttgg ggcgccttcc ttecccatgg    60
cgggacacct ggcttcggat ttgccttct cgcctctcc aggtggtgga ggtgatgggc    120
cagggggggc ggagccgggc tgggttgatc ctcgacctg gctaagcttc caaggccctc    180
ctggagggcc agaatcggg ccggggggtg gcccaggctc tgaggtgtgg gggattcccc    240
catgcccccc gccgtatgag ttctgtgggg ggatggcgta ctgtgggccc caggttgag    300
tggggctagt gcccacaagg ggcttgaga cctctcagcc tgagggcgaa gcaggagtcg    360
gggtggagag caactccgat ggggcctccc cggagccctg caccgtcacc cctggtgccg    420
tgaagctgga gaaggagaag ctggagcaaa acccggagga gtcccaggac atcaaagctc    480
tgcagaaaga actcgagcaa ttgccaagc tctgaagca gaagaggatc accctgggat    540
atacacaggg cgatgtgggg ctaccctgg gggttctatt tgggaaggta ttcagccaaa    600
cgaccatctg ccgctttgag gctctgcagc ttagcttcaa gaacatgtgt aagctcggc    660
ccttgctgca gaagtgggtg gaggaagctg acaacaatga aaatcttcag gagatatgca    720
aagcagaaac cctcgtgcag gcccgaaaga gaaagcgaac cagtatcgag aaccgagtga    780
gaggcaacct ggagaatttg ttctctcagt gcccgaaacc cactctcag cagatcagcc    840
acatcgccca gcagcttggg ctcgagaagg atgtggtccg agtgtggttc tgtaaccggc    900
gccagaaggg caagcgatca agcagcgact atgcacaacg agaggathtt gaggctgctg    960
ggtctccttt ctccagggga ccagtgtcct ttctctcggc ccaggggccc cattttggtg   1020
ccccaggcta tgggagccct cacttcaactg cactgtactc ctgggtccct ttcctgagg   1080
gggaagcctt tccccctgtc tccgtcacca ctctgggctc tcccatgcat tcaaactgag   1140
gtgcctgccc ttctaggaat gggggacagg gggaggggag gagctaggga aagaaaacct   1200
ggagtttgtg ccagggtttt tgggattaag ttcttcattc actaagggaag gaattgggaa   1260
caciaaaggt gggggcaggg gagtttgggg caactggttg gagggaaggt gaagttcaat   1320
gatgctcttg attttaatcc cacatcatgt atcacttttt tcttaataaa agaagcctgg   1380
gacacagtaa aaaaaaaaaa aaaaaaaaaa aaaaaaa   1417

```

-continued

```

<210> SEQ ID NO 7
<211> LENGTH: 360
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 7
Met Ala Gly His Leu Ala Ser Asp Phe Ala Phe Ser Pro Pro Pro Gly
 1          5          10          15
Gly Gly Gly Asp Gly Pro Gly Gly Pro Glu Pro Gly Trp Val Asp Pro
 20          25          30
Arg Thr Trp Leu Ser Phe Gln Gly Pro Pro Gly Gly Pro Gly Ile Gly
 35          40          45
Pro Gly Val Gly Pro Gly Ser Glu Val Trp Gly Ile Pro Pro Cys Pro
 50          55          60
Pro Pro Tyr Glu Phe Cys Gly Gly Met Ala Tyr Cys Gly Pro Gln Val
 65          70          75          80
Gly Val Gly Leu Val Pro Gln Gly Gly Leu Glu Thr Ser Gln Pro Glu
 85          90          95
Gly Glu Ala Gly Val Gly Val Glu Ser Asn Ser Asp Gly Ala Ser Pro
100          105          110
Glu Pro Cys Thr Val Thr Pro Gly Ala Val Lys Leu Glu Lys Glu Lys
115          120          125
Leu Glu Gln Asn Pro Glu Glu Ser Gln Asp Ile Lys Ala Leu Gln Lys
130          135          140
Glu Leu Glu Gln Phe Ala Lys Leu Leu Lys Gln Lys Arg Ile Thr Leu
145          150          155          160
Gly Tyr Thr Gln Ala Asp Val Gly Leu Thr Leu Gly Val Leu Phe Gly
165          170          175
Lys Val Phe Ser Gln Thr Thr Ile Cys Arg Phe Glu Ala Leu Gln Leu
180          185          190
Ser Phe Lys Asn Met Cys Lys Leu Arg Pro Leu Leu Gln Lys Trp Val
195          200          205
Glu Glu Ala Asp Asn Asn Glu Asn Leu Gln Glu Ile Cys Lys Ala Glu
210          215          220
Thr Leu Val Gln Ala Arg Lys Arg Lys Arg Thr Ser Ile Glu Asn Arg
225          230          235          240
Val Arg Gly Asn Leu Glu Asn Leu Phe Leu Gln Cys Pro Lys Pro Thr
245          250          255
Leu Gln Gln Ile Ser His Ile Ala Gln Gln Leu Gly Leu Glu Lys Asp
260          265          270
Val Val Arg Val Trp Phe Cys Asn Arg Arg Gln Lys Gly Lys Arg Ser
275          280          285
Ser Ser Asp Tyr Ala Gln Arg Glu Asp Phe Glu Ala Ala Gly Ser Pro
290          295          300
Phe Ser Gly Gly Pro Val Ser Phe Pro Leu Ala Pro Gly Pro His Phe
305          310          315          320
Gly Thr Pro Gly Tyr Gly Ser Pro His Phe Thr Ala Leu Tyr Ser Ser
325          330          335
Val Pro Phe Pro Glu Gly Glu Ala Phe Pro Pro Val Ser Val Thr Thr
340          345          350
Leu Gly Ser Pro Met His Ser Asn
355          360

```

```

<210> SEQ ID NO 8
<211> LENGTH: 2518

```

-continued

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 8

```

ctattaactt gttcaaaaa gtatcaggag ttgtcaaggc agagaagaga gtgtttgcaa      60
aaggggaaaa gtagtttctt gcctctttaa gactaggact gagagaaaga agaggagaga      120
gaaagaaaagg gagagaagtt tgagccccag gcttaagcct ttccaaaaaa taataataac      180
aatcatcggc ggcggcagga tcggccagag gaggaggaa gcgctttttt tgatcctgat      240
tccagtttgc ctctctcttt ttttcccca aattattctt cgcctgattt tcctcggga      300
gccctgcgct cccgacaccc ccgcccgcct cccctcctcc tctcccccg cccgcgggcc      360
ccccaaagtc ccggccgggc cgagggtcgg cggccgcccg cgggcccggc ccgcgcacag      420
cgcccgatg tacaacatga tggagacgga gctgaagccg ccgggcccgc agcaaaactc      480
ggggggcggc ggcggcaact ccaccgccc ggccgcccgc ggcaaccaga aaaacagccc      540
ggaccgctc aagcggccca tgaatgcctt catggtgtgg tcccggggc agcggcgcaa      600
gatgcccag gagaaccoca agatgcacaa ctccgagatc agcaagcgc tgggcgccga      660
gtgaaaactt ttgtcggaga cggagaagcg gccgttcctc gacgaggcta agcggctgcg      720
agcgtgcac atgaaggagc acccggatta taaataccgg ccccgcgga aaaccaagac      780
gctcatgaag aaggataagt acacgctgcc cggcgggctg ctggccccg gcggcaatag      840
catggcgagc ggggtcgggg tgggcccgg cctgggcccg ggctgaacc agcgcagga      900
cagttacgcy cacatgaacg gctggagcaa cggcagctac agcatgatgc aggaccagct      960
gggctaccog cagcaccocg gcctcaatgc gcacggcgca gcgcagatgc agcccatgca      1020
ccgctacgac gtgagcggcc tgcaagtaca ctccatgacc agctcgcaga cctacatgaa      1080
cggctcggcc acctacagca tgtcctactc gcagcagggc acccctggca tggctcttgg      1140
ctccatgggt tcggtgtgca agtccgaggc cagctccagc cccctgtgg ttacctctc      1200
ctccactcc agggcgcctt gccaggcccg ggacctccgg gacatgatca gcatgtatct      1260
ccccggccc gaggtgcccg aaccgcccgc ccccagcaga cttcacatgt cccagcacta      1320
ccagagcggc ccggtgcccg gcacggccat taacggcaca ctgcccctct cacacatgtg      1380
agggccggac agcgaactgg aggggggaga aatthtcaa gaaaaacgag ggaatggga      1440
ggggtgcaaa agaggagagt aagaaacagc atggagaaaa cccggtacgc tcaaaaagaa      1500
aaaggaaaa aaaaaatccc atcaccoca gcaaatgaca gctgcaaaag agaacacca      1560
tcccatccac actcacgcaa aaaccgcat gccgacaaga aaacttttat gagagagatc      1620
ctggacttct ttttggggga ctatthtgt acagagaaaa cctggggagg gtggggagg      1680
cgggggaatg gacctgtat agatctggag gaaagaaagc tacgaaaaac tttttaaag      1740
ttctagtggc acggtaggag ctttgagga agtttgcaa agtctttacc aataatatt      1800
agagctagtc tccaagcgc gaaaaaatg ttttaattt tgcaagcaac tttgtacag      1860
tatttatcga gataaacatg gcaatcaaaa tgtccattgt ttataagctg agaatttgc      1920
aatatthtcc aaggagagc tcttgctga atthtgatc tgacagtgaa atthaggaca      1980
gttgcaaacg tgaagaagaa aaaattatc aaattggac atthtaattg tttaaaaatt      2040
gtacaaaagg aaaaatttag aataagtact ggcaaccat ctctgtggtc ttgttataaa      2100
agggcaaaag ttttagactg tactaaattt tataacttac tgtaaaagc aaaaatggcc      2160
atgcaggttg acaccgttg taattataa tagctthtgt tcgatccca ctttccattt      2220
tgttcagata aaaaaacca tgaattact gtgttgaaa tathttctta tggtttgtaa      2280

```


-continued

```
tatttctgta aatttattgt gatattttaa ggttttcccc cttttatfff ccgtagttgt 2340
atftttaaag attcggctct gtattatttg aatcagctctg ccgagaatcc atgtatatat 2400
ttgaactaat atcatcctta taacaggtac atfttcaact taagttttta ctccattatg 2460
cacagtttga gataaataaa tttttgaaat atggacactg aaaaaaaaaa aaaaaaaaaa 2518
```

```
<210> SEQ ID NO 9
<211> LENGTH: 317
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
```

```
<400> SEQUENCE: 9
```

```
Met Tyr Asn Met Met Glu Thr Glu Leu Lys Pro Pro Gly Pro Gln Gln
 1          5          10          15
Thr Ser Gly Gly Gly Gly Asn Ser Thr Ala Ala Ala Ala Gly Gly
 20          25          30
Asn Gln Lys Asn Ser Pro Asp Arg Val Lys Arg Pro Met Asn Ala Phe
 35          40          45
Met Val Trp Ser Arg Gly Gln Arg Arg Lys Met Ala Gln Glu Asn Pro
 50          55          60
Lys Met His Asn Ser Glu Ile Ser Lys Arg Leu Gly Ala Glu Trp Lys
 65          70          75          80
Leu Leu Ser Glu Thr Glu Lys Arg Pro Phe Ile Asp Glu Ala Lys Arg
 85          90          95
Leu Arg Ala Leu His Met Lys Glu His Pro Asp Tyr Lys Tyr Arg Pro
100          105          110
Arg Arg Lys Thr Lys Thr Leu Met Lys Lys Asp Lys Tyr Thr Leu Pro
115          120          125
Gly Gly Leu Leu Ala Pro Gly Gly Asn Ser Met Ala Ser Gly Val Gly
130          135          140
Val Gly Ala Gly Leu Gly Ala Gly Val Asn Gln Arg Met Asp Ser Tyr
145          150          155          160
Ala His Met Asn Gly Trp Ser Asn Gly Ser Tyr Ser Met Met Gln Asp
165          170          175
Gln Leu Gly Tyr Pro Gln His Pro Gly Leu Asn Ala His Gly Ala Ala
180          185          190
Gln Met Gln Pro Met His Arg Tyr Asp Val Ser Ala Leu Gln Tyr Asn
195          200          205
Ser Met Thr Ser Ser Gln Thr Tyr Met Asn Gly Ser Pro Thr Tyr Ser
210          215          220
Met Ser Tyr Ser Gln Gln Gly Thr Pro Gly Met Ala Leu Gly Ser Met
225          230          235          240
Gly Ser Val Val Lys Ser Glu Ala Ser Ser Ser Pro Pro Val Val Thr
245          250          255
Ser Ser Ser His Ser Arg Ala Pro Cys Gln Ala Gly Asp Leu Arg Asp
260          265          270
Met Ile Ser Met Tyr Leu Pro Gly Ala Glu Val Pro Glu Pro Ala Ala
275          280          285
Pro Ser Arg Leu His Met Ser Gln His Tyr Gln Ser Gly Pro Val Pro
290          295          300
Gly Thr Ala Ile Asn Gly Thr Leu Pro Leu Ser His Met
305          310          315
```

29

What is claimed is:

1. A mesenchymal stem cell possessing increased developmental potency, said mesenchymal stem cell produced by introducing into said mesenchymal stem cell a nanog gene under conditions to allow expression of said nanog gene,

30

wherein said expression results in higher developmental potency of said mesenchymal stem cell.

* * * * *