

1-1-2010

The weak Stratonovich integral with respect to fractional Brownian motion with Hurst parameter $1/6$

Ivan Nourdin

Anthony Réveillac

Jason Swanson

University of Central Florida

Find similar works at: <https://stars.library.ucf.edu/facultybib2010>


University of Central Florida Libraries <http://library.ucf.edu>

This Article is brought to you for free and open access by the Faculty Bibliography at STARS. It has been accepted for inclusion in Faculty Bibliography 2010s by an authorized administrator of STARS. For more information, please contact STARS@ucf.edu.

Recommended Citation

Nourdin, Ivan; Réveillac, Anthony; and Swanson, Jason, "The weak Stratonovich integral with respect to fractional Brownian motion with Hurst parameter $1/6$ " (2010). *Faculty Bibliography 2010s*. 598.

<https://stars.library.ucf.edu/facultybib2010/598>



Vol. 15 (2010), Paper no. 70, pages 2117–.

Journal URL

<http://www.math.washington.edu/~ejpecp/>

The weak Stratonovich integral with respect to fractional Brownian motion with Hurst parameter $1/6$

Ivan Nourdin*

Université Nancy 1 - Institut de Mathématiques Élie Cartan
B.P. 239, 54506 Vandœuvre-lès-Nancy Cedex, France
nourdin@iecn.u-nancy.fr

Anthony Réveillac[†]

Institut für Mathematik
Humboldt-Universität zu Berlin
Unter des Linden 6, 10099 Berlin, Germany
areveill@mathematik.hu-berlin.de

Jason Swanson[‡]

University of Central Florida
Department of Mathematics
4000 Central Florida Blvd, P.O. Box 161364
Orlando, FL 32816-1364
swanson@mail.ucf.edu

Abstract

Let B be a fractional Brownian motion with Hurst parameter $H = 1/6$. It is known that the symmetric Stratonovich-style Riemann sums for $\int g(B(s)) dB(s)$ do not, in general, converge in probability. We show, however, that they do converge in law in the Skorohod space of càdlàg functions. Moreover, we show that the resulting stochastic integral satisfies a change of variable formula with a correction term that is an ordinary Itô integral with respect to a Brownian motion that is independent of B .

Key words: Stochastic integration; Stratonovich integral; fractional Brownian motion; weak convergence; Malliavin calculus.

AMS 2000 Subject Classification: Primary 60H05; Secondary: 60G15, 60G18, 60J05.

Submitted to EJP on July 6, 2010, final version accepted December 6, 2010.

*Supported in part by the (french) ANR grant 'Exploration des Chemins Rugueux'.

[†]Supported by DFG research center Matheon project E2.

[‡]Supported in part by NSA grant H98230-09-1-0079.

1 Introduction

The Stratonovich integral of X with respect to Y , denoted $\int_0^t X(s) \circ dY(s)$, can be defined as the limit in probability, if it exists, of

$$\sum_{t_j \leq t} \frac{X(t_{j-1}) + X(t_j)}{2} (Y(t_j) - Y(t_{j-1})), \quad (1.1)$$

as the mesh of the partition $\{t_j\}$ goes to zero. Typically, we regard (1.1) as a process in t , and require that it converges uniformly on compacts in probability (ucp).

This is closely related to the so-called symmetric integral, denoted by $\int_0^t X(s) d^\circ Y(s)$, which is the ucp limit, if it exists, of

$$\frac{1}{\varepsilon} \int_0^t \frac{X(s) + X(s + \varepsilon)}{2} (Y(s + \varepsilon) - Y(s)) ds, \quad (1.2)$$

as $\varepsilon \rightarrow 0$. The symmetric integral is an example of the regularization procedure, introduced by Russo and Vallois, and on which there is a wide body of literature. For further details on stochastic calculus via regularization, see the excellent survey article [13] and the many references therein.

A special case of interest that has received considerable attention in the literature is when $Y = B^H$, a fractional Brownian motion with Hurst parameter H . It has been shown independently in [2] and [5] that when $Y = B^H$ and $X = g(B^H)$ for a sufficiently differentiable function $g(x)$, the symmetric integral exists for all $H > 1/6$. Moreover, in this case, the symmetric integral satisfies the classical Stratonovich change of variable formula,

$$g(B^H(t)) = g(B^H(0)) + \int_0^t g'(B^H(s)) d^\circ B^H(s).$$

However, when $H = 1/6$, the symmetric integral does not, in general, exist. Specifically, in [2] and [5], it is shown that (1.2) does not converge in probability when $Y = B^{1/6}$ and $X = (B^{1/6})^2$. It can be similarly shown that, in this case, (1.1) also fails to converge in probability.

This brings us naturally to the notion which is the focus of this paper: the weak Stratonovich integral, which is the limit in law, if it exists, of (1.1). We focus exclusively on the case $Y = B^{1/6}$. For simplicity, we omit the superscript and write $B = B^{1/6}$. Our integrands shall take the form $g(B(t))$, for $g \in C^\infty(\mathbb{R})$, and we shall work only with the uniformly spaced partition, $t_j = j/n$. In this case, (1.1) becomes

$$I_n(g, B, t) = \sum_{j=1}^{\lfloor nt \rfloor} \frac{g(B(t_{j-1})) + g(B(t_j))}{2} \Delta B_j,$$

where $\lfloor x \rfloor$ denotes the greatest integer less than or equal to x , and $\Delta B_j = B(t_j) - B(t_{j-1})$. We show that the processes $I_n(g, B)$ converge in law in $D_{\mathbb{R}}[0, \infty)$, the Skorohod space of càdlàg functions from $[0, \infty)$ to \mathbb{R} . We let $\int_0^t g(B(s)) dB(s)$ denote a process with this limiting law, and refer to this as the weak Stratonovich integral.

The weak Stratonovich integral with respect to B does not satisfy the classical Stratonovich change of variable formula. Rather, we show that it satisfies a change of variable formula with a correction term that is a classical Itô integral. Namely,

$$g(B(t)) = g(B(0)) + \int_0^t g'(B(s)) dB(s) - \frac{1}{12} \int_0^t g'''(B(s)) d[[B]]_s, \quad (1.3)$$

where $[[B]]$ is what we call the signed cubic variation of B . That is, $[[B]]$ is the limit in law of the sequence of processes $V_n(B, t) = \sum_{j=1}^{\lfloor nt \rfloor} \Delta B_j^3$. It is shown in [11] that $[[B]] = \kappa W$, where W is a standard Brownian motion, independent of B , and $\kappa \simeq 2.322$. (See (2.5) for the exact definition of κ .) The correction term in (1.3) is then a standard Itô integral with respect to Brownian motion.

Our precise results are actually somewhat stronger than this, in that we prove the joint convergence of the processes B , $V_n(B)$, and $I_n(g, B)$. (See Theorem 2.13.) We also discuss the joint convergence of multiple sequences of Riemann sums for different integrands. (See Theorem 2.14 and Remark 2.15.)

The work in this paper is a natural follow-up to [1] and [9]. There, analogous results were proven for $B^{1/4}$ in the context of midpoint-style Riemann sums. The results in [1] and [9] were proven through different methods, and in the present work, we combine the two approaches to prove our main results.

Finally, let us stress the fact that, as a byproduct of the proof of (1.3), we show in the present paper that

$$n^{-1/2} \sum_{j=1}^{\lfloor n \cdot \rfloor} g(B(t_{j-1})) h_3(n^{1/6} \Delta B_j) \rightarrow -\frac{1}{8} \int_0^\cdot g'''(B(s)) ds + \int_0^\cdot g(B(s)) d[[B]]_s,$$

in the sense of finite-dimensional distributions on $[0, \infty)$, where $h_3(x) = x^3 - 3x$ denotes the third Hermite polynomial. (See more precisely Theorem 3.7 below. Also see Theorem 3.8.) From our point of view, this result has also its own interest, and should be compared with the recent results obtained in [7, 8], concerning the weighted Hermite variations of fractional Brownian motion.

2 Notation, preliminaries, and main result

Let $B = B^{1/6}$ be a fractional Brownian motion with Hurst parameter $H = 1/6$. That is, B is a centered Gaussian process, indexed by $t \geq 0$, such that

$$R(s, t) = E[B(s)B(t)] = \frac{1}{2}(t^{1/3} + s^{1/3} - |t - s|^{1/3}).$$

Note that $E|B(t) - B(s)|^2 = |t - s|^{1/3}$. For compactness of notation, we will sometimes write B_t instead of $B(t)$. Given a positive integer n , let $\Delta t = n^{-1}$ and $t_j = t_{j,n} = j\Delta t$. We shall frequently have occasion to deal with the quantity $\beta_{j,n} = \beta_j = (B(t_{j-1}) + B(t_j))/2$. In estimating this and similar quantities, we shall adopt the notation $r_+ = r \vee 1$, which is typically applied to nonnegative integers r . We shall also make use of the Hermite polynomials,

$$h_n(x) = (-1)^n e^{x^2/2} \frac{d^n}{dx^n} (e^{-x^2/2}). \tag{2.1}$$

Note that the first few Hermite polynomials are $h_0(x) = 1$, $h_1(x) = x$, $h_2(x) = x^2 - 1$, and $h_3(x) = x^3 - 3x$. The following orthogonality property is well-known: if U and V are jointly normal with $E(U) = E(V) = 0$ and $E(U^2) = E(V^2) = 1$, then

$$E[h_p(U)h_q(V)] = \begin{cases} q!(E[UV])^q & \text{if } p = q, \\ 0 & \text{otherwise.} \end{cases} \tag{2.2}$$

If X is a càdlàg process, we write $X(t-) = \lim_{s \uparrow t} X(s)$ and $\Delta X(t) = X(t) - X(t-)$. The step function approximation to X will be denoted by $X_n(t) = X(\lfloor nt \rfloor/n)$, where $\lfloor \cdot \rfloor$ is the greatest integer function. In this case, $\Delta X_n(t_{j,n}) = X(t_j) - X(t_{j-1})$. We shall frequently use the shorthand notation $\Delta X_j = \Delta X_{j,n} = \Delta X_n(t_{j,n})$. For simplicity, positive integer powers of ΔX_j shall be written without parentheses, so that $\Delta X_j^k = (\Delta X_j)^k$.

The discrete p -th variation of X is defined as

$$V_n^p(X, t) = \sum_{j=1}^{\lfloor nt \rfloor} |\Delta X_j|^p,$$

and the discrete signed p -th variation of X is

$$V_n^{p\pm}(X, t) = \sum_{j=1}^{\lfloor nt \rfloor} |\Delta X_j|^p \operatorname{sgn}(\Delta X_j).$$

For the discrete signed cubic variation, we shall omit the superscript, so that

$$V_n(X, t) = V_n^{3\pm}(X, t) = \sum_{j=1}^{\lfloor nt \rfloor} \Delta X_j^3. \quad (2.3)$$

When we omit the index t , we mean to refer to the entire process. So, for example, $V_n(X) = V_n(X, \cdot)$ refers to the càdlàg process which maps $t \mapsto V_n(X, t)$. We recall the following fact which will be extensively used in this paper.

Remark 2.1. Let $\{\rho(r)\}_{r \in \mathbb{Z}}$ be the sequence defined by

$$\rho(r) = \frac{1}{2}(|r+1|^{1/3} + |r-1|^{1/3} - 2|r|^{1/3}). \quad (2.4)$$

It holds that $\sum_{r \in \mathbb{Z}} |\rho(r)| < \infty$ and $E[\Delta B_i \Delta B_j] = n^{-1/3} \rho(i-j)$ for all $i, j \in \mathbb{N}$.

Let $\kappa > 0$ be defined by

$$\kappa^2 = 6 \sum_{r \in \mathbb{Z}} \rho^3(r) = \frac{3}{4} \sum_{r \in \mathbb{Z}} (|r+1|^{1/3} + |r-1|^{1/3} - 2|r|^{1/3})^3 \simeq 5.391, \quad (2.5)$$

and let W be a standard Brownian motion, defined on the same probability space as B , and independent of B . Define $[[B]]_t = \kappa W(t)$. We shall refer to the process $[[B]]$ as the signed cubic variation of B . The use of this term is justified by Theorem 2.12.

A function $g : \mathbb{R}^d \rightarrow \mathbb{R}$ has *polynomial growth* if there exist positive constants K and r such that $|g(x)| \leq K(1 + |x|^r)$ for all $x \in \mathbb{R}^d$. If k is a nonnegative integer, we shall say that a function g has *polynomial growth of order k* if $g \in C^k(\mathbb{R}^d)$ and there exist positive constants K and r such that $|\partial^\alpha g(x)| \leq K(1 + |x|^r)$ for all $x \in \mathbb{R}^d$ and all $|\alpha| \leq k$. (Here, $\alpha \in \mathbb{N}_0^d = (\mathbb{N} \cup \{0\})^d$ is a multi-index, and we adopt the standard multi-index notation: $\partial_j = \partial / \partial x_j$, $\partial^\alpha = \partial_1^{\alpha_1} \cdots \partial_d^{\alpha_d}$, and $|\alpha| = \alpha_1 + \cdots + \alpha_d$.)

Given $g : \mathbb{R} \rightarrow \mathbb{R}$ and a stochastic process $\{X(t) : t \geq 0\}$, the Stratonovich Riemann sum will be denoted by

$$I_n(g, X, t) = \sum_{j=1}^{\lfloor nt \rfloor} \frac{g(X(t_{j-1})) + g(X(t_j))}{2} \Delta X_j.$$

The phrase “uniformly on compacts in probability” will be abbreviated “ucp.” If X_n and Y_n are càdlàg processes, we shall write $X_n \approx Y_n$ or $X_n(t) \approx Y_n(t)$ to mean that $X_n - Y_n \rightarrow 0$ ucp. In the proofs in this paper, C shall denote a positive, finite constant that may change value from line to line.

2.1 Conditions for relative compactness

The Skorohod space of càdlàg functions from $[0, \infty)$ to \mathbb{R}^d is denoted by $D_{\mathbb{R}^d}[0, \infty)$. Note that $D_{\mathbb{R}^d}[0, \infty)$ and $(D_{\mathbb{R}}[0, \infty))^d$ are not the same. In particular, the map $(x, y) \mapsto x + y$ is continuous from $D_{\mathbb{R}^2}[0, \infty)$ to $D_{\mathbb{R}}[0, \infty)$, but it is not continuous from $(D_{\mathbb{R}}[0, \infty))^2$ to $D_{\mathbb{R}}[0, \infty)$. Convergence in $D_{\mathbb{R}^d}[0, \infty)$ implies convergence in $(D_{\mathbb{R}}[0, \infty))^d$, but the converse is not true.

Note that if the sequences $\{X_n^{(1)}\}, \dots, \{X_n^{(d)}\}$ are all relatively compact in $D_{\mathbb{R}}[0, \infty)$, then the sequence of d -tuples $\{(X_n^{(1)}, \dots, X_n^{(d)})\}$ is relatively compact in $(D_{\mathbb{R}}[0, \infty))^d$. It may not, however, be relatively compact in $D_{\mathbb{R}^d}[0, \infty)$. We will therefore need the following well-known result. (For more details, see Section 2.1 of [1] and the references therein.)

Lemma 2.2. *Suppose $\{(X_n^{(1)}, \dots, X_n^{(d)})\}_{n=1}^{\infty}$ is relatively compact in $(D_{\mathbb{R}}[0, \infty))^d$. If, for each $j \geq 2$, the sequence $\{X_n^{(j)}\}_{n=1}^{\infty}$ converges in law in $D_{\mathbb{R}}[0, \infty)$ to a continuous process, then $\{(X_n^{(1)}, \dots, X_n^{(d)})\}_{n=1}^{\infty}$ is relatively compact in $D_{\mathbb{R}^d}[0, \infty)$.*

Proof. First note that if $x_n \rightarrow x$ in $D_{\mathbb{R}}[0, \infty)$, $y_n \rightarrow y$ in $D_{\mathbb{R}}[0, \infty)$, and x and y have no simultaneous discontinuities, then $x_n + y_n \rightarrow x + y$ in $D_{\mathbb{R}}[0, \infty)$. Thus, if two sequences $\{X_n\}$ and $\{Y_n\}$ are relatively compact in $D_{\mathbb{R}}[0, \infty)$ and every subsequential limit of $\{Y_n\}$ is continuous, then $\{X_n + Y_n\}$ is relatively compact in $D_{\mathbb{R}}[0, \infty)$. The lemma now follows from the fact that a sequence $\{X_n^{(1)}, \dots, X_n^{(d)}\}$ is relatively compact in $D_{\mathbb{R}^d}[0, \infty)$ if and only if $\{X_n^{(k)}\}$ and $\{X_n^{(k)} + X_n^{(\ell)}\}$ are relatively compact in $D_{\mathbb{R}}[0, \infty)$ for all k and ℓ . (See, for example, Problem 3.22(c) in [4].) \square

Our primary criterion for relative compactness is the following moment condition, which is a special case of Corollary 2.2 in [1].

Theorem 2.3. *Let $\{X_n\}$ be a sequence of processes in $D_{\mathbb{R}^d}[0, \infty)$. Let $q(x) = |x| \wedge 1$. Suppose that for each $T > 0$, there exists $\nu > 0$, $\beta > 0$, $C > 0$, and $\theta > 1$ such that $\sup_n E[|X_n(T)|^\nu] < \infty$ and*

$$E[q(X_n(t) - X_n(s))^\beta] \leq C \left(\frac{\lfloor nt \rfloor - \lfloor ns \rfloor}{n} \right)^\theta, \quad (2.6)$$

for all n and all $0 \leq s \leq t \leq T$. Then $\{X_n\}$ is relatively compact.

Of course, a sequence $\{X_n\}$ converges in law in $D_{\mathbb{R}^d}[0, \infty)$ to a process X if $\{X_n\}$ is relatively compact and $X_n \rightarrow X$ in the sense of finite-dimensional distributions on $[0, \infty)$. We shall also need the analogous theorem for convergence in probability, which is Lemma A2.1 in [3]. Note that if $x : [0, \infty) \rightarrow \mathbb{R}^d$ is continuous, then $x_n \rightarrow x$ in $D_{\mathbb{R}^d}[0, \infty)$ if and only if $x_n \rightarrow x$ uniformly on compacts.

Lemma 2.4. *Let $\{X_n\}, X$ be processes with sample paths in $D_{\mathbb{R}^d}[0, \infty)$ defined on the same probability space. Suppose that $\{X_n\}$ is relatively compact in $D_{\mathbb{R}^d}[0, \infty)$ and that for a dense set $H \subset [0, \infty)$, $X_n(t) \rightarrow X(t)$ in probability for all $t \in H$. Then $X_n \rightarrow X$ in probability in $D_{\mathbb{R}^d}[0, \infty)$. In particular, if X is continuous, then $X_n \rightarrow X$ ucp.*

We will also need the following lemma, which is easily proved using the Prohorov metric.

Lemma 2.5. *Let (E, r) be a complete and separable metric space. Let X_n be a sequence of E -valued random variables and suppose, for each k , there exists a sequence $\{X_{n,k}\}_{n=1}^\infty$ such that $\limsup_{n \rightarrow \infty} E[r(X_n, X_{n,k})] \leq \delta_k$, where $\delta_k \rightarrow 0$ as $k \rightarrow \infty$. Suppose also that for each k , there exists Y_k such that $X_{n,k} \rightarrow Y_k$ in law as $n \rightarrow \infty$. Then there exists X such that $X_n \rightarrow X$ in law and $Y_k \rightarrow X$ in law.*

2.2 Elements of Malliavin calculus

In the sequel, we will need some elements of Malliavin calculus that we collect here. The reader is referred to [6] or [10] for any unexplained notion discussed in this section.

We denote by $X = \{X(\varphi) : \varphi \in \mathfrak{H}\}$ an isonormal Gaussian process over \mathfrak{H} , a real and separable Hilbert space. By definition, X is a centered Gaussian family indexed by the elements of \mathfrak{H} and such that, for every $\varphi, \psi \in \mathfrak{H}$,

$$E[X(\varphi)X(\psi)] = \langle \varphi, \psi \rangle_{\mathfrak{H}}.$$

We denote by $\mathfrak{H}^{\otimes q}$ and $\mathfrak{H}^{\odot q}$, respectively, the tensor space and the symmetric tensor space of order $q \geq 1$. Let \mathcal{S} be the set of cylindrical functionals F of the form

$$F = f(X(\varphi_1), \dots, X(\varphi_n)), \quad (2.7)$$

where $n \geq 1$, $\varphi_i \in \mathfrak{H}$ and the function $f \in C^\infty(\mathbb{R}^n)$ is such that its partial derivatives have polynomial growth. The Malliavin derivative DF of a functional F of the form (2.7) is the square integrable \mathfrak{H} -valued random variable defined as

$$DF = \sum_{i=1}^n \frac{\partial f}{\partial x_i}(X(\varphi_1), \dots, X(\varphi_n)) \varphi_i.$$

In particular, $DX(\varphi) = \varphi$ for every $\varphi \in \mathfrak{H}$. By iteration, one can define the m th derivative $D^m F$ (which is an element of $L^2(\Omega, \mathfrak{H}^{\odot m})$) for every $m \geq 2$, giving

$$D^m F = \sum_{i_1, \dots, i_m} \frac{\partial^m f}{\partial x_{i_1} \cdots \partial x_{i_m}}(X(\varphi_1), \dots, X(\varphi_n)) \varphi_{i_1} \otimes \cdots \otimes \varphi_{i_m}.$$

As usual, for $m \geq 1$, $\mathbb{D}^{m,2}$ denotes the closure of \mathcal{S} with respect to the norm $\|\cdot\|_{m,2}$, defined by the relation

$$\|F\|_{m,2}^2 = EF^2 + \sum_{i=1}^m E\|D^i F\|_{\mathfrak{H}^{\otimes i}}^2.$$

The Malliavin derivative D satisfies the following chain rule: if $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is in C_b^1 (that is, the collection of continuously differentiable functions with a bounded derivative) and if $\{F_i\}_{i=1, \dots, n}$ is a vector of elements of $\mathbb{D}^{1,2}$, then $f(F_1, \dots, F_n) \in \mathbb{D}^{1,2}$ and

$$Df(F_1, \dots, F_n) = \sum_{i=1}^n \frac{\partial f}{\partial x_i}(F_1, \dots, F_n) DF_i. \quad (2.8)$$

This formula can be extended to higher order derivatives as

$$D^m f(F_1, \dots, F_n) = \sum_{\nu \in \mathcal{P}_m} C_\nu \sum_{i_1, \dots, i_k=1}^n \frac{\partial^k f}{\partial x_{i_1} \cdots \partial x_{i_k}}(F_1, \dots, F_n) D^{\nu_1} F_{i_1} \otimes \cdots \otimes D^{\nu_k} F_{i_k}, \quad (2.9)$$

where \mathcal{P}_m is the set of vectors $\nu = (\nu_1, \dots, \nu_k) \in \mathbb{N}^k$ such that $k \geq 1$, $\nu_1 \leq \cdots \leq \nu_k$, and $\nu_1 + \cdots + \nu_k = m$. The constants C_ν can be written explicitly as $C_\nu = m! (\prod_{j=1}^m m_j! (j!)^{m_j})^{-1}$, where $m_j = |\{\ell : \nu_\ell = j\}|$.

Remark 2.6. In (2.9), $a \otimes b$ denotes the symmetrization of the tensor product $a \otimes b$. Recall that, in general, the *symmetrization* of a function f of m variables is the function \tilde{f} defined by

$$\tilde{f}(t_1, \dots, t_m) = \frac{1}{m!} \sum_{\sigma \in \mathfrak{S}_m} f(t_{\sigma(1)}, \dots, t_{\sigma(m)}), \quad (2.10)$$

where \mathfrak{S}_m denotes the set of all permutations of $\{1, \dots, m\}$.

We denote by I the adjoint of the operator D , also called the divergence operator. A random element $u \in L^2(\Omega, \mathfrak{H})$ belongs to the domain of I , noted $\text{Dom}(I)$, if and only if it satisfies

$$|E\langle DF, u \rangle_{\mathfrak{H}}| \leq c_u \sqrt{EF^2} \quad \text{for any } F \in \mathcal{S},$$

where c_u is a constant depending only on u . If $u \in \text{Dom}(I)$, then the random variable $I(u)$ is defined by the duality relationship (customarily called ‘‘integration by parts formula’’):

$$E[FI(u)] = E\langle DF, u \rangle_{\mathfrak{H}}, \quad (2.11)$$

which holds for every $F \in \mathbb{D}^{1,2}$.

For every $n \geq 1$, let \mathcal{H}_n be the n th Wiener chaos of X , that is, the closed linear subspace of L^2 generated by the random variables $\{h_n(X(\varphi)) : \varphi \in \mathfrak{H}, |\varphi|_{\mathfrak{H}} = 1\}$, where h_n is the Hermite polynomial defined by (2.1). The mapping

$$I_n(\varphi^{\otimes n}) = h_n(X(\varphi)) \quad (2.12)$$

provides a linear isometry between the symmetric tensor product $\mathfrak{H}^{\otimes n}$ (equipped with the modified norm $\frac{1}{\sqrt{n!}} \|\cdot\|_{\mathfrak{H}^{\otimes n}}$) and \mathcal{H}_n . We set $I_n(f) := I_n(\tilde{f})$ when $f \in \mathfrak{H}^{\otimes n}$. The following duality formula holds:

$$E[FI_n(f)] = E\langle D^n F, f \rangle_{\mathfrak{H}^{\otimes n}}, \quad (2.13)$$

for any element $f \in \mathfrak{H}^{\otimes n}$ and any random variable $F \in \mathbb{D}^{n,2}$. We will also need the following particular case of the classical product formula between multiple integrals: if $\varphi, \psi \in \mathfrak{H}$ and $m, n \geq 1$, then

$$I_m(\varphi^{\otimes m}) I_n(\psi^{\otimes n}) = \sum_{r=0}^{m \wedge n} r! \binom{m}{r} \binom{n}{r} I_{m+n-2r}(\varphi^{\otimes(m-r)} \otimes \psi^{\otimes(n-r)}) \langle \varphi, \psi \rangle_{\mathfrak{H}}^r. \quad (2.14)$$

Finally, we mention that the Gaussian space generated by $B = B^{1/6}$ can be identified with an isonormal Gaussian process of the type $B = \{B(h) : h \in \mathfrak{H}\}$, where the real and separable Hilbert space \mathfrak{H} is defined as follows: (i) denote by \mathcal{E} the set of all \mathbb{R} -valued step functions on $[0, \infty)$, (ii) define \mathfrak{H} as the Hilbert space obtained by closing \mathcal{E} with respect to the scalar product

$$\langle \mathbf{1}_{[0,t]}, \mathbf{1}_{[0,s]} \rangle_{\mathfrak{H}} = E[B(s)B(t)] = \frac{1}{2}(t^{1/3} + s^{1/3} - |t-s|^{1/3}).$$

In particular, note that $B(t) = B(\mathbf{1}_{[0,t]})$. To end up, let us stress that the m th derivative D^m (with respect to B) verifies the Leibniz rule. That is, for any $F, G \in \mathbb{D}^{m,2}$ such that $FG \in \mathbb{D}^{m,2}$, we have

$$D_{t_1, \dots, t_m}^m (FG) = \sum D_J^{|J|} (F) D_{J^c}^{m-|J|} (G), \quad t_i \in [0, T], \quad i = 1, \dots, m, \quad (2.15)$$

where the sum runs over all subsets J of $\{t_1, \dots, t_m\}$, with $|J|$ denoting the cardinality of J . Note that we may also write this as

$$D^m (FG) = \sum_{k=0}^m \binom{m}{k} (D^k F) \otimes (D^{m-k} G). \quad (2.16)$$

2.3 Expansions and Gaussian estimates

A key tool of ours will be the following version of Taylor's theorem with remainder.

Theorem 2.7. *Let k be a nonnegative integer. If $g \in C^k(\mathbb{R}^d)$, then*

$$g(b) = \sum_{|\alpha| \leq k} \partial^\alpha g(a) \frac{(b-a)^\alpha}{\alpha!} + R_k(a, b),$$

where

$$R_k(a, b) = k \sum_{|\alpha|=k} \frac{(b-a)^\alpha}{\alpha!} \int_0^1 (1-u)^k [\partial^\alpha g(a + u(b-a)) - \partial^\alpha g(a)] du$$

if $k \geq 1$, and $R_0(a, b) = g(b) - g(a)$. In particular, $R_k(a, b) = \sum_{|\alpha|=k} h_\alpha(a, b) (b-a)^\alpha$, where h_α is a continuous function with $h_\alpha(a, a) = 0$ for all a . Moreover,

$$|R_k(a, b)| \leq (k \vee 1) \sum_{|\alpha|=k} M_\alpha |b-a|^\alpha,$$

where $M_\alpha = \sup\{|\partial^\alpha g(a + u(b-a)) - \partial^\alpha g(a)| : 0 \leq u \leq 1\}$.

The following related expansion theorem is a slight modification of Corollary 4.2 in [1].

Theorem 2.8. *Recall the Hermite polynomials $h_n(x)$ from (2.1). Let k be a nonnegative integer. Suppose $\varphi : \mathbb{R} \rightarrow \mathbb{R}$ is measurable and has polynomial growth with constants \tilde{K} and r . Suppose $f \in C^{k+1}(\mathbb{R}^d)$ has polynomial growth of order $k+1$, with constants K and r . Let $\xi \in \mathbb{R}^d$ and $Y \in \mathbb{R}$ be jointly normal with mean zero. Suppose that $EY^2 = 1$ and $E\xi_j^2 \leq \nu$ for some $\nu > 0$. Define $\eta \in \mathbb{R}^d$ by $\eta_j = E[\xi_j Y]$. Then*

$$E[f(\xi)\varphi(Y)] = \sum_{|\alpha| \leq k} \frac{1}{\alpha!} \eta^\alpha E[\partial^\alpha f(\xi)] E[h_{|\alpha|}(Y)\varphi(Y)] + R,$$

where $|R| \leq CK|\eta|^{k+1}$ and C depends only on \tilde{K} , r , ν , k , and d .

Proof. Although this theorem is very similar to Corollary 4.2 in [1], we provide here another proof by means of Malliavin calculus.

Observe first that, without loss of generality, we can assume that $\xi_i = X(v_i)$, $i = 1, \dots, d$, and $Y = X(v_{d+1})$, where X is an isonormal process over $\mathfrak{H} = \mathbb{R}^{d+1}$ and where v_1, \dots, v_{d+1} are some adequate vectors belonging in \mathfrak{H} . Since φ has polynomial growth, we can expand it in terms of Hermite polynomials, that is $\varphi = \sum_{q=0}^{\infty} c_q h_q$. Thanks to (2.2), note that $q!c_q = E[\varphi(Y)h_q(Y)]$. We set

$$\widehat{\varphi}_k = \sum_{q=0}^k c_q h_q \quad \text{and} \quad \check{\varphi}_k = \sum_{q=k+1}^{\infty} c_q h_q.$$

Of course, we have

$$E[f(\xi)\varphi(Y)] = E[f(\xi)\widehat{\varphi}_k(Y)] + E[f(\xi)\check{\varphi}_k(Y)].$$

We obtain

$$\begin{aligned} E[f(\xi)\widehat{\varphi}_k(Y)] &= \sum_{q=0}^k \frac{1}{q!} E[\varphi(Y)h_q(Y)] E[f(\xi)h_q(Y)] \\ &= \sum_{q=0}^k \frac{1}{q!} E[\varphi(Y)h_q(Y)] E[f(\xi)I_q(v_{d+1}^{\otimes q})] \quad \text{by (2.12)} \\ &= \sum_{q=0}^k \frac{1}{q!} E[\varphi(Y)h_q(Y)] E[\langle D^q f(\xi), v_{d+1}^{\otimes q} \rangle_{\mathfrak{H}^{\otimes q}}] \quad \text{by (2.13)} \\ &= \sum_{q=0}^k \frac{1}{q!} \sum_{i_1, \dots, i_q=1}^d E[\varphi(Y)h_q(Y)] E \left[\frac{\partial^q f}{\partial x_{i_1} \dots \partial x_{i_q}}(\xi) \right] \prod_{\ell=1}^q \eta_{i_\ell} \quad \text{by (2.9)}. \end{aligned}$$

Since the map $\Phi : \{1, \dots, d\}^q \rightarrow \{\alpha \in \mathbb{N}_0^d : |\alpha| = q\}$ defined by $(\Phi(i_1, \dots, i_q))_j = |\{\ell : i_\ell = j\}|$ is a surjection with $|\Phi^{-1}(\alpha)| = q!/\alpha!$, this gives

$$\begin{aligned} E[f(\xi)\widehat{\varphi}_k(Y)] &= \sum_{q=0}^k \frac{1}{q!} \sum_{|\alpha|=q} \frac{q!}{\alpha!} E[\varphi(Y)h_q(Y)] E[\partial^\alpha f(\xi)] \eta^\alpha \\ &= \sum_{|\alpha| \leq k} \frac{1}{\alpha!} E[\varphi(Y)h_{|\alpha|}(Y)] E[\partial^\alpha f(\xi)] \eta^\alpha. \end{aligned}$$

On the other hand, the identity (2.2), combined with the fact that each monomial x^n can be expanded in terms of the first n Hermite polynomials, implies that $E[Y^{|\alpha|}\check{\varphi}_k(Y)] = 0$ for all $|\alpha| \leq k$. Now, let $U = \xi - \eta Y$ and define $g : \mathbb{R}^d \rightarrow \mathbb{R}$ by $g(x) = E[f(U + xY)\check{\varphi}_k(Y)]$. Since φ (and, consequently, also $\check{\varphi}_k$) and f have polynomial growth, and all derivatives of f up to order $k+1$ have polynomial growth, we may differentiate under the expectation and conclude that $g \in C^{k+1}(\mathbb{R}^d)$. Hence, by Taylor's theorem (more specifically, by the version of Taylor's theorem which appears as Theorem 2.13 in [1]), and the fact that U and Y are independent,

$$\begin{aligned} E[f(\xi)\check{\varphi}_k(Y)] &= g(\eta) = \sum_{|\alpha| \leq k} \frac{1}{\alpha!} \eta^\alpha \partial^\alpha g(0) + R \\ &= \sum_{|\alpha| \leq k} \frac{1}{\alpha!} \eta^\alpha E[\partial^\alpha f(U)] E[Y^{|\alpha|}\check{\varphi}_k(Y)] + R = R, \end{aligned}$$

where

$$|R| \leq \frac{Md^{(k+1)/2}}{k!} |\eta|^{k+1},$$

and $M = \sup\{|\partial^\alpha g(u\eta)| : 0 \leq u \leq 1, |\alpha| = k+1\}$. Note that

$$\partial^\alpha g(u\eta) = E[\partial^\alpha f(U + u\eta Y)Y^{|\alpha|} \check{\varphi}_k(Y)] = E[\partial^\alpha f(\xi - \eta(1-u)Y)Y^{|\alpha|} \check{\varphi}_k(Y)].$$

Hence,

$$\begin{aligned} |\partial^\alpha g(u\eta)| &\leq K\tilde{K}E[(1 + |\xi - \eta(1-u)Y|^r)|Y|^{|\alpha|}(1 + |Y|^r)] \\ &\leq K\tilde{K}E[(1 + 2^r|\xi|^r + 2^r|\eta|^r|Y|^r)(|Y|^{|\alpha|} + |Y|^{|\alpha|+r})]. \end{aligned}$$

Since $|\eta|^2 \leq vd$, this completes the proof. \square

The following special case will be used multiple times.

Corollary 2.9. *Let X_1, \dots, X_n be jointly normal, each with mean zero and variance bounded by $v > 0$. Let $\eta_{ij} = E[X_i X_j]$. If $f \in C^1(\mathbb{R}^{n-1})$ has polynomial growth of order 1 with constants K and r , then*

$$|E[f(X_1, \dots, X_{n-1})X_n^3]| \leq CK\sigma^3 \max_{j < n} |\eta_{jn}|, \quad (2.17)$$

where $\sigma = (EX_n^2)^{1/2}$ and C depends only on r , v , and n .

Proof. Apply Theorem 2.8 with $k = 0$. \square

Finally, the following covariance estimates will be critical.

Lemma 2.10. *Recall the notation $\beta_j = (B(t_{j-1}) + B(t_j))/2$ and $r_+ = r \vee 1$. For any i, j ,*

$$(i) |E[\Delta B_i \Delta B_j]| \leq C\Delta t^{1/3} |j - i|_+^{-5/3},$$

$$(ii) |E[B(t_i) \Delta B_j]| \leq C\Delta t^{1/3} (j^{-2/3} + |j - i|_+^{-2/3}),$$

$$(iii) |E[\beta_i \Delta B_j]| \leq C\Delta t^{1/3} (j^{-2/3} + |j - i|_+^{-2/3}),$$

$$(iv) |E[\beta_j \Delta B_j]| \leq C\Delta t^{1/3} j^{-2/3}, \text{ and}$$

$$(v) C_1 |t_j - t_i|^{1/3} \leq E|\beta_j - \beta_i|^2 \leq C_2 |t_j - t_i|^{1/3},$$

where C_1, C_2 are positive, finite constants that do not depend on i or j .

Proof. (i) By symmetry, we may assume $i \leq j$. First, assume $j - i \geq 2$. Then

$$E[\Delta B_i \Delta B_j] = \int_{t_{i-1}}^{t_i} \int_{t_{j-1}}^{t_j} \partial_{st}^2 R(s, t) dt ds,$$

where $\partial_{st}^2 = \partial_1 \partial_2$. Note that for $s < t$, $\partial_{st}^2 R(s, t) = -(1/9)(t-s)^{-5/3}$. Hence,

$$|E[\Delta B_i \Delta B_j]| \leq C\Delta t^2 |t_{j-1} - t_i|^{-5/3} \leq C\Delta t^{1/3} |j - i|^{-5/3}.$$

Now assume $j - i \leq 1$. By Hölder's inequality, $|E[\Delta B_i \Delta B_j]| \leq \Delta t^{1/3} = \Delta t^{1/3} |j - i|_+^{-5/3}$.

(ii) First note that by (i),

$$|E[B(t_i) \Delta B_j]| \leq \sum_{k=1}^i |E[\Delta B_k \Delta B_j]| \leq C \Delta t^{1/3} \sum_{k=1}^j |k - j|_+^{-5/3} \leq C \Delta t^{1/3}.$$

This proves the lemma when either $j = 1$ or $|j - i|_+ = 1$. To complete the proof of (ii), suppose $j > 1$ and $|j - i| > 1$. Note that if $t > 0$ and $s \neq t$, then

$$\partial_2 R(s, t) = \frac{1}{6} t^{-2/3} - \frac{1}{6} |t - s|^{-2/3} \operatorname{sgn}(t - s).$$

We may therefore write $E[B(t_i) \Delta B_j] = \int_{t_{j-1}}^{t_j} \partial_2 R(t_i, u) du$, giving

$$|E[B(t_i) \Delta B_j]| \leq \Delta t \sup_{u \in [t_{j-1}, t_j]} |\partial_2 R(t_i, u)| \leq C \Delta t^{1/3} (j^{-2/3} + |j - i|_+^{-2/3}),$$

which is (ii).

(iii) This follows immediately from (ii).

(iv) Note that $2\beta_j \Delta B_j = B(t_j)^2 - B(t_{j-1})^2$. Since $EB(t)^2 = t^{1/3}$, the mean value theorem gives $|E[\beta_j \Delta B_j]| \leq C(\Delta t) t_j^{-2/3} = C \Delta t^{1/3} j^{-2/3}$.

(v) Without loss of generality, we may assume $i < j$. The upper bound follows from

$$2(\beta_j - \beta_i) = (B(t_j) - B(t_i)) + (B(t_{j-1}) - B(t_{i-1})),$$

and the fact that $E|B(t) - B(s)|^2 = |t - s|^{1/3}$. For the lower bound, we first assume $i < j - 1$ and write

$$2(\beta_j - \beta_i) = 2(B(t_{j-1}) - B(t_i)) + \Delta B_j + \Delta B_i.$$

For any random variables a, b, c with $a = b + c$ recall that $-E[ac] \leq (E[|a|^2]E[|c|^2])^{1/2}$ leading to

$$E[|b|^2] = E[|a - c|^2] \leq ((E[|a|^2])^{1/2} + (E[|c|^2])^{1/2})^2.$$

Taking the square root in both side of this inequality we get

$$(E[|b|^2])^{1/2} \leq (E[|a|^2])^{1/2} + (E[|c|^2])^{1/2}.$$

Letting, $a = (\beta_j - \beta_i)$, $b = (B(t_{j-1}) - B(t_i))$, and $c = (\Delta B_j + \Delta B_i)/2$ yields

$$(E[|\beta_j - \beta_i|^2])^{1/2} \geq |t_{j-1} - t_i|^{1/6} - \frac{1}{2} (E[|\Delta B_j + \Delta B_i|^2])^{1/2}.$$

Since ΔB_i and ΔB_j are negatively correlated,

$$E[|\Delta B_j + \Delta B_i|^2] \leq E[|\Delta B_j|^2] + E[|\Delta B_i|^2] = 2\Delta t^{1/3}.$$

Thus,

$$(E[|\beta_j - \beta_i|^2])^{1/2} \geq \Delta t^{1/6} |j - 1 - i|^{1/6} - 2^{-1/2} \Delta t^{1/6} \geq C \Delta t^{1/6} |j - i|^{1/6},$$

for some $C > 0$. This completes the proof when $i < j - 1$.

If $i = j - 1$, the conclusion is immediate, since $2(\beta_j - \beta_{j-1}) = B(t_j) - B(t_{j-2})$. □

2.4 Sextic and signed cubic variations

Theorem 2.11. For each $T > 0$, we have $E[\sup_{0 \leq t \leq T} |V_n^6(B, t) - 15t|^2] \rightarrow 0$ as $n \rightarrow \infty$. In particular, $V_n^6(B, t) \rightarrow 15t$ ucp.

Proof. Since $V_n^6(B)$ is monotone, it will suffice to show that $V_n^6(B, t) \rightarrow 15t$ in L^2 for each fixed t . Indeed, the uniform convergence will then be a direct consequence of Dini's theorem. We write

$$V_n^6(B, t) - 15t = \sum_{j=1}^{\lfloor nt \rfloor} (\Delta B_j^6 - 15\Delta t) + 15(\lfloor nt \rfloor / n - t).$$

Since $|\lfloor nt \rfloor / n - t| \leq \Delta t$, it will suffice to show that $E|\sum_{j=1}^{\lfloor nt \rfloor} (\Delta B_j^6 - 15\Delta t)|^2 \rightarrow 0$. For this, we compute

$$\begin{aligned} E \left| \sum_{j=1}^{\lfloor nt \rfloor} (\Delta B_j^6 - 15\Delta t) \right|^2 &= \sum_{i=1}^{\lfloor nt \rfloor} \sum_{j=1}^{\lfloor nt \rfloor} E[(\Delta B_i^6 - 15\Delta t)(\Delta B_j^6 - 15\Delta t)] \\ &= \sum_{i=1}^{\lfloor nt \rfloor} \sum_{j=1}^{\lfloor nt \rfloor} (E[\Delta B_i^6 \Delta B_j^6] - 225\Delta t^2). \end{aligned} \tag{2.18}$$

By Theorem 2.8, if ξ, Y are jointly Gaussian, standard normals, then $E[\xi^6 Y^6] = 225 + R$, where $|R| \leq C|E[\xi Y]|^2$. Applying this with $\xi = \Delta t^{-1/6} \Delta B_i$ and $Y = \Delta t^{-1/6} \Delta B_j$, and using Lemma 2.10(i), gives $|E[\Delta B_i^6 \Delta B_j^6] - 225\Delta t^2| \leq C\Delta t^2 |j - i|_+^{-10/3}$. Substituting this into (2.18), we have

$$E \left| \sum_{j=1}^{\lfloor nt \rfloor} (\Delta B_j^6 - 15\Delta t) \right|^2 \leq C\lfloor nt \rfloor \Delta t^2 \leq Ct\Delta t \rightarrow 0,$$

which completes the proof. \square

Theorem 2.12. As $n \rightarrow \infty$, $(B, V_n(B)) \rightarrow (B, [[B]])$ in law in $D_{\mathbb{R}^2}[0, \infty)$.

Proof. By Theorem 10 in [11], $(B, V_n(B)) \rightarrow (B, \kappa W) = (B, [[B]])$ in law in $(D_{\mathbb{R}}[0, \infty))^2$. By Lemma 2.2, this implies $(B, V_n(B)) \rightarrow (B, [[B]])$ in $D_{\mathbb{R}^2}[0, \infty)$. \square

2.5 Main result

Given $g \in C^\infty(\mathbb{R})$, choose G such that $G' = g$. We then define

$$\int_0^t g(B(s)) dB(s) = G(B(t)) - G(B(0)) + \frac{1}{12} \int_0^t G'''(B(s)) d[[B]]_s. \tag{2.19}$$

Note that, by definition, the change of variable formula (1.3) holds for all $g \in C^\infty$. We shall use the shorthand notation $\int g(B) dB$ to refer to the process $t \mapsto \int_0^t g(B(s)) dB(s)$. Similarly, $\int g(B) d[[B]]$ and $\int g(B) ds$ shall refer to the processes $t \mapsto \int_0^t g(B(s)) d[[B]]_s$ and $t \mapsto \int_0^t g(B(s)) ds$, respectively. Our main result is the following.

Theorem 2.13. *If $g \in C^\infty(\mathbb{R})$, then $(B, V_n(B), I_n(g, B)) \rightarrow (B, [[B]], \int g(B) dB)$ in law in $D_{\mathbb{R}^3}[0, \infty)$.*

We also have the following generalization concerning the joint convergence of multiple sequences of Riemann sums.

Theorem 2.14. *Fix $k \geq 1$. Let $g_j \in C^\infty(\mathbb{R})$ for $1 \leq j \leq k$. Let J_n be the \mathbb{R}^k -valued process whose j -th component is $(J_n)_j = I_n(g_j, B)$. Similarly, define J by $J_j = \int g_j(B) dB$. Then $(B, V_n(B), J_n) \rightarrow (B, [[B]], J)$ in law in $D_{\mathbb{R}^{k+2}}[0, \infty)$.*

Remark 2.15. In less formal language, Theorem 2.14 states that the Riemann sums $I_n(g_j, B)$ converge jointly, and the limiting stochastic integrals are all defined in terms of the same Brownian motion. In other words, the limiting Brownian motion remains unchanged under changes in the integrand. In this sense, the limiting Brownian motion depends only on B , despite being independent of B in the probabilistic sense.

The proofs of these two theorems are given in Section 5.

3 Finite-dimensional distributions

Theorem 3.1. *If $g \in C^\infty(\mathbb{R})$ is bounded with bounded derivatives, then*

$$\left(B, V_n(B), \frac{1}{\sqrt{n}} \sum_{j=1}^{\lfloor n \cdot \rfloor} \frac{g(B(t_{j-1})) + g(B(t_j))}{2} h_3(n^{1/6} \Delta B_j) \right) \rightarrow \left(B, [[B]], \int g(B) d[[B]] \right),$$

in the sense of finite-dimensional distributions on $[0, \infty)$.

The rest of this section is devoted to the proof of Theorem 3.1.

3.1 Some technical lemmas

During the proof of Theorem 3.1, we will need technical results that are collected here. Moreover, for notational convenience, we will make use of the following shorthand notation:

$$\delta_j = \mathbf{1}_{[t_{j-1}, t_j]} \quad \text{and} \quad \varepsilon_j = \mathbf{1}_{[0, t_j]}.$$

For future reference, let us note that by (2.10),

$$\varepsilon_t^{\otimes a} \otimes \varepsilon_s^{\otimes (q-a)} = \binom{q}{a}^{-1} \sum_{\substack{i_1, \dots, i_q \in \{s, t\} \\ |\{j: i_j = s\}| = q-a}} \varepsilon_{i_1} \otimes \dots \otimes \varepsilon_{i_q}. \quad (3.1)$$

Lemma 3.2. *We have*

$$(i) \quad |E[B(r)(B(t) - B(s))]| = |\langle \mathbf{1}_{[0, r]}, \mathbf{1}_{[s, t]} \rangle_{\mathcal{H}}| \leq |t - s|^{1/3} \text{ for any } r, s, t \geq 0;$$

$$(ii) \quad \sup_{0 \leq s \leq T} \sum_{k=1}^{\lfloor nt \rfloor} |E[B(s) \Delta B_k]| = \sup_{0 \leq s \leq T} \sum_{k=1}^{\lfloor nt \rfloor} |\langle \mathbf{1}_{[0, s]}, \delta_k \rangle_{\mathcal{H}}| \underset{n \rightarrow \infty}{=} O(1) \text{ for any fixed } t, T > 0;$$

$$(iii) \sum_{k,j=1}^{\lfloor nt \rfloor} |E(B(t_{j-1})\Delta B_k)| = \sum_{k,j=1}^{\lfloor nt \rfloor} |\langle \varepsilon_{j-1}, \delta_k \rangle_{\mathfrak{H}}| \underset{n \rightarrow \infty}{=} O(n) \text{ for any fixed } t > 0;$$

$$(iv) \sum_{k=1}^{\lfloor nt \rfloor} \left| (E[B(t_{k-1})\Delta B_k])^3 + \frac{1}{8n} \right| = \sum_{k=1}^{\lfloor nt \rfloor} \left| \langle \varepsilon_{k-1}, \delta_k \rangle_{\mathfrak{H}}^3 + \frac{1}{8n} \right| \underset{n \rightarrow \infty}{\longrightarrow} 0 \text{ for any fixed } t > 0;$$

$$(v) \sum_{k=1}^{\lfloor nt \rfloor} \left| (E[B(t_k)\Delta B_k])^3 - \frac{1}{8n} \right| = \sum_{k=1}^{\lfloor nt \rfloor} \left| \langle \varepsilon_k, \delta_k \rangle_{\mathfrak{H}}^3 - \frac{1}{8n} \right| \underset{n \rightarrow \infty}{\longrightarrow} 0 \text{ for any fixed } t > 0.$$

Proof.

(i) We have

$$E(B(r)(B(t) - B(s))) = \frac{1}{2}(t^{1/3} - s^{1/3}) + \frac{1}{2}(|s - r|^{1/3} - |t - r|^{1/3}).$$

Using the classical inequality $||b|^{1/3} - |a|^{1/3}| \leq |b - a|^{1/3}$, the desired result follows.

(ii) Observe that

$$E(B(s)\Delta B_k) = \frac{1}{2n^{1/3}} (k^{1/3} - (k-1)^{1/3} - |k - ns|^{1/3} + |k - ns - 1|^{1/3}).$$

We deduce, for any fixed $s \leq t$:

$$\begin{aligned} \sum_{k=1}^{\lfloor nt \rfloor} |E(B(s)\Delta B_k)| &\leq \frac{1}{2}t^{1/3} + \frac{1}{2n^{1/3}} \left(|(\lfloor ns \rfloor - ns + 1)^{1/3} - (ns - \lfloor ns \rfloor)^{1/3}| \right. \\ &\quad \left. + \sum_{k=1}^{\lfloor ns \rfloor} |(ns + 1 - k)^{1/3} - (ns - k)^{1/3}| + \sum_{k=\lfloor ns \rfloor + 2}^{\lfloor nt \rfloor} |(k - ns)^{1/3} - (k - ns - 1)^{1/3}| \right) \\ &= \frac{1}{2}(t^{1/3} + s^{1/3} + |t - s|^{1/3}) + R_n, \end{aligned}$$

where $|R_n| \leq Cn^{-1/3}$, and C does not depend on s or t . The case where $s > t$ can be obtained similarly. Taking the supremum over $s \in [0, T]$ gives us (ii).

(iii) is a direct consequence of (ii).

(iv) We have

$$\begin{aligned} \left| (E(B(t_{k-1})\Delta B_k))^3 + \frac{1}{8n} \right| &= \frac{1}{8n} (k^{1/3} - (k-1)^{1/3}) \\ &\quad \times \left| (k^{1/3} - (k-1)^{1/3})^2 - 3(k^{1/3} - (k-1)^{1/3}) + 3 \right|. \end{aligned}$$

Thus, the desired convergence is immediately checked by combining the bound $0 \leq k^{1/3} - (k-1)^{1/3} \leq 1$ with a telescoping sum argument.

(v) The proof is very similar to the proof of (iv). \square

Lemma 3.3. *Let $s \geq 1$, and suppose that $\phi \in C^6(\mathbb{R}^s)$ and $g_1, g_2 \in C^6(\mathbb{R})$ have polynomial growth of order 6, all with constants K and r . Fix $a, b \in [0, T]$. Then*

$$\sup_{u_1, \dots, u_s \in [0, T]} \sup_{n \geq 1} \sum_{i_1=1}^{\lfloor na \rfloor} \sum_{i_2=1}^{\lfloor nb \rfloor} \left| E(\phi(B(u_1), \dots, B(u_s))g_1(B(t_{i_1-1}))g_2(B(t_{i_2-1}))I_3(\delta_{i_1}^{\otimes 3})I_3(\delta_{i_2}^{\otimes 3})) \right|$$

is finite.

Proof. Let C denote a constant depending only on T, s, K , and r , and whose value can change from one line to another. Define $f : \mathbb{R}^{s+3} \rightarrow \mathbb{R}$ by

$$f(x) = \phi(x_1, \dots, x_s)g_1(x_{s+1})g_2(x_{s+2})h_3(x_{s+3}).$$

Let $\xi_i = B(u_i)$, $i = 1, \dots, s$; $\xi_{s+1} = B(t_{i_1-1})$, $\xi_{s+2} = B(t_{i_2-1})$, $\xi_{s+3} = n^{1/6}\Delta B_{i_1}$, and $\eta_i = n^{1/6}E[\xi_i\Delta B_{i_2}]$. Applying Theorem 2.8 with $k = 5$, we obtain

$$\begin{aligned} & E(\phi(B(u_1), \dots, B(u_s))g_1(B(t_{i_1-1}))g_2(B(t_{i_2-1}))I_3(\delta_{i_1}^{\otimes 3})I_3(\delta_{i_2}^{\otimes 3})) \\ &= \frac{1}{n}E(\phi(B(u_1), \dots, B(u_s))g_1(B(t_{i_1-1}))g_2(B(t_{i_2-1}))h_3(n^{1/6}\Delta B_{i_1})h_3(n^{1/6}\Delta B_{i_2})) \\ &= \frac{1}{n} \sum_{|\alpha|=3} \frac{6}{\alpha!} E[\partial^\alpha f(\xi)] \eta^\alpha + \frac{R}{n}, \end{aligned}$$

where $|R| \leq C|\eta|^6$.

By Lemma 3.2 (i), we have $|\eta_i| \leq n^{-1/6}$ for any $i \leq s+2$, and $|\eta_{s+3}| \leq 1$ by Lemma 2.10 (i). Moreover, we have

$$\frac{1}{n} \sum_{i_1=1}^{\lfloor na \rfloor} \sum_{i_2=1}^{\lfloor nb \rfloor} |\eta_{s+3}| = \frac{1}{2n} \sum_{i_1=1}^{\lfloor na \rfloor} \sum_{i_2=1}^{\lfloor nb \rfloor} \left| |i_1 - i_2 + 1|^{1/3} + |i_1 - i_2 - 1|^{1/3} - 2|i_1 - i_2|^{1/3} \right| \leq C.$$

Therefore, by taking into account these two facts, we deduce $\frac{1}{n} \sum_{i_1=1}^{\lfloor na \rfloor} \sum_{i_2=1}^{\lfloor nb \rfloor} |R| \leq C$.

On the other hand, if $\alpha \in \mathbb{N}_0^{s+3}$ is such that $|\alpha| = 3$ with $\alpha_{s+3} \neq 0$, we have

$$\begin{aligned} & \frac{1}{n} \sum_{i_1=1}^{\lfloor na \rfloor} \sum_{i_2=1}^{\lfloor nb \rfloor} \frac{6}{\alpha!} |E[\partial^\alpha f(\xi)]| |\eta^\alpha| \\ & \leq \frac{C}{n} \sum_{i_1=1}^{\lfloor na \rfloor} \sum_{i_2=1}^{\lfloor nb \rfloor} \left| |i_1 - i_2 + 1|^{1/3} + |i_1 - i_2 - 1|^{1/3} - 2|i_1 - i_2|^{1/3} \right| \leq C. \end{aligned}$$

Note that $E[\partial^\alpha f(\xi)] < \infty$ since the function $\partial^\alpha f$ has polynomial growth by our assumptions on ϕ , g_1 , and g_2 , and since ξ is a Gaussian vector.

Finally, if $\alpha \in \mathbb{N}_0^{s+3}$ is such that $|\alpha| = 3$ with $\alpha_{s+3} = 0$ then $\partial^\alpha f = \partial^\alpha \hat{f} \otimes h_3$ with $\hat{f} : \mathbb{R}^{s+2} \rightarrow \mathbb{R}$ defined by $\hat{f}(x) = \phi(x_1, \dots, x_s)g_1(x_{s+1})g_2(x_{s+2})$. Hence, applying Theorem 2.8 to $f = \partial^\alpha \hat{f}$ and $\varphi = h_3$ with $k = 2$, we deduce, for $\hat{\eta} \in \mathbb{N}_0^{s+2}$ defined by $\hat{\eta}_i = \eta_i$,

$$\left| E[\partial^\alpha f(\xi)] \right| = \left| E[\partial^\alpha \hat{f}(\xi)h_3(n^{1/6}\Delta B_{i_1})] \right| \leq C|\hat{\eta}|^3 \leq Cn^{-1/2},$$

so that

$$\frac{1}{n} \sum_{i_1=1}^{\lfloor na \rfloor} \sum_{i_2=1}^{\lfloor nb \rfloor} \frac{6}{\alpha!} |E[\partial^\alpha f(\xi)]| |\eta^\alpha| = \frac{1}{n} \sum_{i_1=1}^{\lfloor na \rfloor} \sum_{i_2=1}^{\lfloor nb \rfloor} \frac{6}{\alpha!} |E[\partial^\alpha f(\xi)]| |\hat{\eta}^\alpha| \leq C.$$

The proof of Lemma 3.3 is done. \square

Lemma 3.4. *Let $g, h \in C^q(\mathbb{R})$, $q \geq 1$, having bounded derivatives, and fix $s, t \geq 0$. Set $\varepsilon_t = \mathbf{1}_{[0,t]}$ and $\varepsilon_s = \mathbf{1}_{[0,s]}$. Then $g(B(t))h(B(s))$ belongs in $\mathbb{D}^{q,2}$ and we have*

$$D^q(g(B(t))h(B(s))) = \sum_{a=0}^q \binom{q}{a} g^{(a)}(B(t))h^{(q-a)}(B(s)) \varepsilon_t^{\otimes a} \otimes \varepsilon_s^{\otimes (q-a)}. \quad (3.2)$$

Proof. This follows immediately from (2.16). \square

Lemma 3.5. *Fix an integer $r \geq 1$, and some real numbers $s_1, \dots, s_r \geq 0$. Suppose $\varphi \in C^\infty(\mathbb{R}^r)$ and $g_j \in C^\infty(\mathbb{R})$, $j = 1, 2, 3, 4$, are bounded with bounded partial derivatives. For $i_1, i_2, i_3, i_4 \in \mathbb{N}$, set $\Phi(i_1, i_2, i_3, i_4) := \varphi(B_{s_1}, \dots, B_{s_r}) \prod_{j=1}^4 g_j(B_{t_{i_j-1}})$. Then, for any fixed $a, b, c, d > 0$, the following estimate is in order:*

$$\sup_{n \geq 1} \sum_{i_1=1}^{\lfloor na \rfloor} \sum_{i_2=1}^{\lfloor nb \rfloor} \sum_{i_3=1}^{\lfloor nc \rfloor} \sum_{i_4=1}^{\lfloor nd \rfloor} \left| E \left(\Phi(i_1, i_2, i_3, i_4) I_3(\delta_{i_1}^{\otimes 3}) I_3(\delta_{i_2}^{\otimes 3}) I_3(\delta_{i_3}^{\otimes 3}) I_3(\delta_{i_4}^{\otimes 3}) \right) \right| < \infty. \quad (3.3)$$

Proof. Using the product formula (2.14), we have that $I_3(\delta_{i_3}^{\otimes 3}) I_3(\delta_{i_4}^{\otimes 3})$ equals

$$I_6(\delta_{i_3}^{\otimes 3} \otimes \delta_{i_4}^{\otimes 3}) + 9I_4(\delta_{i_3}^{\otimes 2} \otimes \delta_{i_4}^{\otimes 2}) \langle \delta_{i_3}, \delta_{i_4} \rangle_{\mathfrak{H}} + 18I_2(\delta_{i_3} \otimes \delta_{i_4}) \langle \delta_{i_3}, \delta_{i_4} \rangle_{\mathfrak{H}}^2 + 6 \langle \delta_{i_3}, \delta_{i_4} \rangle_{\mathfrak{H}}^3.$$

As a consequence, we get

$$\begin{aligned} & \sum_{i_1=1}^{\lfloor na \rfloor} \sum_{i_2=1}^{\lfloor nb \rfloor} \sum_{i_3=1}^{\lfloor nc \rfloor} \sum_{i_4=1}^{\lfloor nd \rfloor} \left| E \left(\Phi(i_1, i_2, i_3, i_4) I_3(\delta_{i_1}^{\otimes 3}) I_3(\delta_{i_2}^{\otimes 3}) I_3(\delta_{i_3}^{\otimes 3}) I_3(\delta_{i_4}^{\otimes 3}) \right) \right| \\ & \leq \sum_{i_1=1}^{\lfloor na \rfloor} \sum_{i_2=1}^{\lfloor nb \rfloor} \sum_{i_3=1}^{\lfloor nc \rfloor} \sum_{i_4=1}^{\lfloor nd \rfloor} \left| E \left(\Phi(i_1, i_2, i_3, i_4) I_3(\delta_{i_1}^{\otimes 3}) I_3(\delta_{i_2}^{\otimes 3}) I_6(\delta_{i_3}^{\otimes 3} \otimes \delta_{i_4}^{\otimes 3}) \right) \right| \\ & \quad + 9 \sum_{i_1=1}^{\lfloor na \rfloor} \sum_{i_2=1}^{\lfloor nb \rfloor} \sum_{i_3=1}^{\lfloor nc \rfloor} \sum_{i_4=1}^{\lfloor nd \rfloor} \left| E \left(\Phi(i_1, i_2, i_3, i_4) I_3(\delta_{i_1}^{\otimes 3}) I_3(\delta_{i_2}^{\otimes 3}) I_4(\delta_{i_3}^{\otimes 2} \otimes \delta_{i_4}^{\otimes 2}) \right) \right| \left| \langle \delta_{i_3}, \delta_{i_4} \rangle_{\mathfrak{H}} \right| \\ & \quad + 18 \sum_{i_1=1}^{\lfloor na \rfloor} \sum_{i_2=1}^{\lfloor nb \rfloor} \sum_{i_3=1}^{\lfloor nc \rfloor} \sum_{i_4=1}^{\lfloor nd \rfloor} \left| E \left(\Phi(i_1, i_2, i_3, i_4) I_3(\delta_{i_1}^{\otimes 3}) I_3(\delta_{i_2}^{\otimes 3}) I_2(\delta_{i_3} \otimes \delta_{i_4}) \right) \right| \left| \langle \delta_{i_3}, \delta_{i_4} \rangle_{\mathfrak{H}} \right|^2 \\ & \quad + 6 \sum_{i_1=1}^{\lfloor na \rfloor} \sum_{i_2=1}^{\lfloor nb \rfloor} \sum_{i_3=1}^{\lfloor nc \rfloor} \sum_{i_4=1}^{\lfloor nd \rfloor} \left| E \left(\Phi(i_1, i_2, i_3, i_4) I_3(\delta_{i_1}^{\otimes 3}) I_3(\delta_{i_2}^{\otimes 3}) \right) \right| \left| \langle \delta_{i_3}, \delta_{i_4} \rangle_{\mathfrak{H}} \right|^3 \\ & =: A_1^{(n)} + 9A_2^{(n)} + 18A_3^{(n)} + 6A_4^{(n)}. \end{aligned}$$

(1) First, we deal with the term $A_1^{(n)}$.

$$\begin{aligned} A_1^{(n)} &= \sum_{i_1=1}^{\lfloor na \rfloor} \sum_{i_2=1}^{\lfloor nb \rfloor} \sum_{i_3=1}^{\lfloor nc \rfloor} \sum_{i_4=1}^{\lfloor nd \rfloor} \left| E \left(\Phi(i_1, i_2, i_3, i_4) I_3(\delta_{i_1}^{\otimes 3}) I_3(\delta_{i_2}^{\otimes 3}) I_6(\delta_{i_3}^{\otimes 3} \otimes \delta_{i_4}^{\otimes 3}) \right) \right| \\ &= \sum_{i_1=1}^{\lfloor na \rfloor} \sum_{i_2=1}^{\lfloor nb \rfloor} \sum_{i_3=1}^{\lfloor nc \rfloor} \sum_{i_4=1}^{\lfloor nd \rfloor} \left| E \left(\langle D^6 \left(\Phi(i_1, i_2, i_3, i_4) I_3(\delta_{i_1}^{\otimes 3}) I_3(\delta_{i_2}^{\otimes 3}) \right), \delta_{i_3}^{\otimes 3} \otimes \delta_{i_4}^{\otimes 3} \rangle_{\mathfrak{H}^{\otimes 6}} \right) \right| \end{aligned}$$

When computing the sixth Malliavin derivative $D^6(\Phi(i_1, i_2, i_3, i_4) I_3(\delta_{i_1}^{\otimes 3}) I_3(\delta_{i_2}^{\otimes 3}))$ (using Lemma 3.4), there are three types of terms:

(1a) The first type consists in terms arising when one *only* differentiates $\Phi(i_1, i_2, i_3, i_4)$. By Lemma 3.2 (i), these terms are all bounded by

$$n^{-2} \sum_{i_1=1}^{\lfloor na \rfloor} \sum_{i_2=1}^{\lfloor nb \rfloor} \sum_{i_3=1}^{\lfloor nc \rfloor} \sum_{i_4=1}^{\lfloor nd \rfloor} \left| E \left(\tilde{\Phi}(i_1, i_2, i_3, i_4) I_3(\delta_{i_1}^{\otimes 3}) I_3(\delta_{i_2}^{\otimes 3}) \right) \right|,$$

which is less than

$$cd \sup_{i_3=1, \dots, \lfloor nc \rfloor} \sup_{i_4=1, \dots, \lfloor nd \rfloor} \sum_{i_1=1}^{\lfloor na \rfloor} \sum_{i_2=1}^{\lfloor nb \rfloor} \left| E \left(\tilde{\Phi}(i_1, i_2, i_3, i_4) I_3(\delta_{i_1}^{\otimes 3}) I_3(\delta_{i_2}^{\otimes 3}) \right) \right|.$$

(Here, $\tilde{\Phi}(i_1, i_2, i_3, i_4)$ means a quantity having a similar form as $\Phi(i_1, i_2, i_3, i_4)$.) Therefore, Lemma 3.3 shows that the terms of the first type in $A_1^{(n)}$ well agree with the desired conclusion (3.3).

(1b) The second type consists in terms arising when one differentiates $\Phi(i_1, i_2, i_3, i_4)$ and $I_3(\delta_{i_1}^{\otimes 3})$, but not $I_3(\delta_{i_2}^{\otimes 3})$ (the case where one differentiates $\Phi(i_1, i_2, i_3, i_4)$ and $I_3(\delta_{i_2}^{\otimes 3})$ but not $I_3(\delta_{i_1}^{\otimes 3})$ is, of course, completely similar). In this case, with ρ defined by (2.4), the corresponding terms are bounded either by

$$Cn^{-2} \sum_{i_1=1}^{\lfloor na \rfloor} \sum_{i_2=1}^{\lfloor nb \rfloor} \sum_{i_3=1}^{\lfloor nc \rfloor} \sum_{i_4=1}^{\lfloor nd \rfloor} \sum_{\alpha=0}^2 \left| E \left(\tilde{\Phi}(i_1, i_2, i_3, i_4) I_\alpha(\delta_{i_1}^{\otimes \alpha}) I_3(\delta_{i_2}^{\otimes 3}) \right) \right| |\rho(i_3 - i_1)|,$$

or by the same quantity with $\rho(i_4 - i_1)$ instead of $\rho(i_3 - i_1)$. In order to get the previous estimate, we have used Lemma 3.2 (i) plus the fact that the sequence $\{\rho(r)\}_{r \in \mathbb{Z}}$, introduced in (2.4), is bounded (see Remark 2.1). Moreover, by (2.13) and Lemma 3.2 (i), observe that

$$\left| E \left(\tilde{\Phi}(i_1, i_2, i_3, i_4) I_\alpha(\delta_{i_1}^{\otimes \alpha}) I_3(\delta_{i_2}^{\otimes 3}) \right) \right| = \left| E \left(\langle D^3(\tilde{\Phi}(i_1, i_2, i_3, i_4) I_\alpha(\delta_{i_1}^{\otimes \alpha})), \delta_{i_2}^{\otimes 3} \rangle_{\mathfrak{H}^{\otimes 3}} \right) \right| \leq Cn^{-1},$$

for any $\alpha = 0, 1, 2$. Finally, since

$$\sup_{i_1=1, \dots, \lfloor na \rfloor} \sum_{i_3=1}^{\lfloor nc \rfloor} \sum_{i_4=1}^{\lfloor nd \rfloor} |\rho(i_3 - i_1)| \leq nd \sup_{i_1=1, \dots, \lfloor na \rfloor} \sum_{r \in \mathbb{Z}} |\rho(r)| = Cn$$

(and similarly for $\rho(i_4 - i_1)$ instead of $\rho(i_3 - i_1)$), we deduce that the terms of the second type in $A_1^{(n)}$ also agree with the desired conclusion (3.3).

(1c) The third and last type of terms consist of those that arise when one differentiates $\Phi(i_1, i_2, i_3, i_4)$, $I_3(\delta_{i_1})$ and $I_3(\delta_{i_2})$. In this case, the corresponding terms can be bounded by expressions of the type

$$Cn^{-2} \sum_{i_1=1}^{\lfloor na \rfloor} \sum_{i_2=1}^{\lfloor nb \rfloor} \sum_{i_3=1}^{\lfloor nc \rfloor} \sum_{i_4=1}^{\lfloor nd \rfloor} \sum_{\alpha=0}^2 \sum_{\beta=0}^2 \left| E \left(\tilde{\Phi}(i_1, i_2, i_3, i_4) I_\alpha(\delta_{i_1}^{\otimes \alpha}) I_\beta(\delta_{i_2}^{\otimes \beta}) \right) \right| |\rho(a - i_1)| |\rho(i_2 - a)|$$

with $a = i_3$ or $a = i_4$. Since $\left| E \left(\tilde{\Phi}(i_1, i_2, i_3, i_4) I_\alpha(\delta_{i_1}^{\otimes \alpha}) I_\beta(\delta_{i_2}^{\otimes \beta}) \right) \right|$ is uniformly bounded in n on one hand, and

$$\sum_{i_1=1}^{\lfloor na \rfloor} \sum_{i_2=1}^{\lfloor nb \rfloor} \sum_{i_3=1}^{\lfloor nc \rfloor} |\rho(i_3 - i_1)| |\rho(i_2 - i_3)| \leq nc \left(\sum_{r \in \mathbb{Z}} |\rho(r)| \right)^2 = Cn$$

on the other hand (by Remark 2.1), we deduce that the terms of the third type in $A_1^{(n)}$ also agree with the desired conclusion (3.3).

(2) Second, we focus on the term $A_2^{(n)}$. We have

$$A_2^{(n)} = \sum_{i_1=1}^{\lfloor na \rfloor} \sum_{i_2=1}^{\lfloor nb \rfloor} \sum_{i_3=1}^{\lfloor nc \rfloor} \sum_{i_4=1}^{\lfloor nd \rfloor} \left| E \left(\langle D^4 \left(\Phi(i_1, i_2, i_3, i_4) I_3(\delta_{i_1}^{\otimes 3}) I_3(\delta_{i_2}^{\otimes 3}) \right), \delta_{i_3}^{\otimes 2} \otimes \delta_{i_4}^{\otimes 2} \rangle_{\mathcal{H}^{\otimes 4}} \right) \right| \left| \langle \delta_{i_3}, \delta_{i_4} \rangle_{\mathcal{H}} \right|.$$

When computing the fourth Malliavin derivative $D^4(\Phi(i_1, i_2, i_3, i_4) I_3(\delta_{i_1}^{\otimes 3}) I_3(\delta_{i_2}^{\otimes 3}))$, we have to deal with three types of terms:

(2a) The first type consists in terms arising when one only differentiates $\Phi(i_1, i_2, i_3, i_4)$. By Lemma 3.2 (i), these terms are all bounded by

$$n^{-5/3} \sum_{i_1=1}^{\lfloor na \rfloor} \sum_{i_2=1}^{\lfloor nb \rfloor} \sum_{i_3=1}^{\lfloor nc \rfloor} \sum_{i_4=1}^{\lfloor nd \rfloor} \left| E \left(\tilde{\Phi}(i_1, i_2, i_3, i_4) I_3(\delta_{i_1}^{\otimes 3}) I_3(\delta_{i_2}^{\otimes 3}) \right) \right| |\rho(i_3 - i_4)|,$$

which is less than

$$Cn^{-2/3} \sum_{r \in \mathbb{Z}} |\rho(r)| \sup_{i_3=1, \dots, \lfloor nc \rfloor} \sup_{i_4=1, \dots, \lfloor nd \rfloor} \sum_{i_1=1}^{\lfloor na \rfloor} \sum_{i_2=1}^{\lfloor nb \rfloor} \left| E \left(\tilde{\Phi}(i_1, i_2, i_3, i_4) I_3(\delta_{i_1}^{\otimes 3}) I_3(\delta_{i_2}^{\otimes 3}) \right) \right|.$$

Hence, by Lemma 3.3, we see that the terms of the first type in $A_2^{(n)}$ well agree with the desired conclusion (3.3).

(2b) The second type consists in terms arising when one differentiates $\Phi(i_1, i_2, i_3, i_4)$ and $I_3(\delta_{i_1}^{\otimes 3})$ but not $I_3(\delta_{i_2}^{\otimes 3})$ (the case where one differentiates $\Phi(i_1, i_2, i_3, i_4)$ and $I_3(\delta_{i_2}^{\otimes 3})$ but not $I_3(\delta_{i_1}^{\otimes 3})$ is completely similar). In this case, the corresponding terms can be bounded either by

$$Cn^{-5/3} \sum_{i_1=1}^{\lfloor na \rfloor} \sum_{i_2=1}^{\lfloor nb \rfloor} \sum_{i_3=1}^{\lfloor nc \rfloor} \sum_{i_4=1}^{\lfloor nd \rfloor} \sum_{\alpha=0}^2 \left| E \left(\tilde{\Phi}(i_1, i_2, i_3, i_4) I_\alpha(\delta_{i_1}^{\otimes \alpha}) I_3(\delta_{i_2}^{\otimes 3}) \right) \right| |\rho(i_3 - i_1)| |\rho(i_3 - i_4)|,$$

or by the same quantity with $\rho(i_4 - i_1)$ instead of $\rho(i_3 - i_1)$. By Cauchy-Schwarz inequality, we have

$$\left| E \left(\tilde{\Phi}(i_1, i_2, i_3, i_4) I_\alpha(\delta_{i_1}^{\otimes \alpha}) I_3(\delta_{i_2}^{\otimes 3}) \right) \right| \leq Cn^{-\frac{3+\alpha}{6}} \leq Cn^{-1/2}.$$

Since, moreover, Remark 2.1 entails that

$$\sup_{i_1=1,\dots,[na]} \sum_{i_3=1}^{[nc]} \sum_{i_4=1}^{[nd]} |\rho(i_3 - i_1)| |\rho(i_3 - i_4)| \leq \left(\sum_{r \in \mathbb{Z}} |\rho(r)| \right)^2 = C$$

(and similarly for $\rho(i_4 - i_1)$ instead of $\rho(i_3 - i_1)$), we deduce that the terms of the second type in $A_2^{(n)}$ also agree with the desired conclusion (3.3).

(2c) The third and last type of terms consist of those that arise when one differentiates $\Phi(i_1, i_2, i_3, i_4)$, $I_3(\delta_{i_1})$ and $I_3(\delta_{i_2})$. In this case, the corresponding terms can be bounded by expressions of the type

$$C n^{-5/3} \sum_{i_1=1}^{[na]} \sum_{i_2=1}^{[nb]} \sum_{i_3=1}^{[nc]} \sum_{i_4=1}^{[nd]} \sum_{\alpha=0}^2 \sum_{\beta=0}^2 \left| E \left(\tilde{\Phi}(i_1, i_2, i_3, i_4) I_\alpha(\delta_{i_1}^{\otimes \alpha}) I_\beta(\delta_{i_2}^{\otimes \beta}) \right) \right| \times |\rho(a - i_1)| |\rho(i_2 - b)| |\rho(i_3 - i_4)|.$$

with $a, b \in \{i_3, i_4\}$. Since $\left| E \left(\tilde{\Phi}(i_1, i_2, i_3, i_4) I_\alpha(\delta_{i_1}^{\otimes \alpha}) I_\beta(\delta_{i_2}^{\otimes \beta}) \right) \right|$ is uniformly bounded in n on one hand, and

$$\sum_{i_1=1}^{[na]} \sum_{i_2=1}^{[nb]} \sum_{i_3=1}^{[nc]} \sum_{i_4=1}^{[nd]} |\rho(a - i_1)| |\rho(i_2 - b)| |\rho(i_3 - i_4)| \leq nd \left(\sum_{r \in \mathbb{Z}} |\rho(r)| \right)^3 = Cn$$

on the other hand (still using Remark 2.1), we deduce that the terms of the third type in $A_2^{(n)}$ also agree with the desired conclusion (3.3).

(3) Using exactly the same strategy as in point (2), we can show as well that the terms $A_3^{(n)}$ agree with the desired conclusion (3.3). Details are left to the reader.

(4) Finally, let us focus on the last term, that is $A_4^{(n)}$. We have, using successively the fact that $\sum_{r \in \mathbb{Z}} |\rho(r)|^3 < \infty$ and Lemma 3.3,

$$\begin{aligned} A_4^{(n)} &= \sum_{i_1=1}^{[na]} \sum_{i_2=1}^{[nb]} \sum_{i_3=1}^{[nc]} \sum_{i_4=1}^{[nd]} \left| E \left(\Phi(i_1, i_2, i_3, i_4) I_3(\delta_{i_1}^{\otimes 3}) I_3(\delta_{i_2}^{\otimes 3}) \right) \right| \left| \langle \delta_{i_3}, \delta_{i_4} \rangle_{\mathfrak{H}} \right|^3 \\ &= n^{-1} \sum_{i_1=1}^{[na]} \sum_{i_2=1}^{[nb]} \sum_{i_3=1}^{[nc]} \sum_{i_4=1}^{[nd]} \left| E \left(\Phi(i_1, i_2, i_3, i_4) I_3(\delta_{i_1}^{\otimes 3}) I_3(\delta_{i_2}^{\otimes 3}) \right) \right| |\rho(i_3 - i_4)|^3 \\ &\leq C \sup_{1 \leq i_3 \leq [nc]} \sup_{1 \leq i_4 \leq [nc]} \sum_{i_1=1}^{[na]} \sum_{i_2=1}^{[nb]} \left| E \left(\Phi(i_1, i_2, i_3, i_4) I_3(\delta_{i_1}^{\otimes 3}) I_3(\delta_{i_2}^{\otimes 3}) \right) \right| \leq C. \end{aligned}$$

Hence, the terms $A_4^{(n)}$ agree with the desired conclusion (3.3) and the proof of Lemma 3.5 is now complete. \square

Lemma 3.6. Let $\lambda = (\lambda_1, \dots, \lambda_m) \in \mathbb{R}^m$, $u_1, \dots, u_m > 0$, $u_p > 0$ and suppose $g_1, \dots, g_m \in C^\infty(\mathbb{R})$ are bounded with bounded derivatives. Define $\mathbb{V}_n \in \mathbb{R}^m$ by

$$\mathbb{V}_n := \left(\sum_{i=1}^{[nu_k]} g_k(B(t_{i-1})) I_3(\delta_i^{\otimes 3}) \right)_{k=1, \dots, m},$$

so that

$$\langle \lambda, \mathbb{V}_n \rangle := \sum_{k=1}^m \lambda_k \sum_{i=1}^{\lfloor nu_k \rfloor} g_k(B(t_{i-1})) I_3(\delta_i^{\otimes 3}) \quad (\text{see (3.15) below}). \quad (3.4)$$

Then there exists $C > 0$, independent of n , such that

$$\sup_{j=1, \dots, \lfloor nu_p \rfloor} E(\langle D\langle \lambda, \mathbb{V}_n \rangle, \delta_j \rangle_{\mathfrak{H}}^2) \leq Cn^{-2/3} \quad (3.5)$$

$$\sum_{j=1}^{\lfloor nu_p \rfloor} E(\langle D\langle \lambda, \mathbb{V}_n \rangle, \delta_j \rangle_{\mathfrak{H}}^2) \leq Cn^{-1/3} \quad (3.6)$$

$$\sum_{j=1}^{\lfloor nu_p \rfloor} E(\langle D^2\langle \lambda, \mathbb{V}_n \rangle, \delta_j^{\otimes 2} \rangle_{\mathfrak{H}^{\otimes 2}}^2) \leq Cn^{-2/3}. \quad (3.7)$$

Proof. We have

$$\begin{aligned} \langle D\langle \lambda, \mathbb{V}_n \rangle, \delta_j \rangle_{\mathfrak{H}} &= \sum_{k=1}^m \lambda_k \sum_{i=1}^{\lfloor nu_k \rfloor} g'_k(B(t_{i-1})) I_3(\delta_i^{\otimes 3}) \langle \varepsilon_{i-1}, \delta_j \rangle_{\mathfrak{H}} \\ &\quad + 3 \sum_{k=1}^m \lambda_k \sum_{i=1}^{\lfloor nu_k \rfloor} g_k(B(t_{i-1})) I_2(\delta_i^{\otimes 2}) \langle \delta_i, \delta_j \rangle_{\mathfrak{H}}. \end{aligned} \quad (3.8)$$

Hence, with ρ defined by (2.4),

$$\begin{aligned} &E(\langle D\langle \lambda, \mathbb{V}_n \rangle, \delta_j \rangle_{\mathfrak{H}}^2) \\ &\leq 2m \sum_{k=1}^m \lambda_k^2 \sum_{i, \ell=1}^{\lfloor nu_k \rfloor} |E(g'_k(B(t_{i-1})) g'_k(B(t_{\ell-1})) I_3(\delta_i^{\otimes 3}) I_3(\delta_\ell^{\otimes 3}))| |\langle \varepsilon_{i-1}, \delta_j \rangle_{\mathfrak{H}}| |\langle \varepsilon_{\ell-1}, \delta_j \rangle_{\mathfrak{H}}| \\ &\quad + 18m \sum_{k=1}^m \lambda_k^2 \sum_{i, \ell=1}^{\lfloor nu_k \rfloor} |E(g_k(B(t_{i-1})) g_k(B(t_{\ell-1})) I_2(\delta_i^{\otimes 2}) I_2(\delta_\ell^{\otimes 2}))| |\langle \delta_i, \delta_j \rangle_{\mathfrak{H}}| |\langle \delta_\ell, \delta_j \rangle_{\mathfrak{H}}| \\ &\leq Cn^{-2/3} \sup_{k=1, \dots, m} \sum_{i, \ell=1}^{\lfloor nu_k \rfloor} |E(g'_k(B(t_{i-1})) g'_k(B(t_{\ell-1})) I_3(\delta_i^{\otimes 3}) I_3(\delta_\ell^{\otimes 3}))| \\ &\quad + Cn^{-4/3} \sum_{i, \ell=1}^{\lfloor nu_k \rfloor} |\rho(i-j)| |\rho(\ell-j)| \quad \text{by Lemma 3.2 (i) and Cauchy-Schwarz} \\ &\leq Cn^{-2/3} + Cn^{-4/3} \left(\sum_{r \in \mathbb{Z}} |\rho(r)| \right)^2 \quad \text{by Lemma 3.3 and Remark 2.1} \\ &\leq Cn^{-2/3}, \end{aligned}$$

which is (3.5). Moreover, combining the first inequality of the previous estimate with Lemma 3.2

(ii) and Lemma 3.3, we also have

$$\begin{aligned} & \sum_{j=1}^{\lfloor nu_p \rfloor} E(\langle D\langle \lambda, \mathbb{V}_n \rangle, \delta_j \rangle_{\mathfrak{H}}^2) \\ & \leq Cn^{-1/3} \sup_{k=1, \dots, m} \sum_{i, \ell=1}^{\lfloor nu_k \rfloor} |E(g'_k(B(t_{i-1}))g'_\ell(B(t_{\ell-1}))I_3(\delta_i^{\otimes 3})I_3(\delta_\ell^{\otimes 3}))| \\ & \quad \times \sup_{i=1, \dots, \lfloor nu_k \rfloor} \sum_{j=1}^{\lfloor nu_p \rfloor} |\langle \varepsilon_{i-1}, \delta_j \rangle_{\mathfrak{H}}| + Cn^{-1/3} \left(\sum_{r \in \mathbb{Z}} |\rho(r)| \right)^2 \leq Cn^{-1/3}, \end{aligned}$$

which is (3.6). The proof of (3.7) follows the same lines, and is left to the reader. \square

3.2 Proof of Theorem 3.1

We are now in position to prove Theorem 3.1. For $g : \mathbb{R} \rightarrow \mathbb{R}$, let

$$G_n^-(g, B, t) := \frac{1}{\sqrt{n}} \sum_{j=1}^{\lfloor nt \rfloor} g(B(t_{j-1}))h_3(n^{1/6}\Delta B_j), \quad t \geq 0, \quad n \geq 1.$$

We recall that $h_3(x) = x^3 - 3x$, see (2.1), and the definition (2.3) of $V_n(B, t)$. In particular, observe that

$$V_n(B, t) = G_n^-(1, B, t) + 3n^{-1/3}B(\lfloor nt \rfloor/n). \quad (3.9)$$

Our main theorem which will lead us toward the proof of Theorem 3.1 is the following.

Theorem 3.7. *If $g \in C^\infty(\mathbb{R})$ is bounded with bounded derivatives, then the sequence $(B, G_n^-(1, B), G_n^-(g, B))$ converges to $(B, [[B]], -(1/8) \int g'''(B) ds + \int g(B) d[[B]])$ in the sense of finite-dimensional distributions on $[0, \infty)$.*

Proof. We have to prove that, for any $\ell + m \geq 1$ and any $u_1, \dots, u_{\ell+m} \geq 0$:

$$\begin{aligned} & (B, G_n^-(1, B, u_1), \dots, G_n^-(1, B, u_\ell), G_n^-(g, B, u_{\ell+1}), \dots, G_n^-(g, B, u_{\ell+m})) \\ & \xrightarrow[n \rightarrow \infty]{\text{Law}} \left(B, [[B]]_{u_1}, \dots, [[B]]_{u_\ell}, -\frac{1}{8} \int_0^{u_{\ell+1}} g'''(B(s)) ds + \int_0^{u_{\ell+1}} g(B(s)) d[[B]]_s, \dots, \right. \\ & \quad \left. -\frac{1}{8} \int_0^{u_{\ell+m}} g'''(B(s)) ds + \int_0^{u_{\ell+m}} g(B(s)) d[[B]]_s \right). \end{aligned}$$

Actually, we will prove the following slightly stronger convergence. For any $m \geq 1$, any $u_1, \dots, u_m \geq 0$ and all bounded functions $g_1, \dots, g_m \in C^\infty(\mathbb{R})$ with bounded derivatives, we have

$$\begin{aligned} & (B, G_n^-(g_1, B, u_1), \dots, G_n^-(g_m, B, u_m)) \\ & \xrightarrow[n \rightarrow \infty]{\text{Law}} \left(B, -\frac{1}{8} \int_0^{u_1} g_1'''(B(s)) ds + \int_0^{u_1} g_1(B(s)) d[[B]]_s, \dots, \right. \\ & \quad \left. -\frac{1}{8} \int_0^{u_m} g_m'''(B(s)) ds + \int_0^{u_m} g_m(B(s)) d[[B]]_s \right). \quad (3.10) \end{aligned}$$

Using (2.12), observe that

$$G_n^-(g, B, t) = \sum_{j=1}^{\lfloor nt \rfloor} g(B(t_{j-1})) I_3(\delta_j^{\otimes 3}). \quad (3.11)$$

The proof of (3.10) is divided into several steps, and follows the methodology introduced in [7].

Step 1.- We first prove that:

$$\begin{aligned} & \lim_{n \rightarrow \infty} E \left(G_n^-(g_1, B, u_1), \dots, G_n^-(g_m, B, u_m) \right) \\ &= \left(-\frac{1}{8} \int_0^{u_1} E(g_1'''(B(s))) ds, \dots, -\frac{1}{8} \int_0^{u_m} E(g_m'''(B(s))) ds \right), \\ & \lim_{n \rightarrow \infty} E \left(\left\| (G_n^-(g_1, B, u_1), \dots, G_n^-(g_m, B, u_m)) \right\|_{\mathbb{R}^m}^2 \right) \\ &= \sum_{i=1}^m \left(\kappa^2 \int_0^{u_i} E(g_i^2(B(s))) ds + \frac{1}{64} E \left(\int_0^{u_i} g_i'''(B(s)) ds \right)^2 \right). \end{aligned} \quad (3.12)$$

For g as in the statement of the theorem, we can write, for any fixed $t \geq 0$:

$$\begin{aligned} E \left(G_n^-(g, B, t) \right) &= \sum_{j=1}^{\lfloor nt \rfloor} E \left(g(B(t_{j-1})) I_3(\delta_j^{\otimes 3}) \right) \quad \text{by (3.11)} \\ &= \sum_{j=1}^{\lfloor nt \rfloor} E \left(\langle D^3 g(B(t_{j-1})), \delta_j^{\otimes 3} \rangle_{\mathfrak{H}^{\otimes 3}} \right) \quad \text{by (2.13)} \\ &= \sum_{j=1}^{\lfloor nt \rfloor} E(g'''(B(t_{j-1}))) \langle \varepsilon_{j-1}, \delta_j \rangle_{\mathfrak{H}}^3 \quad \text{by (2.8)} \\ &= -\frac{1}{8n} \sum_{j=1}^{\lfloor nt \rfloor} E(g'''(B(t_{j-1}))) + \sum_{j=1}^{\lfloor nt \rfloor} E(g'''(B(t_{j-1}))) \left(\langle \varepsilon_{j-1}, \delta_j \rangle_{\mathfrak{H}}^3 + \frac{1}{8n} \right) \\ &\xrightarrow{n \rightarrow \infty} -\frac{1}{8} \int_0^t E(g'''(B(s))) ds \quad \text{by Lemma 3.2 (iv)}. \end{aligned}$$

Now, let us turn to the second part of (3.12). We have

$$E \left\| (G_n^-(g_1, B, u_1), \dots, G_n^-(g_m, B, u_m)) \right\|_{\mathbb{R}^m}^2 = \sum_{i=1}^m E(G_n^-(g_i, B, u_i)^2).$$

By the product formula (2.14), we have

$$\begin{aligned} I_3(\delta_j^{\otimes 3}) I_3(\delta_k^{\otimes 3}) &= I_6(\delta_j^{\otimes 3} \otimes \delta_k^{\otimes 3}) + 9I_4(\delta_j^{\otimes 2} \otimes \delta_k^{\otimes 2}) \langle \delta_j, \delta_k \rangle_{\mathfrak{H}} \\ &\quad + 18I_2(\delta_j \otimes \delta_k) \langle \delta_j, \delta_k \rangle_{\mathfrak{H}}^2 + 6 \langle \delta_j, \delta_k \rangle_{\mathfrak{H}}^3. \end{aligned}$$

Thus, for any fixed $t \geq 0$,

$$\begin{aligned}
E(G_n^-(g, B, t)^2) &= \sum_{j,k=1}^{\lfloor nt \rfloor} E \left(g(B(t_{j-1}))g(B(t_{k-1}))I_3(\delta_j^{\otimes 3})I_3(\delta_k^{\otimes 3}) \right) \\
&= \sum_{j,k=1}^{\lfloor nt \rfloor} E \left(g(B(t_{j-1}))g(B(t_{k-1}))I_6(\delta_j^{\otimes 3} \otimes \delta_k^{\otimes 3}) \right) \\
&\quad + 9 \sum_{j,k=1}^{\lfloor nt \rfloor} E \left(g(B(t_{j-1}))g(B(t_{k-1}))I_4(\delta_j^{\otimes 2} \otimes \delta_k^{\otimes 2}) \right) \langle \delta_j, \delta_k \rangle_{\mathfrak{H}} \\
&\quad + 18 \sum_{j,k=1}^{\lfloor nt \rfloor} E \left(g(B(t_{j-1}))g(B(t_{k-1}))I_2(\delta_j \otimes \delta_k) \right) \langle \delta_j, \delta_k \rangle_{\mathfrak{H}}^2 \\
&\quad + 6 \sum_{j,k=1}^{\lfloor nt \rfloor} E \left(g(B(t_{j-1}))g(B(t_{k-1})) \right) \langle \delta_j, \delta_k \rangle_{\mathfrak{H}}^3 \\
&=: A_n + B_n + C_n + D_n.
\end{aligned}$$

We will estimate each of these four terms using the Malliavin integration by parts formula (2.13). For that purpose, we use Lemma 3.4 and the notation of Remark 2.6.

First, we have

$$\begin{aligned}
A_n &= \sum_{j,k=1}^{\lfloor nt \rfloor} E \left(\langle D^6[g(B(t_{j-1}))g(B(t_{k-1}))], \delta_j^{\otimes 3} \otimes \delta_k^{\otimes 3} \rangle_{\mathfrak{H}^{\otimes 6}} \right) \\
&\stackrel{(3.2)}{=} \sum_{j,k=1}^{\lfloor nt \rfloor} \sum_{a=0}^6 \binom{6}{a} E \left(g^{(a)}(B(t_{j-1}))g^{(6-a)}(B(t_{k-1})) \right) \langle \varepsilon_{j-1}^{\otimes a} \otimes \varepsilon_{k-1}^{\otimes (6-a)}, \delta_j^{\otimes 3} \otimes \delta_k^{\otimes 3} \rangle_{\mathfrak{H}^{\otimes 6}} \\
&\stackrel{(3.1)}{=} \sum_{j,k=1}^{\lfloor nt \rfloor} \sum_{a=0}^6 E \left(g^{(a)}(B(t_{j-1}))g^{(6-a)}(B(t_{k-1})) \right) \\
&\quad \times \sum_{\substack{i_1, \dots, i_6 \in \{j-1, k-1\} \\ |\{\ell : i_\ell = j-1\}| = a}} \langle \varepsilon_{i_1} \otimes \dots \otimes \varepsilon_{i_6}, \delta_j^{\otimes 3} \otimes \delta_k^{\otimes 3} \rangle_{\mathfrak{H}^{\otimes 6}}.
\end{aligned}$$

Actually, in the previous double sum with respect to a and i_1, \dots, i_6 , only the following term is

non-negligible:

$$\begin{aligned}
& \sum_{j,k=1}^{\lfloor nt \rfloor} E(g'''(B(t_{j-1}))g'''(B(t_{k-1})))\langle \varepsilon_{j-1}, \delta_j \rangle_{\mathfrak{H}}^3 \langle \varepsilon_{k-1}, \delta_k \rangle_{\mathfrak{H}}^3 \\
&= E \left(\sum_{j=1}^{\lfloor nt \rfloor} g'''(B(t_{j-1}))\langle \varepsilon_{j-1}, \delta_j \rangle_{\mathfrak{H}}^3 \right)^2 \\
&= E \left(-\frac{1}{8n} \sum_{j=1}^{\lfloor nt \rfloor} g'''(B(t_{j-1})) + \sum_{j=1}^{\lfloor nt \rfloor} g'''(B(t_{j-1})) \left(\langle \varepsilon_{j-1}, \delta_j \rangle_{\mathfrak{H}}^3 + \frac{1}{8n} \right) \right)^2 \\
&\xrightarrow{n \rightarrow \infty} \frac{1}{64} E \left(\int_0^t g'''(B(s)) ds \right)^2 \quad \text{by Lemma 3.2 (iv)}.
\end{aligned}$$

Indeed, the other terms in A_n are all of the form

$$\sum_{j,k=1}^{\lfloor nt \rfloor} E(g^{(a)}(B(t_{j-1}))g^{(6-a)}(B(t_{k-1})))\langle \varepsilon_{j-1}, \delta_k \rangle_{\mathfrak{H}} \prod_{i=1}^5 \langle \varepsilon_{x_i-1}, \delta_{y_i} \rangle_{\mathfrak{H}}, \quad (3.13)$$

where x_i and y_i are for j or k . By Lemma 3.2 (iii), we have $\sum_{j,k=1}^{\lfloor nt \rfloor} |\langle \varepsilon_{j-1}, \delta_k \rangle_{\mathfrak{H}}| = O(n)$ as $n \rightarrow \infty$. By Lemma 3.2 (i), $\sup_{j,k=1, \dots, \lfloor nt \rfloor} \prod_{i=1}^5 |\langle \varepsilon_{x_i-1}, \delta_{y_i} \rangle_{\mathfrak{H}}| = O(n^{-5/3})$ as $n \rightarrow \infty$. Hence, the quantity in (3.13) tends to zero as $n \rightarrow \infty$. We have proved

$$A_n \xrightarrow{n \rightarrow \infty} \frac{1}{64} E \left(\int_0^t g'''(B(s)) ds \right)^2.$$

Using the integration by parts formula (2.13) as well as Lemma 3.4, we have similarly that

$$\begin{aligned}
|B_n| &\leq 9 \sum_{j,k=1}^{\lfloor nt \rfloor} \sum_{a=0}^4 \binom{4}{a} |E(g^{(a)}(B(t_{j-1}))g^{(4-a)}(B(t_{k-1})))\langle \varepsilon_{j-1}^{\otimes a} \otimes \varepsilon_{k-1}^{\otimes(4-a)}, \delta_j^{\otimes 2} \otimes \delta_k^{\otimes 2} \rangle_{\mathfrak{H}^{\otimes 4}} \langle \delta_j, \delta_k \rangle_{\mathfrak{H}}| \\
&\leq Cn^{-4/3} \sum_{j,k=1}^{\lfloor nt \rfloor} |\langle \delta_j, \delta_k \rangle_{\mathfrak{H}}| \quad \text{by Lemma 3.2 (i)} \\
&= Cn^{-5/3} \sum_{j,k=1}^{\lfloor nt \rfloor} |\rho(j-k)| \leq Cn^{-2/3} \sum_{r \in \mathbb{Z}} |\rho(r)| = Cn^{-2/3} \xrightarrow{n \rightarrow \infty} 0,
\end{aligned}$$

with ρ defined by (2.4).

Using similar computations, we also have

$$|C_n| \leq Cn^{-1/3} \sum_{r=-\infty}^{\infty} \rho^2(r) = Cn^{-1/3} \xrightarrow{n \rightarrow \infty} 0,$$

while

$$\begin{aligned}
D_n &= \frac{6}{n} \sum_{j,k=1}^{\lfloor nt \rfloor} E(g(B(t_{j-1}))g(B(t_{k-1})))\rho^3(j-k) \\
&= \frac{6}{n} \sum_{r \in \mathbb{Z}} \sum_{j=1 \vee (1-r)}^{\lfloor nt \rfloor \wedge (\lfloor nt \rfloor - r)} E(g(B(t_{j-1}))g(B(t_{j+r-1})))\rho^3(r) \\
&\xrightarrow{n \rightarrow \infty} 6 \sum_{r \in \mathbb{Z}} \rho^3(r) \int_0^t E(g^2(B(s))) ds = \kappa^2 \int_0^t E(g^2(B(s))) ds,
\end{aligned}$$

the previous convergence being obtained as in the proof of (3.27) below. Finally, we have obtained

$$E(G_n^-(g, B, t)^2) \xrightarrow{n \rightarrow \infty} \kappa^2 \int_0^t E(g^2(B(s))) ds + \frac{1}{64} E \left(\int_0^t g'''(B(s)) ds \right)^2, \quad (3.14)$$

and the proof of (3.12) is done.

Step 2.- By Step 1, the sequence $(B, G_n^-(g_1, B, u_1), \dots, G_n^-(g_m, B, u_m))$ is tight in $D_{\mathbb{R}}[0, \infty) \times \mathbb{R}^m$. Consider a subsequence converging in law to some limit denoted by

$$(B, G_{\infty}^-(g_1, B, u_1), \dots, G_{\infty}^-(g_m, B, u_m))$$

(for convenience, we keep the same notation for this subsequence and for the sequence itself). Recall \mathbb{V}_n , defined in Lemma 3.6, and note that by (3.11), we have

$$\mathbb{V}_n := (G_n^-(g_1, B, u_1), \dots, G_n^-(g_m, B, u_m)), \quad n \in \mathbb{N} \cup \{\infty\}. \quad (3.15)$$

Let us also define

$$\begin{aligned}
\mathbb{W} := & \left(-\frac{1}{8} \int_0^{u_1} g_1'''(B(s)) ds + \int_0^{u_1} g_1(B(s)) d[[B]]_s, \dots, \right. \\
& \left. -\frac{1}{8} \int_0^{u_m} g_m'''(B(s)) ds + \int_0^{u_m} g_m(B(s)) d[[B]]_s \right).
\end{aligned}$$

We have to show that, conditioned on B , the laws of \mathbb{V}_{∞} and \mathbb{W} are the same.

Let $\lambda = (\lambda_1, \dots, \lambda_m)$ denote a generic element of \mathbb{R}^m and, for $\lambda, \mu \in \mathbb{R}^m$, write $\langle \lambda, \mu \rangle$ for $\sum_{i=1}^m \lambda_i \mu_i$. We consider the conditional characteristic function of \mathbb{W} given B :

$$\Phi(\lambda) := E \left(e^{i \langle \lambda, \mathbb{W} \rangle} \mid B \right). \quad (3.16)$$

Recall that $[[B]] = \kappa W$, where W is a Brownian motion independent of B . Hence, if we condition on B , then \mathbb{W} has the same law as $\mu + N$, where $\mu \in \mathbb{R}^m$ has components $\mu_k := -(1/8) \int_0^{u_k} g_k'''(B(s)) ds$, and N is a centered Gaussian random vector in \mathbb{R}^m with covariance matrix $Q = (q_{ij})$ given by

$$q_{ij} := \kappa^2 \int_0^{u_i \wedge u_j} g_i(B(s)) g_j(B(s)) ds.$$

Thus, $\Phi(\lambda) = e^{i\langle \lambda, \mu \rangle - \frac{1}{2}\langle \lambda, Q\lambda \rangle}$. The point is that Φ is the unique solution of the following system of PDEs (see [12]):

$$\frac{\partial \varphi}{\partial \lambda_p}(\lambda) = \varphi(\lambda) \left(i\mu_p - \sum_{k=1}^m \lambda_k q_{pk} \right), \quad p = 1, \dots, m, \quad (3.17)$$

where the unknown function $\varphi : \mathbb{R}^m \rightarrow \mathbb{C}$ satisfies the initial condition $\varphi(0) = 1$. Hence, we have to show that, for every random variable ξ of the form $\psi(B(s_1), \dots, B(s_r))$, with $\psi : \mathbb{R}^r \rightarrow \mathbb{R}$ belonging to $C_b^\infty(\mathbb{R}^r)$ and $s_1, \dots, s_r \geq 0$, we have

$$\begin{aligned} \frac{\partial}{\partial \lambda_p} E \left(e^{i\langle \lambda, \mathbb{V}_\infty \rangle} \xi \right) &= -\frac{i}{8} \int_0^{u_p} E \left(g_p'''(B(s)) \xi e^{i\langle \lambda, \mathbb{V}_\infty \rangle} \right) ds \\ &\quad - \kappa^2 \sum_{k=1}^m \lambda_k \int_0^{u_p \wedge u_k} E \left(g_p(B(s)) g_k(B(s)) \xi e^{i\langle \lambda, \mathbb{V}_\infty \rangle} \right) ds \end{aligned} \quad (3.18)$$

for all $p \in \{1, \dots, m\}$.

Step 3.- Since (\mathbb{V}_∞, B) is defined as the limit in law of (\mathbb{V}_n, B) on one hand, and \mathbb{V}_n is bounded in L^2 on the other hand, note that

$$\frac{\partial}{\partial \lambda_p} E \left(e^{i\langle \lambda, \mathbb{V}_\infty \rangle} \xi \right) = \lim_{n \rightarrow \infty} \frac{\partial}{\partial \lambda_p} E \left(e^{i\langle \lambda, \mathbb{V}_n \rangle} \xi \right).$$

Let us compute $\frac{\partial}{\partial \lambda_p} E \left(e^{i\langle \lambda, \mathbb{V}_n \rangle} \xi \right)$. We have

$$\frac{\partial}{\partial \lambda_p} E \left(e^{i\langle \lambda, \mathbb{V}_n \rangle} \xi \right) = i E \left(G_n^-(g_p, B, u_p) e^{i\langle \lambda, \mathbb{V}_n \rangle} \xi \right). \quad (3.19)$$

Moreover, see (3.11) and use (2.13), for any $t \geq 0$:

$$\begin{aligned} E \left(G_n^-(g, B, t) e^{i\langle \lambda, \mathbb{V}_n \rangle} \xi \right) &= \sum_{j=1}^{\lfloor nt \rfloor} E \left(g(B(t_{j-1})) I_3(\delta_j^{\otimes 3}) e^{i\langle \lambda, \mathbb{V}_n \rangle} \xi \right) \\ &= \sum_{j=1}^{\lfloor nt \rfloor} E \left(\langle D^3(g(B(t_{j-1})) e^{i\langle \lambda, \mathbb{V}_n \rangle} \xi), \delta_j^{\otimes 3} \rangle_{\mathfrak{H}^{\otimes 3}} \right). \end{aligned} \quad (3.20)$$

The first three Malliavin derivatives of $g(B(t_{j-1})) e^{i\langle \lambda, \mathbb{V}_n \rangle} \xi$ are respectively given by

$$\begin{aligned} &D(g(B(t_{j-1})) e^{i\langle \lambda, \mathbb{V}_n \rangle} \xi) \\ &= g'(B(t_{j-1})) e^{i\langle \lambda, \mathbb{V}_n \rangle} \xi \varepsilon_{j-1} + i g(B(t_{j-1})) e^{i\langle \lambda, \mathbb{V}_n \rangle} \xi D\langle \lambda, \mathbb{V}_n \rangle \\ &\quad + g(B(t_{j-1})) e^{i\langle \lambda, \mathbb{V}_n \rangle} D\xi, \end{aligned}$$

$$\begin{aligned} &D^2(g(B(t_{j-1})) e^{i\langle \lambda, \mathbb{V}_n \rangle} \xi) \\ &= g''(B(t_{j-1})) e^{i\langle \lambda, \mathbb{V}_n \rangle} \xi \varepsilon_{j-1}^{\otimes 2} + 2i g'(B(t_{j-1})) e^{i\langle \lambda, \mathbb{V}_n \rangle} \xi D\langle \lambda, \mathbb{V}_n \rangle \otimes \varepsilon_{j-1} \\ &\quad + 2g'(B(t_{j-1})) e^{i\langle \lambda, \mathbb{V}_n \rangle} D\xi \otimes \varepsilon_{j-1} - g(B(t_{j-1})) e^{i\langle \lambda, \mathbb{V}_n \rangle} \xi D\langle \lambda, \mathbb{V}_n \rangle^{\otimes 2} \\ &\quad + 2i g(B(t_{j-1})) e^{i\langle \lambda, \mathbb{V}_n \rangle} D\xi \otimes D\langle \lambda, \mathbb{V}_n \rangle + i g(B(t_{j-1})) e^{i\langle \lambda, \mathbb{V}_n \rangle} \xi D^2\langle \lambda, \mathbb{V}_n \rangle \\ &\quad + g(B(t_{j-1})) e^{i\langle \lambda, \mathbb{V}_n \rangle} D^2\xi, \end{aligned}$$

and

$$\begin{aligned}
& D^3(g(B(t_{j-1}))e^{i\langle\lambda, \mathbb{V}_n\rangle}\xi) \\
&= g'''(B(t_{j-1}))e^{i\langle\lambda, \mathbb{V}_n\rangle}\xi \varepsilon_{j-1}^{\otimes 3} + 3ig''(B(t_{j-1}))e^{i\langle\lambda, \mathbb{V}_n\rangle}\xi \varepsilon_{j-1}^{\otimes 2} \otimes D\langle\lambda, \mathbb{V}_n\rangle \\
&\quad + 3g''(B(t_{j-1}))e^{i\langle\lambda, \mathbb{V}_n\rangle} \varepsilon_{j-1}^{\otimes 2} \otimes D\xi - 3g'(B(t_{j-1}))e^{i\langle\lambda, \mathbb{V}_n\rangle}\xi D\langle\lambda, \mathbb{V}_n\rangle^{\otimes 2} \otimes \varepsilon_{j-1} \\
&\quad + 6ig'(B(t_{j-1}))e^{i\langle\lambda, \mathbb{V}_n\rangle} D\xi \otimes D\langle\lambda, \mathbb{V}_n\rangle \otimes \varepsilon_{j-1} \\
&\quad - ig(B(t_{j-1}))e^{i\langle\lambda, \mathbb{V}_n\rangle}\xi D\langle\lambda, \mathbb{V}_n\rangle^{\otimes 3} - 3g(B(t_{j-1}))e^{i\langle\lambda, \mathbb{V}_n\rangle} D\langle\lambda, \mathbb{V}_n\rangle^{\otimes 2} \otimes D\xi \\
&\quad + ig(B(t_{j-1}))e^{i\langle\lambda, \mathbb{V}_n\rangle}\xi D^3\langle\lambda, \mathbb{V}_n\rangle + g(B(t_{j-1}))e^{i\langle\lambda, \mathbb{V}_n\rangle} D^3\xi \\
&\quad + 3ig(B(t_{j-1}))e^{i\langle\lambda, \mathbb{V}_n\rangle} D^2\xi \otimes D\langle\lambda, \mathbb{V}_n\rangle + 3ig(B(t_{j-1}))e^{i\langle\lambda, \mathbb{V}_n\rangle} D\xi \otimes D^2\langle\lambda, \mathbb{V}_n\rangle \\
&\quad + 3g'(B(t_{j-1}))e^{i\langle\lambda, \mathbb{V}_n\rangle} D^2\xi \otimes \varepsilon_{j-1} + 3ig'(B(t_{j-1}))e^{i\langle\lambda, \mathbb{V}_n\rangle}\xi \varepsilon_{j-1} \otimes D^2\langle\lambda, \mathbb{V}_n\rangle \\
&\quad - 3g(B(t_{j-1}))e^{i\langle\lambda, \mathbb{V}_n\rangle}\xi D\langle\lambda, \mathbb{V}_n\rangle \otimes D^2\langle\lambda, \mathbb{V}_n\rangle.
\end{aligned} \tag{3.21}$$

Let us compute the term $D^3\langle\lambda, \mathbb{V}_n\rangle$. Recall that

$$\langle\lambda, \mathbb{V}_n\rangle = \sum_{k=1}^m \lambda_k G_n^-(g_k, B, u_k) = \sum_{k=1}^m \lambda_k \sum_{\ell=1}^{\lfloor nu_k \rfloor} g_k(B(t_{\ell-1})) I_3(\delta_\ell^{\otimes 3}).$$

Combining the Leibniz rule (2.15) with $D(I_q(f^{\otimes q})) = qI_{q-1}(f^{\otimes(q-1)})f$ for any $f \in \mathfrak{H}$, we have

$$\begin{aligned}
D^3\langle\lambda, \mathbb{V}_n\rangle = \sum_{k=1}^m \lambda_k \sum_{\ell=1}^{\lfloor nu_k \rfloor} \left[g_k'''(B(t_{\ell-1})) I_3(\delta_\ell^{\otimes 3}) \varepsilon_{\ell-1}^{\otimes 3} + 9g_k''(B(t_{\ell-1})) I_2(\delta_\ell^{\otimes 2}) \varepsilon_{\ell-1}^{\otimes 2} \otimes \delta_\ell \right. \\
\left. + 18g_k'(B(t_{\ell-1})) I_1(\delta_\ell) \varepsilon_{\ell-1} \otimes \delta_\ell^{\otimes 2} + 6g_k(B(t_{\ell-1})) \delta_\ell^{\otimes 3} \right].
\end{aligned} \tag{3.22}$$

Combining relations (3.19), (3.20), (3.21), and (3.22) we obtain the following expression:

$$\begin{aligned}
\frac{\partial}{\partial \lambda_p} E \left(e^{i\langle\lambda, \mathbb{V}_n\rangle} \xi \right) &= iE \left(e^{i\langle\lambda, \mathbb{V}_n\rangle} \xi \sum_{j=1}^{\lfloor nu_p \rfloor} g_p'''(B(t_{j-1})) \langle \varepsilon_{j-1}, \delta_j \rangle_{\mathfrak{H}}^3 \right) \\
&\quad - 6E \left(e^{i\langle\lambda, \mathbb{V}_n\rangle} \xi \sum_{j=1}^{\lfloor nu_p \rfloor} \sum_{k=1}^m \lambda_k \sum_{\ell=1}^{\lfloor nu_k \rfloor} g_p(B(t_{j-1})) g_k(B(t_{\ell-1})) \langle \delta_\ell, \delta_j \rangle_{\mathfrak{H}}^3 \right) + i \sum_{j=1}^{\lfloor nu_p \rfloor} r_{j,n},
\end{aligned} \tag{3.23}$$

with

$$\begin{aligned}
r_{j,n} &= i \sum_{k=1}^m \lambda_k \sum_{\ell=1}^{\lfloor nu_k \rfloor} E \left(g_p(B(t_{j-1})) g_k'''(B(t_{\ell-1})) I_3(\delta_\ell^{\otimes 3}) e^{i\langle\lambda, \mathbb{V}_n\rangle} \xi \right) \langle \varepsilon_{\ell-1}, \delta_j \rangle_{\mathfrak{H}}^3 \\
&\quad + 9i \sum_{k=1}^m \lambda_k \sum_{\ell=1}^{\lfloor nu_k \rfloor} E \left(g_p(B(t_{j-1})) g_k''(B(t_{\ell-1})) I_2(\delta_\ell^{\otimes 2}) e^{i\langle\lambda, \mathbb{V}_n\rangle} \xi \right) \langle \varepsilon_{\ell-1}, \delta_j \rangle_{\mathfrak{H}}^2 \langle \delta_\ell, \delta_j \rangle_{\mathfrak{H}} \\
&\quad + 18i \sum_{k=1}^m \lambda_k \sum_{\ell=1}^{\lfloor nu_k \rfloor} E \left(g_p(B(t_{j-1})) g_k'(B(t_{\ell-1})) I_1(\delta_\ell) e^{i\langle\lambda, \mathbb{V}_n\rangle} \xi \right) \langle \delta_\ell, \delta_j \rangle_{\mathfrak{H}}^2 \langle \varepsilon_{\ell-1}, \delta_j \rangle_{\mathfrak{H}}
\end{aligned}$$

$$\begin{aligned}
& + 3iE \left(g_p''(B(t_{j-1}))e^{i\langle \lambda, \mathbb{V}_n \rangle} \xi \langle D\langle \lambda, \mathbb{V}_n \rangle, \delta_j \rangle_{\mathfrak{H}} \right) \langle \varepsilon_{j-1}, \delta_j \rangle_{\mathfrak{H}}^2 \\
& + 3E \left(g_p''(B(t_{j-1}))e^{i\langle \lambda, \mathbb{V}_n \rangle} \langle D\xi, \delta_j \rangle_{\mathfrak{H}} \right) \langle \varepsilon_{j-1}, \delta_j \rangle_{\mathfrak{H}}^2 \\
& - 3E \left(g_p'(B(t_{j-1}))e^{i\langle \lambda, \mathbb{V}_n \rangle} \xi \langle D\langle \lambda, \mathbb{V}_n \rangle, \delta_j \rangle_{\mathfrak{H}}^2 \right) \langle \varepsilon_{j-1}, \delta_j \rangle_{\mathfrak{H}} \\
& + 6iE \left(g_p'(B(t_{j-1}))e^{i\langle \lambda, \mathbb{V}_n \rangle} \langle D\langle \lambda, \mathbb{V}_n \rangle, \delta_j \rangle_{\mathfrak{H}} \langle D\xi, \delta_j \rangle_{\mathfrak{H}} \right) \langle \varepsilon_{j-1}, \delta_j \rangle_{\mathfrak{H}} \\
& + 3E \left(g_p'(B(t_{j-1}))e^{i\langle \lambda, \mathbb{V}_n \rangle} \langle D^2\xi, \delta_j^{\otimes 2} \rangle_{\mathfrak{H}^{\otimes 2}} \right) \langle \varepsilon_{j-1}, \delta_j \rangle_{\mathfrak{H}} \\
& - iE \left(g_p(B(t_{j-1}))e^{i\langle \lambda, \mathbb{V}_n \rangle} \xi \langle D\langle \lambda, \mathbb{V}_n \rangle, \delta_j \rangle_{\mathfrak{H}}^3 \right) \\
& - 3E \left(g_p(B(t_{j-1}))e^{i\langle \lambda, \mathbb{V}_n \rangle} \langle D\langle \lambda, \mathbb{V}_n \rangle, \delta_j \rangle_{\mathfrak{H}}^2 \langle D\xi, \delta_j \rangle_{\mathfrak{H}} \right) \\
& + 3iE \left(g_p'(B(t_{j-1}))e^{i\langle \lambda, \mathbb{V}_n \rangle} \xi \langle D^2\langle \lambda, \mathbb{V}_n \rangle, \delta_j^{\otimes 2} \rangle_{\mathfrak{H}^{\otimes 2}} \right) \langle \varepsilon_{j-1}, \delta_j \rangle_{\mathfrak{H}} \\
& - 3E \left(g_p(B(t_{j-1}))e^{i\langle \lambda, \mathbb{V}_n \rangle} \xi \langle D\langle \lambda, \mathbb{V}_n \rangle, \delta_j \rangle_{\mathfrak{H}} \langle D^2\langle \lambda, \mathbb{V}_n \rangle, \delta_j^{\otimes 2} \rangle_{\mathfrak{H}^{\otimes 2}} \right) \\
& + 3iE \left(g_p(B(t_{j-1}))e^{i\langle \lambda, \mathbb{V}_n \rangle} \langle D\xi, \delta_j \rangle_{\mathfrak{H}} \langle D^2\langle \lambda, \mathbb{V}_n \rangle, \delta_j^{\otimes 2} \rangle_{\mathfrak{H}^{\otimes 2}} \right) \\
& + 3iE \left(g_p(B(t_{j-1}))e^{i\langle \lambda, \mathbb{V}_n \rangle} \langle D\langle \lambda, \mathbb{V}_n \rangle, \delta_j \rangle_{\mathfrak{H}} \langle D^2\xi, \delta_j^{\otimes 2} \rangle_{\mathfrak{H}^{\otimes 2}} \right) \\
& + E \left(g_p(B(t_{j-1}))e^{i\langle \lambda, \mathbb{V}_n \rangle} \langle D^3\xi, \delta_j^{\otimes 3} \rangle_{\mathfrak{H}^{\otimes 3}} \right) \\
& = \sum_{a=1}^{15} R_{j,n}^{(a)}. \tag{3.24}
\end{aligned}$$

Assume for a moment (see Steps 4 to 8 below) that

$$\sum_{j=1}^{\lfloor nu_p \rfloor} r_{j,n} \xrightarrow{n \rightarrow \infty} 0. \tag{3.25}$$

By Lemma 3.2 (iv) and since $e^{i\langle \lambda, \mathbb{V}_n \rangle}$, ξ and g_p''' are bounded, we have

$$\left| E \left(e^{i\langle \lambda, \mathbb{V}_n \rangle} \xi \sum_{j=1}^{\lfloor nu_p \rfloor} g_p'''(B(t_{j-1})) \langle \varepsilon_{j-1}, \delta_j \rangle_{\mathfrak{H}}^3 \right) - E \left(e^{i\langle \lambda, \mathbb{V}_n \rangle} \xi \times \frac{(-1)}{8n} \sum_{j=1}^{\lfloor nu_p \rfloor} g_p'''(B(t_{j-1})) \right) \right| \rightarrow 0.$$

Moreover, by Lebesgue bounded convergence, we have that

$$\left| E \left(e^{i\langle \lambda, \mathbb{V}_n \rangle} \xi \times \frac{(-1)}{8n} \sum_{j=1}^{\lfloor nu_p \rfloor} g_p'''(B(t_{j-1})) \right) - E \left(e^{i\langle \lambda, \mathbb{V}_n \rangle} \xi \times \frac{(-1)}{8} \int_0^{u_p} g_p'''(B(s)) ds \right) \right| \rightarrow 0.$$

Finally, note that $(B, \mathbb{V}_n) \rightarrow (B, \mathbb{V}_\infty)$ in law in $D_{\mathbb{R}}[0, \infty) \times \mathbb{R}^m$. By the Skorohod representation theorem (see, for example, Theorem 2.1.8 in [4]), we may construct a version of this sequence which converges a.s., so that again by Lebesgue bounded convergence, we have

$$E \left(e^{i\langle \lambda, \mathbb{V}_n \rangle} \xi \times \frac{(-1)}{8} \int_0^{u_p} g_p'''(B(s)) ds \right) \rightarrow E \left(e^{i\langle \lambda, \mathbb{V}_\infty \rangle} \xi \times \frac{(-1)}{8} \int_0^{u_p} g_p'''(B(s)) ds \right).$$

Putting these convergences together, we obtain:

$$E \left(e^{i\langle \lambda, \mathbb{V}_n \rangle} \xi \sum_{j=1}^{\lfloor nu_p \rfloor} g_p'''(B(t_{j-1})) \langle \varepsilon_{j-1}, \delta_j \rangle_{\mathfrak{H}}^3 \right) \rightarrow E \left(e^{i\langle \lambda, \mathbb{V}_\infty \rangle} \xi \times \frac{(-1)}{8} \int_0^{u_p} g_p'''(B(s)) ds \right). \quad (3.26)$$

Similarly, let us show that

$$\begin{aligned} 6E \left(e^{i\langle \lambda, \mathbb{V}_n \rangle} \xi \sum_{j=1}^{\lfloor nu_p \rfloor} \sum_{\ell=1}^{\lfloor nu_k \rfloor} g_p(B(t_{j-1})) g_k(B(t_{\ell-1})) \langle \delta_\ell, \delta_j \rangle_{\mathfrak{H}}^3 \right) \\ \rightarrow \kappa^2 E \left(e^{i\langle \lambda, \mathbb{V}_\infty \rangle} \xi \times \int_0^{u_p \wedge u_k} g_p(B(s)) g_k(B(s)) ds \right). \end{aligned} \quad (3.27)$$

We have, see (2.4) for the definition of ρ :

$$\begin{aligned} 6 \sum_{j=1}^{\lfloor nu_p \rfloor} \sum_{\ell=1}^{\lfloor nu_k \rfloor} g_p(B(t_{j-1})) g_k(B(t_{\ell-1})) \langle \delta_\ell, \delta_j \rangle_{\mathfrak{H}}^3 \\ = \frac{6}{n} \sum_{j=1}^{\lfloor nu_p \rfloor} \sum_{\ell=1}^{\lfloor nu_k \rfloor} g_p(B(t_{j-1})) g_k(B(t_{\ell-1})) \rho^3(\ell - j) \\ = \frac{6}{n} \sum_{r=1-\lfloor nu_p \rfloor}^{\lfloor nu_k \rfloor - 1} \rho^3(r) \sum_{j=1 \vee (1-r)}^{\lfloor nu_p \rfloor \wedge (\lfloor nu_k \rfloor - r)} g_p(B(t_{j-1})) g_k(B(t_{r+j-1})). \end{aligned} \quad (3.28)$$

For each fixed integer $r > 0$ (the case $r \leq 0$ being similar), we have

$$\begin{aligned} \left| \frac{1}{n} \sum_{j=1 \vee (1-r)}^{\lfloor nu_p \rfloor \wedge (\lfloor nu_k \rfloor - r)} g_p(B(t_{j-1})) g_k(B(t_{r+j-1})) - \frac{1}{n} \sum_{j=1 \vee (1-r)}^{\lfloor nu_p \rfloor \wedge (\lfloor nu_k \rfloor - r)} g_p(B(t_{j-1})) g_k(B(t_{j-1})) \right| \\ \leq C \|g_p\|_\infty \sup_{1 \leq j \leq \lfloor nu_p \rfloor} |g_k(B(t_{r+j-1})) - g_k(B(t_{j-1}))| \\ \leq C \|g_p\|_\infty \|g_k'\|_\infty \sup_{s \in [0, u_p]} |B(s + r/n) - B(s)| \xrightarrow[n \rightarrow \infty]{\text{a.s.}} 0, \end{aligned}$$

since the Brownian paths are uniformly continuous on compact intervals. Hence, for all fixed $r \in \mathbb{Z}$,

$$\frac{1}{n} \sum_{j=1 \vee (1-r)}^{\lfloor nu_p \rfloor \wedge (\lfloor nu_k \rfloor - r)} g_p(B(t_{j-1})) g_k(B(t_{r+j-1})) \xrightarrow[n \rightarrow \infty]{\text{a.s.}} \int_0^{u_p \wedge u_k} g_p(B(s)) g_k(B(s)) ds.$$

By combining a bounded convergence argument with (3.28) (observe in particular that $\kappa^2 = 6 \sum_{r \in \mathbb{Z}} \rho^3(r) < \infty$), we deduce that

$$6 \sum_{j=1}^{\lfloor nu_p \rfloor} \sum_{\ell=1}^{\lfloor nu_k \rfloor} g_p(B(t_{j-1})) g_k(B(t_{\ell-1})) \langle \delta_\ell, \delta_j \rangle_{\mathfrak{H}}^3 \xrightarrow[n \rightarrow \infty]{\text{a.s.}} \kappa^2 \int_0^{u_p \wedge u_k} g_p(B(s)) g_k(B(s)) ds.$$

Since $(B, \mathbb{V}_n) \rightarrow (B, \mathbb{V}_\infty)$ in $D_{\mathbb{R}}[0, \infty) \times \mathbb{R}^m$, we deduce that

$$\left(\mathbb{V}_n, \xi, 6 \sum_{j=1}^{\lfloor nu_p \rfloor} \sum_{\ell=1}^{\lfloor nu_k \rfloor} g_p(B(t_{j-1})) g_k(B(t_{\ell-1})) \langle \delta_\ell, \delta_j \rangle_{\mathfrak{H}}^3 \right)_{k=1, \dots, m} \xrightarrow{\text{Law}} \left(\mathbb{V}_\infty, \xi, \kappa^2 \int_0^{u_p \wedge u_k} g_p(B(s)) g_k(B(s)) ds \right)_{k=1, \dots, m} \text{ in } \mathbb{R}^d \times \mathbb{R} \times \mathbb{R}^m.$$

By boundedness of $e^{i\langle \lambda, \mathbb{V}_n \rangle}$, ξ and g_i , we have that (3.27) follows. Putting (3.25), (3.26), and (3.27) into (3.23), we deduce (3.18).

Now, it remains to prove (3.25).

Step 4.- Study of $R_{j,n}^{(5)}$, $R_{j,n}^{(8)}$ and $R_{j,n}^{(15)}$ in (3.24). Let $k \in \{1, 2, 3\}$. Since

$$D^k \xi = \sum_{i_1, \dots, i_k=1}^r \frac{\partial^k \psi}{\partial s_{i_1} \dots \partial s_{i_k}} (B_{s_1}, \dots, B_{s_r}) \mathbf{1}_{[0, s_{i_1}]} \otimes \dots \otimes \mathbf{1}_{[0, s_{i_k}]},$$

with $\psi \in C_b^\infty(\mathbb{R}^r)$, we have $\sum_{j=1}^{\lfloor nt \rfloor} |\langle D^k \xi, \delta_j^{\otimes k} \rangle_{\mathfrak{H}}| \leq Cn^{-(k-1)/3}$ by Lemma 3.2 (i) and (ii). Moreover, $|\langle \varepsilon_{j-1}, \delta_j \rangle_{\mathfrak{H}}| \leq n^{-1/3}$ by Lemma 3.2 (i). Hence, $\sum_{j=1}^{\lfloor nt \rfloor} |R_{j,n}^{(p)}| = O(n^{-2/3}) \xrightarrow{n \rightarrow \infty} 0$ for $p \in \{5, 8, 15\}$.

Step 5.- Study of $R_{j,n}^{(2)}$ and $R_{j,n}^{(3)}$ in (3.24). We can write, using Lemma 3.2 (i), Cauchy-Schwarz inequality and the definition (2.4) of ρ :

$$\begin{aligned} & \sum_{j=1}^{\lfloor nu_p \rfloor} |R_{j,n}^{(3)}| \\ & \leq 18 \sum_{k=1}^m |\lambda_k| \sum_{j=1}^{\lfloor nu_p \rfloor} \sum_{\ell=1}^{\lfloor nu_k \rfloor} \left| E \left(g_p(B(t_{j-1})) g'_k(B(t_{\ell-1})) I_1(\delta_\ell) e^{i\langle \lambda, \mathbb{V}_n \rangle} \xi \right) \right| |\langle \delta_\ell, \delta_j \rangle_{\mathfrak{H}}^2| |\langle \varepsilon_{\ell-1}, \delta_j \rangle_{\mathfrak{H}}| \\ & \leq Cn^{-7/6} \sum_{j=1}^{\lfloor nu_p \rfloor} \sum_{\ell=1}^{\lfloor nu_k \rfloor} \rho(\ell - j)^2 \leq Cn^{-1/6} \sum_{r \in \mathbb{Z}} \rho(r)^2 = Cn^{-1/6} \xrightarrow{n \rightarrow \infty} 0. \end{aligned}$$

Concerning $R_{j,n}^{(2)}$, we can write similarly:

$$\begin{aligned} & \sum_{j=1}^{\lfloor nu_p \rfloor} |R_{j,n}^{(2)}| \\ & \leq 9 \sum_{k=1}^m |\lambda_k| \sum_{j=1}^{\lfloor nu_p \rfloor} \sum_{\ell=1}^{\lfloor nu_k \rfloor} \left| E \left(g_p(B(t_{j-1})) g''_k(B(t_{\ell-1})) I_2(\delta_\ell^{\otimes 2}) e^{i\langle \lambda, \mathbb{V}_n \rangle} \xi \right) \right| |\langle \delta_\ell, \delta_j \rangle_{\mathfrak{H}}| |\langle \varepsilon_{\ell-1}, \delta_j \rangle_{\mathfrak{H}}^2| \\ & \leq Cn^{-4/3} \sum_{j=1}^{\lfloor nu_p \rfloor} \sum_{\ell=1}^{\lfloor nu_k \rfloor} |\rho(\ell - j)| \leq Cn^{-1/3} \sum_{r \in \mathbb{Z}} |\rho(r)| = Cn^{-1/3} \xrightarrow{n \rightarrow \infty} 0. \end{aligned}$$

Step 6.- Study of $R_{j,n}^{(1)}$, $R_{j,n}^{(6)}$, $R_{j,n}^{(10)}$, and $R_{j,n}^{(12)}$. First, let us deal with $R_{j,n}^{(1)}$. In order to lighten the notation, we set $\tilde{\xi}_{j,\ell} = g_p(B(t_{j-1}))g_k'''(B(t_{\ell-1}))\xi$. Using $I_3(\delta_\ell^{\otimes 3}) = I_2(\delta_\ell^{\otimes 2})I_1(\delta_\ell) - 2n^{-1/3}I_1(\delta_\ell)$ (see e.g. Proposition 1.1.2 from [10]) and then integrating by parts through (2.11), we get

$$\begin{aligned} E\left(e^{i\langle \lambda, \mathbb{V}_n \rangle} \tilde{\xi}_{j,\ell} I_3(\delta_\ell^{\otimes 3})\right) &= E\left(e^{i\langle \lambda, \mathbb{V}_n \rangle} \tilde{\xi}_{j,\ell} I_2(\delta_\ell^{\otimes 2}) I_1(\delta_\ell)\right) - 2n^{-1/3} E\left(e^{i\langle \lambda, \mathbb{V}_n \rangle} \tilde{\xi}_{j,\ell} I_1(\delta_\ell)\right) \\ &= E\left(e^{i\langle \lambda, \mathbb{V}_n \rangle} I_2(\delta_\ell^{\otimes 2}) \langle D\tilde{\xi}_{j,\ell}, \delta_\ell \rangle_{\mathfrak{H}}\right) \\ &\quad + iE\left(e^{i\langle \lambda, \mathbb{V}_n \rangle} I_2(\delta_\ell^{\otimes 2}) \tilde{\xi}_{j,\ell} \langle D\langle \lambda, \mathbb{V}_n \rangle, \delta_\ell \rangle_{\mathfrak{H}}\right). \end{aligned}$$

Due to Lemma 3.2 (i) and Cauchy-Schwarz inequality, we have

$$\sup_{\ell=1,\dots,[nu_k]} \sup_{j=1,\dots,[nu_p]} \left| E\left(e^{i\langle \lambda, \mathbb{V}_n \rangle} I_2(\delta_\ell^{\otimes 2}) \langle D\tilde{\xi}_{j,\ell}, \delta_\ell \rangle_{\mathfrak{H}}\right) \right| \leq Cn^{-2/3}.$$

By (3.5) and Cauchy-Schwarz inequality, we also have

$$\sup_{\ell=1,\dots,[nu_k]} \sup_{j=1,\dots,[nu_p]} \left| E\left(e^{i\langle \lambda, \mathbb{V}_n \rangle} I_2(\delta_\ell^{\otimes 2}) \tilde{\xi}_{j,\ell} \langle D\langle \lambda, \mathbb{V}_n \rangle, \delta_\ell \rangle_{\mathfrak{H}}\right) \right| \leq Cn^{-2/3}.$$

Hence, combined with Lemma 3.2 (i) and (ii), we get:

$$\begin{aligned} \sum_{j=1}^{[nu_p]} |R_{j,n}^{(1)}| &\leq \sum_{k=1}^m |\lambda_k| \sum_{j=1}^{[nu_p]} \sum_{\ell=1}^{[nu_k]} \left| E\left(g_p(B(t_{j-1}))g_k'''(B(t_{\ell-1}))I_3(\delta_\ell^{\otimes 3})e^{i\langle \lambda, \mathbb{V}_n \rangle} \xi\right) \right| \left| \langle \varepsilon_{\ell-1}, \delta_j \rangle_{\mathfrak{H}} \right|^3 \\ &\leq Cn^{-1/3} \sup_{\ell=1,\dots,[nu_k]} \sum_{j=1}^{[nu_p]} \left| \langle \varepsilon_{\ell-1}, \delta_j \rangle_{\mathfrak{H}} \right| \leq Cn^{-1/3} \xrightarrow{n \rightarrow \infty} 0. \end{aligned}$$

Now, let us concentrate on $R_{j,n}^{(6)}$. Since $e^{i\langle \lambda, \mathbb{V}_n \rangle}$, ξ , and g_p' are bounded, we have that

$$\begin{aligned} \sum_{j=1}^{[nu_p]} |R_{j,n}^{(6)}| &\leq C \sum_{j=1}^{[nu_p]} E\left(\langle D\langle \lambda, \mathbb{V}_n \rangle, \delta_j \rangle_{\mathfrak{H}}^2\right) |\langle \varepsilon_{j-1}, \delta_j \rangle_{\mathfrak{H}}| \\ &\leq Cn^{-1/3} \sum_{j=1}^{[nu_p]} E\left(\langle D\langle \lambda, \mathbb{V}_n \rangle, \delta_j \rangle_{\mathfrak{H}}^2\right) \quad \text{by Lemma 3.2 (i)} \\ &\leq Cn^{-2/3} \xrightarrow{n \rightarrow \infty} 0 \quad \text{by (3.6)}. \end{aligned}$$

Similarly,

$$\begin{aligned} \sum_{j=1}^{[nu_p]} |R_{j,n}^{(12)}| &\leq 3 \sum_{j=1}^{[nu_p]} \left| E\left(g_p(B(t_{j-1}))e^{i\langle \lambda, \mathbb{V}_n \rangle} \xi \langle D\langle \lambda, \mathbb{V}_n \rangle, \delta_j \rangle_{\mathfrak{H}} \langle D^2\langle \lambda, \mathbb{V}_n \rangle, \delta_j^{\otimes 2} \rangle_{\mathfrak{H}^{\otimes 2}}\right) \right| \\ &\leq C \sum_{j=1}^{[nu_p]} \left(E\left(\langle D\langle \lambda, \mathbb{V}_n \rangle, \delta_j \rangle_{\mathfrak{H}}^2\right) + E\left(\langle D^2\langle \lambda, \mathbb{V}_n \rangle, \delta_j^{\otimes 2} \rangle_{\mathfrak{H}^{\otimes 2}}^2\right) \right) \\ &\leq Cn^{-1/3} \xrightarrow{n \rightarrow \infty} 0 \quad \text{by (3.6) and (3.7)}. \end{aligned}$$

For $R_{j,n}^{(10)}$, we can write:

$$\begin{aligned} \sum_{j=1}^{\lfloor nu_p \rfloor} |R_{j,n}^{(10)}| &\leq 3 \sum_{j=1}^{\lfloor nu_p \rfloor} \left| E \left(g_p(B(t_{j-1})) e^{i\langle \lambda, \mathbb{V}_n \rangle} \langle D \langle \lambda, \mathbb{V}_n \rangle, \delta_j \rangle_{\mathfrak{H}}^2 \langle D \xi, \delta_j \rangle_{\mathfrak{H}} \right) \right| \\ &\leq C n^{-1/3} \sum_{j=1}^{\lfloor nu_p \rfloor} \left| E \left(\langle D \langle \lambda, \mathbb{V}_n \rangle, \delta_j \rangle_{\mathfrak{H}}^2 \right) \right| \quad \text{by Lemma 3.2 (i)} \\ &\leq C n^{-2/3} \xrightarrow{n \rightarrow \infty} 0 \quad \text{by (3.6)}. \end{aligned}$$

Step 7.- Study of $R_{j,n}^{(4)}$, $R_{j,n}^{(7)}$, $R_{j,n}^{(11)}$, $R_{j,n}^{(13)}$, and $R_{j,n}^{(14)}$.

Using (3.8), and then Cauchy-Schwarz inequality and Lemma 3.2 (i), we can write

$$\begin{aligned} &\sum_{j=1}^{\lfloor nu_p \rfloor} |R_{j,n}^{(4)}| \\ &\leq 3 \sum_{j=1}^{\lfloor nu_p \rfloor} \left| E \left(g_p''(B(t_{j-1})) e^{i\langle \lambda, \mathbb{V}_n \rangle} \xi \langle D \langle \lambda, \mathbb{V}_n \rangle, \delta_j \rangle_{\mathfrak{H}} \right) \right| \langle \varepsilon_{j-1}, \delta_j \rangle_{\mathfrak{H}}^2 \\ &\leq 3 \sum_{k=1}^m |\lambda_k| \sum_{j=1}^{\lfloor nu_p \rfloor} \sum_{i=1}^{\lfloor nu_k \rfloor} \left| E \left(g_p''(B(t_{j-1})) g_k'(B(t_{i-1})) e^{i\langle \lambda, \mathbb{V}_n \rangle} \xi I_3(\delta_i^{\otimes 3}) \right) \right| \left| \langle \varepsilon_{i-1}, \delta_j \rangle_{\mathfrak{H}} \right| \langle \varepsilon_{j-1}, \delta_j \rangle_{\mathfrak{H}}^2 \\ &\quad + 9 \sum_{k=1}^m |\lambda_k| \sum_{j=1}^{\lfloor nu_p \rfloor} \sum_{i=1}^{\lfloor nu_k \rfloor} \left| E \left(g_p''(B(t_{j-1})) g_k(B(t_{i-1})) e^{i\langle \lambda, \mathbb{V}_n \rangle} \xi I_2(\delta_i^{\otimes 2}) \right) \right| \left| \langle \delta_i, \delta_j \rangle_{\mathfrak{H}} \right| \langle \varepsilon_{j-1}, \delta_j \rangle_{\mathfrak{H}}^2 \\ &\leq C n^{-1/6} \sup_{1 \leq i \leq \lfloor nu_k \rfloor} \sum_{j=1}^{\lfloor nu_p \rfloor} \left| \langle \varepsilon_{i-1}, \delta_j \rangle_{\mathfrak{H}} \right| + C n^{-4/3} \sum_{j=1}^{\lfloor nu_p \rfloor} \sum_{i=1}^{\lfloor nu_k \rfloor} |\rho(i-j)| \leq C n^{-1/6} \xrightarrow{n \rightarrow \infty} 0. \end{aligned}$$

Using the same arguments, we show that $\sum_{j=1}^{\lfloor nu_p \rfloor} |R_{j,n}^{(7)}| \xrightarrow{n \rightarrow \infty} 0$ and $\sum_{j=1}^{\lfloor nu_p \rfloor} |R_{j,n}^{(14)}| \xrightarrow{n \rightarrow \infty} 0$.

Differentiating in (3.8), we get

$$\begin{aligned} \langle D^2 \langle \lambda, \mathbb{V}_n \rangle, \delta_j^{\otimes 2} \rangle_{\mathfrak{H}^{\otimes 2}} &= \sum_{k=1}^m \lambda_k \sum_{i=1}^{\lfloor nu_k \rfloor} g_k''(B(t_{i-1})) I_3(\delta_i^{\otimes 3}) \langle \varepsilon_{i-1}, \delta_j \rangle_{\mathfrak{H}}^2 \\ &\quad + 6 \sum_{k=1}^m \lambda_k \sum_{i=1}^{\lfloor nu_k \rfloor} g_k'(B(t_{i-1})) I_2(\delta_i^{\otimes 2}) \langle \varepsilon_{i-1}, \delta_j \rangle_{\mathfrak{H}} \langle \delta_i, \delta_j \rangle_{\mathfrak{H}} \\ &\quad + 6 \sum_{k=1}^m \lambda_k \sum_{i=1}^{\lfloor nu_k \rfloor} g_k(B(t_{i-1})) I_1(\delta_i) \langle \delta_i, \delta_j \rangle_{\mathfrak{H}}^2. \end{aligned}$$

Hence, using Cauchy-Schwarz inequality and Lemma 3.2 (i)-(ii), we can write

$$\begin{aligned}
& \sum_{j=1}^{\lfloor nu_p \rfloor} |R_{j,n}^{(11)}| \\
& \leq 3 \sum_{j=1}^{\lfloor nu_p \rfloor} \left| E(g'_p(B(t_{j-1})) e^{i\langle \lambda, \mathbb{V}_n \rangle} \xi \langle D^2 \langle \lambda, \mathbb{V}_n \rangle, \delta_j^{\otimes 2} \rangle_{\mathfrak{H}^{\otimes 2}}) \right| \langle \varepsilon_{j-1}, \delta_j \rangle_{\mathfrak{H}} \\
& \leq 3 \sum_{k=1}^m |\lambda_k| \sum_{j=1}^{\lfloor nu_p \rfloor} \sum_{i=1}^{\lfloor nu_k \rfloor} \left| E(g'_p(B(t_{j-1})) g''_k(B(t_{i-1})) e^{i\langle \lambda, \mathbb{V}_n \rangle} \xi I_3(\delta_i^{\otimes 3})) \right| \langle \varepsilon_{i-1}, \delta_j \rangle_{\mathfrak{H}}^2 |\langle \varepsilon_{j-1}, \delta_j \rangle_{\mathfrak{H}}| \\
& + 18 \sum_{k=1}^m |\lambda_k| \sum_{j=1}^{\lfloor nu_p \rfloor} \sum_{i=1}^{\lfloor nu_k \rfloor} \left| E(g'_p(B(t_{j-1})) g'_k(B(t_{i-1})) e^{i\langle \lambda, \mathbb{V}_n \rangle} \xi I_2(\delta_i^{\otimes 2})) \right| \\
& \quad \times |\langle \varepsilon_{i-1}, \delta_j \rangle_{\mathfrak{H}}| |\langle \varepsilon_{j-1}, \delta_j \rangle_{\mathfrak{H}}| |\langle \delta_i, \delta_j \rangle_{\mathfrak{H}}| \\
& + 18 \sum_{k=1}^m |\lambda_k| \sum_{j=1}^{\lfloor nu_p \rfloor} \sum_{i=1}^{\lfloor nu_k \rfloor} \left| E(g'_p(B(t_{j-1})) g_k(B(t_{i-1})) e^{i\langle \lambda, \mathbb{V}_n \rangle} \xi I_1(\delta_i)) \right| |\langle \varepsilon_{j-1}, \delta_j \rangle_{\mathfrak{H}}| \langle \delta_i, \delta_j \rangle_{\mathfrak{H}}^2 \\
& \leq C n^{-1/6} \sup_{1 \leq i \leq \lfloor nu_k \rfloor} \sum_{j=1}^{\lfloor nu_p \rfloor} |\langle \varepsilon_{i-1}, \delta_j \rangle_{\mathfrak{H}}| + C n^{-4/3} \sum_{j=1}^{\lfloor nu_p \rfloor} \sum_{i=1}^{\lfloor nu_k \rfloor} |\rho(i-j)| \\
& \quad + C n^{-5/6} \sum_{j=1}^{\lfloor nu_p \rfloor} \sum_{i=1}^{\lfloor nu_k \rfloor} |\rho(i-j)| \\
& \leq C n^{-1/6} \xrightarrow{n \rightarrow \infty} 0.
\end{aligned}$$

Using the same arguments, we show that $\sum_{j=1}^{\lfloor nu_p \rfloor} |R_{j,n}^{(13)}| \xrightarrow{n \rightarrow \infty} 0$.

Step 8.- Now, we consider the last term in (3.24), that is $R_{j,n}^{(9)}$. Since $e^{i\langle \lambda, \mathbb{V}_n \rangle}$, ξ , and g_p are bounded, we can write

$$\begin{aligned}
\left| \sum_{j=1}^{\lfloor nu_p \rfloor} R_{j,n}^{(9)} \right| & \leq C \sum_{j=1}^{\lfloor nu_p \rfloor} E(|\langle D \langle \lambda, \mathbb{V}_n \rangle, \delta_j \rangle_{\mathfrak{H}}|^3) \\
& \leq C \sum_{j=1}^{\lfloor nu_p \rfloor} E(\langle D \langle \lambda, \mathbb{V}_n \rangle, \delta_j \rangle_{\mathfrak{H}}^2) + E(\langle D \langle \lambda, \mathbb{V}_n \rangle, \delta_j \rangle_{\mathfrak{H}}^4).
\end{aligned}$$

In addition we have, see (3.8), that

$$\begin{aligned}
& \sum_{j=1}^{\lfloor nu_p \rfloor} E(\langle D\langle \lambda, \mathbb{V}_n \rangle, \delta_j \rangle_{\mathfrak{H}}^4) \\
& \leq 8m^3 \sum_{j=1}^{\lfloor nu_p \rfloor} \sum_{k=1}^m \lambda_k^4 \sum_{i_1=1}^{\lfloor nu_k \rfloor} \sum_{i_2=1}^{\lfloor nu_k \rfloor} \sum_{i_3=1}^{\lfloor nu_k \rfloor} \sum_{i_4=1}^{\lfloor nu_k \rfloor} E \left(\prod_{a=1}^4 \langle \varepsilon_{t_{i_a}}, \delta_j \rangle_{\mathfrak{H}} g'_k(B(t_{i_a-1})) I_3(\delta_{i_a}^{\otimes 3}) \right) \\
& \quad + 648m^3 \sum_{j=1}^{\lfloor nu_p \rfloor} \sum_{k=1}^m \lambda_k^4 \sum_{i_1=1}^{\lfloor nu_k \rfloor} \sum_{i_2=1}^{\lfloor nu_k \rfloor} \sum_{i_3=1}^{\lfloor nu_k \rfloor} \sum_{i_4=1}^{\lfloor nu_k \rfloor} E \left(\prod_{a=1}^4 \langle \delta_{i_a}, \delta_j \rangle_{\mathfrak{H}} g_k(B(t_{i_a-1})) I_2(\delta_{i_a}^{\otimes 2}) \right) \\
& \leq C \sum_{k=1}^m \lambda_k^4 \left[n^{-4/3} \sum_{j=1}^{\lfloor nu_p \rfloor} \sum_{i_1=1}^{\lfloor nu_k \rfloor} \sum_{i_2=1}^{\lfloor nu_k \rfloor} \sum_{i_3=1}^{\lfloor nu_k \rfloor} \sum_{i_4=1}^{\lfloor nu_k \rfloor} \left| E \left(\prod_{a=1}^4 g'_k(B(t_{i_a-1})) I_3(\delta_{i_a}^{\otimes 3}) \right) \right| \right. \\
& \quad \left. + \sum_{j=1}^{\lfloor nu_p \rfloor} \sum_{i_1=1}^{\lfloor nu_k \rfloor} \sum_{i_2=1}^{\lfloor nu_k \rfloor} \sum_{i_3=1}^{\lfloor nu_k \rfloor} \sum_{i_4=1}^{\lfloor nu_k \rfloor} \prod_{a=1}^4 |\langle \delta_{i_a}, \delta_j \rangle_{\mathfrak{H}}| \left| E \left(\prod_{a=1}^4 g_k(B(t_{i_a-1})) I_2(\delta_{i_a}^{\otimes 2}) \right) \right| \right].
\end{aligned}$$

By Lemma 3.5 we have that

$$\sum_{i_1=1}^{\lfloor nu_k \rfloor} \sum_{i_2=1}^{\lfloor nu_k \rfloor} \sum_{i_3=1}^{\lfloor nu_k \rfloor} \sum_{i_4=1}^{\lfloor nu_k \rfloor} \left| E \left(\prod_{a=1}^4 g'_k(B(t_{i_a-1})) I_3(\delta_{i_a}^{\otimes 3}) \right) \right| \leq C,$$

so that

$$n^{-4/3} \sum_{j=1}^{\lfloor nu_p \rfloor} \sum_{i_1=1}^{\lfloor nu_k \rfloor} \sum_{i_2=1}^{\lfloor nu_k \rfloor} \sum_{i_3=1}^{\lfloor nu_k \rfloor} \sum_{i_4=1}^{\lfloor nu_k \rfloor} \left| E \left(\prod_{a=1}^4 g'_k(B(t_{i_a-1})) I_3(\delta_{i_a}^{\otimes 3}) \right) \right| \leq Cn^{-1/3}.$$

On the other hand, by Cauchy-Schwarz inequality, we have

$$\left| E \left(\prod_{a=1}^4 g_k(B(t_{i_a-1})) I_2(\delta_{i_a}^{\otimes 2}) \right) \right| \leq C,$$

so that, with ρ defined by (2.4),

$$\begin{aligned}
& \sum_{j=1}^{\lfloor nu_p \rfloor} \sum_{i_1=1}^{\lfloor nu_k \rfloor} \sum_{i_2=1}^{\lfloor nu_k \rfloor} \sum_{i_3=1}^{\lfloor nu_k \rfloor} \sum_{i_4=1}^{\lfloor nu_k \rfloor} \prod_{a=1}^4 |\langle \delta_{i_a}, \delta_j \rangle_{\mathfrak{H}}| \left| E \left(\prod_{a=1}^4 g_k(B(t_{i_a-1})) I_2(\delta_{i_a}^{\otimes 2}) \right) \right| \\
& \leq Cn^{-4/3} \sum_{j=1}^{\lfloor nu_p \rfloor} \sum_{i_1=1}^{\lfloor nu_k \rfloor} |\rho(i_1 - j)| \times \sum_{i_2=1}^{\lfloor nu_k \rfloor} |\rho(i_2 - j)| \times \sum_{i_3=1}^{\lfloor nu_k \rfloor} |\rho(i_3 - j)| \times \sum_{i_4=1}^{\lfloor nu_k \rfloor} |\rho(i_4 - j)| \\
& \leq Cn^{-1/3} \left(\sum_{r \in \mathbb{Z}} |\rho(r)| \right)^4 = Cn^{-1/3}.
\end{aligned}$$

As a consequence, combining the previous estimates with (3.6), we have shown that

$$\left| \sum_{j=1}^{\lfloor nu_p \rfloor} R_{j,n}^{(9)} \right| \leq Cn^{-1/3} \xrightarrow{n \rightarrow \infty} 0,$$

and the proof of Theorem 3.7 is done. \square

Theorem 3.8. *If $g \in C^\infty(\mathbb{R})$ is bounded with bounded derivatives, then the sequence $(B, G_n^+(1, B), G_n^+(g, B))$ converges to $(B, [[B]], (1/8) \int g'''(B) ds + \int g(B) d[[B]])$ in the sense of finite-dimensional distributions on $[0, \infty)$, where*

$$G_n^+(g, B, t) := \frac{1}{\sqrt{n}} \sum_{j=1}^{\lfloor nt \rfloor} g(B(t_j)) h_3(n^{1/6} \Delta B_j), \quad t \geq 0, \quad n \geq 1.$$

Proof. The proof is exactly the same as the proof of Theorem 3.7, except ε_{j-1} must be everywhere replaced ε_j , and Lemma 3.2 (v) must be used instead of Lemma 3.2 (iv). \square

Proof of Theorem 3.1. We begin by observing the following general fact. Suppose U and V are càdlàg processes adapted to a filtration under which V is a semimartingale. Similarly, suppose \tilde{U} and \tilde{V} are càdlàg processes adapted to a filtration under which \tilde{V} is a semimartingale. If the processes (U, V) and (\tilde{U}, \tilde{V}) have the same law, then $U(0)V(0) + \int_0^\cdot U(s-) dV(s)$ and $\tilde{U}(0)\tilde{V}(0) + \int_0^\cdot \tilde{U}(s-) d\tilde{V}(s)$ have the same law. This is easily seen by observing that these integrals are the limit in probability of left-endpoint Riemann sums.

Now, let G_n^- and G_n^+ be as defined previously in this section. Define

$$G^-(g, B, t) = -\frac{1}{8} \int_0^t g'''(B(s)) ds + \int_0^t g(B(s)) d[[B]]_s,$$

$$G^+(g, B, t) = \frac{1}{8} \int_0^t g'''(B(s)) ds + \int_0^t g(B(s)) d[[B]]_s.$$

Let $\mathbf{t} = (t_1, \dots, t_d)$, where $0 \leq t_1 < \dots < t_d$. Let

$$G_n^-(g, B, \mathbf{t}) = (G_n^-(g, B, t_1), \dots, G_n^-(g, B, t_d)),$$

and similarly for G_n^+ , G^- , and G^+ . By Theorems 2.12, 3.7, and 3.8, the sequence

$$\{(B, V_n(B), G_n^-(g, B, \mathbf{t}), G_n^+(g, B, \mathbf{t}))\}_{n=1}^\infty$$

is relatively compact in $D_{\mathbb{R}^2}[0, \infty) \times \mathbb{R}^d \times \mathbb{R}^d$. By passing to a subsequence, we may assume it converges in law in $D_{\mathbb{R}^2}[0, \infty) \times \mathbb{R}^d \times \mathbb{R}^d$ to $(B, [[B]], X, Y)$, where $X, Y \in \mathbb{R}^d$.

By Theorems 2.12 and 3.7, $\{(B, V_n(B), G_n^-(g, B, \mathbf{t}))\}$ is relatively compact in $D_{\mathbb{R}^2}[0, \infty) \times \mathbb{R}^d$, and converges in the sense of finite-dimensional distributions to $(B, [[B]], G^-(g, B, \mathbf{t}))$. It follows that $(B, V_n(B), G_n^-(g, B, \mathbf{t})) \rightarrow (B, [[B]], G^-(g, B, \mathbf{t}))$ in law in $D_{\mathbb{R}^2}[0, \infty) \times \mathbb{R}^d$. Hence, $(B, [[B]], X)$ and $(B, [[B]], G^-(g, B, \mathbf{t}))$ have the same law in $D_{\mathbb{R}^2}[0, \infty) \times \mathbb{R}^d$. By the general fact we observed at the beginning of the proof, $(G^-(g, B), X)$ and $(G^-(g, B), G^-(g, B, \mathbf{t}))$ have the same law. This can be seen, for example, by letting

$$U(t) = \begin{pmatrix} -\frac{1}{8}g'''(B(t)) & g(B(t)) & 0 \\ 0 & 0 & X \end{pmatrix}, \quad \tilde{U}(t) = \begin{pmatrix} -\frac{1}{8}g'''(B(t)) & g(B(t)) & 0 \\ 0 & 0 & G^-(g, B, \mathbf{t}) \end{pmatrix},$$

$$V(t) = \tilde{V}(t) = \begin{pmatrix} t \\ [[B]]_t \\ 1 \end{pmatrix}.$$

In particular, $(G^-(g, B, \mathbf{t}), X)$ and $(G^-(g, B, \mathbf{t}), G^-(g, B, \mathbf{t}))$ have the same law. But this implies $G^-(g, B, \mathbf{t}) - X$ has the same law as the zero random variable, which gives $G^-(g, B, \mathbf{t}) - X = 0$ a.s.

We have thus shown that $X = G^-(g, B, \mathbf{t})$ a.s. Similarly, $Y = G^+(g, B, \mathbf{t})$ a.s. It follows that

$$(B, V_n(B), G_n^-(g, B), G_n^+(g, B)) \rightarrow (B, [[B]], G^-(g, B), G^+(g, B)),$$

and therefore

$$\left(B, V_n(B), \frac{G_n^-(g, B) + G_n^+(g, B)}{2} \right) \rightarrow \left(B, [[B]], \frac{G^-(g, B) + G^+(g, B)}{2} \right),$$

in the sense of finite-dimensional distributions on $[0, \infty)$, which is what was to be proved. \square

4 Moment bounds

The following four moment bounds are central to our proof of relative compactness in Theorem 2.13.

Theorem 4.1. *There exists a constant C such that*

$$E|V_n(B, t) - V_n(B, s)|^4 \leq C \left(\frac{\lfloor nt \rfloor - \lfloor ns \rfloor}{n} \right)^2,$$

for all n, s , and t .

Proof. The calculations in the proof of Theorem 10 in [11] show that

$$E \left| \sum_{j=\lfloor ns \rfloor + 1}^{\lfloor nt \rfloor} \Delta B_j^3 \right|^{2p} \leq C_p \left(\frac{\lfloor nt \rfloor - \lfloor ns \rfloor}{n} \right)^p,$$

for all n, s , and t . \square

Theorem 4.2. *Let $g \in C^1(\mathbb{R})$ have compact support. Fix $T > 0$ and let c and d be integers such that $0 \leq t_c < t_d \leq T$. Then*

$$E \left| \sum_{j=c+1}^d g(\beta_j) \Delta B_j^5 \right|^2 \leq C \|g\|_{1,\infty}^2 \Delta t^{1/3} |t_d - t_c|^{4/3},$$

where $\|g\|_{1,\infty} = \|g\|_\infty + \|g'\|_\infty$, and C depends only on T .

Proof. Note that

$$E \left| \sum_{j=c+1}^d g(\beta_j) \Delta B_j^5 \right|^2 = \sum_{i=c+1}^d \sum_{j=c+1}^d E_{ij}, \tag{4.1}$$

where $E_{ij} = E[g(\beta_i) \Delta B_i^5 g(\beta_j) \Delta B_j^5]$. Let $K = \|g\|_{1,\infty}$, and define $f : \mathbb{R}^3 \rightarrow \mathbb{R}$ by $f(x) = K^{-2} g(x_1) g(x_2) x_3^5$. Note that f has polynomial growth of order 1 with constants $\tilde{K} = 1$ and $r = 5$.

Let $\xi_1 = \beta_i$, $\xi_2 = \beta_j$, $\xi_3 = \Delta t^{-1/6} \Delta B_i$, $Y = \Delta t^{-1/6} \Delta B_j$, and $\varphi(y) = y^5$. Then $E_{ij} = K^2 \Delta t^{5/3} E[f(\xi)\varphi(Y)]$. By Theorem 2.8 with $k = 0$, $|E[f(\xi)\varphi(Y)]| \leq C|\eta|$, where $\eta_j = E[\xi_j Y]$. Using Lemma 2.10, we have

$$\begin{aligned} |\eta_1| &\leq C\Delta t^{1/6}(j^{-2/3} + |j-i|_+^{-2/3}), \\ |\eta_2| &\leq C\Delta t^{1/6}j^{-2/3}, \\ |\eta_3| &\leq C|j-i|_+^{-5/3}. \end{aligned}$$

Hence,

$$|E_{ij}| = K^2 \Delta t^{5/3} |E[f(\xi)\varphi(Y)]| \leq CK^2(\Delta t^{11/6}(j^{-2/3} + |j-i|_+^{-2/3}) + \Delta t^{5/3}|j-i|_+^{-5/3}).$$

Substituting this into (4.1) gives

$$\begin{aligned} E \left| \sum_{j=c+1}^d g(\beta_j) \Delta B_j^5 \right|^2 &\leq CK^2(\Delta t^{11/6}(d-c)^{4/3} + \Delta t^{5/3}(d-c)) \\ &\leq CK^2 \Delta t^{5/3} (d-c)^{4/3} = CK^2 \Delta t^{1/3} |t_d - t_c|^{4/3}, \end{aligned}$$

which completes the proof. \square

Theorem 4.3. Let $g \in C^2(\mathbb{R})$ have compact support. Fix $T > 0$ and let c and d be integers such that $0 \leq t_c < t_d \leq T$. Then

$$E \left| \sum_{j=c+1}^d g(\beta_j) \Delta B_j^3 \right|^2 \leq \|g\|_{2,\infty}^2 |t_d - t_c|,$$

where $\|g\|_{2,\infty} = \|g\|_\infty + \|g'\|_\infty + \|g''\|_\infty$, and C depends only on T .

Proof. Note that

$$E \left| \sum_{j=c+1}^d g(\beta_j) \Delta B_j^3 \right|^2 = \sum_{i=c+1}^d \sum_{j=c+1}^d E_{ij}, \quad (4.2)$$

where $E_{ij} = E[g(\beta_i) \Delta B_i^3 g(\beta_j) \Delta B_j^3]$. Let $K = \|g\|_{2,\infty}$, and define $f : \mathbb{R}^3 \rightarrow \mathbb{R}$ by $f(x) = K^{-2} g(x_1) g(x_2) x_3^3$. Note that f has polynomial growth of order 2 with constants $\tilde{K} = 1$ and $r = 3$.

Let $\xi_1 = \beta_i$, $\xi_2 = \beta_j$, $\xi_3 = \Delta t^{-1/6} \Delta B_i$, $Y = \Delta t^{-1/6} \Delta B_j$, and $\varphi(y) = y^3$. Then $E_{ij} = K^2 \Delta t E[f(\xi)\varphi(Y)]$. By Theorem 2.8 with $k = 1$, $E[f(\xi)\varphi(Y)] = \eta_1 E[\partial_1 f(\xi)] + \eta_2 E[\partial_2 f(\xi)] + R$, where $|R| \leq C(|\eta_3| + |\eta|^2)$. By (2.17), if $j = 1$ or $j = 2$, $|E[\partial_j f(\xi)]| \leq C(|E[\xi_1 \xi_3]| + |E[\xi_2 \xi_3]|)$. Therefore, using $|\eta_3|^2 \leq |\eta_3|$ and $|ab| \leq |a|^2 + |b|^2$,

$$|E_{ij}| \leq CK^2 \Delta t (|\eta_3| + |\eta_1|^2 + |\eta_2|^2 + |E[\xi_1 \xi_3]|^2 + |E[\xi_2 \xi_3]|^2).$$

Using Lemma 2.10, we have

$$\begin{aligned} |E[\xi_2 \xi_3]| &\leq C\Delta t^{1/6}(i^{-2/3} + |j-i|_+^{-2/3}), \\ |E[\xi_1 \xi_3]| &\leq C\Delta t^{1/6}i^{-2/3}. \end{aligned}$$

Together with the estimates from the proof of Theorem 4.2, this gives

$$|E_{ij}| \leq CK^2(\Delta t^{4/3}(i^{-4/3} + j^{-4/3} + |j - i|_+^{-4/3}) + \Delta t|j - i|_+^{-5/3}).$$

Substituting this into (4.2) gives

$$E \left| \sum_{j=c+1}^d g(\beta_j) \Delta B_j^3 \right|^2 \leq CK^2 \Delta t (d - c) = CK^2 |t_d - t_c|,$$

which completes the proof. \square

Theorem 4.4. *Suppose $g \in C^3(\mathbb{R})$ has compact support. Fix $T > 0$ and let c and d be integers such that $0 \leq t_c < t_d \leq T$. Then*

$$E \left| \sum_{j=c+1}^d (g(\beta_j) - g(\beta_c)) \Delta B_j^3 \right|^2 \leq C |t_d - t_c|^{4/3},$$

where C depends only on g and T .

Proof. Let $Y_j = g(\beta_j) - g(\beta_c)$, and note that

$$E \left| \sum_{j=c+1}^d Y_j \Delta B_j^3 \right|^2 = \sum_{i=c+1}^d \sum_{j=c+1}^d E_{ij}, \quad (4.3)$$

where $E_{ij} = E[Y_i \Delta B_i^3 Y_j \Delta B_j^3]$. For fixed i, j , define $f : \mathbb{R}^4 \rightarrow \mathbb{R}$ by

$$f(x) = \left(\frac{g(x_1 + \sigma_i x_2) - g(x_1)}{\sigma_i} \right) \left(\frac{g(x_1 + \sigma_j x_3) - g(x_1)}{\sigma_j} \right) x_4^3,$$

where $\sigma_j^2 = E|\beta_j - \beta_c|^2$. Note that f has polynomial growth of order 2 with constants K and r that do not depend on i or j .

Let $\xi_1 = \beta_c$, $\xi_2 = \sigma_i^{-1}(\beta_i - \beta_c)$, $\xi_3 = \sigma_j^{-1}(\beta_j - \beta_c)$, $\xi_4 = \Delta t^{-1/6} \Delta B_i$, $Y = \Delta t^{-1/6} \Delta B_j$, and $\varphi(y) = y^3$. Note that $E_{ij} = \sigma_i \sigma_j \Delta t E[f(\xi) \varphi(Y)]$, so that by Lemma 2.10(v),

$$|E_{ij}| \leq C \Delta t^{4/3} |i - c|^{1/6} |j - c|^{1/6} |E[f(\xi) \varphi(Y)]|. \quad (4.4)$$

By Theorem 2.8 with $k = 1$, $E[f(\xi) \varphi(Y)] = 3 \sum_{k=1}^3 \eta_k E[\partial_k f(\xi)] + R$, where $|R| \leq C(|\eta_4| + |\eta|^2)$. Using $|ab| \leq |a|^2 + |b|^2$ and the fact that $|\eta_j|^2 \leq |\eta|^2$, this gives

$$|E[f(\xi) \varphi(Y)]| \leq C \left(\sum_{k=1}^3 |E[\partial_k f(\xi)]|^2 + |\eta_4| + |\eta|^2 \right).$$

By (2.17), for each $k \leq 3$, $|E[\partial_k f(\xi)]| \leq C \sum_{\ell=1}^3 |E[\xi_\ell \xi_4]|$. Therefore, since $\eta_j = E[\xi_j Y]$, we have

$$|E[f(\xi) \varphi(Y)]| \leq C \left(|E[\xi_4 Y]| + \sum_{k=1}^3 (|E[\xi_k Y]|^2 + |E[\xi_k \xi_4]|^2) \right). \quad (4.5)$$

To estimate these covariances, first note that $d - c = n(t_d - t_c) \leq nT$. Hence, $\Delta t = n^{-1} \leq C(d - c)^{-1}$. Now, using Lemma 2.10,

$$\begin{aligned} |E[\xi_1 Y]| &\leq C\Delta t^{1/6}|j - c|^{-2/3} \leq C|d - c|^{-1/6}|j - c|^{-2/3} \leq C|j - c|^{-5/6}, \\ |E[\xi_2 Y]| &\leq C|i - c|^{-1/6}(|j - c|^{-2/3} + |j - i|_+^{-2/3}), \\ |E[\xi_3 Y]| &\leq C|j - c|^{-1/6}|j - c|^{-2/3} = C|j - c|^{-5/6}, \\ |E[\xi_4 Y]| &\leq C|j - i|_+^{-5/3}. \end{aligned}$$

Similarly,

$$\begin{aligned} |E[\xi_1 \xi_4]| &\leq C\Delta t^{1/6}|i - c|^{-2/3} \leq C|d - c|^{-1/6}|i - c|^{-2/3} \leq C|i - c|^{-5/6}, \\ |E[\xi_2 \xi_4]| &\leq C|i - c|^{-1/6}|i - c|^{-2/3} = C|i - c|^{-5/6}, \\ |E[\xi_3 \xi_4]| &\leq C|j - c|^{-1/6}(|i - c|^{-2/3} + |j - i|_+^{-2/3}). \end{aligned}$$

Substituting these estimates into (4.5) and using (4.4) gives

$$\begin{aligned} |E_{ij}| &\leq C\Delta t^{4/3}(|i - c|^{1/6}|j - c|^{1/6}|j - i|_+^{-5/3} \\ &\quad + |i - c|^{1/6}|j - c|^{-3/2} + |i - c|^{-1/6}|j - c|^{-7/6} + |i - c|^{-1/6}|j - c|^{1/6}|j - i|_+^{-4/3} \\ &\quad + |i - c|^{-3/2}|j - c|^{1/6} + |i - c|^{-7/6}|j - c|^{-1/6} + |i - c|^{1/6}|j - c|^{-1/6}|j - i|_+^{-4/3}). \end{aligned}$$

We can simplify this to

$$\begin{aligned} |E_{ij}| &\leq C\Delta t^{4/3}(|i - c|^{1/6}|j - c|^{1/6}|j - i|_+^{-4/3} \\ &\quad + |i - c|^{1/6}|j - c|^{-7/6} + |j - c|^{1/6}|j - i|_+^{-4/3} \\ &\quad + |i - c|^{-7/6}|j - c|^{1/6} + |i - c|^{1/6}|j - i|_+^{-4/3}). \end{aligned}$$

Using $|ab| \leq |a|^2 + |b|^2$, this further simplifies to

$$\begin{aligned} |E_{ij}| &\leq C\Delta t^{4/3}(|i - c|^{1/3}|j - i|_+^{-4/3} + |j - c|^{1/3}|j - i|_+^{-4/3} \\ &\quad + |i - c|^{1/6}|j - c|^{-7/6} + |i - c|^{-7/6}|j - c|^{1/6}). \end{aligned}$$

We must now make use of (4.3). Note that

$$\begin{aligned} \Delta t^{4/3} \sum_{i=c+1}^d \sum_{j=c+1}^d |i - c|^{1/3}|j - i|_+^{-4/3} &\leq C\Delta t^{4/3} \sum_{i=c+1}^d |i - c|^{1/3} \\ &\leq C\Delta t^{4/3}(d - c)^{4/3} = C|t_d - t_c|^{4/3}. \end{aligned}$$

Similarly,

$$\Delta t^{4/3} \sum_{j=c+1}^d \sum_{i=c+1}^d |j - c|^{1/3}|j - i|_+^{-4/3} \leq C|t_d - t_c|^{4/3}.$$

Also,

$$\begin{aligned} \Delta t^{4/3} \sum_{i=c+1}^d \sum_{j=c+1}^d |i-c|^{1/6} |j-c|^{-7/6} &\leq C \Delta t^{4/3} \sum_{i=c+1}^d |i-c|^{1/6} \leq C \Delta t^{4/3} (d-c)^{7/6} \\ &\leq C \Delta t^{4/3} (d-c)^{4/3} = C |t_d - t_c|^{4/3}, \end{aligned}$$

and similarly,

$$\Delta t^{4/3} \sum_{j=c+1}^d \sum_{i=c+1}^d |i-c|^{-7/6} |j-c|^{1/6} \leq C |t_d - t_c|^{4/3}.$$

It follows, therefore, that $\sum_{i=c+1}^d \sum_{j=c+1}^d |E_{ij}| \leq C |t_d - t_c|^{4/3}$. By (4.3), this completes the proof. \square

5 Proof of main result

Lemma 5.1. *If $g \in C^1(\mathbb{R})$ has compact support, then $\sum_{j=1}^{\lfloor nt \rfloor} g(\beta_j) \Delta B_j^5 \rightarrow 0$ ucp.*

Proof. Let $X_n(g, t) = \sum_{j=1}^{\lfloor nt \rfloor} g(\beta_j) \Delta B_j^5$. Fix $T > 0$ and let $0 \leq s < t \leq T$ be arbitrary. Then

$$X_n(g, t) - X_n(g, s) = \sum_{j=c+1}^d g(\beta_j) \Delta B_j^5,$$

where $c = \lfloor ns \rfloor$ and $d = \lfloor nt \rfloor$. By Theorem 4.2,

$$E|X_n(g, t) - X_n(g, s)|^2 \leq C \Delta t^{1/3} |t_d - t_c|^{4/3} \leq C |t_d - t_c|^{5/3} = C \left(\frac{\lfloor nt \rfloor - \lfloor ns \rfloor}{n} \right)^{5/3},$$

where C depends only on g and T . This verifies condition (2.6) of Theorem 2.3. By Theorem 4.2, $\sup_n E|X_n(g, T)|^2 \leq CT^{4/3} < \infty$. Hence, by Theorem 2.3, $\{X_n(g)\}$ is relatively compact in $D_{\mathbb{R}}[0, \infty)$. By Lemma 2.4, it will therefore suffice to show that $X_n(g, t) \rightarrow 0$ in probability for each fixed t . But this follows easily by taking $s = 0$ above, which gives $E|X_n(g, t)|^2 \leq C \Delta t^{1/3}$ and completes the proof. \square

We recall that we shall write $X_n(t) \approx Y_n(t)$ to mean that $X_n - Y_n \rightarrow 0$ uniformly on compacts in probability.

Lemma 5.2. *If $g \in C^6(\mathbb{R})$ has compact support, then*

$$I_n(g', B, t) \approx g(B(t)) - g(B(0)) + \frac{1}{12} \sum_{j=1}^{\lfloor nt \rfloor} g'''(\beta_j) \Delta B_j^3.$$

Proof. Fix $a, b \in \mathbb{R}$. Let $x = (a + b)/2$ and $h = (b - a)/2$. By Theorem 2.7,

$$\begin{aligned} g(b) - g(a) &= (g(x+h) - g(x)) - (g(x-h) - g(x)) \\ &= \sum_{j=1}^6 g^{(j)}(x) \frac{h^j}{j!} - \sum_{j=1}^6 g^{(j)}(x) \frac{(-h)^j}{j!} + R_1(x, h) - R_1(x, -h) \\ &= \sum_{\substack{j=1 \\ j \text{ odd}}}^6 \frac{1}{j!2^{j-1}} g^{(j)}(x)(b-a)^j + R_1(x, h) - R_1(x, -h) \\ &= g'(x)(b-a) + \frac{1}{24}g'''(x)(b-a)^3 + \frac{1}{5!2^4}g^{(5)}(x)(b-a)^5 + R_2(a, b), \end{aligned}$$

where $R_2(a, b) = R_1(x, h) - R_1(x, -h)$ and

$$R_1(x, h) = \frac{h^6}{5!} \int_0^1 (1-u)^6 [g^{(6)}(x+uh) - g^{(6)}(x)] du.$$

Similarly,

$$\begin{aligned} \frac{g'(a) + g'(b)}{2} - g'(x) &= \frac{1}{2}(g'(x+h) - g'(x)) + \frac{1}{2}(g'(x-h) - g'(x)) \\ &= \frac{1}{2} \sum_{j=1}^5 g^{(j+1)}(x) \frac{h^j}{j!} + \frac{1}{2} \sum_{j=1}^5 g^{(j+1)}(x) \frac{(-h)^j}{j!} + R_4(a, b) \\ &= \frac{1}{8}g'''(x)(b-a)^2 + \frac{1}{4!2^4}g^{(5)}(x)(b-a)^4 + R_4(a, b), \end{aligned}$$

where $R_4(a, b) = R_3(x, h) + R_3(x, -h)$ and

$$R_3(x, h) = \frac{h^5}{4!} \int_0^1 (1-u)^5 [g^{(6)}(x+uh) - g^{(6)}(x)] du.$$

Combining these two expansions gives

$$g(b) - g(a) = \frac{g'(a) + g'(b)}{2}(b-a) - \frac{1}{12}g'''(x)(b-a)^3 + \gamma g^{(5)}(x)(b-a)^5 + R_6(a, b),$$

where $\gamma = (5!2^4)^{-1} - (4!2^4)^{-1}$ and

$$R_6(a, b) = R_2(a, b) - R_4(a, b)(b-a).$$

Note that $R_6(a, b) = h(a, b)(b-a)^6$, where

$$|h(a, b)| \leq C \sup_{0 \leq u \leq 1} |g^{(6)}(x+uh) - g^{(6)}(x)|.$$

Taking $a = B(t_{j-1})$ and $b = B(t_j)$ gives

$$\begin{aligned} g(B(t_j)) - g(B(t_{j-1})) &= \frac{g'(B(t_{j-1})) + g'(B(t_j))}{2} \Delta B_j - \frac{1}{12}g'''(\beta_j) \Delta B_j^3 + \gamma g^{(5)}(\beta_j) \Delta B_j^5 \\ &\quad + h(B(t_{j-1}), B(t_j)) \Delta B_j^6 \end{aligned}$$

Recall that $B_n(t) = B(\lfloor nt \rfloor/n)$, so that

$$g(B(t)) - g(B(0)) = I_n(g', B, t) - \frac{1}{12} \sum_{j=1}^{\lfloor nt \rfloor} g'''(\beta_j) \Delta B_j^3 + \varepsilon_n(g, t),$$

where

$$\varepsilon_n(g, t) = \gamma \sum_{j=1}^{\lfloor nt \rfloor} g^{(5)}(\beta_j) \Delta B_j^5 + \sum_{j=1}^{\lfloor nt \rfloor} h(B(t_{j-1}), B(t_j)) \Delta B_j^6 + g(B(t)) - g(B_n(t)).$$

It will therefore suffice to show that $\varepsilon_n(g, t) \rightarrow 0$ ucp.

By the continuity of g and B , $g(B(t)) - g(B_n(t)) \rightarrow 0$ uniformly on compacts, with probability one. By Lemma 5.1, since $g^{(5)} \in C^1(\mathbb{R})$, $\gamma \sum_{j=1}^{\lfloor nt \rfloor} g^{(5)}(\beta_j) \Delta B_j^5 \rightarrow 0$ ucp. It remains only to show that

$$\sum_{j=1}^{\lfloor nt \rfloor} h(B(t_{j-1}), B(t_j)) \Delta B_j^6 \rightarrow 0 \quad \text{ucp.} \quad (5.1)$$

Fix $T > 0$. Let $\{n(k)\}_{k=1}^{\infty}$ be an arbitrary sequence of positive integers. By Theorem 2.11, we may find a subsequence $\{m(k)\}_{k=1}^{\infty}$ and a measurable subset $\Omega^* \subset \Omega$ such that $P(\Omega^*) = 1$, $t \mapsto B(t, \omega)$ is continuous for all $\omega \in \Omega^*$, and

$$\sum_{j=1}^{\lfloor m(k)t \rfloor} \Delta B_{j, m(k)}(\omega)^6 \rightarrow 15t, \quad (5.2)$$

as $k \rightarrow \infty$ uniformly on $[0, T]$ for all $\omega \in \Omega^*$. Fix $\omega \in \Omega^*$. We will show that

$$\sum_{j=1}^{\lfloor m(k)t \rfloor} h(B(t_{j-1}^{m(k)}, \omega), B(t_j^{m(k)}, \omega)) \Delta B_{j, m(k)}(\omega)^6 \rightarrow 0,$$

as $k \rightarrow \infty$ uniformly on $[0, T]$, which will complete the proof.

For this, it will suffice to show that

$$\sum_{j=1}^{\lfloor m(k)T \rfloor} |h(B(t_{j-1}^{m(k)}, \omega), B(t_j^{m(k)}, \omega))| \Delta B_{j, m(k)}(\omega)^6 \rightarrow 0,$$

as $k \rightarrow \infty$. We begin by observing that, by (5.2), there exists a constant L such that $\sum_{j=1}^{\lfloor m(k)T \rfloor} \Delta B_{j, m(k)}(\omega)^6 < L$ for all k . Now let $\varepsilon > 0$. Since g has compact support, $g^{(6)}$ is uniformly continuous. Hence, there exists $\delta > 0$ such that $|b - a| < \delta$ implies $|h(a, b)| < \varepsilon/L$ for all t . Moreover, there exists k_0 such that $k \geq k_0$ implies $|\Delta B_{j, m(k)}(\omega)| < \delta$ for all $1 \leq j \leq \lfloor m(k)T \rfloor$. Hence, if $k \geq k_0$, then

$$\sum_{j=1}^{\lfloor m(k)T \rfloor} |h(B(t_{j-1}^{m(k)}, \omega), B(t_j^{m(k)}, \omega))| \Delta B_{j, m(k)}(\omega)^6 < \frac{\varepsilon}{L} \sum_{j=1}^{\lfloor m(k)T \rfloor} \Delta B_{j, m(k)}(\omega)^6 < \varepsilon,$$

which completes the proof. \square

As a corollary to this result, we find that although we do not have a bound on the second moments of $I_n(g, B)$, we can instead approximate $I_n(g, B)$, in the ucp sense, by processes whose second moments can be uniformly bounded.

Corollary 5.3. *If $g \in C^6(\mathbb{R})$ has compact support, then there exists a sequence of càdlàg processes X_n such that $I_n(g', B, t) \approx X_n(t)$ and, for any $T > 0$,*

$$\sup_n \sup_{t \in [0, T]} E|X_n(t)|^2 < \infty.$$

Proof. This follows immediately from Lemma 5.2 and Theorem 4.3 by taking $X_n(t) = g(B(t)) - g(B(0)) + \frac{1}{12} \sum_{j=1}^{\lfloor nt \rfloor} g'''(\beta_j) \Delta B_j^3$. \square

Lemma 5.4. *If $g \in C^6(\mathbb{R})$ has compact support, then $\{I_n(g', B)\}$ is relatively compact in $D_{\mathbb{R}}[0, \infty)$.*

Proof. Define

$$\begin{aligned} X_n(t) &:= \frac{1}{12} \sum_{j=1}^{\lfloor nt \rfloor} g'''(\beta_j) \Delta B_j^3, \\ Y(t) &:= g(B(t)) - g(B(0)) \\ \varepsilon_n(t) &:= I_n(g', B, t) - Y(t) - X_n(t). \end{aligned}$$

Since $(x, y, z) \mapsto x + y + z$ is a continuous function from $D_{\mathbb{R}^3}[0, \infty)$ to $D_{\mathbb{R}}[0, \infty)$, it will suffice to show that $\{(X_n, Y, \varepsilon_n)\}$ is relatively compact in $D_{\mathbb{R}^3}[0, \infty)$. By Lemma 5.2, $\varepsilon_n \rightarrow 0$ ucp, and therefore in $D_{\mathbb{R}}[0, \infty)$. Hence, by Lemma 2.2, it will suffice to show that $\{X_n\}$ is relatively compact in $D_{\mathbb{R}}[0, \infty)$.

For this, we apply Theorem 2.3 with $\beta = 4$. Fix $T > 0$ and let $0 \leq s < t \leq T$. Let $c = \lfloor ns \rfloor$ and $d = \lfloor nt \rfloor$. Note that $q(a + b)^4 \leq C(|a|^2 + |b|^4)$. Hence, since g has compact support and, therefore, g''' is bounded,

$$\begin{aligned} E[q(X_n(t) - X_n(s))^4] &= E \left[q \left(\frac{1}{12} \sum_{j=c+1}^d g'''(\beta_j) \Delta B_j^3 \right)^4 \right] \\ &\leq CE \left| \sum_{j=c+1}^d (g'''(\beta_j) - g'''(\beta_c)) \Delta B_j^3 \right|^2 + CE \left| \sum_{j=c+1}^d g'''(\beta_c) \Delta B_j^3 \right|^4 \\ &\leq CE \left| \sum_{j=c+1}^d (g'''(\beta_j) - g'''(\beta_c)) \Delta B_j^3 \right|^2 + CE \left| \sum_{j=c+1}^d \Delta B_j^3 \right|^4. \end{aligned}$$

Since $g''' \in C^3(\mathbb{R})$, we may apply Theorems 4.4 and 4.1, which give

$$E[q(X_n(t) - X_n(s))^4] \leq C|t_d - t_c|^{4/3} + C|t_d - t_c|^2 \leq C \left(\frac{\lfloor nt \rfloor - \lfloor ns \rfloor}{n} \right)^{4/3},$$

which verifies condition (2.6) of Theorem 2.3. As above,

$$\begin{aligned} E|X_n(T)|^2 &\leq CE \left| \sum_{j=1}^{\lfloor nT \rfloor} (g'''(\beta_j) - g'''(\beta_c)) \Delta B_j^3 \right|^2 + CE \left| \sum_{j=1}^{\lfloor nT \rfloor} \Delta B_j^3 \right|^2 \\ &\leq CE \left| \sum_{j=1}^{\lfloor nT \rfloor} (g'''(\beta_j) - g'''(\beta_c)) \Delta B_j^3 \right|^2 + C \left(E \left| \sum_{j=1}^{\lfloor nT \rfloor} \Delta B_j^3 \right|^4 \right)^{1/2} \\ &\leq CT^{4/3} + CT. \end{aligned}$$

Hence, $\sup_n E|X_n(T)|^2 < \infty$. By Theorem 2.3, $\{X_n\}$ is relatively compact, completing the proof. \square

Lemma 5.5. *If $g \in C^9(\mathbb{R})$ has compact support, then*

$$I_n(g', B, t) \approx g(B(t)) - g(B(0)) + \frac{1}{12\sqrt{n}} \sum_{j=1}^{\lfloor nt \rfloor} \frac{g'''(B(t_{j-1})) + g'''(B(t_j))}{2} h_3(n^{1/6} \Delta B_j).$$

Proof. Using the Taylor expansions in the proof of Lemma 5.2, together with Lemma 5.1, we have

$$\sum_{j=1}^{\lfloor nt \rfloor} g'''(\beta_j) \Delta B_j^3 \approx \sum_{j=1}^{\lfloor nt \rfloor} \frac{g'''(B(t_{j-1})) + g'''(B(t_j))}{2} \Delta B_j^3.$$

By Lemma 5.2, since $h_3(x) = x^3 - 3x$, it therefore suffices to show that

$$n^{-1/3} \sum_{j=1}^{\lfloor nt \rfloor} \frac{g'''(B(t_{j-1})) + g'''(B(t_j))}{2} \Delta B_j = n^{-1/3} I_n(g''', B, t) \approx 0.$$

Since $g''' \in C^6(\mathbb{R})$, this follows from Lemma 5.4, Corollary 5.3, and Lemma 2.4. \square

Proof of Theorem 2.13. We first assume that g (and also G) has compact support. By Lemma 5.5 and Theorem 3.1, we need only show that $\{(B, V_n(B), I_n(g, B))\}$ is relatively compact in $D_{\mathbb{R}^3}[0, \infty)$. By Lemma 2.2, it will suffice to show that $\{I_n(g, B)\}$ is relatively compact in $D_{\mathbb{R}}[0, \infty)$. But this follows from Lemma 5.4, completing the proof when g has compact support.

Now consider general g . Let

$$\Xi_n = (B, V_n(B), I_n(g, B)) \quad \text{and} \quad \Xi = (B, [[B]], \int g(B) dB).$$

For $T > 0$, define $\Xi_n^T(t) = \Xi_n(t)1_{\{t < T\}}$ and $\Xi^T(t) = \Xi(t)1_{\{t < T\}}$. Using the definition of the Skorohod metric r on $D_{\mathbb{R}^d}[0, \infty)$ (formally given by (3.5.2) in [4] that we will not recall for simplicity), it holds that

$$r(x, y) \leq \int_0^\infty e^{-u} (\sup_{t \geq 0} |x(t \wedge u) - y(t \wedge u)| \wedge 1) du,$$

for any elements x and y in $D_{\mathbb{R}^d}[0, \infty)$. Hence, if two càdlàg functions x and y agree on the interval $[0, T)$, then $r(x, y) \leq \int_T^\infty e^{-u} du = e^{-T}$. Hence, by Lemma 2.5, it will suffice to show that $\Xi_n^T \rightarrow \Xi^T$ in law, where $T > 0$ is fixed.

Let $H : D_{\mathbb{R}^3}[0, \infty) \rightarrow \mathbb{R}$ be continuous and bounded, with $M = \sup |H(x)|$. Define $X_n = H(\Xi_n^T)$, so that it will suffice to show that $X_n \rightarrow H(\Xi^T)$ in law. For each $k > 0$, choose $G_k \in C^6(\mathbb{R})$ with compact support such that $G_k = G$ on $[-k, k]$. Let $g_k = G'_k$,

$$\tilde{\Xi}_{n,k} = (B, V_n(B), I_n(g_k, B)), \quad \tilde{\Xi}_k = (B, [[B]], \int g_k(B) dB),$$

$X_{n,k} = H(\tilde{\Xi}_{n,k}^T)$ and $Y_k = H(\tilde{\Xi}_k^T)$. Note that since $I_n(g_k, B) = I_n(g, B)$ on $\{\omega \in \Omega : \sup_{0 \leq t \leq T} |B(t)(\omega)| < k\}$, we have $E|X_n - X_{n,k}| \leq \delta_k$, where

$$\delta_k = 2MP \left(\sup_{0 \leq t \leq T} |B(t)| \geq k \right).$$

Also note that that $\delta_k \rightarrow 0$ as $k \rightarrow \infty$. Since G_k has compact support, we have already proven that $X_{n,k} \rightarrow Y_k$ in law. Hence, by Lemma 2.5, the sequences $\{X_n\}$ and $\{Y_k\}$ are both convergent in law, and they have the same limit. Thus, to show that $X_n \rightarrow H(\Xi^T)$ in law, it will suffice to show that $Y_k \rightarrow H(\Xi^T)$ in law. However, it is an immediate consequence of (2.19) that $\tilde{\Xi}_k^T \rightarrow \Xi^T$ ucp, which completes the proof. \square

Proof of Theorem 2.14. As in the proof of Theorem 2.13, $\{(B, V_n(B), J_n)\}$ is relatively compact. Let (B, X, Y) be any subsequential limit. By Theorem 2.12, $X = \kappa W$, where W is a standard Brownian motion, independent of B . Hence, $(B, X, Y) = (B, [[B]], Y)$. Fix $j \in \{1, \dots, k\}$. By Theorem 2.13, $(B, [[B]], Y_j)$ has the same law as $(B, [[B]], \int g_j(B) dB)$. Using the general fact we observed at the beginning of the proof of Theorem 3.1, together with (2.19) and the definition of $[[B]]$, this implies $(\int g_j(B) dB, Y_j)$ and $(\int g_j(B) dB, \int g_j(B) dB)$ have the same law. Hence, $Y_j = \int g_j(B) dB$ a.s., so $(B, X, Y) = (B, [[B]], J)$. \square

Acknowledgments

We would like to thank the referee for his or her careful and thorough reading, and the many comments and suggestions which helped us to improve the presentation of the paper.

References

- [1] K. Burdzy and J. Swanson. A change of variable formula with Itô correction term. *Ann. Probab.*, 38(5):1817–1869, 2010. <http://www.ams.org/mathscinet-getitem?mr=MR2722787>
- [2] P. Cheridito and D. Nualart. Stochastic integral of divergence type with respect to fractional Brownian motion with Hurst parameter $H \in (0, \frac{1}{2})$. *Ann. Inst. H. Poincaré Probab. Statist.*, 41(6):1049–1081, 2005. <http://www.ams.org/mathscinet-getitem?mr=MR2172209>
- [3] P. Donnelly and T. G. Kurtz. A countable representation of the Fleming-Viot measure-valued diffusion. *Ann. Probab.*, 24(2):698–742, 1996. <http://www.ams.org/mathscinet-getitem?mr=MR1419488>
- [4] S. N. Ethier and T. G. Kurtz. *Markov processes: Characterization and convergence*. Wiley Series in Probability and Mathematical Statistics: Probability and Mathematical Statistics. John Wiley & Sons Inc., New York, 1986. <http://www.ams.org/mathscinet-getitem?mr=MR0838085>
- [5] M. Gradinaru, I. Nourdin, F. Russo, and P. Vallois. m -order integrals and generalized Itô’s formula: the case of a fractional Brownian motion with any Hurst index. *Ann. Inst. H. Poincaré Probab. Statist.*, 41(4):781–806, 2005. <http://www.ams.org/mathscinet-getitem?mr=MR2144234>
- [6] P. Malliavin. *Stochastic analysis*, volume 313 of *Fundamental Principles of Mathematical Sciences*. Springer-Verlag, Berlin, 1997. <http://www.ams.org/mathscinet-getitem?mr=MR1450093>

- [7] I. Nourdin and D. Nualart. Central limit theorems for multiple Skorohod integrals. *J. Theoret. Probab.*, 23(1):39–64, 2010. <http://www.ams.org/mathscinet-getitem?mr=MR2591903>
- [8] I. Nourdin, D. Nualart, and C. A. Tudor. Central and non-central limit theorems for weighted power variations of fractional Brownian motion. *Ann. Inst. H. Poincaré Probab. Statist.*, 46(4), 1055–1079, 2010. Math. Review not available.
- [9] I. Nourdin and A. Réveillac. Asymptotic behavior of weighted quadratic variations of fractional Brownian motion: the critical case $H = 1/4$. *Ann. Probab.*, 37(6):2200–2230, 2009. <http://www.ams.org/mathscinet-getitem?mr=MR2573556>
- [10] D. Nualart. *The Malliavin calculus and related topics*. Probability and its Applications (New York). Springer-Verlag, Berlin, 2006. <http://www.ams.org/mathscinet-getitem?mr=MR2200233>
- [11] D. Nualart and S. Ortiz-Latorre. Central limit theorems for multiple stochastic integrals and Malliavin calculus. *Stochastic Process. Appl.*, 118(4):614–628, 2008. <http://www.ams.org/mathscinet-getitem?mr=MR2394845>
- [12] A. Réveillac. Convergence of finite-dimensional laws of the weighted quadratic variations process for some fractional Brownian sheets. *Stoch. Anal. Appl.*, 27(1):51–73, 2009. <http://www.ams.org/mathscinet-getitem?mr=MR2473140>
- [13] F. Russo and P. Vallois. Elements of stochastic calculus via regularization. In *Séminaire de Probabilités XL*, volume 1899 of *Lecture Notes in Math.*, pages 147–185. Springer, Berlin, 2007. <http://www.ams.org/mathscinet-getitem?mr=MR2409004>