

1-1-2006

## Identification and characterization of Rv3281 as a novel subunit of a biotin-dependent Acyl-CoA carboxylase in *Mycobacterium tuberculosis* H37Rv

Tae-Jin Oh  
*University of Central Florida*

Jaiyanth Daniel  
*University of Central Florida*

Hwa-Jung Kim  
*University of Central Florida*

Tatiana D. Sirakova  
*University of Central Florida*

Pappachan E. Kolattukudy  
Find similar works at <https://stars.library.ucf.edu/facultybib2000>  
*University of Central Florida* Libraries <http://library.ucf.edu>

This Article is brought to you for free and open access by the Faculty Bibliography at STARS. It has been accepted for inclusion in Faculty Bibliography 2000s by an authorized administrator of STARS. For more information, please contact [STARS@ucf.edu](mailto:STARS@ucf.edu).

### Recommended Citation

Oh, Tae-Jin; Daniel, Jaiyanth; Kim, Hwa-Jung; Sirakova, Tatiana D.; and Kolattukudy, Pappachan E., "Identification and characterization of Rv3281 as a novel subunit of a biotin-dependent Acyl-CoA carboxylase in *Mycobacterium tuberculosis* H37Rv" (2006). *Faculty Bibliography 2000s*. 6483.  
<https://stars.library.ucf.edu/facultybib2000/6483>

# Identification and Characterization of Rv3281 as a Novel Subunit of a Biotin-dependent Acyl-CoA Carboxylase in *Mycobacterium tuberculosis* H37Rv\*

Received for publication, October 31, 2005, and in revised form, December 5, 2005. Published, JBC Papers in Press, December 14, 2005, DOI 10.1074/jbc.M511761200

Tae-Jin Oh, Jaiyanth Daniel, Hwa-Jung Kim, Tatiana D. Sirakova, and Pappachan E. Kolattukudy<sup>1</sup>

From the Burnett College of Biomedical Sciences, University of Central Florida, Orlando, Florida 32816

*Mycobacterium tuberculosis* produces a large number of structurally diverse lipids generated from the carboxylation products of acetyl-CoA and propionyl-CoA. A biotin-dependent acyl-CoA carboxylase was purified from *M. tuberculosis* H37Rv by avidin affinity chromatography, and the three major protein components were determined by N-terminal sequencing to be the 63-kDa  $\alpha$ 3-subunit (AccA3, Rv3285), the 59-kDa  $\beta$ 5-subunit (AccD5, Rv3280), and the 56-kDa  $\beta$ 4-subunit (AccD4, Rv3799). A minor protein of about 24 kDa that co-purified with the above subunits was identified by matrix-assisted laser desorption/ionization-time of flight mass spectrometry to be the product of Rv3281 that is located immediately downstream of the open reading frame encoding the  $\beta$ 5-subunit. This protein displays identity over a short stretch of amino acids with the recently discovered  $\epsilon$ -subunits of *Streptomyces coelicolor*, suggesting that it might be an  $\epsilon$ -subunit of the mycobacterial acyl-CoA carboxylase. To test this hypothesis, the carboxylase subunits were expressed in *Escherichia coli* and purified. Acyl-CoA carboxylase activity was successfully reconstituted for the first time from purified subunits of the acyl-CoA carboxylase of *M. tuberculosis*. The reconstituted  $\alpha$ 3- $\beta$ 5 showed higher activity with propionyl-CoA than with acetyl-CoA, and the addition of the  $\epsilon$ -subunit stimulated the carboxylation by 3.2- and 6.3-fold, respectively. The  $\alpha$ 3- $\beta$ 4 showed very low activity with the above substrates but carboxylated long chain acyl-CoA. This  $\epsilon$ -subunit contains five sets of tandem repeats at the N terminus that are required for maximal enhancement of carboxylase activity. The Rv3281 open reading frame is co-transcribed with Rv3280 in the mycobacterial cell, and the level of  $\epsilon$ -protein was highest during the log phase and decreased during the stationary phase.

Biotin-dependent carboxylases are involved in the synthesis of malonyl-CoA and methylmalonyl-CoA. Most biotin-dependent enzymes contain three functional components: the biotin carboxylase (BC),<sup>2</sup> the biotin carboxyl carrier protein (BCCP), and the carboxyltransferase (CT) (1). The acyl-CoA carboxylation reaction occurs in two steps in two separate subsites of the enzyme. The first partial reaction involves the fixation of CO<sub>2</sub> on biotin and requires the cooperation of BC and

BCCP components; the biotin group is moved to interact with the BC component, resulting in the formation of carboxyl biotin. This carboxyl biotin then swings out to the carboxyltransferase component, resulting in the formation of the carboxylated product (2).

Acyl-CoA carboxylases from *Mycobacterium tuberculosis* and *Mycobacterium bovis* BCG have been purified previously and shown to have both propionyl-CoA carboxylase (PCC) and acetyl-CoA carboxylase (ACC) activities (3). *Mycobacterium smegmatis* PCC was reported to be composed of two subunits with the biotin being associated with the heavier subunit (4, 5), as found also in *M. tuberculosis* (3). *M. tuberculosis* has three *accA* genes annotated as an  $\alpha$ -subunit that contains the BC and BCCP domains and six *accD* genes annotated as  $\beta$ -subunits that comprise the CT domain (6). It is not known which of these genes are expressed in *M. tuberculosis*. The acyl-CoA selectivity of the CT domain in the different  $\beta$ -subunits also remains unclear. Isolation of the various subunits and reconstitution of active carboxylase would be required to examine the catalytic activities of the various gene products. However, there are no reports of reconstitution of active acyl-CoA carboxylase from purified subunits of the *M. tuberculosis* acyl-CoA carboxylase.

Recently, a new group of acyl-CoA carboxylases was identified in *S. coelicolor* that contained a third type of subunit, designated  $\epsilon$ . The  $\epsilon$ -subunit considerably enhanced the basal activity obtained with the  $\alpha$ - and  $\beta$ -subunits from this organism (7, 8). It is not known whether the mycobacterial carboxylase belongs to this new group. In the genome of *M. tuberculosis*, an open reading frame (ORF) corresponding to an  $\epsilon$ -subunit has not been identified, and the purified carboxylase from the pathogen has not been thoroughly examined for subunit composition. We report the identification of an  $\epsilon$ -like ORF adjacent to the  $\beta$ 5 ORF (Rv3280) and present evidence for the expression of this ORF in *M. tuberculosis*. MALDI-TOF MS analysis of a 24-kDa protein that was purified with the native carboxylase by avidin-affinity chromatography showed it to be the  $\epsilon$ -subunit of the carboxylase. We also showed that ACCA3, ACCD4, and ACCD5 are the major subunits of the native carboxylase. Reconstitution of active carboxylase from expressed and purified subunits showed that the  $\epsilon$ -subunit stimulates carboxylation by  $\alpha$ 3- $\beta$ 5, demonstrating that *M. tuberculosis* has a novel carboxylase similar to that in *S. coelicolor*.

## EXPERIMENTAL PROCEDURES

**Materials**—Sodium [<sup>14</sup>C]bicarbonate (50 Ci mol<sup>-1</sup>) was purchased from American Radiolabeled Chemicals Inc., and all other chemicals and reagents were of the highest grade and were purchased from Sigma and Fisher. Nucleotide primers were synthesized by Integrated DNA Technologies, Inc.

**Bacterial Strains, Culture, and Transformation Conditions**—*Escherichia coli* strain DH5 $\alpha$  was used for routine subcloning and was transformed according to Sambrook *et al.* (9). *E. coli* BL21 Star (DE3) and *E. coli* Rosetta (DE3) were used for expression of recombinant proteins

\* This work was supported by Grants AI46582 and AI35272 from the National Institutes of Health. The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

<sup>1</sup> To whom correspondence should be addressed: Burnett College of Biomedical Sciences, University of Central Florida, BMS 136, 4000 Central Florida Blvd., Orlando, FL 32816-2364; Tel.: 407-823-1206; Fax: 407-823-0956; E-mail: pk@mail.ucf.edu.

<sup>2</sup> The abbreviations used are: BC, biotin carboxylase; BCCP, biotin carboxyl carrier protein; CT, carboxyltransferase; PCC, propionyl-CoA carboxylase; ACC, acetyl-CoA carboxylase; ORF, open reading frame; MALDI-TOF MS, matrix-assisted laser desorption/ionization time-of-flight mass spectrometry; RT, reverse transcriptase; SC, specificity constant; CAPS, 3-(cyclohexylamino)propanesulfonic acid.

# A Novel $\epsilon$ -Subunit in Mycobacterial Acyl-CoA Carboxylase

**TABLE 1**  
Primers used for this study

Primer name	Sequence <sup>a</sup> (5'→3' direction)	Comments
AccA3F	<u>GGCATATG</u> GCTAGTCACGCCGGCT	Upstream of <i>accA3</i> , NdeI site added
AccA3R	GGAGATCTT <u>TACTTGATC</u> TCGGCG	Downstream of <i>accA3</i> , BglII site added
AccD4F	<u>GGCATATG</u> ACCGTCACCGAGCCG	Upstream of <i>accD4</i> , NdeI site added
AccD4R	GGCTCGAGTCAGGCCGCTGCTTGC	Downstream of <i>accD4</i> , XhoI site added
AccD5F	<u>GGCATATG</u> ACAAGCGTTACCGAC	Upstream of <i>accD5</i> , NdeI site added
AccD5R	GGCTCGAGTCACAGGGGCACGTT	Downstream of <i>accD5</i> , XhoI site added
1ENF	<u>CATATGGGA</u> ACGTGCCCTGTG	Upstream <sup>1</sup> M of Rv3281, NdeI site added
1ECR	GGATCCTCGGCGCATGTGCGTC	Downstream of Rv3281, BamHI site added
102EF	<u>GGCATATG</u> ACCGAGAAGCCGCTGC	Positions <sup>102</sup> V of Rv3281, NdeI site added
102ER	GGCTCGAGTCATCGGCGCATGTGCGTC	Downstream of Rv3281, XhoI site added
D5-227F	CTTCGTGTACCGCCAGCAGCTGG	Internal primer of <i>accD5</i> , RT-PCR
E-139R	CTGCCGGATTGTTCTCGTCTC	Internal primer of Rv3281, RT-PCR

<sup>a</sup> Endonuclease restriction sites added to sequences are underlined.

(10). All media were purchased from Difco. Host strains for cloning and expression experiments were grown on Luria Bertani (LB) broth or agar. Ampicillin, chloramphenicol, and kanamycin were added when required at final concentrations of 100, 34, and 50  $\mu\text{g ml}^{-1}$ , respectively.

**Growth Conditions, Protein Production, and Preparation of Cell-free Extracts**—For expression of heterologous proteins, *E. coli* strains harboring the appropriate plasmids were grown at 37 °C in LB medium in the presence of the corresponding antibiotics for plasmid maintenance. Overnight cultures were diluted 1:100 in fresh medium, grown to  $A_{600}$  0.8, and induced with 1 mM isopropyl- $\beta$ -D-thiogalactopyranoside for 4 h at the same temperature with shaking. *E. coli* BL21 transformed with pCY216 (11), which expresses the *E. coli* biotin ligase (BirA), was grown under the same conditions as above and induced by the addition of arabinose to a final concentration of 0.5%. The cells were harvested, washed, and resuspended in 1 $\times$  equilibration/wash buffer, pH 7.0, containing 50 mM sodium phosphate and 300 mM NaCl and were disrupted by sonication using a Branson Sonifier 450 (Branson Ultrasonics Corp., Danbury, CT). Cell debris was removed by centrifugation, and the supernatant was used as the cell-free extract.

**Gel Electrophoresis and Western Blot Analysis**—Cell-free extracts and purified proteins were analyzed by SDS-PAGE (12) using a Bio-Rad minigel apparatus. Protein concentration was determined by the method of Bradford (13). To detect His-tagged protein and biotin-containing proteins, Western blot analyses were carried out with His probe (H-3) (Santa Cruz Biotechnology, Inc., Santa Cruz, CA) and peroxidase-conjugated avidin (Pierce), respectively. For Western blotting, cell extracts or purified proteins were subjected to SDS-PAGE and electroblotted onto a polyvinylidene difluoride membrane by using Trans-blot Cell from Bio-Rad.


**Antibody Production and  $\epsilon$ -Protein Expression in *M. tuberculosis***—The purified 177-residue  $\epsilon$ -subunit was resolved on SDS-PAGE and used for raising antibodies in rabbit by Harlan Bioproducts for Science, Inc (Madison, WI) according to their standard 73-day protocol. *M. tuberculosis* H37Rv cultures were grown aerobically in roller cultures in 7H9 medium at 37 °C. Equal quantities of cell lysates from 3-, 5-, 7-, and 9-day-old cultures were resolved on 12% SDS-PAGE and were transferred to Immobilon polyvinylidene difluoride membrane. The blot was probed with antisera raised as described above, and the bands were quantitated on an AlphaImager Gel Doc system.

**Avidin Affinity Chromatography and Electrophoresis**—Acyl-CoA carboxylase was purified by avidin affinity chromatography as described previously (3). *M. tuberculosis* H37Rv inactivated by  $\gamma$  irradiation (obtained from Colorado State University, TB Research Materials and Vaccine Testing Contract NIAID, National Institutes of Health, NO1 AI-753230) were disrupted by a French pressure cell operating at 15,000 p.s.i. (passed four times) in 100 mM potassium phosphate (pH 7.0) containing 1.0 mM dithioerythritol, 1.0 mM EDTA, and 10% glycerol. Just before each passage, 50  $\mu\text{l}$  of 0.1 M phenylmethylsulfonyl fluoride in isopropyl alcohol was added to

inhibit proteases. The homogenate was centrifuged at 30,000  $\times g$  for 30 min, and the supernatant was then centrifuged at 105,000  $\times g$  for 90 min. The final supernatant was mixed with a monomeric avidin-Sepharose affinity resin; the mixture was stirred slowly for 1 h, and the gel was poured into a column. The gel was washed with 5 bed volumes of the same buffer at 0.5 ml/min, and a linear gradient of 0–0.2 mM *d*-biotin in the phosphate buffer in a total volume of 300 ml was applied. The fractions (5.0 ml) containing the highest carboxylase activity were pooled and concentrated by ultrafiltration using Amicon PM-3 membrane. To determine the N-terminal protein sequence, the separated proteins from 7.5% SDS-PAGE were transferred on Immobilon polyvinylidene difluoride membrane (Millipore) in 10 mM CAPS buffer (pH 11.0) with 10% methanol. After staining with Coomassie Blue, N-terminal sequence of each protein band was determined by automated Edman degradation on a protein sequencer (Cleveland Clinic, Cleveland, OH). The  $\epsilon$ -like proteins were resolved on 15% SDS-PAGE and analyzed by MALDI-TOF MS following tryptic digestion.

**Gene Cloning and Plasmid Constructions**—DNA corresponding to each carboxylase subunit ORF was amplified using *Pfu* Turbo Hotstart DNA polymerase (Stratagene), and expression was performed using the pET expression vector (Novagen). The primers used for amplification of the genes *Rv3285* ( $\alpha 3$ ), *Rv3799c* ( $\beta 4$ ), *Rv3280* ( $\beta 5$ ), and *Rv3281* ( $\epsilon$ ) are listed in Table 1. All of the PCR products were cloned using the PCR-BluntII TOPO vector into *E. coli* TOP10 competent cells. These inserts were released by digestion as described by Table 1 and subcloned in pET16b or pET28a (+) vectors. *Rv3285*, *Rv3799c*, and *Rv3280* were expressed in pET16b, and *Rv3281* was expressed in pET28a (+). In these vectors, the ORFs were directionally cloned and expressed as His-tagged fusion proteins in *E. coli* BL21 Star (DE3) or *E. coli* Rosetta (DE3).

**Purification of Expressed Proteins**—The *E. coli* strain BL21 star (DE3) containing the plasmids pTJA3 ( $\alpha 3$ ), pTJD4 ( $\beta 4$ ), and pTJD5 ( $\beta 5$ ) were induced with isopropyl- $\beta$ -D-thiogalactopyranoside for protein overexpression as described above. Following lysis, the 16,000  $\times g$  supernatants were mixed with TALON resin (BD Biosciences) at room temperature for 30 min to allow the polyhistidine-tagged protein to bind the resin. The His-tagged fusion proteins were recovered by elution with 100 mM imidazole for the  $\alpha 3$ - and  $\beta 4$ -subunits or 500 mM imidazole for the  $\beta 5$ -subunits. For biotinylation of the cloned  $\alpha 3$ -subunit, *E. coli* cell lysate containing the expressed  $\alpha 3$ -subunit was incubated with an extract of *E. coli* cells expressing the cloned *E. coli* biotin ligase (BirA) from the pCY216 plasmid at 30 °C for 3 h (11). Biotinylation of the  $\alpha 3$ -subunit was confirmed by Western blotting with avidin-horseradish peroxidase conjugate (Santa Cruz Biotechnology). The *E. coli* strain Rosetta (DE3) was used for expression of pTJE1 and pTJE102. The His-tagged  $\epsilon$ -proteins were insoluble and were solubilized with 6 M guanidine-HCl from the 16,000  $\times g$  pellet obtained from cell lysate. The solubilized proteins were bound to TALON resin, which was pre-equilibrated in 6 M guanidine-HCl, and were purified under




**FIGURE 1. Determination of subunit composition and domain organization of *M. tuberculosis* acyl-CoA carboxylase.** A, identification of the subunits of the biotin-dependent acyl-CoA carboxylase. Acyl-CoA carboxylase was purified by single-step avidin affinity chromatography, and the purified complex was resolved on 7.5% SDS-PAGE. The N-terminal amino acid sequence of the 63- ( $\alpha$ 3), 59- ( $\beta$ 5), and 56- ( $\beta$ 4) kDa bands were determined by Edman degradation. The 24-kDa band ( $\epsilon$ ) was resolved on 15% SDS-PAGE, and the amino acid sequence was determined by MALDI-TOF MS following tryptic digestion. Lane 1, molecular weight protein marker; lane 2, avidin-purified acyl-CoA carboxylase complex on 15% SDS-PAGE; lane 3, molecular weight protein marker; and lane 4, 7.5% SDS-PAGE of avidin-purified acyl-CoA carboxylase. B, domain organization and amino acid sequence similarity of *M. tuberculosis* and *S. coelicolor* acyl-CoA carboxylase. Sequence similarities are shown by similarly shaded boxes. *AccA3*, Rv3285, acyl-CoA carboxylase  $\alpha$ -subunit from *M. tuberculosis*; *AccD5*, Rv3280, propionyl-CoA carboxylase  $\beta$ -subunit from *M. tuberculosis*; *Acc/PccE*, Rv3281, acyl-CoA carboxylase  $\epsilon$ -subunit from *M. tuberculosis*; *AccA2*, acyl-CoA carboxylase  $\alpha$ -subunit from *S. coelicolor*; *AccB*, acetyl-CoA carboxylase  $\beta$ -subunit from *S. coelicolor*; *PccB*, propionyl-CoA carboxylase  $\beta$ -subunit from *S. coelicolor*; *AccE*, acetyl-CoA carboxylase  $\epsilon$ -subunit from *S. coelicolor*; *PccE*, propionyl-CoA carboxylase  $\epsilon$ -subunit from *S. coelicolor*; ATP/CO<sub>3</sub><sup>-</sup>, biotin carboxylation domain that binds ATP and CO<sub>2</sub> fixation, respectively; CB, putative carboxybiotin-binding domain; AcC and PcC, carboxyltransferase domain that binds acetyl-CoA and propionyl-CoA, respectively.

denaturing conditions according to the manufacturer's protocol. Following elution with 150 mM imidazole, the guanidine-HCl was removed, and the purified  $\epsilon$ -proteins were refolded by stepwise dialysis. The first step of dialysis was performed against a solution containing 2 M guanidine-HCl, 100 mM potassium phosphate (pH 8.0), and 10% glycerol for 12 h at 4 °C followed by two rounds of dialysis against 100 mM potassium phosphate (pH 8.0) and 10% glycerol for 12 h at 4 °C.

**Reconstitution of Carboxylase Activity**—The purified  $\epsilon$ -subunit in dilute form (100  $\mu$ g/ml) was initially combined with a dilute (100  $\mu$ g/ml) preparation of either the  $\beta$ 4- or the  $\beta$ 5-subunits, after which the  $\beta$ - $\epsilon$  subunits were combined with a dilute preparation of the biotinylated  $\alpha$ 3-subunit. Following this, the reconstituted enzyme was concentrated 5–7-fold by ultrafiltration in Centrprep YM-10 filters (Millipore) at 700  $\times$  g for 6 h at 4 °C.

**Acyl-CoA Carboxylase Assay**—Enzyme activities of the *in vitro* reconstituted subunits obtained by mixing purified proteins were measured by following the incorporation of label from H<sup>14</sup>CO<sub>3</sub><sup>-</sup> into acid-stable reaction material (14). The reaction mixture contained 100 mM potassium phosphate, pH 8.0, 300  $\mu$ g of bovine serum albumin, 2 mM ATP, 10 mM MgCl<sub>2</sub>, 5 mM NaH<sup>14</sup>CO<sub>3</sub> (50 Ci mol<sup>-1</sup>), 2 mM acetyl-CoA, 2 mM propionyl-CoA, and 200–600  $\mu$ g of reconstituted enzyme in a total reaction volume of 100  $\mu$ l. The reaction was initiated by the addition of NaH<sup>14</sup>CO<sub>3</sub>, allowed to proceed at 30 °C for 3 h, and stopped with 150  $\mu$ l of 6 M HCl. Following evaporation, the residue was resuspended in 100  $\mu$ l of water, and the <sup>14</sup>C radioactivity was determined by liquid scintillation counting. To determine the effect of inhibitors on acyl-CoA carboxylase activity, the reconstituted enzyme was incubated with the inhibitor for 30 min at room temperature prior to the assay described above.



**FIGURE 2.  $\epsilon$  transcript and protein expression level in *M. tuberculosis* cultures grown *in vitro*.** *A*, semi-quantitative RT-PCR showing expression of the  $\epsilon$  ORF in *M. tuberculosis*. *Lane 1*, PCR product obtained with internal primers 102EF and 1ECR (Table 1); *lane 2*, PCR product obtained with primers spanning the  $\beta 5$ - $\epsilon$  ORF junction (D5-227F and E-139R, Table 1). *B*, Western blot of total cell lysates from *M. tuberculosis* H37Rv cultures grown at 37 °C. Equal quantities of lysates from each time point were resolved on 12% SDS-PAGE, transferred to Immobilon polyvinylidene difluoride membrane, and probed with rabbit antisera raised against purified  $\epsilon$ -subunit.

## RESULTS

**Avidin Purification and Identification of the Native Carboxylase Subunits**—The carboxylases purified from *M. tuberculosis* H37Rv by avidin affinity chromatography showed activity with both acetyl-CoA (12.3 nmol/mg/min) and propionyl-CoA (20.6 nmol/mg/min) as substrates. When the purified native carboxylase preparation from *M. tuberculosis* was resolved on 7.5% SDS-PAGE, three proteins with apparent molecular masses of 63, 59, and 56 kDa were observed. The N-terminal amino acid sequence determination revealed them as being the products of *Rv3285* (AccA3,  $\alpha 3$ ; ASHAGSRIARISK, lacking the terminal methionine), *Rv3280* (AccD5,  $\beta 5$ ; TSVTDRSAHSAER, lacking the terminal methionine), and *Rv3799c* (AccD4,  $\beta 4$ ; VTEPVL, lacking 2 N-terminal amino acids) (Fig. 1A).

We expressed the  $\alpha 3$ -,  $\beta 4$ -, and  $\beta 5$ -subunits as His-tagged fusion proteins in *E. coli*. The biotinylated  $\alpha 3$ -subunit was purified and mixed with the purified  $\beta 4$ - or  $\beta 5$ -subunit and assayed for carboxylation of acetyl-CoA and propionyl-CoA. The reconstituted enzyme showed only very low activity but appeared to support the conclusion that the  $\beta 5$ -subunit preferred propionyl-CoA. The low activity observed with the purified subunits made us wonder whether we were missing another component that might be required


for effective carboxylation. At this time, a novel class of carboxylase that requires an additional  $\epsilon$ -subunit for activity was described in *S. coelicolor* (7, 8). We used the most conserved segments of the four  $\epsilon$ -subunits found in *Streptomyces* to search the mycobacterial genome. We discovered that a segment of a hypothetical protein downstream of *accD5* showed homology to the conserved region of the  $\epsilon$ -subunit of *Streptomyces*. To determine whether the native carboxylase contained an  $\epsilon$ -subunit protein, the purified native carboxylase preparation was resolved by 15% SDS-PAGE. In addition to  $\alpha 3$ ,  $\beta 4$ , and  $\beta 5$ , a 24-kDa band was observed (Fig. 1A). MALDI-TOF MS analysis following trypsin digestion revealed a sequence, VSGT-NEVSDGNE, which we identified as the product of the *Rv3281* ORF lacking 17 N-terminal amino acids. The presence of a trypsin cleavage site immediately upstream of this peptide may explain the absence of the N-terminal end. Interestingly, *Rv3281* is located immediately downstream of the ORF encoding the AccD5 ( $\beta 5$ , *Rv3280*), as shown in Fig. 1B. Since this suggested a potential role for *Rv3281* gene product in carboxylation involving the  $\beta 5$ -subunit, we examined the role of *Rv3281* in acyl-CoA carboxylation.

**$\epsilon$  Transcript and Protein Expression in *M. tuberculosis***—To test whether the putative  $\epsilon$ -subunit of *M. tuberculosis* is expressed in the

pathogen, semiquantitative RT-PCR analysis of the transcripts was done. RT-PCR data clearly showed that the ORF encoding the  $\epsilon$ -subunit is transcribed in the mycobacterial cell. RT-PCR analysis with primers spanning the  $\beta 5$ - $\epsilon$  junction yielded a product expected if both ORFs were transcribed as one unit (Fig. 2A). We assayed the  $\epsilon$ -protein expression level in cultures of *M. tuberculosis* using rabbit antisera raised against the  $\epsilon$ -protein expressed in *E. coli*. The protein level was maximal at 5 days of growth when the culture was in its exponential growth phase and declined when the culture entered late log and stationary phase (Fig. 2B).

**In Vitro Reconstitution of Carboxylase Activity with the Purified Subunits**—To examine the possible role of Rv3281 in the carboxylation reaction involving the  $\alpha 3$ -,  $\beta 4$ -, and  $\beta 5$ -subunits, we expressed  $\alpha 3$ -,  $\beta 4$ -,  $\beta 5$ -, and  $\epsilon$ -subunits as His-tagged fusion proteins in *E. coli* and purified them (Fig. 3A). Biotinylation of the expressed  $\alpha 3$ -subunit was performed with the expressed *E. coli* biotin ligase, and successful biotinylation of the  $\alpha 3$ -subunit was confirmed by Western blotting using the avidin-horseradish peroxidase conjugate (data not shown). The biotinylated  $\alpha 3$ -subunit was purified and mixed with the purified  $\beta 4$ - or  $\beta 5$ -subunit and assayed for carboxylation of acetyl-CoA or propionyl-CoA. In our initial reconstitution experiments, the expressed and purified subunits were mixed together after concentration of individual subunits. These preparations showed very low acyl-CoA carboxylase activities. When we analyzed the concentrated subunits by Superose-12 gel filtration chromatography, we found that the individual concentrated subunits were in aggregated forms; aggregation would prevent interaction with other subunits and thus hinder successful reconstitution. Attempts to mix the aggregated forms under denaturing conditions in 6 M urea followed by renaturation of the subunits by stepwise dialysis yielded extremely low carboxylase activity. Therefore, individual subunits were purified by cobalt affinity chromatography, and the dilute subunit preparations were mixed prior to concentration. The purified  $\epsilon$ -subunit in dilute form was initially combined with the dilute solution of either the  $\beta 4$ - or the  $\beta 5$ -subunits before the addition of a dilute preparation of the biotinylated  $\alpha 3$ -subunit followed by concentration by ultrafiltration. Such a reconstitution protocol yielded high levels of carboxylase activity with the  $\alpha 3$ - $\beta 5$ - $\epsilon$  when acetyl-CoA or propionyl-CoA was used as the substrate. As shown in Fig. 3B, the reconstituted  $\alpha 3$ - $\beta 5$  showed nearly 3-fold higher activity with propionyl-CoA when compared with acetyl-CoA. However, when reconstituted with the  $\epsilon$ -subunit, the  $\alpha 3$ - $\beta 5$ - $\epsilon$  gave 6.3-fold increased ACC activity. The PCC activity increased only 3.2-fold with the inclusion of the  $\epsilon$ -subunit. The reconstituted  $\alpha 3$ - $\beta 4$  and  $\alpha 3$ - $\beta 4$ - $\epsilon$  showed very low carboxylase activity with either substrate.  $\alpha 3$ - $\beta 4$  and  $\alpha 3$ - $\beta 5$  catalyzed carboxylation of long chain fatty acyl-CoA, and avidin completely inhibited this carboxylation. This carboxylation was not affected by the addition of the  $\epsilon$ -subunit (Fig. 3, B and C). These results indicate that the  $\epsilon$ -subunit interacts with only the  $\beta 5$ -subunit and modulates its carboxylase activity for acetyl-CoA and propionyl-CoA.

**Characterization of the  $\epsilon$ -Subunit (Rv3281)**—An analysis of the amino acid sequence of the  $\epsilon$ -subunit revealed a low level of identity at the C terminus with the reported  $\epsilon$ -subunit of *S. coelicolor* (7, 8). As shown in Fig. 4A, the *M. tuberculosis*  $\epsilon$ -subunit has five sets of the repeated amino acids ((N/A)EVSDGNETNNP) at the N terminus that are not found in the *S. coelicolor*  $\epsilon$ -subunits (CAC21625 and CAA19984). To determine whether the N-terminal repeated amino acids were important for activity, we cloned a truncated version of the  $\epsilon$ -subunit that contained only the C-terminal portion that showed homology with *S. coelicolor*  $\epsilon$ -subunit. When the full-length ( $\epsilon$ ) or truncated ( $\epsilon_{\text{mut}}$ ) versions of the  $\epsilon$ -subunit were used for reconstitution with



**FIGURE 3. Purification of *M. tuberculosis* acyl-CoA carboxylase subunits expressed in *E. coli* and reconstitution of acyl-CoA carboxylase activity with purified subunits.** A, each subunit was expressed in *E. coli* and purified by cobalt affinity chromatography. The  $\alpha 3$ -subunit was biotinylated using biotin ligase (pCY216) prior to purification. Lane 1, Mw, molecular weight protein marker; lane 2, purified  $\epsilon$ -subunit (full-length) from *E. coli*; lane 3, purified  $\alpha 3$ -subunit from *E. coli*; lane 4, purified  $\beta 4$ -subunit from *E. coli*; and lane 5, purified  $\beta 5$ -subunit from *E. coli*. B, influence of  $\epsilon$  on the carboxylase activities of  $\alpha 3$ - $\beta 4$  and  $\alpha 3$ - $\beta 5$  with acetyl-CoA (white bars), propionyl-CoA (black bars), and long chain acyl-CoA (shaded bars) substrates was measured under optimal conditions. C, carboxylase activities of  $\alpha 3$ - $\beta 4$  (white bars) and  $\alpha 3$ - $\beta 4$ - $\epsilon$  (black bars) with acetyl-CoA, propionyl-CoA, and long chain acyl-CoA substrates.

the  $\alpha 3$ - and  $\beta 5$ -subunits and assayed for acyl-CoA carboxylase activities with acetyl-CoA or propionyl-CoA, the truncated  $\epsilon$ -subunit showed very much less stimulation of the carboxylase activity when compared with the full-length  $\epsilon$ -subunit, indicating that N-terminal repeats in the *M. tuberculosis*  $\epsilon$ -subunit are important for carboxylase activity (Fig. 4B).






FIGURE 5. **Determination of optimal stoichiometry for reconstitution of acyl-CoA carboxylase activity.** Purified subunits were mixed in the indicated ratios prior to concentration. The molar ratios of subunits are indicated as A:B:C where A represents the ratio of the  $\alpha$ -subunit, B represents the ratio of the  $\beta$ -subunit, and C represents the ratio of the  $\epsilon$ -subunit. Acyl-CoA carboxylase activity with acetyl-CoA (white bars) and propionyl-CoA (black bars) substrates were measured after concentration. Results presented are the average of at least three independent experiments with a standard error.

$\epsilon$ -subunit increased the SC of the  $\alpha$ 3- $\beta$ 5 for acetyl-CoA by 5.5-fold, whereas the SC for propionyl-CoA was increased only 1.4-fold.

**Effect of Inhibitors on the Reconstituted Carboxylase**—Incubation of the purified native carboxylase with avidin completely inhibited the acetyl-CoA and propionyl-CoA carboxylation reactions, showing that it is a biotin-dependent acyl-CoA carboxylase (data not shown). When the purified reconstituted  $\alpha$ 3- $\beta$ 5 and  $\alpha$ 3- $\beta$ 5- $\epsilon$  were incubated with avidin, the ACC and PCC activities were also completely inhibited as expected (Fig. 8). Acetyl-CoA carboxylation by  $\alpha$ 3- $\beta$ 5 and  $\alpha$ 3- $\beta$ 5- $\epsilon$  was inhibited by 90% with 100  $\mu$ M diclofop when compared with the solvent ( $\text{Me}_2\text{SO}$ ) control. In contrast, propionyl-CoA carboxylase activity showed only 30% inhibition (Fig. 8).

## DISCUSSION

Malonyl-CoA and methylmalonyl-CoA are products of the carboxylation of acetyl-CoA and propionyl-CoA, respectively, by an acyl-CoA carboxylase. Such acyl-CoA carboxylases, in most organisms, are composed of the  $\alpha$ -subunit, which contains the BC and BCCP domains and the  $\beta$ -subunit, which contains the CT domain. In the *M. tuberculosis* genome, where the genes involved in lipid metabolism are abundant, there are three genes encoding the putative  $\alpha$ -subunits and six genes encoding the putative  $\beta$ -subunits of the acyl-CoA carboxylases (6). It is not known which ORFs encode the subunits of the native acyl-CoA carboxylase. We identified the three major proteins in our avidin-purified *M. tuberculosis* acyl-CoA carboxylase preparation as the  $\alpha$ 3-,  $\beta$ 4-, and  $\beta$ 5-subunits by N-terminal sequencing. Recently, a new group of acyl-CoA carboxylases was discovered in *S. coelicolor*, which contained a third type of subunit designated as  $\epsilon$  in addition to the  $\alpha$ - and  $\beta$ -subunits (7, 8). Adjacent to the ORF encoding the  $\beta$ 5-subunit, an ORF that has not been annotated in the mycobacterial genome (*Rv3281*) was found, the product of which showed homology with the recently identified  $\epsilon$ -subunit of *Streptomyces*. We also identified a novel 24-kDa protein in the native carboxylase preparation and determined that it was the product of *Rv3281*.

To understand the nature of the biotin-dependent enzymes, we analyzed the amino acid sequence similarity between each similarly functioning domain of the *M. tuberculosis* carboxylase with other reported




FIGURE 6. **Time course of acyl-CoA carboxylase activity with the reconstituted subunits.** A, time dependence of ACC measured with  $\alpha$ 3- $\beta$ 5 ( $\diamond$ ) and  $\alpha$ 3- $\beta$ 5- $\epsilon$  ( $\blacklozenge$ ). B, time dependence of PCC measured with  $\alpha$ 3- $\beta$ 5 ( $\diamond$ ) and  $\alpha$ 3- $\beta$ 5- $\epsilon$  ( $\blacklozenge$ ). Results presented are the average of at least three independent experiments with standard error.

ACC and PCC protein sequences. As the biotin-dependent enzymes share similar reaction mechanisms, these enzymes also share a high degree of amino acid sequence similarity (data not shown). As shown in Fig. 1B, the domain organization of *M. tuberculosis* ACC/PCC is similar to that of the PCC and ACC of *S. coelicolor* (7, 8). The first half of *M. tuberculosis* AccA3 is composed of the BC that catalyzes the ATP-driven carboxylation of biotin to form carboxybiotin, and the amino acid sequence displays a high degree of homology with the N-terminal domain of the  $\alpha$ -subunit (AccA2) of ACC and PCC from *S. coelicolor*. The other half of *M. tuberculosis* AccA3 is the BCCP domain that is found in all biotin-dependent enzymes, and this domain functions in transferring  $\text{CO}_2$  from one subsite to another, allowing carboxylation, and the amino acid sequence shows high homology with the C-terminal domain of the  $\alpha$ -subunit (AccA2) of ACC and PCC from *S. coelicolor*. The amino acid sequence of *M. tuberculosis* AccA3 showed 68% identity with AccA2 of *S. coelicolor*. The *M. tuberculosis* AccD5 is most similar to the *S. coelicolor* PccB (67% identity), and the *M. tuberculosis* AccD4 also has similarity to PccB (49% identity) and AccB (44% identity) of *S. coelicolor*.

There are significant differences between the *M. tuberculosis*  $\epsilon$ -subunit and the *S. coelicolor*  $\epsilon$ -subunits. The *M. tuberculosis* genome contains only one ORF encoding an  $\epsilon$ -subunit, whereas in the case of *S. coelicolor*, there are two  $\epsilon$ -subunits, AccE and PccE, which interact specifically with the AccB and PccB, respectively. The *M. tuberculosis*  $\epsilon$ -subunit is larger (177 amino acids) than its counterparts in *S. coelicolor* (65–69 amino acids) and contains a set of five tandem repeats of 12



FIGURE 7. Substrate concentration dependence of acetyl-CoA and propionyl-CoA carboxylase activities of  $\alpha 3$ - $\beta 5$  and  $\alpha 3$ - $\beta 5$ - $\epsilon$ . Acetyl-CoA (A), propionyl-CoA (B), bicarbonate (C), ATP (D), and  $MgCl_2$  (E) dependence of the carboxylation was measured with reconstituted  $\alpha 3$ - $\beta 5$  and  $\alpha 3$ - $\beta 5$ - $\epsilon$ .  $\diamond$  and  $\blacklozenge$ , ACC activity of  $\alpha 3$ - $\beta 5$  ( $\diamond$ ) and  $\alpha 3$ - $\beta 5$ - $\epsilon$  ( $\blacklozenge$ ).  $\blacktriangledown$  and  $\blacktriangle$ , PCC activity of  $\alpha 3$ - $\beta 5$  ( $\blacktriangledown$ ) and  $\alpha 3$ - $\beta 5$ - $\epsilon$  ( $\blacktriangle$ ). Results presented are the average of at least three independent experiments with a standard error.




TABLE 2 Kinetic parameters for purified and reconstituted acyl-CoA carboxylase subunits of *M. tuberculosis* expressed in *E. coli*

$K_m$  ( $\mu M$ ),  $V_{max}$  (pmol/mg/min), and SC ( $V_{max}/K_m$ ) are shown.

Substrates	Complexes						
	$\alpha 3\beta 5$			$\alpha 3\beta 5\epsilon$			
	$K_m$	$V_{max}$	SC	$K_m$	$V_{max}$	SC	
ACC	Acetyl-CoA	232.0	302.9	1.3	377.7	2709.0	7.2
	Bicarbonate	584.4	438.5	0.8	631.0	2669.0	4.2
	ATP	75.2	339.3	4.5	289.8	2431.0	8.4
PCC	Propionyl-CoA	105.4	1095.0	10.4	233.0	3442.0	14.8
	Bicarbonate	1752.0	1493.0	0.9	545.6	3802.0	7.0
	ATP	120.7	1099.0	9.1	215.3	3923.0	18.2

amino acids each at the N terminus, which is not seen in the *S. coelicolor*  $\epsilon$ -subunits. We determined that the N-terminal tandem repeats were important for activity and probably stabilize the protein-protein interactions between the  $\alpha 3$ -,  $\beta 5$ -, and  $\epsilon$ -subunits. Although the  $\alpha 3$ - $\beta 5$ - $\epsilon$  of *M. tuberculosis* belongs to the new group of acyl-CoA carboxylases found in *S. coelicolor*, there were substantial differences in the kinetic parameters between the two. The SC of the *M. tuberculosis*  $\alpha 3$ - $\beta 5$ - $\epsilon$  for acetyl-CoA was nearly 2-fold lower than the value for propionyl-CoA (Table 2). In contrast, the SC of the *S. coelicolor* ACC for acetyl-CoA

was nearly 3-fold lower than that of the PCC for propionyl-CoA, and no activity was detected with the PCC when acetyl-CoA was used as substrate (8). Furthermore, the reconstituted *M. tuberculosis* acyl-CoA carboxylase yielded the highest activity when the ratio of the  $\alpha 3$ -,  $\beta 5$ -, and  $\epsilon$ -subunits was 1:1:2, whereas in the case of *S. coelicolor*, near maximal velocity was obtained when the ratio of the subunits was 1:1:3 for AccE and 1:1:9 for PccE (8).

Mutagenesis studies on related bacteria suggest that *accD4* is involved in carboxylation of long chain acids required for mycolic acid

synthesis, consistent with the finding that it was an essential gene in *M. smegmatis* and *M. tuberculosis* (15–17). In agreement with these reports, the reconstituted  $\alpha 3$ - $\beta 4$  of *M. tuberculosis* carboxylated long chain fatty acyl-CoA in an  $\epsilon$ -independent manner. The mycobacterial cell wall is rich in unusual lipids that are known to play critical roles in pathogenesis. Methylmalonyl-CoA is the unique precursor of the multiple methyl-branched lipids that have been shown to function as virulence factors, and its synthesis is a potential drug target. It is expected that the mycobacterium may require higher levels of *n*-fatty acids dur-

ing its exponential growth phase and, as it enters the stationary phase, higher levels of branched fatty acids may be required for modifications of the cell wall. Our data indicated that expression of the  $\epsilon$ -protein in the mycobacterial cell reaches its highest level during the exponential growth phase and decreases during the stationary phase. During the exponential phase, when *n*-acyl lipids are synthesized from malonyl-CoA, the ACC activity of the carboxylase needs to be increased, and that may be done by increased levels of the  $\epsilon$ -subunit, which appears to favor the ACC activity. When branched lipids are required during the stationary phase, the PCC activity of the carboxylase may be favored by decreasing the expression of the  $\epsilon$ -subunit.

Commercial herbicides such as diclofop, haloxyfop, and sethoxydim are potent inhibitors of ACC in plants and have been shown to act on the CT domain (18). Our studies with the reconstituted  $\alpha 3$ - $\beta 5$ - $\epsilon$  showed that diclofop inhibited the ACC activity almost completely. Since there is no report on the inhibition of PCC activity with such inhibitors, we examined whether the PCC activity of our reconstituted carboxylase was sensitive to inhibition by diclofop. We found that the PCC activity of the  $\alpha 3$ - $\beta 5$ - $\epsilon$  was inhibited by a much lesser degree by diclofop, suggesting that the inhibitor may distinguish the binding sites for acetyl-CoA and propionyl-CoA on the CT domain of the  $\beta 5$ -subunit. Further studies with inhibitors of the  $\alpha 3$ - $\beta 5$ - $\epsilon$  may provide leads for drug development.

*M. tuberculosis* and *M. bovis* have three  $\alpha$ -subunit genes and six  $\beta$ -subunit genes, but *Mycobacterium leprae*, which is known to have lost a substantial part of the original mycobacterial genome, has one  $\alpha$ -subunit gene, three  $\beta$ -subunit genes, and one  $\epsilon$ -subunit gene (19). The organization of the ORFs for the  $\alpha 3$ -,  $\beta 5$ -, and  $\epsilon$ -subunits and that of the biotin ligase (*birA*) in the genomes of *M. tuberculosis* H37Rv, *M. tuberculosis* CDC1551, *M. bovis*, *M. leprae*, and *M. smegmatis* is nearly identical in all these organisms and, in each case, the  $\epsilon$ -subunit is





FIGURE 8. Effect of inhibitors on reconstituted carboxylases. Inhibition of acyl-CoA carboxylase activity was measured by preincubating the purified  $\alpha 3$ - $\beta 5$  and  $\alpha 3$ - $\beta 5$ - $\epsilon$  with avidin (1 unit) and diclofop (100  $\mu$ M). Activity obtained under optimal substrate concentrations is represented as mean  $\pm$  S.D. from three independent measurements. ACC activity was measured in  $\alpha 3$ - $\beta 5$  (white bars) and  $\alpha 3$ - $\beta 5$ - $\epsilon$  (light shaded bars), and PCC activity was measured in  $\alpha 3$ - $\beta 5$  (dark shaded bars) and  $\alpha 3$ - $\beta 5$ - $\epsilon$  (black bars).

FIGURE 9. Organization of the carboxylase gene cluster in *M. tuberculosis* H37Rv and other mycobacterial species. The percentages of amino acid (aa) identity of *M. tuberculosis* H37Rv subunits with other respective mycobacterial subunits are indicated in bold. The sizes in amino acids are given in regular numerals above the respective subunit.



## A Novel $\epsilon$ -Subunit in Mycobacterial Acyl-CoA Carboxylase

located immediately downstream of the  $\beta$ 5-subunit (Fig. 9). This conservation indicates that the  $\epsilon$ -subunit probably plays an important role in the carboxylation reaction in all these species. Therefore, further study of its role may lead to deeper understanding of the acyl-CoA carboxylases in mycobacteria.

*Acknowledgment*—We thank Prof. John E. Cronan, Jr. for kindly providing us the pCY216 plasmid.

### REFERENCES

1. Cronan, J. E., Jr., and Waldrop, G. L. (2002) *Prog. Lipid Res.* **41**, 407–435
2. Attwood, P. V., and Wallace, J. C. (2002) *Acc. Chem. Res.* **35**, 113–120
3. Rainwater, D. L., and Kolattukudy, P. E. (1982) *J. Bacteriol.* **151**, 905–911
4. Henrikson, K. P., and Allen, S. H. G. (1979) *J. Biol. Chem.* **254**, 5888–5891
5. Hasse, F. C., Henrikson, K. P., Treble, D. H., and Allen, S. H. G. (1982) *J. Biol. Chem.* **257**, 11994–11999
6. Cole, S. T., Brosch, R., Parkhill, J., Garnier, T., Churcher, C., Harris, D., Gordon, S. V., Eiglmeier, K., Gas, S., Barry, C. E., III, Tekaia, F., Badcock, K., Basham, D., Brown, D., Chillingworth, T., Connor, R., Davies, R., Devlin, K., Feltwell, T., Gentles, S., Hamlin, N., Holroyd, S., Hornsby, T., Jagels, K., Krogh, A., McLean, J., Moule, S., Murphy, L., Oliver, K., Osborne, J., Quail, M. A., Rajandream, M. A., Rogers, J., Rutter, S., Seeger, K., Skelton, J., Squares, R., Squares, S., Sulston, J. E., Taylor, K., Whitehead, S., and Barrell, B. G. (1998) *Nature* **393**, 537–544
7. Rodríguez, E., Banchio, C., Diacovich, L., Bibb, M. J., and Gramajo, H. (2001) *Appl. Environ. Microbiol.* **67**, 4166–4176
8. Diacovich, L., Peirú, S., Kurth, D., Rodríguez, E., Podestá, F., Khosla, C., and Gramajo, H. (2002) *J. Biol. Chem.* **277**, 31228–31236
9. Sambrook, J., Fritsch, E. F., and Maniatis, T. (1989) *Molecular Cloning: A Laboratory Manual*, 2nd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY
10. Studier, F. W., and Moffatt, B. A. (1986) *J. Mol. Biol.* **189**, 113–130
11. Chapman-Smith, A., Turner, D. L., Cronan, J. E., Jr., Morris, T. W., and Wallace, J. (1994) *Biochem. J.* **302**, 881–887
12. Laemmli, U. K. (1970) *Nature* **227**, 680–685
13. Bradford, M. M. (1976) *Anal. Biochem.* **72**, 248–254
14. Hunaiti, A. R., and Kolattukudy, P. E. (1982) *Arch. Biochem. Biophys.* **216**, 362–371
15. Portevin, D., de Sousa-D'Auria, C., Montrozier, H., Houssin, C., Stella, A., Laneelle, M. A., Bardou, F., Guilhot, C., and Daffe, M. (2005) *J. Biol. Chem.* **280**, 8862–8874
16. Gande, R., Gibson, K. J., Brown, A. K., Krumbach, K., Dover, L. G., Sahm, H., Shioyama, S., Oikawa, T., Besra, G. S., and Eggeling, L. (2004) *J. Biol. Chem.* **279**, 44847–44857
17. Sassetti, C. M., Boyd, D. H., and Rubin, E. J. (2003) *Mol. Microbiol.* **48**, 77–84
18. Zhang, H., Tweel, B., and Tong, L. (2004) *Proc. Natl. Acad. Sci.* **101**, 5910–5915
19. Cole, S. T., Eiglmeier, K., Parkhill, J., James, K. D., Thomson, N. R., Wheeler, P. R., Honore, N., Garnier, T., Churcher, C., Harris, D., Mungall, K., Basham, D., Brown, D., Chillingworth, T., Connor, R., Davies, R. M., Devlin, K., Duthoy, S., Feltwell, T., Fraser, A., Hamlin, N., Holroyd, S., Hornsby, T., Jagels, K., Lacroix, C., Maclean, J., Moule, S., Murphy, L., Oliver, K., Quail, M. A., Rajandream, M. A., Rutherford, K. M., Rutter, S., Seeger, K., Simon, S., Simmonds, M., Skelton, J., Squares, R., Squares, S., Stevens, K., Taylor, K., Whitehead, S., Woodward, J. R., and Barrell, B. G. (2001) *Nature* **409**, 1007–1011

**Identification and Characterization of Rv3281 as a Novel Subunit of a Biotin-dependent Acyl-CoA Carboxylase in *Mycobacterium tuberculosis* H37Rv**  
Tae-Jin Oh, Jaiyanth Daniel, Hwa-Jung Kim, Tatiana D. Sirakova and Pappachan E. Kolattukudy

*J. Biol. Chem.* 2006, 281:3899-3908.

doi: 10.1074/jbc.M511761200 originally published online December 14, 2005

---

Access the most updated version of this article at doi: [10.1074/jbc.M511761200](https://doi.org/10.1074/jbc.M511761200)

Alerts:

- [When this article is cited](#)
- [When a correction for this article is posted](#)

[Click here](#) to choose from all of JBC's e-mail alerts

This article cites 17 references, 8 of which can be accessed free at <http://www.jbc.org/content/281/7/3899.full.html#ref-list-1>