



---

Proceedings of the Second Russia–China International Meeting on the  
Central Asian Orogenic Belt (September 6–12, 2017, Irkutsk, Russia)

---

## EVOLUTION OF THE NORTHERN ALXA BLOCK IN THE PALEOZOIC: CONSTRAINTS FROM GEOCHRONOLOGY, GEOCHEMICAL CHARACTERISTICS AND ZIRCON HF ISOTOPES OF GRANITOIDS

Lei Zhang<sup>1</sup>, Tao Wang<sup>1</sup>, Jian-jun Zhang<sup>1</sup>, Xing-jun Shi<sup>2</sup>, Ying Tong<sup>1</sup>

<sup>1</sup>*Institute of Geology, Chinese Academy of Geological Sciences, Beijing 100037, China*

<sup>2</sup>*Faculty of City and Environmental Science, Xinyang Normal University, Xinyang 464000, China*

---

**For citation:** Zhang L., Wang T., Zhang J.-J., Shi X.-J., Tong Y., 2017. Evolution of the northern Alxa block in the Paleozoic: constraints from geochronology, geochemical characteristics and zircon Hf isotopes of granitoids. *Geodynamics & Tectonophysics* 8 (3), 613–614. doi:10.5800/GT-2017-8-3-0307.

---

The Alxa block is situated to the south of the CAOB, situated to the east of the Tarim block and west of the NCC. Voluminous intrusive and extrusive rocks outcrop in the northern Alxa block and adjacent southern CAOB. Most of them are thought to be related to the closure of the Paleo-Asia Ocean and subsequent collision [Wu, 1993; Wu *et al.*, 1998; Zhang *et al.*, 2013; Dan *et al.*, 2016].

Traditionally, the northern Alxa block is proposed to consist of two tectono-magmatic belts, the Zongnaishan-Shalazhashan belt to the north (the north belt) and the Yabulai-Nurugong-Honggueryulin belt to the south (the south belt) (e.g. [Wang *et al.*, 1992, 1994; Wu,

1993; Zheng *et al.*, 2014]). Based on previous studies and this study, zircon Hf isotope of granitoids in the north belt are generally depleted ( $\epsilon\text{Hf}(t)$  is higher than CHUR) and the zircon Hf isotope of granitoids in the south belt are more enriched ( $\epsilon\text{Hf}(t)$  is negative). This indicates that the granitoids in the two belts have different sources: the source for the north belt granitoids is generally juvenile and the source for the south belt granitoids consists of more old materials. Therefore, the north belt could not be a part of the Alxa block [Shi *et al.*, 2014a, 2014b; Zhang *et al.*, 2015].

According to the newly obtained geochronological data, most granitoids in the north belt are generated in

the Late Carboniferous to the beginning of the Triassic (e.g. [Zhang et al., 2013; Shi et al., 2014a, 2014b]), and granitoids in the south belt formed from the late stage of Early Paleozoic to the Late Permian (e.g. [Li, 2006; Dan et al., 2016]). It is noted that the a few Late Carboniferous to Early Permian Amushan Formation occurred

in both belts. Possibly, subduction of the ocean plate between the two belts could start before the late stage of Early Paleozoic and the closure of the ocean could be finished before the Late Carboniferous. The pervasive granitoids in this area would be formed in a collisional setting.

## REFERENCES

- Dan W., Li X.H., Wang Q., Wang X.C., Wyman D.A., Liu Y., 2016. Phanerozoic amalgamation of the Alxa block and North China craton: Evidence from Paleozoic granitoids, U-Pb geochronology and Sr-Nd-Pb-Hf-O isotope geochemistry. *Gondwana Research* 32, 105–121. <https://doi.org/10.1016/j.gr.2015.02.011>.
- Li J.J., 2006. Regional metallogenic system of Alashan Block in the Inner Mongolia Autonomous region. A Dissertation Submitted to China University of Geoscience for Doctoral Degree (in Chinese with English Abstract).
- Shi X.J., Wang T., Zhang L., Castro A., Xiao X.C., Tong Y., Zhang J.J., Guo L., Yang Q.D., 2014a. Timing, petrogenesis and tectonic setting of the Late Paleozoic gabbro–granodiorite–granite intrusions in the Shalazhashan of northern Alxa: Constraints on the southernmost boundary of the Central Asian Orogenic Belt. *Lithos* 208–209, 158–177. <https://doi.org/10.1016/j.lithos.2014.08.024>.
- Shi X.J., Zhang L., Wang T., Xiao X.C., Tong Y., Zhang J.J., Geng J.Z., Ye K., 2014b. Geochronology and geochemistry of the intermediate-acid intrusive rocks from Zongnaishan area in northern Alxa, Inner Mongolia, and their tectonic implications. *Acta Petrologica et Mineralogica* 33 (6), 989–1007 (in Chinese with English abstract).
- Wang T.Y., Wang J.R., Wang S.Z., 1992. The identification of the Engerwusu mélangé in northern Alxa area and its tectonic implications. *Journal of Lanzhou University (Natural Sciences)* 28 (2), 194–196 (in Chinese).
- Wang T.Y., Wang S.Z., Wang J.R., 1994. Formation and Evolution of the Paleozoic Continental Crust in the Alxa Area. Lanzhou University Press, Lanzhou, 213 p. (in Chinese).
- Wu T.R., 1993. Tectonic units and their fundamental characteristics on the northern margin of the Alxa block. *Acta Geologica Sinica* 67, 97–108 (in Chinese with English abstract).
- Wu T.R., He G.Q., Zhang C., 1998. On Paleozoic tectonics in the Alxa region. *Acta Geologica Sinica* 72 (3), 256–263.
- Zhang J.J., Wang T., Zhang L., Tong Y., Shi X.J., Guo L., Zhang Z.C., Yang Q.D., Huang W., Zhao J.X., Ye K., 2015. Tracking deep crust by zircon xenocrysts within igneous rocks from the northern Alxa region, China: Constraint on the southernmost boundary of the Central Asian Orogenic Belt. *Journal of Asian Earth Sciences* 108, 150–169. <https://doi.org/10.1016/j.jseaes.2015.04.019>.
- Zhang W., Wu T.R., Feng J.C., Zheng R.G., He Y.K., 2013. Time constraints for the closing of the Paleo-Asian ocean in the Northern Alxa region: evidence from Wuliji granites. *Science China Earth Sciences* 56 (1), 153–164. <https://doi.org/10.1007/s11430-012-4435-y>.
- Zheng R.G., Wu T.R., Zhang W., Xu C., Meng Q.P., Zhang Z.Y., 2014. Late Paleozoic subduction system in the northern margin of the Alxa block, Altaids: Geochronological and geochemical evidences from ophiolites. *Gondwana Research* 25 (2), 842–858. <https://doi.org/10.1016/j.gr.2013.05.011>.