

Systematic and zoogeographical characteristics of the oribatid mite fauna (Acari: Oribatida) of Vietnam

Đặc điểm cấu trúc phân loại và địa động vật của khu hệ ve giáp (Acari: Oribatifa) ở Việt Nam

Research article

Vu Quang Manh^{1*}, Dao Duy Trinh², Nguyen Hai Tien³, Nguyen Huy Tri¹, Lai Thu Hien¹, Ha Tra My¹, Do Thi Duyen^o

¹Hanoi National University of Education, 136 Xuan Thuy, Hanoi, Vietnam; ²Vietnam National University Hanoi, 144 Xuan Thuy, Hanoi, Vietnam; ³Thai Binh University of Medicine and Pharmacy, 373 Ly Bon, Thai Binh, Vietnam; ^oInstitute of Ecology and Bioresources, Hanoi, Vietnam

The article is a synthesis of the studies on oribatid mites carried out in Vietnam during the period of 1980-2013, and is based on the oribatid materials obtained throughout the country. The oribatid mite fauna (Acari: Oribatida) of Vietnam is diversified, and has high specialization. It is diverse by the number of superfamilies, families, genera and species recorded. However, the number of genera per family, as well as the number of species and subspecies per genus, is not high. 43.75% and 37.50% of the total 64 families and subfamilies consist of one and of 2-3 genera, respectively. The only one family Oppiidae Grandjean, 1954 consists of 23 genera. The majority of the genera, 68.10% of the total, are represented by one species. The only two genera are represented by more than 10 species, namely *Galumna* Heyden, 1826 and *Pergalumna* Grandjean, 1936, with 13 and 11 species, respectively. The main zoogeographical characteristics of the oribatid mite fauna of Vietnam are the Oriental species, representing 60.30% of the total number. It also includes the elements of the Palaearctic – Oriental (12.2%), the Cosmopolitan (10.6%), the Afrotropical (Ethiropical) – Oriental (6.9%), the Australian – Oriental (5.0%), the Neotropical – Oriental (3.8%), the Nearctic – Oriental (0.9%), and the Pacific – Oriental (0.3%).

Để đánh giá cấu trúc phân loại và đặc điểm địa động vật của khu hệ ve giáp Việt Nam (Acari: Oribatida), trên cơ sở mẫu vật nghiên cứu thu từ toàn lãnh thổ quốc gia, công trình đã tổng hợp và phân tích các kết quả nghiên cứu về ve giáp trong giai đoạn 1980-2013. Khu hệ động vật ve giáp Việt Nam có tính chuyên biệt cao, và rất đa dạng về số lượng họ, giống và loài xác định được. Tuy nhiên số lượng giống trong 1 họ, cũng như số lượng loài trong 1 giống lại không cao. 43,75% và 37,50% của 64 họ và phân họ, tương ứng chỉ xác định được có 2 và 3 giống. Duy nhất có họ Oppiidae Grandjean, 1954 ghi nhận được 23 giống. 68,10% tổng số giống, chỉ xác định được 1 loài. Duy nhất có 2 giống ghi nhận được hơn 10 loài, là Galumna Heyden, 1826 và Pergalumna Grandjean, 1936, tương ứng có 13 và 11 loài. Đặc điểm địa động vật cơ bản của khu hệ ve giáp Việt Nam là tính chất Đông phương (Oriental), với 60,30% tổng số loài xác định được. Tính chất địa động vật của nó còn bao gồm các yếu tố sau: Cổ bắc - Đông phương (Palaearctic-Oriental, 12,2% tổng số loài xác định được), Toàn cầu (Cosmopolite, 10,6%), Nhiệt đới Phi châu - Đông phương (Afrotropical (Ethiropical)-Oriental, 6,9%), Úc châu - Đông phương (Australian-Oriental, 5,0%), Tân nhiệt đới - Đông phương (Neotropical-Oriental, 3,8%), Vùng cực - Đông phương (Nearctic-Oriental, 0,9%), và Thái Bình Dương - Đông phương (Pacific-Oriental, 0,3%).

Keywords: oribatida, systematic, zoogeography, Vietnam

1. INTRODUCTION

Oribatid mites (Acarina: Oribatida) are small arachnids (Microarthropoda), comprising the main component of microarthropod populations in the soil ecosystem, and are an important component in all biological processes taking place in soil. They disperse bacteria and fungi, both externally on their body surface, or by feeding, with subsequent survival of spores during passage through their alimentary tracts. Others are intermediate hosts for important tapeworm parasites (Cestoda) of animals (Ghilarov & Krivolutsky 1975, Behan-Pelletier 1999) [4, 13].

The study on oribatid mites of Vietnam started in 1967 by Hungarian oribatologists (Balogh & Mahunka 1967) [3]. It continued in the 1980's, and has obtained the important results (Golosova 1983 & 1984, Vu Quang Manh 1985, Jeleva & Vu 1987, Mahunka 1987, 1988 & 1989) [14, 16, 17, 18, 28]. However, because of the diverse fauna of the tropical soil ecosystem of Vietnam the studies on oribatid mites of Vietnam are still not enough sufficient (Krivolutsky et al. 1997, Vu & Nguyen 2000, Vu Quang Manh 2007, 2012 & 2015, Dao Duy Trinh et al. 2010, Vu et al. 2010, Ermilov et al. 2011, Ermilov & Vu 2012, Nguyen Hai Tien & Vu Quang Manh 2012, Nguyen Huy Tri et al. 2014, Fernandez et al. 2014 & 2015) [5, 9, 11, 12, 15, 20, 30, 31, 32, 34, 35, 36].

In recent years, the Russian colleague Sergey G. Ermilov and his collaborators have made a significant contributions to the knowledge of the oribatid mite fauna of Southeast and Southwest of Vietnam. However, many of their oribatid specimens have been obtained by unofficial lines, and Ermilov rarely collected materials by himself in the field. Their data on the geographical locations and natural conditions of Vietnam are founded on incorrect and sometime on illegal sources. In particular, the administrative maps of the socialist republic of Vietnam in their publications are introduced fully false, lacking islands and the island regions of the Vietnam (Ermilov et al. 2012, Ermilov & Anichkin 2014, Ermilov 2015, Minor & Ermilov 2015) [7, 8, 10, 19]. Therefore, the data on Vietnam's oribatid mites given by the Russian colleague Sergey G. Ermilov and his coauthors must be checked and revised carefully, before consulting them. Even so, although lacking of knowledge and right source of Vietnam's nature, this is an encouraging attempt by Russian colleague on study of oribatid mites of Vietnam, and it showed up the importance of the soil oribatid mites of Vietnam.

This article is a synthesis of the studies on oribatid mites carried out in Vietnam during the period of 1980-2013, and is based on the oribatid materials obtained throughout the country. Its aim is to characterize the systematic and zoogeographical characteristics of the oribatid mite fauna of Vietnam.

2. MATERIALS AND METHODS

2.1 Study area

During the period of 1980-2013, oribatid specimens obtained throughout Vietnam, including fifty sites located in

27 provinces, and from all of the 8 natural geographical regions of Vietnam, namely Northwest, Northeast, Red River Delta, North Central Coast, South Central Coast, Central Highlands, Southeast, and Mekong River Delta (Vu, Tailard 1994) [37]. Fifty study sites were classified into six main types, as follows: Natural forest, Human-disturbed forest, Grassland and scrub, Grassland, Cultivated land with perennial and annual plants, and Agricultural land with annual plants (Vietnam Ministry of Agriculture and Rural Development 2006) [26]. The study soils were arranged in a six main groups: Coastal saline-acid soil, Acid alluvial soil, Neutral alluvial soil, Ferralitic reddish brown soil, Ferralitic brownish soil derived from limestone, Reddish brown soils derived from basic and intermediate magmatic rocks (Vietnam National Institute for Soils and Fertilizers 2002) [27].

2.2 Sampling and extraction

Soil samples were taken according to soil vertical layers, as follows: Forest litter, and soil vertical layers of 0-10cm, 10-20cm, 20-30cm. Soil samples were taken by rectangular metal sampler, with (0,25m²) surface area, 20 cm in depth and with both ends open. Forest litter samples were taken from the total forest litter covered an area of (0,25m²). Modifications of Berlese-Tullgren funnels were used for extraction of oribatid mites from the obtained materials. An extraction lasted seven days in the laboratory at normal air condition of 25-30°C. Extracted oribatid mites were preserved in 70° ethanol, sorted and counted (Edwards 1991) [6].

2.3. Identification and data analysis

Oribatida materials are identified mainly after Balogh (1992), Gilyarov & Krivolutsky (1975), Balogh & Balogh (1988, 1989, 2002), Norton & Behan-Pelletier (2009), Schatz et al. (2011), and Subias (2013) [1, 2, 13, 22, 23, 24]. A zoogeographical characteristic of the oribatid mite fauna of Vietnam is analyzed according to the World zoogeographical regions, consisting NE. Nearctic, PA. Palaearctic, NO. Neotropical, AF. Afrotropical (Ethiopian), OR. Oriental, AU. Australian, PC. Pacific, and AN (Illies 1974, Udvardy 1975) [25].

3. RESULT AND DISCUSSION

3.1. Species biodiversity

Table 1 presents the species diversity and an analysis of a systematic characteristics of the oribatid mite fauna of Vietnam, consulting to classifications of Gilyarov & Krivolutsky (1975), Balogh & Balogh (2002), Norton & Behan-Pelletier (2009), Schatz et al. (2011), and Subias (2013) [2, 13, 22, 23, 24]. In the table introduced are also a number of superfamilies per order, of families per superfamily, of genera per family, as well as a number of species per genus.

Table 1. Systematic characteristics of the oribatid mite fauna of Vietnam

No	Infraorder	Superfamily	Family	Genus	Species & Sub-species		
I	ENARTHRO TA Grandjean, 1947	1.HYPOCHTHONOID EA Berlese, 1910	1. Hypochthoni- idae Berlese, 1910	1. <i>Malacoangelia</i> Berlese, 1913	1		
				2. <i>Eohypochthonius</i> Jacot, 1938	2		
				3. <i>Eniochthonius</i> Grandjean, 1933	1		
				4. <i>Cosmochthonius</i> Berlese, 1910	1		
			2. PROTHOPLOPHORI DEA Ewing, 1917	2. Cosmochthoni- idae Grandjean, 1947	5. <i>Sphaerochthonius</i> Berlese, 1910	1	
				3. Sphaerochtho- niidae Grandjean, 1947	6. <i>Apoplophora</i> Aoki, 1980	2	
				4. Prothoplophori- dae Ewing, 1917	7. <i>Arthrophlophora</i> Berlese, 1910	1	
		II	MIXONOMATA Grandjean, 1969	3. PHTHIRACAROIDEA Perty, 1841	5. Steganacaridae Niedbala, 1986	8. <i>Arphthiracarus</i> Niedbala, 1994	1
						9. <i>Atropacarus</i> Ewing, 1917	3
						10. <i>Austrophthiracarus</i> Balogh et Mahunka, 1978	1
						11. <i>Plonaphacarus</i> Niedbala, 1986	2
	12. <i>Hoplophorella</i> Berlese, 1923				4		
	13. <i>Phthiracarus</i> Perty, 1841				1		
	14. <i>Indotritia</i> Jacot, 1929				1		
	15. <i>Oribotritia</i> Jacot, 1924				1		
	16. <i>Sabahtritia</i> Mahunka, 1987				1		
	17. <i>Rhysotritia</i> Markel & Meyer, 1959				6		
	18. <i>Haplacarus</i> Wallwork, 1962				1		
	19. <i>Javacarus</i> Balogh, 1961			1			
	20. <i>Lohmannia</i> Michael, 1898			1			
	21. <i>Meristacarus</i> Grandjean, 1934			1			
	22. <i>Mixacarus</i> Balogh, 1958			2			
	23. <i>Papilacarus</i> Kunst, 1959			8			
	24. <i>Vepracarus</i> Aoki, 1965			1			
	4. EUPHTHIRACAROID EA Jacot, 1930			7. Oribotritiidae Grandjean, 1954	25. <i>Trimalaconothrus</i> Berlese, 1916	1	
				8. Euphthiracari- dae Jacot, 1930	26. <i>Nothrus</i> Koch, 1836	4	
		9. Lohmanniidae Berlese, 1916	27. <i>Afronothrus</i> Wallwork, 1961	1			
	5. LOHMANNIOIDEA Berlese 1916	10. Malaconothri- dae Berlese, 1916	28. <i>Allonothrus</i> Hammer, 1953	1			
		11. Nothridae Ber- lese, 1896	29. <i>Archegozetes</i> Grandjean, 1931	1			
		12. Trhypochtho- niidae Willmann, 1931					
	6. CROTONIOIDEA Thorell, 1876						

No	Infraorder	Superfamily	Family	Genus	Species & Sub-species
III	DESMONOMA TA Woolley, 1973	7.	13. Epilohmanniidae Oudemans, 1923	30. <i>Epolihmannia</i> Berlese, 1910	5
		EPILOHMANNIOIDEA Oudemans, 1923			
		8.	14. Nanhermanniidae Sellnick, 1928	31. <i>Cosmohermannia</i> Aoki & Yoshida, 1970	1
		NANHERMANNIOIDEA Sellnick, 1928			
				32. <i>Cyrthermannia</i> Balogh, 1958	1
				33. <i>Masthermannia</i> Berlese, 1913	1
				34. <i>Nanhermannia</i> Berlese, 1913	1
				35. <i>Phyllhermannia</i> Berlese, 1916	3
				36. <i>Hermannella</i> Berlese, 1908	1
				37. <i>Neolides</i> Heyden, 1826	1
				38. <i>Platyliodes</i> Berlese, 1916	1
				39. <i>Adrodamaeus</i> (Paschoal, 1984)	1
				40. <i>Hexachaetoniella</i> Paschoal, 1987	1
				41. <i>Pedrocortesella</i> Hammer, 1961	2
				42. <i>Belba</i> Heyden, 1826	1
				43. <i>Metabelba</i> Grandjean, 1936	1
		IV	BRACHYPYLIN A Hull, 1918	9. HERMANNIOIDEA Sellnick, 1928	15. Hermanniidae Sellnick, 1928
10. HERMANNIELLOIDEA Grandjean, 1934	16. Hermanniellidae Grandjean, 1934				
11. NEOLIODOIDEA Grandjean, 1954	17. Neoliodidae Grandjean, 1954				
12. GYMNODAMAEOIDEA Grandjean, 1954	18. Gymnodamaeidae Grandjean, 1954				
13. PLATEREMAEOIDEA Traegardh, 1931	19. Pedrocortesellidae Paschoal, 1987				
14. DAMAEOIDEA Berlese, 1896	20. Damaeidae Berlese, 1896				
15. CEPHEOIDEA Berlese, 1896	21. Cepheidae Berlese, 1896				
	22. Microtegeidae Balogh, 1972				
	23. Eremaeozetidae Piffel, 1972				
16. MICROZETOIDEA Grandjean, 1936	24. Microzetidae Grandjean, 1936				
				44. <i>Sphodrocephus</i> Woolley & Higgins, 1963	1
				45. <i>Microtegeus</i> Berlese, 1916	4
				46. <i>Eremaeozetes</i> Berlese, 1913	1
				47. <i>Berlesezetes</i> Mahunka, 1980	2
				48. <i>Caucasiozetes</i> Shtanchaeva, 1984	1
				49. <i>Kaszabozetes</i> Mahunka, 1988	1
				50. <i>Schalleriella</i> Balogh, 1962	1
		51. <i>Roynortonia</i> Ermilov, 2011	1		
		52. <i>Austroeremulus</i> Mahunka, 1985	1		
		53. <i>Eremulus</i> Berlese, 1908	3		
		54. <i>Mahunkana</i> Kocak and Kemal, 2008	1		
		55. <i>Fosseremus</i> Grandjean, 1954	1		
		56. <i>Gressittolus</i> Balogh, 1970	1		
		57. <i>Eremobelba</i> Berlese, 1908	6		
		58. <i>Fenestrella</i> Mahunka, 1987	1		
		59. <i>Heterobelba</i> Berlese, 1913	1		
		25. Amerobelbidae Grandjean, 1954			
		26. Eremulidae Grandjean, 1965			
		27. Damaeolidae Grandjean, 1965			
		28. Eremobelbidae Balogh, 1961			
		29. Heterobelbidae Balogh, 1961			
		17. AMEROBELBOIDEA Grandjean, 1954			

No	Infraorder	Superfamily	Family	Genus	Species & Sub-species
			30. Basilobelbidae Balogh, 1961	60. <i>Basilobelba</i> Balogh, 1958	1
	18.				
	ZETORCHESTOIDEA		31. Zetorchestidae Michael, 1898	61. <i>Zetorchestes</i> Berlese, 1888	2
	Balogh, 1961				
	19. GUSTAVIOIDEA		32. Astegistidae Balogh, 1961	62. <i>Cultroribula</i> Berlese, 1908	2
	Oudemans, 1900				
			33. Peloppiidae Balogh, 1943	63. <i>Ceratoppia</i> Berlese, 1908	1
				64. <i>Furcoppia</i> Balogh et Mahunka, 1969	2
	20. CARABODOIDEA		34. Carabodidae C.L.Koch, 1837	65. <i>Aokiella</i> Balogh et Mahunka, 1967	2
	C. L. Koch, 1837			66. <i>Austrocarabodes</i> Hammer, 1966	3
				67. <i>Chistyakovella</i> Ermilov, Aoki, Anichkin, 2013	1
				68. <i>Gibbicepheus</i> Balogh, 1958	1
				69. <i>Pentabodes</i> P.Balogh, 1984	1
				70. <i>Carabodes (Phyllocarabodes)</i> Balogh et Mahunka, 1969	1
			35. Nipobodidae Aoki, 1959	71. <i>Leobodes</i> Aoki, 1965	1
	21.				
	TECTOCEPHEOIDEA		36. Tectocephidae Grandjean, 1954	72. <i>Tectocephus</i> Berlese, 1896	2
	A Grandjean, 1954			73. <i>Tegezozetes</i> Berlese, 1913	1
			37. Otocephidae Balogh, 1961	74. <i>Otocephus (Acrotocephus)</i> Aoki, 1965	4
				75. <i>Archegotocephus (Megalotocephus)</i> Mahunka, 1988	1
				76. <i>Dolicheremaeus</i> Jacot, 1938	8
				77. <i>Eurostocephus</i> Aoki, 1965	1
	22. OPPIOIDEA		38. Eremellidae Balogh, 1961	78. <i>Eremella</i> Berlese, 1913	1
	Grandjean, 1954		39. Granuloppiidae Balogh, 1983	79. <i>Gigantoppia</i> Mahunka, 2008	1
				80. <i>Granuloppia</i> Balogh, 1958	1
			40. Oppiidae Grandjean, 1954	81. <i>Acroppia</i> Balogh, 1883	1
				82. <i>Arcoppia</i> Hammer, 1977	6
				83. <i>Belloppia</i> Hammer, 1968	1
				84. <i>Brachioppiella</i> Hammer, 1962	1
				85. <i>Congoppia</i> Balogh, 1963	1
				86. <i>Cryptoppia</i> Csiszár, 1961	1
				87. <i>Hammerella</i> Ermilov, Shtanchaeva, Subias, Anichkin, 2012	1
				88. <i>Helioppia</i> Balogh, 1983	1

No	Infraorder	Superfamily	Family	Genus	Species & Sub-species
				89. <i>Karenella</i> Hammer, 1962	1
				90. <i>Kokoppia</i> Balogh, 1983	1
				91. <i>Lanceoppia</i> Subías, 1989	1
				92. <i>Lasiobelba</i> Aoki, 1959	2
				93. <i>Lineoppia</i> J & P. Balogh, 1983	1
				94. <i>Multioppia</i> Hammer, 1961	1
				95. <i>Neoamerioppia</i> Subías, 1989	1
				96. <i>Oppia</i> Wallwork, 1961	2
				97. <i>Oppiella</i> Jacot, 1937	1
				98. <i>Oxybrachioppia</i> Subías, 1989	1
				99. <i>Pseudoamerioppia</i> Subías, 1989	1
				100. <i>Pulchroppia</i> Hammer, 1979	5
				101. <i>Ramusella</i> Hammer, 1962	3
				102. <i>Ramuselloppia</i> Subías & Rodríguez, 1986	1
				103. <i>Striatoppia</i> Balogh, 1958	3
				104. <i>Taiwanoppia</i> Tseng, 1982	1
				105. <i>Suctobelba</i> Paoli, 1908	3
		23. TRICOZETOIDEA Ewing, 1917	41. Suctobelbidae Jacot, 1938	106. <i>Suctobelbila</i> Jacot, 1937	1
				107. <i>Suctobelbella</i> Jacot, 1937	8
		24. LIMNOZETOIDEA Thor, 1937	42. Limnozetestidae Grandjean, 1954	108. <i>Limnozetes</i> Hull, 1916	1
		25. CYMBAEREMAEOIDEA Sellnick, 1928	43. Cymbaeremaeidae Sellnick, 1928	109. <i>Scapheremaeus</i> Berlese, 1910	4
V	PORONOTA Balogh J. et Balogh P. 2002 (?)	26. IDIOZETOIDEA Aoki, 1976	44. Idiozetidae Aoki, 1976	110. <i>Idiozetes</i> Aoki, 1976	1
		27. ORIPODOIDEA Jacot, 1925	45. Chaunoproctidae Balogh, 1961	111. <i>Chaunoproctus</i> Pearce, 1906	1
			46. Parakalummidae Grandjean, 1936	112. <i>Neoribates</i> Berlese, 1914	3
			47. Mochlozetidae Grandjean, 1960	113. <i>Unguizetes</i> Sellnick, 1925	4
				114. <i>Uracrobates</i> Balogh et Mahunka, 1967	1
			48. Haplozetidae Grandjean, 1936 (Xylobatidae J. Balogh et Balogh P., 1984)	115. <i>Brasilobates</i> Pérez-Íñigo y Baggio, 1980	1
				116. <i>Perxylobates</i> Hammer, 1972	6
				117. <i>Setoxylobates</i> Balogh et Mahunka, 1967	1
				118. <i>Vilhenabates</i> Balogh, 1963	1

No	Infraorder	Superfamily	Family	Genus	Species & Sub-species
				119. <i>Xylobates</i> Jacot, 1925	6
49.	Protoribatidae		Balogh Balogh J. Balogh et P. Balogh, 1984	120. <i>Protoribates</i> Berlese, 1908	3
		(1) Protoribatinae	Balogh Balogh J. Balogh et P. Balogh, 1984 (Subfamily)	121. <i>Liebstadia</i> Oudemans, 1906	1
		(2) Liebstadiinae	Balogh Balogh J. Balogh et P. Balogh, 1984 (Subfamily)		
50.	Oribatulidae		Thor, 1929	122. <i>Cordiozetes</i> Mahunka, 1983	1
				123. <i>Oribatula</i> Berlese, 1896	1
				124. <i>Sellnickia</i> Oudemans, 1927	1
				125. <i>Zygoribatula</i> Berlese, 1916	3
51.	Haplozetidae		Grandjean, 1936	126. <i>Cosmobates</i> Balogh, 1959	1
				127. <i>Indoribates</i> Willmann, 1935	2
				128. <i>Magnobates</i> Hammer, 1967	1
				129. <i>Peloribates</i> Berlese, 1908	7
				130. <i>Rostrozetes (Trachyribates)</i> Sellnick, 1925	5
52.	Schelorbitidae		Grandjean, 1953	131. <i>Euscheloribates</i> Kunst, 1958	1
				132. <i>Fijibates</i> Hammer, 1971	1
				133. <i>Ischeloribates</i> Corpuz-Raros, 1980	1
				134. <i>Nanobates</i> Balogh et Mahunka, 1980	1
				135. <i>Philoribates</i> L.A.Corpuz-Raros, 1980	1
				136. <i>Rhabdoribates</i> Aoki, 1967	1
				137. <i>Scheloribates</i> Berlese, 1908	8
				138. <i>Tuberemaeus</i> Sellnick, 1930	2
53.	Oripodidae		Jacot, 1925	139. <i>Cosmopirnodus</i> Balogh, 1970	1
				140. <i>Oripoda</i> Bank, 1904	1
				141. <i>Subpirnodus</i> Mahunka, 1988	1
				142. <i>Truncopes</i> Grandjean, 1956	1
54.	Birobatidae		J. Balogh et P. Balogh, 1984	143. <i>Brachyoripoda</i> Balogh, 1970	1
55.	Mycobatidae		Grandjean, 1954	144. <i>Allozetes</i> Berlese, 1913	1

No	Infraorder	Superfamily	Family	Genus	Species & Sub-species
		28. CERATOZETOIDEA Jacot, 1925	56. Ceratozetidae Jacot, 1925	145. <i>Ceratozetes</i> Berlese, 1908	2
				146. <i>Fuscozetes</i> Sellnick, 1928	1
			57. Austrachipteriidae Luxton, 1985	147. <i>Austrachipteria</i> Balogh et Mahunka, 1966	1
				148. <i>Lamellobates</i> Hammer, 1958	4
				149. <i>Paralamellobates</i> Bhaduri y Raychaudhuri, 1968	3
			58. Punctori- batidae Thor, 1937	150. <i>Punctoribates</i> Subías, Kahwash y Ruiz, 1990	1
		29. ORIBATELLOIDEA Jacot, 1925	59. Oribatellidae Jacot, 1925	151. <i>Novoribatella</i> Engelbrecht, 1986	1
				152. <i>Oribatella</i> Banks, 1895	3
			60. Achipteridae Thor, 1929	153. <i>Achipteria</i> Berlese, 1885	1
				154. <i>Parachipteria</i> Hammen, 1952	1
		30. GALUMNOIDEA Jacot, 1925	61. Galumnellidae Piffel, 1970	155. <i>Galumnella</i> Berlese, 1917	2
			62. Galumnidae Jacot, 1925	156. <i>Allogalumna</i> Grandjean, 1936	1
				157. <i>Dimidiogalumna</i> Engelbrecht, 1972	1
				158. <i>Galumna</i> Heyden, 1826	13
				159. <i>Globogalumna</i> P. y G. Palogh, 1990	1
				160. <i>Leptogalumna</i> Balogh, 1960	1
				161. <i>Neogalumna</i> Hammer, 1973	1
				162. <i>Pergalumna</i> Grandjean, 1936	11
				163. <i>Trichogalumna</i> Balogh, 1960	3
Total	5 infraorders	30 superfamilies	62 families (2 subfamilies)	163 genera	320 species, subspecies

Up to 2014, the oribatid mite fauna of Vietnam is represented by 320 species (including 4 subspecies), belonging to 163 genera, 62 families (not including two subfamilies), and 30 superfamilies (Balogh, Balogh 2002, Norton, Behan-Pelletier 2009, Schatz et al. 2011, and Subías 2013). Among the three hundred and twenty species (320 species, 100%) recorded from Vietnam, one hundred and twenty species (120 species, 37.50% of the total number), were described new for science. One hundred and eleven species (111 species, 34.68% of the total number), have been found only in Vietnam, and are probably endemic species. One hundred and fifty five (155) species, representing 48.44% of the total oribatid fauna, were new record for Vietnam (Table 1). After Subías (2013) [24], the World oribatid fauna comprised 10,342 species and subspecies, belonging to 1,249 genera and 163 families. In comparison with the World oribatid mite fauna the one of Vietnam occupies

3.09% (320 vs. 10,342 species), 13.05% (163 vs. 1,249 genera), and 38.03% (62 vs. 163 families) of the World fauna (Vu Quang Manh 2013, 2015 [33, 34]).

In general, the oribatid mite fauna of Vietnam is very diverse, with a high number of species probably endemic. However, these data still not completely.

3.2. Systematic characteristics

Table 2 presents the oribatid systematic structure according to the number of genera per family. Almost all of the families consist of 1-3 genera, 43.75% and 37.50%, respectively, of the total 64 families and subfamilies. The families consisting of 4-5, and of 6-10 species are few, only 10.94% and 6.25%, respectively. Only one family consists of more

than 10 species, the family Oppiidae Grandjean, 1954 consisting of 23 genera (Table 1 and 2).

Table 2. Systematic characteristics of the oribatid mite fauna according to the number of genera per family

Total	Number of genera per family				
	1	2-3	4-5	6-10	>10
64 families and subfamilies	28	24	7	4	1
100 %	43.75	37.50	10.94	6.25	1.56

Table 3 presents systematic characteristics of the oribatid mite fauna according to the number of species per genus. Almost all of the genera consist of one species, 68.10% of the 163 genera. The genera comprising of 2-3, 4-5 and 6-

10 species are few, only 18.41%, 6.13% and 6.13%, respectively, of the total of 163 genera. Only two genera comprise of more than 10 species, namely Galumna Heyden, 1826 and Pergalumna Grandjean, 1936, with 13 and 11 species, respectively (Table 1 and 3).

Table 3. Systematic characteristics of the oribatid mite fauna according to the number of species per genus

Total	Number of species per genus				
	1	2-3	4-5	6-10	>10
163 genera	111	30	10	10	2
100%	68.10	18.41	6.13	6.13	1.23

In general, according to the number of superfamilies, of families, of genera, as well as of species and subspecies recorded, the systematic characteristics of the oribatid mite fauna of Vietnam is diversified. However, the number of genera per family, as well as the number of species per genus are not high.

3.3. Zoogeographical characteristics

Table 2 presents the oribatid systematic structure according to the number of genera Vietnam is completely distributed in the in the Southeast Asia. Hammer and Wallwork (1979) considered that the source for the oribatid mite fauna of the south Pacific is probably Southeast Asia and that dispersal has occurred over seas, via island “stepping stones”. However, Vietnam is a highly interesting region from a zoogeographical aspect. Although the whole country is mainly in the Oriental region, there is distinct differentiation between northern and southern parts of the country, and even between different subportions of the northern or the southern part. Northern Vietnam is closer to the south Chinese mountain range that makes it possible for some Palaeartic elements to infiltrate into this area, while Southern Vietnam is closer to the Pacific region.

On the basis of the zoogeographical data provided by Balogh (1961, 1972), Hammer (1972), Gilyarov & Krivolutsky (1975), Hammer & Wallwork (1979), Karpinen & Krivolutsky (1982), Karpinen et al. (1986, 1987), Balogh & Balogh (1988, 1989, 2002, 2012), Aoki (1999), Yin Wenying et al. (2000), Corpuz-Raros (2005), Subias (2013) [1, 2, 3, 13, 24], the results of the analysis of a zoogeographical characteristics of the oribatid fauna of Vietnam is presented in the Figure 1.




Figure 1. Zoogeographical characteristics of the oribatid mite fauna of Vietnam

Legends: (1) Oriental region: OR, (2) Palaeartic – Oriental: PA, (3) Cosmopolitan: CO, (4) Afrotropical (Ethiopic): AF, (5) Australian – Oriental: AU, (6) Neotropical – Oriental: NO, (7) Nearctic – Oriental: NE, (8) Pacific – Oriental: PC.

The result shows that, the oribatid mite fauna of Vietnam includes all eight zoogeographical elements. These eight zoogeographical elements can be arranged according to decreasing representation of species, as follows: (1) Oriental region (OR) occupies 60.3% of the total 320 species, with 193 species registered, (2) Palaeartic – Oriental (PA) - 12.2%, with 39 species, (3) Cosmopolitan (CO) - 10.6%, with 34 species, (4) Afrotropical (Ethiopic) – Oriental (AF) - 6.9%, with 22 species, (5) Australian – Oriental (AU) - 5.0%, with 16 species, (6) Neotropical – Oriental (NO) - 3.8%, with 12 species, (7) Nearctic – Oriental (NE) - 0.9%, with 3 species, (8) Pacific - Oriental (PC) – 0.3%, with 1 species (Figure 1).

The main zoogeographical component of the oribatid mite fauna of Vietnam are the Oriental species, representing 60.3% of the total 320 species, with 193 recorded. This oribatid mite fauna has high specialization, with 111 species (34.68% of the total) recorded only from Vietnam (probably endemic species, i.e. conditional endemics). Besides

that, the oribatid mite fauna of Vietnam includes also the Palaearctic - Oriental elements, with 39 species recorded, and cosmopolitan elements - 34 species, representing 12.2% and 10.6% of the total, respectively. The other categories are represented by a smaller number of species. There is now Antarctic element recorded in the oribatid mite fauna of Vietnam.

4. CONCLUSIONS

The oribatid mite fauna (Acari: Oribatida) of Vietnam is diversified, and has high specialization. It is diverse by the number of superfamilies, families, genera and species recorded. However, the number of genera per family, as well as the number of species and subspecies per genus, is not high. 43.75% and 37.50% of the total 64 families and subfamilies consist of one and of 2-3 genera, respectively. The only one family Oppiidae Grandjean, 1954 consists of 23 genera. The majority of the genera, 68.10% of the total, are represented by one species. The only two genera are represented by more than 10 species, namely *Galumna* Heyden, 1826 and *Pergalumna* Grandjean, 1936, with 13 and 11 species, respectively.

The main zoogeographical characteristics of the oribatid mite fauna of Vietnam are the Oriental species.

Acknowledgements: This research is funded by Vietnam National Foundation for Science and Technology Development (NAFOSTED) under grant number 106.14-2012.46. The first author thanks Prof. B. Georgiev and Assoc. Prof. Anna Ganeva for their assistance during his study visit, November-December 2015, at the Institute of Biodiversity and Ecosystem Research (IBER: BAS) Sofia, Bulgarian.

5. References

- [1] Balogh J, P. Balogh 1992. The Oribatid Genera of the World, HNHM Press, Budapest, V.1, 263p, V.2, 375p.
- [2] Balogh J., P. Balogh 2002. Identification Keys to the Oribatid Mites of the Extra-Holarctic Regions. Well-Press Publishing Limited. Budapest, V. I, 451p. and V.II, 504p.
- [3] Balogh J., S. Mahunka 1967. New oribatids (Acari, Oribatei) from Vietnam.- Acta Zoologica Hungarica, 13(1-2): 39-74.
- [4] Behan-Pelletier V. 1999. Oribatid mite biodiversity in agroecosystems role for bioindication.- Agricultural Ecosystem and Environment, 74: 411-423.
- [5] Đào Duy Trinh, Trịnh Thị Thu, Vu Quang Manh, 2010: Dẫn liệu về thành phần loài, phân bố và địa động vật của khu hệ ve giáp ở VQG Xuân Sơn, Phú Thọ.- Tạp chí Khoa học ĐHQG Hà Nội, 26(1): 49-56.
- [6] Edwards C. A. 1991. The assessment of population of soil-inhabiting invertebrates.- Agricultural Ecosystem and Environment, 34: 145-176.
- [7] Ermilov SG, 2015, A list of oribatid mites (Acari, Oribatida) of Vietnam, Zookeys. 2015; (546): 61-85.
- [8] Ermilov S. G., Anichkin A. E., 2014. Taxonomy study of oribatid mites (Acari, Oribatida) of Bu Dup - Nui Ba National Park (Southern Vietnam).- Zootaxa, 3834(1): 1-86.
- [9] Ermilov S.G., Vu Q. M., 2012. Two new species of oribatid mite (Acari: Oribatida) from Phong Nha - Ke Bang National Park of Central Vietnam. International Journal of Acarology 38, 2, 160-167.
- [10] Ermilov SG, Anichkin AE, Wu D., 2012. Oribatid mites from Bu Gia Map National Park (Southern Vietnam), with description of a new species of *Dolichcheremaeus* (Tetracondylidae)(Acari: Oribatida). Genus. 23(4):591-601.
- [11] Ermilov S.G., Vu Q. M., Trinh T. T. and Dao D. T., 2011. *Percylobates thanhoaensis*, a new species of oribatid mite from Vietnam (Acari: Oribatida: Haplozetidae). International Journal of Acarology, 37, 2, 161-166
- [12] Fernandez N., Theron P., Rollard C., Casillo E. R., 2014: Oribatid mites from deep soils of Hòn Chông limestone hills, Vietnam: the Family Lohmanniidae (Acari: Oribatida), with the descriptions of *Bedoslohmannia annea* n.gen., n. sp., and *Paulianaqcarus vietnamese* n. sp. Zoosystema 36(4), 771-787.
- [13] Ghilarov M., D. Krivolutsky 1975. Identification of Soil mites Sarcoptiformes, Nauka, Moscow, 491p. (in Russ.).
- [14] Jeleva M., M. Q. Vu 1987. New Oribatids (Oribatei, Acari) from the Northern part of Vietnam. Acta Zoologica Bulgarica, 33, 10-18.
- [15] Krivolutskiy et al. 1997 (Криволицкий Д. А., Ву Куанг Мань, Фан Тхе Вьет 1997). Панцирных клещей (Acari: Oribatei) Вьетнама. Тропическая медицина, Т. I, Тропцентр, Наука, Москва - Ханой, 152-167 (In Russ.).
- [16] Mahunka S., 1987. A survey of the Oribatid (Acari) fauna of Vietnam. I., Annals. Hist.-nat. Mus. natn. Hung., 79, 259-279.
- [17] Mahunka S., 1988. A survey of the Oribatid (Acari) fauna of Vietnam. II, Act. Zool. Hung., 34 (2-3), 215-246.
- [18] Mahunka S., 1989. A survey of the Oribatid (Acari) fauna of Vietnam. III, Folia Entomologica Hungarica, L, 47-59.
- [19] Minor M. A., Ermilov, S. G. 2015. Effects of topography on soil and litter mites (Acari: Oribatida, Mesostigmata) in a tropical monsoon forest in Southern Vietnam. Experimental and Applied Acarology, 67(3), 357-372.
- [20] Nguyễn Hải Tiến, Vũ Quang Mạnh, 2012: Thành phần loài ve giáp (Acari: Oribatida) ở vườn quốc gia Phong Nha – Kẻ Bàng, Quảng Bình. Tạp chí Khoa học ĐHQGHN, Khoa học Tự nhiên và Công nghệ, 28(2012), 125-134.
- [21] Nguyễn Huy Trí, Lại Thu Hiền, Nguyễn Hải Tiến, Vũ Quang Mạnh, 2014: Cấu trúc quần xã ve giáp (Acari: Oribatida) như yếu tố chỉ thị sinh học sự biến đổi sinh cảnh ở Vườn Quốc gia Cát Bà, thành phố Hải Phòng. Tạp chí Khoa học, ĐHQG Hà Nội, 268-274.

- [22] Norton R. A., V. Behan-Pelletier 2009. Chapter 15. Suborder Oribatida. 430-564pp.. in: KRANTZ, G.W.; WALTER, D.E. (eds) A manual of acarology. Third edition. Texas Tech University Press, Lubbock Texas.
- [23] Schatz H. et al. 2011. Suborder Oribatida van der Hammen, 1968. In: Zhang Z. -Q. (ed.) 2011. Animal biodiversity: an outline of higher-level classification and survey of taxonomic richness. *Zootaxa*, 3148: 141–148.
- [24] Subias L. S. 2013. Listado sistemático, sinonímico y biogeográfico de los ácaros oribátidos (Acariformes: Oribatida) del mundo (excepto fósiles) Graellsia 60 (número extraordinario). 1-570p. (<http://www.ucm.es/info/zoo/Artropodos/Catalogo.pdf>).
- [25] Udvardy M. D. F. 1975. A classification of the biogeographic provinces of the World. IUCN. Morges.
- [26] Vietnam Ministry of Agriculture and Rural Development 2006. Forest area and forest land not used for forestry planning. Agricultural Publishing House, 152 p. Hanoi (In Vietnamese).
- [27] Vietnam National Institute for Soils and fertilizers 2002. The basic information of main soil units of Vietnam. Thegioi Publishers. Hanoi. 1-158.
- [28] Vietnam Soil Society 2000. The Soils of Vietnam. Agricultural Publishing House, Hanoi. 1-412 (In Vietnamese).
- [29] Vu Quang Manh 1985 (Ву Куанг Манх). Фаунистично - екологично исследование върху орибатеите (Acari: Oribatei) в северната част на Виетнам.- Дисертация Канд. Биолог. Науки, София, 1-175 (in Bulg.).
- [30] Vũ Quang Mạnh, 1990. Chân khớp bé (Microarthropoda) trong quần lạc động vật đất ở Việt Nam. Tạp chí Sinh học, 12, 1, 3-10.
- [31] Vu Q. M., Nguyen T. T. 2000. Microarthropod community structures (Oribatei and Collembola) in Tam Dao National Park. Journal of Biosciences. 25(4): 379-386.
- [32] Vũ Quang Mạnh, 2007. Động vật chí Việt Nam, T. 21: Bộ Ve giáp Oribatida, Nxb KHKТ, 1-355.
- [33] Vu Q. M. 2012. Oribatid soil mite (Acari: Oribatida) of northern Vietnam: Species distribution and densities according to soil and habitat type. The Pan-Pacific Entomologist, 87(4), 209-222.
- [34] Vu Quang Manh, 2013. The Oribatida (Acari: Oribatida) fauna of Vietnam. Institute of Biodiversity and Ecosystem Research (IBER: BAS), DSc. Thesis, 205p. (www.ecolab.bas.bg/main/Members/gpv/Vu.../Vu_Manh_Thesis_all.pdf).
- [35] Vu Quang Manh, 2015. The oribatid mite fauna (Acari: Oribatida) of Vietnam - Systematics, Zoogeography and Formation, PENSOFT Publishers, Sofia-Moscow. 1-212.
- [36] Vu Q. M., Nguyen T. T. 2000. Microarthropod community structures (Oribatei and Collembola) in Tam Dao National Park. Journal of Biosciences. 25(4): 379-386.
- [37] Vu Quang Manh, Ermilov S.G., Dao Duy Trinh, 2010: Two new species of Oribatid Mites (Acari: Oribatida) from Vietnam. Tạp chí Sinh học, 32(3), 12-19.
- [38] Vu T. L., C. Taillard 1994. Atlas of Vietnam (Cartographic material), RECLUS, Montpellier, Paris, 1-421 pp.