

Bioaccumulation of organochlorine pesticides (OCPs) in molluscs and fish at the Sai Gon - Dong Nai estuary

Tích lũy sinh học hóa chất bảo vệ thực vật clo hữu cơ trong nhuyễn thể và cá tại cửa sông Sài Gòn - Đồng Nai

Research article

Nguyen Xuan Tong^{1,2}, Tran Thi Thu Huong³, Duong Thi Thuy^{1,4*}, Mai Huong⁵, Duong Trong Khang², Huynh Cong Luc², Pham Thi Loan², Le Thi Phuong Quynh⁶

¹Graduate University of Science and Technology, Viet Nam Academy of Science and Technology (VAST); ²Institute for Environmental Science, Engineering and Management, Industrial University of Ho Chi Minh City; ³Faculty of Environmental, Ha Noi University of Mining and Geology; ⁴Institute of Environmental Technology, VAST; ⁵University of Science and Technology of Ha Noi, VAST; ⁶Institute of Natural Product Chemistry, VAST, 18 Hoang Quoc Viet, Cau Giay, Ha Noi, Vietnam

The aim of this study is to assess the biological accumulation of pesticide residues in aquatic organisms in Sai Gon - Dong Nai (SG-DN) estuary. Fish and mollusks were collected directly at the Soai Rap and Long Tau estuary of the SG-DN river system, washed and separated for taking the tissue. The organochlorine compounds from the tissue were then extracted and analyzed by gas chromatography system. The results showed that, the concentration of OCPs in *Tegillarca granosa*, *Meretrix lyrata*, *Margaritifera auricularia* and *Bostrychus sinensis* varied from 6.4 to 59.9 µg/kg, 7.2 to 322 µg/kg, 4.5 to 62.1 µg/kg and 2.9 to 114.3 µg/kg fresh weight, respectively. In general, molluscs species that accumulate more heptachlor, aldrin, endrin or dieldrin tend to accumulate less DDT (dichlorodiphenyltrichloroethane). Endosulfan was the most commonly found in three bivalve mollusks while DDTs (1.5–75.2 µg/kg, averaging 8.7 µg/kg weight) was the most popular OCPs in the fish (*Bostrychus sinensis*) samples. In DDT group, the p,p'-DDT metabolite accounted for the largest percentage, reaching 50% of total DDTs. In HCH (Hexachlorocyclohexane) group, β-HCH isomer was predominant in almost samples.

Mgr đích ccc nghiên cch này là đánh giá tích lũy sinh hhn ccn thuu trr sâu trong các sinh vvn sshn dưnh nưnh ttr khu vvu ccu sông Sài Gòn - ĐĐn Nai (SG-DN). Cá và nhuyyu thh đưuy llr trry tiyy iicủa sông Soài RRà và Lòng Tàu thuu hh thhu sông SG-DN, đư-D rư- ss-D và tách llc phhh mô thht. Các hhc chh clo hho cơ sau đó đưud tách chiü và phân tích bbch hh thhh ssh ký khí. Kết quả nghiên cứu cho thấy, dư lượng OCPs tích tụ trong sò điệp (*Tegillarca granosa*), ngao (*Meretrix lyrata*), trai nước ngọt (*Margaritifera auricularia*) và cá bớp (*Bostrychus sinensis*) dao động tương ứng từ 6,4 đến 59,9 µg/kg, 7,2 đến 322 µg/kg, 4,5 đến 62,1 µg/kg và 2,9 đến 114,3 µg/kg trọng lượng tươi. Nhìn chung, loài nhuyễn thể nào tích lũy nhiều heptachlor, aldrin, endrin hoặc dieldrin có xu hướng tích lũy ít DDT (Dichlorodiphenyltrichloroethane). Endosulfan là nhóm thuốc được tìm thấy nhiều nhất trong các loài nhuyễn thể nghiên cứu. Ngược lại, nhóm DDT lại phổ biến ở cá *Bostrychus sinensis* (1.5–75.2 µg/kg, trung bình 8.7 µg/kg trọng lượng). Dạng p,p'-DDT trong nhóm DDT chiếm tỷ lệ cao nhất, tới 50% DDT tổng. Trong khi đó, đồng dạng β-HCH của nhóm HCH (Hexachlorocyclohexane) chiếm đa số trong hầu hết các mẫu.

Keywords: biological accumulation, estuarine, fish, molluscs, pesticide

1. Introduction

The Sai Gon – Don Nai (SG-DN) River system with a large estuarine area is the main water sources for agriculture and the daily activities of people living along the river banks in Ho Chi Minh City and the neighbouring provinces (Ba Ria Vung Tau, Dong Nai, Long An and Binh Duong). However, rapid development in this region raised concerns about local environment and ecological integrity (Anh, 2003; Nguyen et al, 2007). Large amounts of untreated municipal and industrial wastewater, as well as runoff from landfills carrying numerous toxic complex contaminants are released directly into the river. The increase in population and productive activities, as well as use of pesticides without logical in agricultural production increased pesticide residues in the river. Since many years, this estuary has been served as one of the main areas for large-scale aquaculture. Many species of fishes and molluscs such as oysters, clam etc. were cultivated in here. These species have a long growing cycle, ranging from 8 to 10 months and need to live in a clean environment without contamination. Therefore, the protection of river water quality is an important task for sustainable development, especially when demand for water supply and use of fishery products increases rapidly.

Organochlorine pesticides (OCPs) which are large groups of crop protection chemicals have been widely used for various applications during the past several decades. Due to their persistent, semi-volatile, toxic properties and highly bio-accumulation especially in lipophilic substances and tissues, organochlorines distribute ubiquitously in the global environment and can be detected at all levels of food chains, including humans. Many OCPs are classified into persistent organic pollutants (POPs), according to Stockholm formula (UNEP, 2011). These compounds affect seriously public health of humans and wildlife because they alter normal functions of endocrine (endocrine disrupter pesticides) (Mnif et al, 2011), interfere with female hormonal function, leading to negative effects on the reproductive systems (Bretveld et al, 2006), have the close relationship with neurodegeneration diseases (Dana, 2008), many popular cancers such as breast cancer (Wolff et al, 2007; Bhatnagar et al, 2002), non Hodgkin’s lymphoma (Cantor et al, 1992), and other types of cancer (Alavanja et al 2013).

Although the use of many organochlorine compounds such as Lindane, DDT, Endrin, Dieldrin etc. has been prohibited according to Circular No. 21/2013/MARD because of the long half-life and difficulty for controlling persistent residue (MARD, 2013). However, pesticide contamination still occurs and is difficult to handle, including the issue of OCPs bioaccumulation in the environment. Some reports published recently focus on the assessment of OCPs and heavy metal pollution in sediments, rivers, seas and estuaries along Vietnam such as: Mekong River Delta (Nguyen et al, 2006), Sai Gon - Dong Nai River (Nguyen et al, 2007), Tien estuary (Nguyen et al, 2017), Thi Vai Estuary and Can Gio Mangrove Forest (Costa-Böddeker, 2016), Nha Trang Bay (Hoang et al, 2015), Red River Delta and along the coast of North Vietnam (Dang et al, 1998). Until now, there is a few of studies investigating the toxic OCPs bioaccumulation in *Tegillarca granosa*, *Meretrix lyrata*,

Margaritifera auricularia and *Bostrychus sinensis* cultured in the SG-DN estuary. Therefore, the aim of this work is to investigate levels of OCPs residues and their isomer including of the groups such as: HCHs, DDTs, Aldrin, Dieldrin, Heptachlor, Endosulfans in these species in SG-DN estuary area.

2. Material and methods

2.1. Study area

The current study was conducted in the Long Tau and Soai Rap estuary belonging to the Sai Gon-Dong Nai estuary, located in Ho Chi Minh city, Vietnam (Figure 1). The estuary with funnel shape has a total catchment area of 43,500 km², including areas contiguous with the Ba Ria - Vung Tau, Dong Nai, Long An and Binh Duong provinces. The river catchment has two distinct seasons: the dry season is from December to April and the rainy season is from May to November, with total annual rainfall from 170.8 – 212.4 mm. There are four main rivers (Soai Rap, Long Tau, Thi Vai and Cai Mep) collective adding water to the SG-DN River system, before coming to the East Sea with the total annual discharge of approximately 34 billion m³ (Le et al, 2015). Currently, the area of the paddy field in the catchment of the SG-DN river system is around 18,000 hectares.




Figure 1. The SG-DN estuary and sampling areas

2.2. Sampling

2.2.1. Sample identification

For sample identification, all the samples are classified according to their morphological characteristics and explicit enumeration to the David's category (1969): scallops,

clams, mussel and fish. The bivalve molluscs (clams, oysters and clams) are brushed with a brush, washed with water to remove the adhering detritus.

2.2.2. Sampling

Thirty-eight sites surrounding the boundary of the Long Tau and Soai Rap estuary belong to the SG-DN system were selected for the current study (Figure 1). Samplings were conducted in two seasons (rainy season: May 2017, dry season: October and November 2017). The sampling process is similar to the procedure of two authors Kožul (2008) and Mohamed (2012). The experimental samples including: scallops, mussel, clams and fish were taken directly and randomly by hand. All the sample were immediately kept in plastic boxes and stored in an ice-chest at 5°C and transported to the laboratory for analysis. The group of bivalve molluscs was weighed, classified, peeled off the shells to collect the meat. Their tissues were homogenized and stored in the dark glass bottles at -5°C for further analysis (Kožul et al, 2008; Shreadah et al, 2012).

2.3. Procedure for sample treatment and analysis

The processing analysis is followed the steps presented in the Figure 2 (Kožul, 2008; Shreadah, 2012).




Figure 2. The procedure for analyzing OCPs in experimental samples

2.4. Statistical analysis

All experiments were done in triplicate and the data were calculated as mean ± SD (standard deviation) and drawn by the software SigmaPlot. Statistical significance was accepted at a level of p < 0.05.

3. Results and discussion

3.1. The analysis results of OCPs component

In this study, the individual numbers of the analysed species *M. auricularia*, *T. granosa*, *M. lyrata* and *B. sinensis* were 8, 9, 13 and 8, respectively. Table 1 shows the average mean (mean), the standard error of the mean (SEM) and the varied values of OCPs in all analysed samples. According to Vietnamese standard, 12 different common components of OCPs and pesticides residues were detected with high contents in the both three bivalve molluscs and fish *B. sinensis* species collected in the study area (MARD, 2013; 2015). The bioaccumulation of OCPs in the survey species is significantly different. Almost all individuals of *Bostrychus sinensis* did not accumulate endosulfan II; many individuals of *Tegillarca granosa* did not accumulate α-HCH and δ-HCH; *Meretrix lyrata* individuals did not accumulate aldrin, sulphate endosulfan and *Margaritifera auricularia* did not accumulate aldrin.

The highest total OCP accumulation in *M. lyrata* was 114.317 µg/kg, while the lowest in *M. auricularia* varied from 4.472 to 62.114 µg/kg by wet weight. The analysis results showed that the highest total OCP accumulation in *Meretrix lyrata* was 114,317 µg / kg, while the lowest in *Margaritifera auricularia* varied from 4,472 to 62,114 µg kg by wet weight. However, the endosulfan group was found in all three bivalve molluscs with the content of 1.5 - 75.2 µg/kg and the DDT group was the most common OCPs in the fish *B. sinensis* with the content of 8.7 µg/kg by wet weight. The high accumulation of HCHs in fish sample indicated the pollution in the environment and the risk to human health.

Table 1. Organochlorine pesticides detected (µg/kg) in all sample

Species	<i>Margaritifera auricularia</i>	<i>Tegillarca granosa</i>	<i>Meretrix lyrata</i>	<i>Bostrychus sinensis</i>
Sample size	(N = 8)	(N = 13)	(N = 9)	(N = 8)
Total HCHs	6.527	5.005	7.160	4.939
SEM	1.914	0.925	4.344	1.509
Range	2.001 – 16.591	1.706 – 12.149	1.742 – 41.832	0.804 – 13.788
α-HCH	1.759	0.717	1.515	0.970
SEM	0.492	0.288	0.905	0.367
Range	0.745 – 4.128	0 – 3.695	0 – 8.695	0 – 3.132

Species	<i>Margaritifera auricularia</i>	<i>Tegillarca granosa</i>	<i>Meretrix lyrata</i>	<i>Bostrychus sinensis</i>
Sample size	(N = 8)	(N = 13)	(N = 9)	(N = 8)
β -HCH	3.107	2.076	3.330	2.246
SEM	1.098	0.482	2.058	0.783
Range	0.817 – 10.012	0 – 6.455	0.586 – 19.704	0.221 – 5.458
δ -HCH	0.686	0.889	0.389	1.224
SEM	0.313	0.445	0.195	0.483
Range	0 – 2.502	0 – 4.702	0 – 1.587	0 – 3.961
γ -HCH	0.976	1.323	1.925	0.500
SEM	0.245	0.392	1.247	0.199
Range	0.429 – 2.102	0 – 3.922	0.182 – 11.845	0 – 1.707
Total DDTs	8.240	6.083	6.801	16.662
SEM	2.432	1.673	2.455	8.749
Range	0.542 – 17.729	1.038 – 20.094	2.030 – 24.475	1.532 – 75.224
<i>p,p'</i> -DDE	1.060	1.075	0.991	2.084
SEM	0.525	0.394	0.475	0.683
Range	0 – 4.094	0 – 4.721	0 – 4.419	0 – 4.596
<i>p,p'</i> -DDD	4.560	3.107	1.899	4.245
SEM	1.752	1.058	0.851	1.613
Range	0.001 – 10.856	0.109 – 12.746	0 – 8.350	0.564 – 12.603
<i>p,p'</i> -DDT	2.620	1.942	3.910	10.333
SEM	0.81	0.653	1.175	7.417
Range	0.442 – 6.640	0 – 9.017	1.336 – 11.706	0.464 – 61.944
Total endosulfans	4.591	14.482	27.831	9.674
SEM	1.188	2.952	24.02	15.836
Range	0.534 – 9.862	0.813 – 30.615	0.205 – 219.491	0 – 47.061
Endosulfan I	2.405	1.415	1.059	1.737
SEM	1.199	0.661	0.454	0.7
Range	0 – 8.854	0 – 8.075	0.118 – 3.537	0 – 5.682
Endosulfan II	0.456	2.620	0.531	0.164
SEM	0.13	0.962	0.164	0.109
Range	0 – 1.143	0.639 – 13.460	0 – 1.409	0 – 0.765
Endosulfan Sulfat	1.730	10.446	26.241	7.772
SEM	0.868	3.098	23.707	5.637
Range	0.105 – 5.743	0 – 29.679	0 – 215.414	0 – 46.507
Total OCPs	28.020	36.992	51.585	35.750
SEM	7.105	4.488	34.026	13.326
Range	4.472 – 62.114	6.360 – 59.897	7.181 – 321.996	2.925 – 114.317
Heptachlor	1.360	3.516	7.738	1.003
SEM	0.315	0.755	3.853	0.256
Range	0.178 – 2.917	0 – 8.121	0 – 30.524	0 – 2.056
Aldrin	1.002	1.713	0.458	1.245
SEM	0.547	0.474	0.337	0.43
Range	0 – 4.380	0 – 5.421	0 – 3.055	0 – 3.815
Dieldrin	4.626	2.793	0.201	1.743
SEM	3.441	1.756	0.074	0.564
Range	0.055 – 27.816	0 – 23.157	0 – 0.557	0 – 3.822
Endrin	1.675	3.359	1.936	0.589
SEM	1.252	1.317	0.916	0.217
Range	0.039 – 10.391	0 – 17.351	0 – 8.341	0 – 1.580

* SEM: Standard error of the mean

3.2. The analysis results of the organochlorine derivative groups

Figure 3 illustrates the content of seven types of pesticides affecting 4 studied species: *M. auricularia*, *T. granosa*, *M. lyrata*, and *B. sinensis*. In general, it can be seen that endosulfans were the most predominant compounds in three molluscs species; followed by DDT, HCH, heptachlor and the rest substances. However, the content of substances

high or low is very different in different species. The contents of endosulfans in *M. lyrata*, *T. granosa*, *B. sinensis* and *M. auricularia* respectively were 27,831 $\mu\text{g}/\text{kg}$, 14,482 $\mu\text{g}/\text{kg}$, 9,674 $\mu\text{g}/\text{kg}$ and 4,591 $\mu\text{g}/\text{kg}$, respectively. Followed by the content of DDT was 6,083 $\mu\text{g}/\text{kg}$ in *T. granosa* and the lowest content in this species is the HCH group. In contrast, in fish *B. sinensis*, DDTs were the most predominant OCPs (16,662 $\mu\text{g}/\text{kg}$), followed by endosulfans and HCH derivatives. However, in *M. lyrata*, heptachlor had the relatively equal amount to HCHs and DDTs.




Figure 3. Total OCPs and organochlorine groups in the samples

Figure 4 shows contents ($\mu\text{g}/\text{kg}$) of five HCH derivatives determined in the study species. The highest bioaccumulation of total HCHs was observed in *M. lyrata* (7,159 $\mu\text{g}/\text{kg}$), followed by *M. auricularia* (6,528 $\mu\text{g}/\text{kg}$), *B. sinensis* (5,94 $\mu\text{g}/\text{kg}$) and the lowest value of 5 $\mu\text{g}/\text{kg}$ for *T. granosa*. In this group, β -HCH d accounted for more than 45.5% of all HCHs found in the survey species. It is interesting to note that *M. auricularia* and *M. lyrata* contained the similar content of β -HCH (about 3.33 $\mu\text{g}/\text{kg}$), higher than *B. sinensis* (2,246 $\mu\text{g}/\text{kg}$) and *T. granosa* (2,076 $\mu\text{g}/\text{kg}$). Anh finally, the content of α -HCH was the largest within HCHs group in *M. auricularia* (1.759 $\mu\text{g}/\text{kg}$), while γ -HCH was predominant in *M. lyrata* (nearly 2 $\mu\text{g}/\text{kg}$), δ -HCH in *B. sinensis* (1.224 $\mu\text{g}/\text{kg}$). This result is different from the previous report (Pham et al, 2011), in which HCHs were detected in fish samples collected in Minh Dai commune, Phu Tho province with concentrations varied from 12 to 78 ng/g wet weight and the average value was 31 ng/g.




Figure 4. HCHs and compositions in the samples

Figure 5 illustrates the level of three DDT derivatives including p,p'-DDE, p,p'-DDD and p,p'-DDT observed in the study species. The highest bioaccumulation of DDTs total was observed in *B. sinensis* (16,662 $\mu\text{g}/\text{kg}$), followed by *M. auricularia* (8.24 $\mu\text{g}/\text{kg}$), *M. lyrata* (6,8 $\mu\text{g}/\text{kg}$), and the lowest value of only 6 $\mu\text{g}/\text{kg}$ was in *T. granosa*. In general, the content of p,p'-DDT metabolite accounted for the largest percentage, reaching 50% of DDTs total in all samples. In particular, the ratio p,p'-DDT/DDTs of *B. sinensis* was highest (62%), followed by *M. lyrata* (57%). In contrast, p,p'-DDD was more predominant in *Margaritifera auricularia* and *Tegillarca granosa*, accounting for to 50% of DDTs total. This indicates the high stability of the stock compounds (p, p'-DDT

and p, p'-DDD) in the aquatic environment compared with other metabolites.




Figure 5. DDTs and compositions in the samples

Figure 6 shows the levels of endosulfans total and the form of derivatives in 4 studied species. The highest bioaccumulation of endosulfan total was observed in *M. lyrata* (up to 27,831 $\mu\text{g}/\text{kg}$), followed by *T. granosa* (14,481 $\mu\text{g}/\text{kg}$), *B. sinensis* (9,673 $\mu\text{g}/\text{kg}$) and the lowest value of only 4.591 $\mu\text{g}/\text{kg}$ was in *M. auricularia*. In figure 5, except *M. auricularia* samples, endosulfan sulphate tended to be more active than other endosulfan derivatives in the remaining three species. It accounted for the largest percentage, reaching up to 80-90% of endosulfans total and this value in *M. lyrata* (26,241 $\mu\text{g}/\text{kg}$), *T. granosa* (10,441 $\mu\text{g}/\text{kg}$) and *B. sinensis* is 7,772 $\mu\text{g}/\text{kg}$, respectively. For *M. auricularia*, total endosulfans was slightly lower than endosulfan I, this value is 1.73 $\mu\text{g}/\text{kg}$ compared with 2.4 $\mu\text{g}/\text{kg}$ endosulfan I. According to Dang et al (2001), in the canals in Hanoi city, p,p'-DDE always accounts for the largest proportion, up to 70-80% of DDT in freshwater mussel (*Angulyagra* sp.).




Figure 6. Endosulfans and compositions in the samples

The study of bioaccumulation OCPs in some species of bivalve molluscs and fish in the SG-DN estuary showed that there was a difference in the level of bioaccumulation in each species. In general, molluscs species tended to accumulate heptachlor, aldrin, endrin or dieldrin more than DDTs. This result is different from the report of Dang et al (2001) and Marta (2015). (Commendatore et al, 2015) showed that organochlorine bioaccumulation processes through bottom sediment resuspension were found in surface sediment and bivalve molluscs (29.4–206.0 ng/g dry weight); whereas imposex incidence was only 15% in the gastropod *Pareuthria plumbea*. Moreover, the presence of p,p'-DDT and the p,p'-DDE ratio likely reflect the high

stability of the stock compounds and their isomer in the aquatic environment compared with other metabolites. The OCP compounds in bivalve molluscs showed different patterns. This result may be due to different input pathways such as OCPs being incorporated into organisms directly from the water column's suspended particles while being transformed in sediments. According to these authors (Dang et al, 1998; Commendatore, 2015), the organochlorinated pollution was clearly related to the migration of contaminants and the ability to self-disintegrate pesticides by environmental factors such as: temperature, atmosphere, transport etc.

4. Conclusion

This study demonstrates the high accumulation levels of organochlorine pesticides in different species of bivalve molluscs cultured in the Sai Gon – Dong Nai estuary compared with Vietnam standard 10:2008/MONRE. Endosulfan was the most commonly found in three bivalve molluscs (4,591-27,831 µg/kg, average 14,144 µg/kg by wet weight) while DDTs (6,124-16,662 µg/kg, average 9,4565 µg/kg by wet weight) was the most popular OCPs in the fish (*B. sinensis*) samples. In DDT group, the p,p'-DDT metabolite accounted for the largest percentage, reaching 50% of total DDTs. In HCH (Hexachlorocyclohexane) group, β-HCH isomer was predominant in almost samples. However, in order to have more consistent and scientific assessments, it is necessary to combine with the study of OCPs deposit in water, sediment and other aquatic parameters to limit the potential for bioaccumulation of the pesticides in these species.

5. References

[1] Anh MT, Chi DHL, Vinh NN, Loan TTC, Triet LM, Slooten KB, Tarradellas J (2003) Micropollutants in the sediment of the SaiGon–DongNai River: situation and ecological risks. *Chimia*, 57, 537–541.

[2] Canadian Cancer Society (2013) Cosmetic Pesticides. Information Brief. Available from: <https://www.cancer.ca/~media/cancer.ca/AB/get%20involved/take%20action/CosmeticPesticides-InformationBrief-AB>.

[3] Cantor KP, Blair A, Everett G, Gibson R, Burmeister LF (1992) Pesticides and other agricultural risk factors for non-Hodgkin's lymphoma among men in Iowa and Minnesota. *Cancer Res*, 52, 2447–55.

[4] Circular No. 21/2013/MARD: The promulgating the list of pesticides permitted for use, limited use, forbidden to use and the additional list of plant varieties permitted for production and trading in Viet Nam.

[5] Circular No. 31/2015/MARD: Regulations on monitoring residues of toxic substances in animals and seafood products.

[6] Cohn B, Wolff M, Cirillo P, Sholtz R (2007) DDT and Breast Cancer in Young Women: New Data on the Significance of Age at Exposure. *Environmental Health Perspectives*, 115(10), 1410-1414.

[7] Hancock DB, Martin ER, Mayhew GM, Stajich JM,

Jewett R, Stacy MA, Scott BL, Vance JM, Scott WK (2008) Pesticide exposure and risk of Parkinson's disease: A family-based case-control study. *BMC Neurology*, 8(6). doi:10.1186/1471-2377-8-6

[8] Dang DN, Carvalho FP, Nguyen MA, Nguyen QT, Nguyen THY, Villeneuve JP, Cattini C (2001) Chlorinated pesticides and PCBs in sediments and molluscs from freshwater canals in the Hanoi region. *Environmental Pollution*, 112(3), 311-320.

[9] Dang DN, Nguyen MA, Nguyen CH, Luu VD, Carvalho FP, Villeneuve JP, Cattini C (1998) Organochlorine pesticides and PCBs in the Red River Delta, North Vietnam. *Marine Pollution Bulletin*, 36(9), 742-749.

[10] Nicol D (1969) The Number of Living Species of Molluscs. *Systematic Zoology*, 18(2), 251-254

[11] Hoang TD, Kunzmann A (2015) The Sediment Load and Deposition by River Discharge and Their Relation to Organochlorine Pesticides Pollutants in the Sediment Bottom of Nha Trang Bay, Vietnam. *Ocean Sci. J*, 50(2), 455-466.

[12] Darija K, Snježana HR, Zorana KG, Jere V (2008) Levels of Polychlorinated Biphenyls and Organochlorine pesticides in Mediterranean blue mussel from the Adriatic sea. *Organohalogen Compounds*, Volume 70, pp. 001243.

[13] Le Duc An (2015) Coastal Vietnam, structure and natural resources. Natural Science and Technology Publisher (in Vietnamese).

[14] Commendatore MG, Franco MA, Costa GP, Castro IB, Fillmann G, Bigatii G, Esteves JL, Nievas ML (2015) Butyltins, polyaromatic hydrocarbons, organochlorine pesticides, and polychlorinated biphenyls in sediments and bivalve mollusks in a mid-latitude environment from the patagonian coastal zone. *Environmental Toxicology and Chemistry*, 34(12), 2750–2763.

[15] Mathur V, Bhatnagar P, Sharma RG, Acharya V, Sexana R. (2002) Breast Cancer incidence and exposure to pesticides among young women originating from Jaipur. *Environment International*, 28, 331-336.

[16] Alavanja MCR, Ross MK, Bonner MR (2013) Increased Cancer Burden Among Pesticide Applicators and Others Due to Pesticide Exposure. *J CLIN*, 63, 120–142.

[17] Minh NH, Someya M, Minh TB, Kunisue T, Watanabe M, Tanabe S, Viet PH, Tuyen BC (2004) Persistent organochlorine residues in human breast milk from Hanoi and Ho Chi Minh City, Vietnam: Contamination, accumulation kinetics, and risk assessment for infants. *Environ Pollut*, 129, 431–441.

[18] Mnif W, Hassine AIH, Bouaziz A, Bartegi A, Thomas O, Roig Bendo (2011) Effect of endocrine disruptor pesticides: a review. *Int J Environ Res Public Health*, 8, 2265–2203.

[19] Shreadah MA, Said TO, Othman IM, Eiman M. I. Fathallah MEM (2012) Polychlorinated Biphenyls and Chlorinated Pesticides in Sediments along the

- Semi-Closed Areas of Alexandria, Egypt. *Journal of Environmental Protection* 3, 141-149.
- [20] Nguyen HM, Tu BM, Iwata H, Kajiwara N, Kunisue T, Takahashi S, Pham HV, Bui CT, Tanabe S (2007) Persistent Organic Pollutants in Sediments from Sai Gon–Dong Nai River Basin, Vietnam: Levels and Temporal Trends. *Arch. Environ. Contam. Toxicol.* 52, 458–465.
- [21] Nguyen HM, Tu BM, Kajiwara N, Kunisue T, Hisato Iwata, Pham HV, Nguyen PCT, Bui CT, Tanabe S (2006) Contamination by polybrominated diphenyl ethers and persistent organochlorines in catfish and feed from Mekong river delta, Vietnam. *Environmental Toxicology and Chemistry*, 25(10), 2700–2709.
- [22] Nguyen VH, Hoang TQD, Nguyen HP (2017) Metal speciation in sediment and bioaccumulation in *Meretrix lyrata* in the Tien Estuary in Vietnam. *Environ Monit Assess*, 189(6), 299.
- [23] Pham MH, Zita S, Tu BM, Pham HV, Renaud FG (2011) Pesticide pollution in agricultural areas of Northern Vietnam: Case study in Hoang Liet and Minh Dai communes. *Environmental Pollution*, 159, 3344-3350.
- [24] Bretveld RW, Thomas CMG, Scheepers PTJ, Zielhuis AG, Roeleveld N (2006) Pesticide exposure: the hormonal function of the female reproductive system disrupted? *Reproductive Biology and Endocrinology*, 4, 30.
- [25] Costa-Böddeker S, Hoelzmann P, Le XT, Hoang DH, Hoang AN, Richter O, Schwalb A (2016) Ecological risk assessment of a coastal zone in Southern Vietnam: Spatial distribution and content of heavy metals in water and surface sediments of the Thi Vai Estuary and Can Gio. *Marine Pollution Bulletin* 114(2), 1141-1151.
- [26] UNEP (2011) Stockholm Convention on Persistent Organic Pollutants.
- [27] Vietnam standard 10:2008/MONRE: National technical regulation on coastal water quality.