

Chemical composition and antimicrobial activity of essential oil from *Pyrethrum pulchrum* Ledeb.

Uugangerel Erdenetsogt¹, Chojjamts Gotov¹, Kerstin Voigt², Stefan Bartram³, Wilhelm Boland³, Enkhmaa Dagvadorj^{1*}

¹School of Bio-Medicine, Mongolian National University of Medical Sciences, Zorig St., Ulaanbaatar 14210, Mongolia

²Leibniz Institute for Natural Product Research and Infection Biology, Hans Knöll Institute (HKI), Adolf-Reichwein Str. 23, Jena 07745, Germany

³Max Planck Institute for Chemical Ecology, Hans-Knöll Str. 8, Jena 07745, Germany

* Corresponding author: enkhmaa.d@mnums.edu.mn; [ORCID ID:0000-0003-0138-4790](https://orcid.org/0000-0003-0138-4790)

Received: 01 December 2018; revised: 07 January 2019; accepted: 08 January 2019

ABSTRACT

The chemical composition and antimicrobial activity of the essential oil from the aerial parts of *Pyrethrum pulchrum* Ledeb. were investigated. Dried plant material was hydro-distilled yielding 0.1% of essential oil. The oil was analyzed by GC-MS techniques. Fifty-five compounds were identified representing 99.7% of the total oil composition. Camphor was the predominant compound (33.9%) followed by linalool (21.1%) and α -pinene (9.0%). The antimicrobial activity of the oil was determined using the disk diffusion method against Gram-positive bacteria (*Bacillus subtilis*, *Staphylococcus aureus* and *Enterococcus faecalis*), Gram-negative bacteria (*Pseudomonas aeruginosa* and *Escherichia coli*), *Mycobacterium vaccae* and fungi (*Candida albicans*, *Sporidiobolus salmonicolor* and *Penicillium notatum*). The essential oil of *P. pulchrum* displays an intermediate activity against selected bacteria.

Keywords: Asteraceae, *Pyrethrum pulchrum*, *Tanacetum*, essential oil, camphor

INTRODUCTION

The Asteraceae also known as Compositae or sunflower family, comprises of the largest family of flowering plants with over 1900 genera and ca. 40000 species [1]. The genus *Tanacetum* Ledeb., formerly *Pyrethrum* Zinn, is a large, poorly defined classification group in the Asteraceae containing 150–200 species distributed over West Asia and Europe [2, 3]. Many species of this genera have traditionally been used as a spicy additive for food, in cosmetics and as herbal remedies due to their biologically active compounds [4]. Moreover, *Tanacetum* has been used as medicinal plants for over 2000 years [5] and it is a known remedy for the treatment of many diseases, including women's ailments, psoriasis, toothache, insect bites, rheumatism, asthma, vertigo and migraine prophylaxis [6, 7], while *Pyrethrum* is used for treating various inflammatory disorders, wound, anthrax, bone fractures as well as reducing fever [8, 9].

The strong and aromatic odor of the *Tanacetum*

species is mainly due to high concentrations of volatile terpene constituents in their essential oil, especially in their leaves and flowers [10]. Camphor, 1,8-cineole, α -thujone, carvone, thymol, *trans*-sabinyl acetate, borneol, caryophyllene oxide, (E)-myroxide, sabinene, bornyl acetate, isopulegone, artemisia ketone, limonene, and camphene were identified as the main constituents of *Tanacetum* species essential oils [3, 11]. It is well known, this genus is also found to contain sesquiterpene lactones, a large group of molecules with several biological activity [11]. Several studies have shown that essential oils and extracts of the genus *Tanacetum* exhibit anti-inflammatory, anticancer, antibacterial, antifungal, anthelmintic, insecticidal and antiprotozoal effects [12–14].

According to literature, six species present in Mongolia that belong to two different genera of *Tanacetum* and *Pyrethrum*, namely *Tanacetum tanacetoides*, *T. vulgare*, *Pyrethrum pulchrum*, *P. lanuginosum*, *P. alatavicum* and *P. changaicum* [15]. *P. pulchrum* is

an impressive flowering plant with about 15-45 cm in height, growing on large-stoned screes at the snow line, usually on glacier moraines in the mountain regions. To our knowledge, no previous studies have been reported neither on the phytochemical analysis of *P. pulchrum* nor on its biological activity. This is the first report on the chemical composition and antimicrobial activity of the essential oil from *P. pulchrum* growing in Mongolia.

EXPERIMENTAL

Plant material: Aerial parts of *P. pulchrum* were collected in the Mongolian Altai mountain region of Govi-Altai province, in July 2012. Plants were dried in shade at ambient temperature. Plant material was authenticated by Prof., ScD, Ch.Sanchir, Institute of General and Experimental Biology of the Mongolian Academy of Sciences.

Isolation of the essential oil: For essential oil extraction, 25 grams of milled plant material were extracted for 2 h in a Clevenger apparatus using hydro-distillation process.

Gas Chromatography-Mass Spectrometry (GC-MS) analysis: The oil was analyzed using a ThermoQuest Trace gas chromatography mass spectrometry system (TRACE 2000 series, Thermo Finnigan, Bremen, Germany) equipped with a ZB-5 capillary column (15 m×0.25 mm, film thickness 0.25 μm, Phenomenex, Torrance, USA). The essential oil was diluted in *n*-hexane at a ratio of 1 to 10. The samples were then injected and separated under programmed conditions to achieve complete separation of the essential oil. The oven temperature was kept at 40 °C for 2 min and then ramped up to 280 °C at 10 °C min⁻¹ followed by a heating to 320 °C with a rate of 30 °C min⁻¹. The injector temperature was set to 220 °C and helium served as

carrier gas with a flow rate of 1.5 mL min⁻¹ and a split ratio of 1:10. Mass spectra were recorded from 33-401 Da (m/z) in full scan mode representing the total ion current chromatogram (TIC, Figure 1).

The relative composition of the essential oil was calculated based on GC peak area. Series of C₈-C₄₀ of *n*-alkanes were injected for the calculation of retention indices (RI) of individual components [16]. Proposals for the identity of the separated compounds were obtained by matching their recorded mass spectra and RIs with library mass spectra and RIs from NIST, Adams, and Massfinder databases [17-19].

Antimicrobial activity assay: *P. pulchrum*, essential oil antimicrobial activity was investigated against Gram-negative bacteria, Gram-positive bacteria and fungi in accordance with reports by Krieg *et al.* [20].

In essence: *Bacillus subtilis* (6633), *Staphylococcus aureus* (134/94 MRSA; 511), *Escherichia coli* (458), *Enterococcus faecalis* (1528, VRSA), *Pseudomonas aeruginosa* (SG137, K799/6) and *Mycobacterium vaccae* (10670) were cultivated on standard I nutrient agar (NA I) in Petri dishes at 37 °C. Antifungal bioassays were conducted on yeast morphology agar (YMA) at 37 °C for *C. albicans* and on malt agar (MA) at 30 °C against *Sporobolomyces salmonicolor* (549) and *Penicillium notatum*. After inoculation, a disc (9 mm in diameter) was removed from the center of the Petri dish and 50 μL of the test solution was added to the cavity. After 18 h of incubation at the respective temperatures the inhibiting areola were measured. Ciprofloxacin (5 μg·ml⁻¹ in deionized water) and amphotericin B (10 μg·ml⁻¹ in DMSO/MeOH 1:1) were used as reference compounds against bacterial and fungal strains, respectively.




Fig. 1. Typical TIC-GC/MS chromatogram of *P. pulchrum* oil separated on a semi - polar ZB - 5 capillary column. Compounds are listed by their peak number in Table 1.

RESULTS AND DISCUSSION

The yield of essential oil obtained by hydro-distillation from aerial parts of *P. pulchrum* was 0.1%. The literature data indicate that the essential oil content in the genus *Tanacetum* ranges between 0.1% to 6.98% [5, 12, 21]. Our result is close to those reported for *T. audibertii* (0.1%) [11] and *T. chilliophyllum* (0.06-0.16%) [22]. The general chemical profiling of the essential oils, the identity and percentage content of the individual components are summarized in Table 1. The GC-MS analyses of *P. pulchrum* essential oil provided the separation of 55 components, representing 99.7% of the oil. In particular, camphor (33.9%), linalool (21.1%), α -pinene (9.0%), (*Z*)- γ -curcumyl 2-methylbutyrate (4.8%), pentylcurcume (3.2%), camphene (2.9%), tricosane (2.8%) and nerolidol (2.6%) were present as

the main constituents.

Generally, essential oils are comprised of two or three major components in relatively high concentrations (20-95%) and other components present in trace levels. On the other hand, quantity and quality of essential oil composition depended considerably on the type of variety, growth stage, time of collection and climatic conditions of the habitat. Table 2 shows a comparison between the major components of *P. pulchrum* essential oil and some species of *Tanacetum*. A comparison of the data presented in this paper with those in the literature for other species of *Tanacetum* shows that there are qualitative and quantitative differences in the levels of some of the compounds present. Our results indicate that the camphor, linalool and α -pinene are three major compounds of *P. pulchrum* essential oil.

Table 1. Constituents of the essential oil from *Pyrethrum pulchrum*

No	RI _{exp.}	RI _{db.}	Compound	RPA, %	No	RI _{exp.}	RI _{db.}	Compound	RPA, %
1	928	932	α -Pinenez	9.0	30	1455	1454	(<i>E</i>)- β -Farnesene	0.24
2	942	946	Camphene	2.85	31	1467	1466	Dehydrosesquicineol	0.17
3	971	974	β -Pinene	1.40	32	1472	1474	β -Chamigrene	0.01
4	990	981	2-Pentylfuran	0.40	33	1477	1484	Germacrene D	0.07
5	1001	1002	α -Phellandrene	1.03	34	1492	1494	α -Selinene	1.80
6	1003	1004	2-methyl-, 2-methylpropyl ester	0.14	35	1563	1561	(<i>E</i>)-Nerolidol	2.63
7	1013	1014	α -Terpinene	0.18	36	1579	1582	Caryophyllene oxide	0.25
8	1022	1020	<i>p</i> -Cymene	0.64	37	1623	1626	Benzophenone	0.06
9	1025	1025	Limonene	0.31	38	1634	1639	allo-Aromadendrene epoxide	0.10
10	1038	1032	(<i>E</i>)- β -Ocimene	0.27	39	1683	1673	α -Bisabolol	0.70
11	1044	1036	Benzene acetaldehyde	0.14	40	1726	1730	Chamazulene	0.52
12	1057	1054	γ -Terpinene	0.22	41	1738	1742	(2 <i>E</i> ,6 <i>E</i>)-Farnesol	0.07
13	1072	1067	<i>cis</i> -Linalool oxide	0.15	42	1747	1741	α -Cyperone	0.54
14	1087	1084	<i>trans</i> -Linalool oxide	0.52	43	1845	1844	2-Pentadecanone, 6,10,14-trimethyl	0.86
15	1100	1095	Linalool	21.12	44	1926	1821	Methyl hexadecanoate	0.17
16	1103	1100	Methyl butyl-2-methyl butyrate-2	0.61	45	1952	1951	Pentylcurcumene	3.21
17	1108	1103	Methyl butyl isovalerate-2	0.19	46	2010	2011	(<i>Z</i>)- γ -Curcumyl 2-methylbutyrate	4.82
18	1139	1141	Camphor	33.94	47	2024	1701	Arglabin	0.57
19	1162	1165	Borneol	0.67	48	2091	2095	Methyl linoleate	0.02
20	1175	1174	Terpinen-4-ol	0.43	49	2098	2100	Heneicosane	1.24
21	1190	1186	α -Terpineol	0.18	50	2139	2132	Linoleic acid	0.78
22	1235	1232	Thymol, methyl ether	0.05	51	2199	2200	Docosane	0.08
23	1243	1244	Geranial	0.06	52	2299	2300	Tricosane	2.78
24	1295	1289	Thymol	0.16	53	2400	2400	Tetracosane	0.15
25	1317	1315	(2 <i>E</i> ,4 <i>E</i>)-Decadienal	0.08	54	2499	2500	Pentacosane	1.36
26	1358	1356	Eugenol	0.01	55	2699	2700	Heptacosane	0.20
27	1375	1373	<i>n</i> -Decanoic acid	1.06					
28	1382	1379	Geranyl acetate	0.38				SUM	99.67
29	1415	1421	(<i>E</i>)- β -Caryophyllene	0.08					

Retention indices (RI) measured on a ZB-5 column; exp: experimental values, db: values from databases; RPA: relative peak area.

Table 2. Major components of essential oils from *P. pulchrum* and some species of *Tanacetum*

Species	Major constituents	References
<i>P. pulchrum</i>	camphor - 33.9%, linalool - 21.1%, α -pinene - 9.0%	our result
<i>T. parthenium</i>	camphor - 45.1%, chrysanthenyl acetate - 21.5%, camphene - 9.6%	[23]
<i>T. parthenium</i>	camphor - 53.8%, <i>trans</i> - β -farnesene - 8.3%	[24]
<i>T. punctatum</i>	camphor - 45.5%, <i>trans</i> - β -farnesene - 7.4%	[24]
<i>T. chiliophyllum</i>	camphor - 32.5%, 1,8-cineol - 16.1%	[14]
<i>T. armenum</i>	camphor - 27%, 1,8-cineol - 11%	[25]
<i>T. vulgare</i>	camphor - 22.3-41.4%, 1,8-cineol - 10.5-26.4%	[26]
<i>T. pinnatum</i>	camphor - 23.2%, α -pinene - 8.5%, camphene - 7.7%	[27]
<i>T. angulatum</i>	1,8-cineol - 75.3%, camphor - 8.1%	[22]
<i>T. praeteritum</i>	borneol - 28%, 1,8-cineol - 12%	[3]
<i>T. audibertii</i>	artemisia ketone - 39.8%, <i>trans</i> -linalool oxide - 32%	[11]
<i>T. messicyticu</i>	thujone - 51%	[3]
<i>T. argyrophyllum</i>	<i>cis</i> -thujone - 69.9%	[13]
<i>T. balsamita</i>	carvone - 52%	[25]
<i>T. longifolium</i>	eudesmol - 22.5%, 1,4-dimethyl azulene - 13.5%, germacrone - 8.2%	[28]

Table 3. Antimicrobial activity of essential oil from *P. pulchrum*

Microorganism	Inhibition zone (mm)		
	Essential oil	Ciprofloxacin	Amphotericin B
<i>Bacillus subtilis</i> (6633)	11	29	-
<i>Staphylococcus aureus</i> (511)	11	19	-
<i>Staphylococcus aureus</i> (134/94, MRSA)	-	-	-
<i>Enterococcus faecalis</i> (1528, VRSA)	14	16	-
<i>Escherichia coli</i> (458)	0	24/32	-
<i>Pseudomonas aeruginosa</i> (K977/61)	17	28/35	-
<i>Pseudomonas aeruginosa</i> (SG 137)	-	25	-
<i>Mycobacterium vaccae</i> (10670)	19	22	-
<i>Sporidiobolus salmonicolor</i>	-	-	19
<i>Candida albicans</i>	12	-	20
<i>Penicillium notatum</i>	-	-	19

One of the major constituents of the *P. pulchrum* is camphor. There are many pharmaceutical applications for camphor such as topical analgesic, antiseptic, antispasmodic, antipruritic, anti-inflammatory, anti-infective and cough suppressant [29]. Linalool, identified as the other major constituent in the essential oil of *P. pulchrum*, is a well-known sedative [30].

In addition, in the essential oil of *P. pulchrum*, compounds with an interesting biological activity, such as arglabin and chamazulene were detected with a relative amount of 0.57% and 0.52%, respectively.

Arglabin belongs to the guaianolide, the class of sesquiterpene lactones, previously isolated from *Artemisia* species [31]. Arglabin shows a promising antitumor activity against different tumor cell lines [32]. Chamazulene is an aromatic sesquiterpene found in a variety of plants including chamomile (*Matricaria*

chamomilla), wormwood (*Artemisia absinthium*) and yarrow (*Achillea millefolium*) [33]. Chamazulene has anti-inflammatory properties in vivo and inhibits the CYP1A2 enzyme involved in leukotriene biosynthesis [34].

During the last decades, antimicrobial plant products have gained special interest because of the resistance to antibiotics that some microorganisms have acquired, the increasing popular concern about the safety of food and the potential impact of synthetic additives on health. One of the novel ways to reduce the proliferation of microorganisms is the use of essential oils [35]. Thus, the antimicrobial activities of the essential oil of *P. pulchrum* were studied against eight bacterial and three fungi strains (Table 3).

Antimicrobial activity was detectable in six out of 11 cases (*Bacillus subtilis*, *Staphylococcus aureus*,

Enterococcus faecalis, *Pseudomonas aeruginosa*, *Mycobacterium vaccae* and *Candida albicans*). The results of the antimicrobial tests of the essential oil from the aerial parts displayed an intermediate effect [20, 36] on the Gram-negative bacteria *P. aeruginosa* (K799/61) and *M. vaccae* (10670). On the other hand, the essential oil exhibited a weak activity against Gram-positive bacteria, *B. subtilis* (6633), *S. aureus* (511) and *E. faecalis* (VRSA), and against the fungus *C. albicans*. No inhibition was observed against *S. aureus* 134/94 (MRSA), *E. coli* (458), *P. aeruginosa* (SG137) and the fungi *S. salmonicolor* and *P. notatum*.

Several studies have also reported that *T. parthenium* essential oils which are rich in camphor, prevent the growth of some bacteria such as *S. aureus*, *S. epidermidis*, *S. flexneri*, *Klebsiella pneumonia*, *E. coli* (25923 and 157), methicillin resistant *S. aureus*, *B. subtilis*, *S. saprophyticus* and *P. aeruginosa* [23, 24, 37]. However, other publications reported that the essential oils of *T. parthenium* (from various regions) were resistant to *E. coli*, *B. subtilis* and *S. aureus* [5, 38].

As already mentioned, natural habitat of plant species was having a bigger impact on the chemical composition and even the biological activity of the essential oils.

CONCLUSION

In the current study, the chemical composition and antimicrobial activity of the essential oil of *P. pulchrum* from Mongolia was investigated for the first time. The yield of essential oil obtained by hydro-distillation from *P. pulchrum* was 0.1 percent of the dried weight. In total, 55 components were identified representing 99.7% of the oil. Camphor (33.9%), linalool (21.1%), α -pinene (9.0%), (*Z*)- γ -curcumyl-2-methylbutyrate (4.8%), pentycurcumene (3.2%), camphene (2.9%), tricosane (2.8%), nerolidol (2.6%), α -selinene (1.8%) and β -pinene (1.4%) were the main components. The essential oil from *P. pulchrum* exhibited an intermediate antimicrobial effect on bacteria *P. aeruginosa* (K799/61) and *M. vaccae* (10670).

REFERENCE

1. The Plant List: Compositae. Royal botanic gardens kew and Missouri botanic garden. Retrieved 07 December 2018.
2. Brown A.M.G., Edwards C.M., Hartman T.P.V., Marshall J.A., et al. (1999) Sexual hybrids of *Tanacetum*: Biochemical, cytological and pharmacological characterization. *J. Exp. Bot.*, **50**(333), 435-444. doi:10.1093/jxb/50.333.435
3. Kumar V., Tyagi D. (2013) Chemical composition and biological activities of essential oils of genus *Tanacetum* - A review. *J. Pharmacogn. Phytochem.*, **2**(3), 159-163.
4. Rohloff J., Mordal R., Dragland S. (2004) Chemotypical variation of tansy (*Tanacetum vulgare* L.) from 40 different locations in Norway. *J. Agric. Food Chem.*, **52**(6), 1742-1748. doi:10.1021/jf0352430
5. Izadi Z., Esna-Ashari M., Piri K., Davoodi P. (2010) Chemical composition and antimicrobial activity of feverfew (*Tanacetum parthenium*) essential oil. *Int. J. Agric. Biol.*, **12**(5), 759-763.
6. Esmaeili M.A., Sonboli A., Ayyari Noushabadi M. (2010) Antioxidant and protective properties of six *Tanacetum* species against hydrogen peroxide-induced oxidative stress in K562 cell line: A comparative study. *Food Chem.*, **121**(1), 148-155. doi:10.1016/j.foodchem.2009.12.022
7. Ernst E., Pittler M.H. (2000) The efficacy and safety of feverfew (*Tanacetum parthenium* L.): An update of a systematic review. *Public Health Nutr.*, **3**(4a), 509-514. doi:10.1017/S1368980000000598
8. Volodya Ts. (2014) *Methods for the clinical usage of Mongolian Medicinal plants*. Ulaanbaatar. ISBN: 978-99929-0-431-3 (in Mongolian)
9. Khaidav T., Menshikova T.L. (1978) *Medicinal Plants in Mongolian Traditional Medicine*. state publishing house, Ulaanbaatar (In Mongolian)
10. Abad M.J., Bermejo P., Villar A. (1995) An approach to the genus *Tanacetum* L. (Compositae): Phytochemical and pharmacological review. *Phyther. Res.*, **9**(2), 79-92. doi:10.1002/ptr.2650090202
11. Maxia A., Sanna C., Piras A., Porcedda S., Falconieri D., et al. (2015) Chemical composition and biological activity of *Tanacetum audibertii* (Req.) DC. (Asteraceae), an endemic species of Sardinia Island, Italy. *Ind. Crops Prod.*, **65**, 472-476. doi:10.1016/j.indcrop.2014.10.039
12. Mohsenzadeh F., Chehregani A., Amiri H. (2011) Chemical composition, antibacterial activity and cytotoxicity of essential oils of *Tanacetum parthenium* in different developmental stages. *Pharm. Biol.*, **49**(9), 920-926. doi:10.3109/13880209.2011.556650
13. Akpulat H.A., Tepe B., Sokmen A., Daferera D., and Polissiou M. (2005) Composition of the essential oils of *Tanacetum argyrophyllum* (C. Koch) Tvetz. var. *argyrophyllum* and *Tanacetum parthenium* (L.) Schultz Bip. (Asteraceae) from Turkey. *Biochem. Syst. Ecol.*, **33**(5), 511-516. doi:10.1016/j.bse.2004.10.006
14. Mohammad bagher Rezaee, Kamkar Jaimand, M.N. (2013) Chemical composition of the essential oil of three *Tanacetum* species from North-West of Iran. *J. Med. Plants By-products*, **2**(2012), 79-82.
15. Grubov V.I. (1982) *Key to the Vascular Plants of Mongolia*, Nauka, Leningrad.
16. Van den Dool H., Kratz P.D. (1963) A Generalization of retention index system including linear temperature programmed gas-liquid partition chromatography. *J Chromatogr*, **11**(4), 463-71.
17. NIST. NIST/EPA/NIH Mass Spectral Library. Retention Index Library, 2014.

18. Adams R.P. (2007) *Identification of essential oil components by gas chromatography/mass spectrometry*, 4th ed., Allured Pub Corp., USA. ISBN-13: 978-1932633214
19. Hochmuth D.H. (2010). MassFinder 4.21, Hochmuth Scientific Consulting, Hamburg, Germany.
20. Krieg R., Jortzik E., Goetz A. A., Blandin S., et al. (2017) Arylmethylamino steroids as antiparasitic agents. *Nat. Commun.*, **8**, 14478 [doi:10.1038/ncomms14478](https://doi.org/10.1038/ncomms14478)
21. Mirjalili M.H., Salehi P., Sonboli A., Mohammadi Vala M. (2007) Essential oil composition of feverfew (*Tanacetum parthenium*) in wild and cultivated populations from Iran. *Chem. Nat. Compd.*, **43**(2), 218-220. [doi:10.1007/s10600-007-0085-2](https://doi.org/10.1007/s10600-007-0085-2)
22. Polatoğlu K., Demirci B., Demirci F., Gören N., Can Başer K.H. (2012) Biological activity and essential oil composition of two new *Tanacetum chiliophyllum* (Fisch. & Mey.) Schultz Bip. var. *chiliophyllum* chemotypes from Turkey. *Ind. Crops Prod.*, **39**, 97-105. [doi:10.1016/j.indcrop.2012.02.005](https://doi.org/10.1016/j.indcrop.2012.02.005)
23. Izadi Z., Aghaalkhani M., Esna-Ashari M., Davoodi P. (2013) Determining chemical composition and antimicrobial activity of feverfew (*Tanacetum parthenium* L.) essential oil on some microbial strains. *Zahedan J. Res. Med. Sci. J.*, **15**(6), 8-13.
24. Shafaghat A., Ghorban-Dadras O., Mohammad hosseini M., Akhavan M., Shafaghatlonbar M., Panahi A. (2017) A comparative study on chemical composition and antimicrobial activity of essential oils from *Tanacetum parthenium* (L.) Schultz. Bip. and *Tanacetum punctatum* (Desr.) Grierson. Leaves from Iran. *J. Essent. Oil Bearing Plants*, **20**(4), 1143-1150. [doi:10.1080/0972060X.2017.1383859](https://doi.org/10.1080/0972060X.2017.1383859)
25. Can Baser K.H., Demirci B., Tabanca N., Ozek T., Goren N. (2001) Composition of the essential oils of *Tanacetum armenum* (DC.) Schultz Bip., *T. balsamita* L., *T. chiliophyllum* (Fisch. & Mey.) Schultz Bip. var. *chiliophyllum* and *T. haradjani* (Rech. fil.) Grierson and the enantiomeric distribution of camphor and carvone. *Flavour Fragr. J.*, **16**(3), 195-200. [doi:10.1002/ffj.977](https://doi.org/10.1002/ffj.977)
26. Mockute, D., Judzentiene, A. (2004) Composition of the essential oils of *Tanacetum vulgare* L. growing wild in Vilnius district (Lithuania). *J. Essent. Oil Res.*, **16**(6), 550-553. [doi:10.1080/10412905.2004.9698795](https://doi.org/10.1080/10412905.2004.9698795)
27. Esmaeili A., Amiri H., Mmaterial P. (2011) The in vitro antioxidant and antibacterial activities of *Tanacetum pinnatum* boiss.grow in Iran. *Bulg. Chem. Commun.*, **43**(4), 532-537.
28. Kumar V., Sharma Y. (2016) Chemical composition and antibacterial activity of essential oils of *Tanacetum longifolium*. *Int. J. Curr. Microbiol. Appl. Sci.*, **5**(10), 836-841. [doi:10.20546/ijcmas.2016.510.091](https://doi.org/10.20546/ijcmas.2016.510.091)
29. Mihaylova H., Fakih H. (2013) Cherubism - A case report. *Rentgenol. i Radiol.*, **52**(3), 203-208.
30. Elisabetsky E. (2002) Chapter 11 Traditional medicines and the new paradigm of psychotropic drug action. *Adv. Phytomedicine*, **1**(C), 133-144. [doi:10.1016/S1572-557X\(02\)80020-4](https://doi.org/10.1016/S1572-557X(02)80020-4)
31. Adekenov S.M., Mukhametzhano M.N., Kagarlitskii A.D., Kupriyanov A.N. (1982) Arglabin - a new sesquiterpene lactone from *Artemisia glabella*. *Chem. Nat. Compd.*, **18**(5), 623-62.
32. Lone S.H., Bhat K.A., Khuroo M.A. (2015) Arglabin: From isolation to antitumor evaluation. *Chem. Biol. Interact.*, **240**, 180-198. [doi:10.1016/j.cbi.2015.08.015](https://doi.org/10.1016/j.cbi.2015.08.015)
33. Conti B., Canale A., Bertoli A., Gozzini F., P.L. (2010) Essential oil composition and larvicidal activity of six Mediterranean aromatic plants against the mosquito *Aedes albopictus* (Diptera: Culicidae). *Parasitol. Res.*, **107**(6), 1455-61.
34. Safayhi H., Sabieraj J., Sailer E.R., Ammon H.P. (1994) Chamazulene: An antioxidant-type inhibitor of leukotriene B4 formation. *Planta Medica.*, **60**, 410-413.
35. Bagci E., Kursat M., Kocak A., Gur S. (2008) Composition and antimicrobial activity of the essential oils of *Tanacetum balsamita* L. subsp. *balsamita* and *T. chiliophyllum* (Fisch. et Mey.) Schultz Bip. var. *chiliophyllum* (Asteraceae) from Turkey. *J. Essent. Oil Bearing Plants*, **11**(5), 476-484. [doi:10.1080/0972060X.2008.10643656](https://doi.org/10.1080/0972060X.2008.10643656)
36. Bauer A.W., Kirby W.M.M., Sherris J.C., Turck K. (1966) Antibiotic Susceptibility test by a standardized single disk method. *Am. J Clin. Pathol.*, **45**, 493-496.
37. Polatoglu K., Demirci F., Demirci B., Gören N., Başer K.H.C. (2010) Antibacterial activity and the variation of *Tanacetum parthenium* (L.) Schultz Bip. essential oils from Turkey. *J. Oleo Sci.*, **59**(4), 177-184. [doi:10.5650/jos.59.177](https://doi.org/10.5650/jos.59.177)
38. Saharkhiz M.J., Satari Morteza., Goudarzi Gholam Reza., Omid Beygi. (2008). Assesment of antibacterial properties of *Tanacetum Parthenium* L. essential oil. *Iran. J. Med. Aromat. Plants*, **24**(1), 47-55.