

RESEARCH ON THE INFLUENCE OF CHEMICAL FERTILIZERS ON TEMPORARY SHORT-EXPLOITATION GRASSLAND

CROITORU IRINA MARINELA, CROITORU ALIN, MILUȚ MARIUS

University of Craiova, Faculty of Agriculture and Horticulture

e-mail: croitoruimarinela@yahoo.com

Keywords: *mixtures, chemical fertilizers, production*

ABSTRACT

Averaged over the two years of experimentation (2012-2013) separate influence of the mixture of species on short-exploitation temporary grassland production was reduced, meaning that the production of the 4 mixtures were close.

Whatever the mixture, the unfertilized variant, considered control, has been acquired very low production of only 2.44 t ha^{-1} d.m. Through fertilization with 100 kg/ha N (background of $50 \text{ kg/ha P}_2\text{O}_5$ and $50 \text{ kg/ha K}_2\text{O}$) were made 5.41 t ha^{-1} d.m., which represents an increase of 2.97 t ha^{-1} or in the relative numbers by 122%.

The dose of 120 kg/ha N applied in two rounds (80 kg in the spring + 40 kg after first harvest) increased production by 2.08 t ha^{-1} , the amount harvested per unit area being 4.52 t ha^{-1} d.m.

INTRODUCTION

To improve highly degraded grasslands can give good results only radical reconstruction, which consists of grubbing and sowing a seed mixture by perennial forage grasses, in other words, setting up a temporary meadows.

Currently in Europe there is a tendency of using mixtures with small number of species or monocultures, a consequence of high levels of nitrogen fertilization which favors a low number of species, of most production transformation in semisilage and increasing of animals number per unit area (Vântu V. și colab., 2004).

An important element to be taken into account when making mixtures for temporary meadows is the way of use of the future grasslands. It up mixture to be used only by mowing, grazing only or in mixed case. Usually mixtures exploited as hay are simple, containing 2-3 species and are sown in arable land for a short period of exploitation (2-3 years). The other two are sown to the place of permanent grassland degraded, on fallow land and contain a total of 5-6 species being exploited 4-6 years or more.

Formixtures designed as pastures used by mixed regime should be used both high and medium-sized species and with low height, ensuring a balance between floors of vegetation.

Currently, when increased requirements for animal products is necessary to intensify fodder production, which can not be achieved only by natural reserves of soil nutrients assimilable. It is necessary to supplement the reserves of nutrients in the soil by applying organic or chemical fertilizers, without which no temporary meadows reveal remarkable productive potential.

In the hills of Oltenia, low soil fertility is a real limiting factor for crop temporary meadows. Therefore, in this area, characterized by very poor and highly acidic soils, fertilizers and amendments are sometimes obligatory for emergence and persistence of eutrophic species by grasses and legumes (Ionescu I., 2001, 2003).

MATERIAL AND METHOD

The experience was placed in the first decade of March in 2012, on a plan area, in subdivided plots with two factors.

Experimental factors were:

Factor A - the mixture with 4 graduations:

- $a_1 = Lolium\ perenne + Trifolium\ pratense$
- $a_2 = Festuca\ pratensis + Trifolium\ pratense$
- $a_3 = Lolium\ perenne + Festuca\ pratensis$
- $a_4 = Lolium\ perenne + Festuca\ pratensis + Trifolium\ pratense$

In the first three mixtures, each of the two species participated in equal proportion (50% each). In the fourth mixture, the proportion of grasses was of 25% each (total 50%) and 50% of legumes.

For seeding were used following varieties:

- Calibra variety for *Lolium perenne*;
- Laura variety for *Festuca pratensis*;
- Nike variety for *Trifolium pratense*.

Factor B - dose of fertilizer, with graduations:

- $b_1 = 0$ - unfertilized
- $b_2 = 100$ kg/ha N
- $b_3 = 80$ kg /ha N in the sprigtime + 40 kg/ha N after first harvest

At the graduations b_2 and b_3 the doses of nitrogen had the background of 50 kg/ha P_2O_5 and 50 kg/ha K_2O .

RESULTS AND DISCUSSIONS

Dry matter production averaged over two years (2012-2013)

Averaged over the two years of experimentation (2012-2013) separate influence of the mixture of species on short-exploitation temporary grassland production was reduced, meaning that the production of the 4 mixtures were close (table 1.).

Table 1

Separate influence of the mixture on the production of short-exploitation temporary meadows from Preajba–Gorj (average 2012-2013, t ha⁻¹ d.m.)

No	Mixture	Yield (t ha ⁻¹ d.m.)	%	Difference	Significance
1	<i>Lolium perenne</i> + <i>Trifolium pratense</i>	4,46	100	-	Control
2	<i>Festuca pratensis</i> + <i>Trifolium pratense</i>	4,60	103	0, 14	-
3	<i>Lolium perenne</i> + <i>Festuca pratensis</i>	3,22	72	- 1, 24	00
4	<i>Lolium perenne</i> + <i>Festuca pratensis</i> + <i>Trifolium pratense</i>	4,23	95	- 0,23	-

DL 5 % = 0,77 t ha⁻¹ d.m.; DL 1 % = 1,09 t ha⁻¹ d.m.; DL 0,1 % = 1,54 t ha⁻¹ d.m

Relative to the amount of dry matter made from the mixture by *Lolium perenne* + *Trifolium pratense* (4.46 t ha⁻¹), which is considered a control, the other three mixtures were obtained the following results:

- the mixture *Festuca pratensis* + *Trifolium pratense* gave an increase of 3% or, in absolute numbers of 0.14 t ha⁻¹, obviously insignificant, the average being 4.60 t ha⁻¹ d.m.;
- mixtures of *Lolium perenne* + *Festuca pratensis* and *Lolium perenne* + *Festuca pratensis* + *Trifolium pratense* gave lower yields than the control, of 1.24 t ha⁻¹ (distinct significant negative difference) and 0.23 t ha⁻¹ (insignificant negative difference).

As is evident from these data, the three mixtures in which he participated red clover ranged from a quantitative perspective, on a higher plane and very close in terms of productive capacity.

Mixture consisting only of grasses (*Lolium perenne* + *Festuca pratensis*) proved less indicated being not differing from the others in terms of quantity.

Consequently, in the case of mixtures of short duration, similar to those by medium and long, similar yields are obtained if in their composition are used species well adapted to the area. This has great practical significance since, facilitates the procurement of seeds, allowing for different formulas, depending on the supply of seed material, with about the same results.

Spectacular differences in the production of short duration grasslands were determined by chemical fertilizers (Table 2).

Table 2

Separate influence of of chemical fertilizers on the production of short-exploitation temporary meadows from Preajba – Gorj (average 2012-2013, t ha⁻¹ d.m.)

Nr. crt.	Agrofond (kg ha ⁻¹)	Yield (t ha ⁻¹ d.m.)	%	Difference	Significance
1	0	2,44	100	-	Control
2	100 N, 50 P ₂ O ₅ , 50 K ₂ O	5,41	222	2,97	***
3	80 N + 40 N, 50 P ₂ O ₅ , 50 K ₂ O	4,52	185	2,08	***

DL 5 % = 0,26 t ha⁻¹ d.m.; DL 1 % = 0,37 t ha⁻¹ d.m.; DL 0,1 % = 0,47 t ha⁻¹ d.m.

On average, regardless of mixture the unfertilized variant, considered as control, achieved a very low production of only 2.44 t ha⁻¹ d.m. Through fertilization with 100 kg / ha N (background of 50 kg / ha P₂O₅ and 50 kg / ha K₂O) were obtained 5.41 t / ha d.m, which represents an increase of 2.97 t ha⁻¹ or in the relative numbers of 122%.

The dose of 120 kg/ha N applied in two rounds (80 kg in the spring + 40 kg after first harvest) increased production by 2.08 t ha⁻¹, the amount harvested per unit area being 4.52 t ha⁻¹ d.m. At the both treatments, increases obtained were very significant.

Results emphasize the urgent need for fertilizer use as a way to obtain normal yields on very poor and acidic soils of the Carpathian area of Oltenia. It should also be pointed out that use of moderate nitrogen dose, springfully applied, gave better results than taking fractional, increasingly frequent phenomenon in the area, due to climate change manifested by long periods of drought accompanied by high temperatures.

Special Efficacy of fertilizers can be better highlighted if we take into account the mixture of species (Table 3).

The table recorded real yields, determined on each variant and not average data, as presented in the two previous tables. From the data it appears that under the 4 mixtures fertilization, without exception, were obtained compared with unfertilized control, very significant increases. They ranged from 1.85 t ha⁻¹ and 3.17 t ha⁻¹. For example, the mixture of *Festuca pratensis* with *Trifolium pratense* without fertilizers, harvested (average over 2 years) 2.75 t ha⁻¹ d.m. Treatment with 100 kg / ha N, 50 kg / ha P₂O₅, 50 kg / ha K₂O increased production by 3.12 t ha⁻¹ and treatment with 80 + 40 kg / ha N (background phospho-potassic) with 2.43 t ha⁻¹, yields per hectare being 5.87 and respectively 5.18 t d.m.

In all four mixtures tested growth caused by dose of 100 kg / ha, fully given in spring was superior to treatment with growth recorded when nitrogen was applied in two rounds.

Table 3

The combined influence of chemical fertilizers with the mixture on the production of short-exploitation temporary meadows from Preajba – Gorj (average 2012-2013, t ha⁻¹d.m.)

Nr. crt.	Agrofond (kg ha ⁻¹)	Mixture	Yield (t ha ⁻¹ d.m.)	%	Difference	Significance
1	0	<i>Lolium perenne</i> +	2,88	100	-	Control
2	100 N, 50 P ₂ O ₅ , 50 K ₂ O	<i>Trifolium pratense</i>	5,69	197	3,21	***
3	80 N + 40 N, 50 P ₂ O ₅ , 50 K ₂ O		4,82	167	1,94	***
4	0	<i>Festuca pratensis</i>	2,75	100	-	Control
5	100 N, 50 P ₂ O ₅ , 50 K ₂ O	+ <i>Trifolium pratense</i>	5,87	213	3,12	***
6	80 N + 40 N, 50 P ₂ O ₅ , 50 K ₂ O		5,18	188	2,43	***
7	0	<i>Lolium perenne</i> +	1,67	100	-	Control
8	100 N, 50 P ₂ O ₅ , 50 K ₂ O	<i>Festuca pratensis</i>	4,47	268	2,80	***
9	80 N + 40 N, 50 P ₂ O ₅ , 50 K ₂ O		3,52	211	1,85	***
10	0	<i>Lolium perenne</i> +	2,46	100	-	Control
11	100 N, 50 P ₂ O ₅ , 50 K ₂ O	<i>Festuca pratensis</i> + <i>Trifolium pratense</i>	5,63	229	3,17	***
12	80 + 40 N, 50 P ₂ O ₅ , 50 K ₂ O		4,57	186	2,11	***

DL 5 % = 0,53 t ha⁻¹ d.m.; DL 1 % = 0,71 t ha⁻¹ d.m.; DL 0,1 % = 0,95 t ha⁻¹ d.m.

The combined influence of mixture with chemical fertilizers is recorded in Table 4.

Also in this table appear concrete production, harvested on the variants. Statistical calculations showed that within each of the 3 agrofunds differences between the control mixture (*Lolium perenne* + *Trifolium pratense*) and the other three mixtures were insignificant, or at the most significant.

Unfertilized *Festuca pratensis* with *Trifolium pratense* mixture gave a small negative difference towards the control, 0.13 t ha⁻¹, the other two variants of fertilization (100 + 40 N or 80 N) proved superiority against the witness with 0.18 and respectively 0.36 t ha⁻¹.

The other two mixtures gave production minus both to unfertilized and at the two fertilized variants. Note that in the case of mixtures of *Lolium perenne* + *Festuca pratensis*, negative differences to control mixture were considered significant by 1.21 to 1.30 t ha⁻¹ d.m.

More important are the differences that can be observed between the yields of the 4 mixtures at different graduations of fertilizer factor and comparing them with the calculated values of the limit differences.

Looking at the table it follows that the yields of dry matter of the 12 variants ranged from 1.67 t ha⁻¹ (the mixture only with the grass species, unfertilized) and 5.87 t ha⁻¹ in mixture *Festuca pratensis* with *Trifolium pratense* fertilized with 100 kg / ha N, 50 kg / ha P₂O₅, 50 kg / ha K₂O. Between these two variants there is a remarkable difference very significant: 5.87 - 1.67 = 4.2 t ha⁻¹.

Table 4

The combined influence of mixture with the chemical fertilizer on the production of short-exploitation temporary meadows from Preajba – Gorj (average 2012-2013, t ha⁻¹d.m.)

Nr. crt.	Mixture	Agrofond (kg ha ⁻¹)	Yield (t ha ⁻¹ d.m.)	%	Difference	Significance
1	<i>Lolium perenne</i> + <i>Trifolium pratense</i>	0	2,88	100	-	Control
2	<i>Festuca pratensis</i> + <i>Trifolium pratense</i>		2,75	95	-0,13	-
3	<i>Lolium perenne</i> + <i>Festuca pratensis</i>		1,67	58	-1,21	0
4	<i>Lolium perenne</i> + <i>Festuca pratensis</i> + <i>Trifolium pratense</i>		2,46	85	-0,42	-
5	<i>Lolium perenne</i> + <i>Trifolium pratense</i>	100 N, 50 P ₂ O ₅ , 50 K ₂ O	5,69	100	-	Control
6	<i>Festuca pratensis</i> + <i>Trifolium pratense</i>		5,87	103	0,18	-
7	<i>Lolium perenne</i> + <i>Festuca pratensis</i>		4,47	78	-1,22	0
8	<i>Lolium perenne</i> + <i>Festuca pratensis</i> + <i>Trifolium pratense</i>		5,63	99	-0,06	-
9	<i>Lolium perenne</i> + <i>Trifolium pratense</i>	80 + 40 N, 50 P ₂ O ₅ , 50 K ₂ O	4,82	100	-	Control
10	<i>Festuca pratensis</i> + <i>Trifolium pratense</i>		5,18	107	0,36	-
11	<i>Lolium perenne</i> + <i>Festuca pratensis</i>		3,52	73	-1,30	0
12	<i>Lolium perenne</i> + <i>Festuca pratensis</i> + <i>Trifolium pratense</i>		4,57	95	-0,25	-

DL 5 % = 0,89 t ha⁻¹ d.m.; DL 1 % = 1,23 t ha⁻¹ d.m.; DL 0,1 % = 1,71 t ha⁻¹ d.m.

Such differences may be found in this experience, given that six variants have achieved yields of 4-5 t ha⁻¹, very close to the maximum. These are the three mixtures of grass + red clover fertilised with nitrogen along with phosphorus and potassium.

From these data it is clear the importance of correct choice of cultivated mixtures meaning use of species well adapted to the area and especially the use of appropriate fertilization which can make the difference between a high and low production, to the standards of a modern agriculture.

CONCLUSIONS

1. Averaged over the two years of experimentation (2012-2013) separate influence of the mixture of species on short-exploitation temporary grassland production was reduced, meaning that the production of the 4 mixtures were close.

2. On average, regardless of mixture the unfertilized variant, considered as control, achieved a very low production of only 2.44 t / ha d.m.

3. Through fertilization with 100 kg / ha N (background of 50 kg / ha P₂O₅ and 50 kg / ha K₂O) were obtained 5.41 t / ha d.m, which represents an increase of 2.97 t / ha or in the relative numbers of 122%.

4. The dose of 120 kg/ha N applied in two rounds (80 kg in the spring + 40 kg after first harvest) increased production by 2.08 t ha⁻¹, the amount harvested per unit area being 4.52 t ha⁻¹d.m. At the both treatments, increases obtained were very significant.

5. Results emphasize the urgent need for fertilizer use as a way to obtain normal yields on very poor and acidic soils of the Carpathian area of Oltenia. It should also be pointed out that use of moderate nitrogen dose, springfully applied, gave better results than taking fractional, increasingly frequent phenomenon in the area, due to climate change manifested by long periods of drought accompanied by high temperatures.

REFERENCES

1. **Ionescu I., 2003** – *Pajiștile temporare în zona subcarpatică a Olteniei*, Edit. Sitech, Craiova;
2. **Ionescu I., 2001** – *Cultura pajiștilor, curs*, Edit. Sitech;
3. **Ionescu I., 2001** – *Pajiștile permanente din nordul Olteniei*, Edit. Universitaria, Craiova;
4. **Pavel C., 1973** – *Pajiștile din zona subcarpatică a Olteniei*, Edit. Scrisul Românesc, Craiova;
5. **Vintu V. și colab., 2004** – *Cultura pajiștilor și a plantelor furajere*, Edit. Ion Ionescu de la Brad, Iași;