



Easton, S., Bartley, D. J., Hotchkiss, E., Hodgkinson, J. E., Pinchbeck, G. L. and Matthews, J. B. (2016) Use of a multiple choice questionnaire to assess UK prescribing channels' knowledge of helminthology and best practice surrounding anthelmintic use in livestock and horses. *Preventive Veterinary Medicine*, 128, pp. 70-77.

There may be differences between this version and the published version. You are advised to consult the publisher's version if you wish to cite from it.

<http://eprints.gla.ac.uk/201451/>

Deposited on: 23 October 2019

Enlighten – Research publications by members of the University of Glasgow_
<http://eprints.gla.ac.uk>

1 **Use of a multiple choice questionnaire to assess UK prescribing channels'**
2 **knowledge of helminthology and best practice surrounding anthelmintic use**
3 **in livestock and horses**

4

5 Stephanie Easton ^a, David J Bartley ^a, Emily Hotchkiss ^a, Jane E Hodgkinson^b, Gina L
6 Pinchbeck ^c, Jacqueline B Matthews ^{a,*}

7

8 *^aMoredun Research Institute, Pentlands Science Park, Midlothian, EH26 0PZ, UK*

9 *^bVeterinary Parasitology, Institute of Infection and Global Health, University of Liverpool,*
10 *Liverpool, L69 7ZJ, UK*

11 *^cVeterinary Epidemiology, Institute of Infection and Global Health, University of Liverpool,*
12 *Liverpool, L69 7ZJ, UK*

13

14 *Corresponding author.

15

16 Address: Moredun Research Institute, Pentlands Science Park, Edinburgh EH26 0PZ, UK.

17 Tel.: + 44 (0)131 445 5111; fax: + 44 (0)131 445 6111.

18 *E-mail address:* jacqui.matthews@moredun.ac.uk (J.B. Matthews).

19

20

21

22

23 **Keywords: livestock, equines, helminths, anthelmintics, prescribing**

24 **ABSTRACT**

25 Grazing livestock and equines are at risk of infection from a variety of helminths, for which
26 the primary method of control has long been the use of anthelmintics. Anthelmintic
27 resistance is now widespread in a number of helminth species across the globe so it is
28 imperative that best practice control principles be adopted to delay the further spread of
29 resistance. It is the responsibility of all who prescribe anthelmintics (in the UK, this being
30 veterinarians, suitably qualified persons (SQPs) and pharmacists) to provide adequate
31 information on best practice approaches to parasite control at the point of purchase. Poor
32 uptake of best practice guidelines at farm level has been documented; this could be due to a
33 lack of, or inappropriate, advice at the point of anthelmintics purchase. Therefore, the aim
34 here was to evaluate levels of basic knowledge of helminthology, best practice guidelines and
35 dispensing legislation among veterinarians and SQPs in the UK, through a Multiple Choice
36 Question (MCQ) test, that was distributed online via targeted emails and social media sites.
37 For each respondent, the percentage correct was determined (for the MCQ test overall and for
38 subsections) and the results analysed initially using parametric and non-parametric statistics
39 to compare differences between prescribing channels. The results showed that channels
40 generally performed well; veterinarians achieved a mean total percentage correct of 79.7%
41 (range 34.0-100%) and SQPs, a mean total percentage correct of 75.8% (range 38.5-100%)
42 ($p=0.051$). The analysis indicated that veterinarians performed better in terms of knowledge
43 of basic helminthology ($p=0.001$), whilst the SQP group performed better on legislation type
44 questions ($p=0.032$). There was no significant difference in knowledge levels of best practice
45 between the two channels. Multivariable linear regression analysis showed that veterinarians
46 and those answering equine questions only performed significantly better than those
47 answering all questions. Based on information gaps identified by analysis of individual
48 questions, a number of areas for improvement in knowledge transfer to both channels are
49 suggested to improve the quality of advice at the point of anthelmintics purchase.

50 **1. Introduction**

51 All grazing livestock and equines are at risk of infection from a variety of helminths, and
52 these parasites can have a substantial negative impact on animal productivity, health and
53 welfare (Love et al., 1999; Charlier et al., 2014). For the last half century, the main method of
54 controlling helminth infections has been the application of anthelmintics. Until recently, for
55 nematodes, three broad spectrum anthelmintic classes were primarily used for this purpose:
56 benzimidazoles, imidazothiazoles/tetrahydropyrimidines and macrocyclic lactones (Prichard,
57 1990). There are also a variety of flukicide products licenced for the treatment and control of
58 trematode infections (Fairweather and Boray, 1999). Recently, two new anthelmintic
59 compounds were licenced and marketed for use in the control of nematode species in sheep in
60 several countries. These compounds are monepantel (Zolvix®, Elanco Animal Health),
61 which is an amino-acetonitrile derivative compound (Kaminsky et al., 2008), and derquantel
62 (a spiroindole), which is combined with the avermectin, abamectin (Startect®, Zoetis) (Little
63 et al., 2011). Resistance (including multi-class resistance) to the traditional classes of
64 anthelmintic is widespread in nematodes of cattle, small ruminants and equids (Bartley, 2011;
65 Sutherland and Leathwick, 2011; Kaplan and Vidyashankar, 2012; Papadopoulos et al., 2012;
66 Matthews, 2014) and there have been reports of resistance to the commonly used, most broad
67 spectrum flukicide, triclabendazole (Fairweather, 2011; Flanagan et al., 2011). Moreover, the
68 first reports of resistance to monepantel are already published (Scott et al., 2013; Mederos et
69 al., 2014). With regard to pigs, resistance has been reported to benzimidazoles and
70 imidazothiazoles/ tetrahydropyrimidines in the EU (Gerwert et al., 2002); however, there is a
71 lack of published data on anthelmintic resistance status in worms this host species.

72 In the UK, the two new classes of anthelmintic can be dispensed only by veterinarians or
73 by pharmacists on veterinary prescription (Prescription Only Medicine – Veterinarian, POM-
74 V). All the other classes are categorised under a Prescription Only Medicine – Veterinary,
75 Pharmacist, Suitably Qualified Person (POM-VPS) label and can be dispensed by Registered

76 Qualified Persons, the types of which are Veterinarians, Pharmacists and Suitably Qualified
77 Persons (SQPs). A SQP must possess a qualification awarded by the Animal Medicines
78 Training Regulatory Authority (AMTRA; <http://www.amtra.org.uk/>), a body appointed under
79 the Veterinary Medicines Regulations by the UK Secretary of State. There are a number of
80 SQP categories in relation to the animals for which they can prescribe; there are various
81 combinations, each of which is assigned a lettered code to describe an individual's permit in
82 the prescribing and supply of medicines (Table 1). Prescribers can be located at a variety of
83 premises such as veterinary surgeries, feed merchants, pharmacies and online. It is the
84 responsibility of prescribers to provide information on current best practice approaches to
85 parasite management at the point of purchase. In the UK, these are described in industry
86 guidelines such as the Sustainable Control Of Parasites in Sheep (SCOPS,
87 <http://www.scops.org.uk/> (Abbott et al., 2012)) and Control Of Worms Sustainably (COWS,
88 <http://www.cattleparasites.org.uk/> (EBLEX, 2010)). No such guidelines exist for horses in
89 the UK, but similar principles apply as laid out in guidelines of the American Association of
90 Equine Practitioners (<http://www.aaep.org/info/parasite-control-guidelines>). For pigs,
91 guidelines are described by the Responsible Use of Medicines in Animals Alliance
92 (<http://www.ruma.org.uk/pigs/anthelmintics-pigs/>). A number of farmer and horse owner
93 surveys indicate that there has been a relatively poor uptake of the guidelines (Morgan and
94 Coles, 2010; McMahon et al., 2013). Risk factors highlighted as important in preserving
95 anthelmintic efficacy are not widely implemented, particular examples being the use of
96 effective quarantine or in the calculation of accurate dose rates (Barton et al., 2006; Relf et al.,
97 2012). It is imperative that best practice control principles be adopted by farmers and horse
98 owners to delay further dissemination of anthelmintic resistance and to preserve efficacy of
99 the currently effective products.

100 In 2013, the British Veterinary Association (BVA) lobbied the UK Veterinary Medicines
101 Directorate (VMD) to make changes to the Veterinary Medicines Regulations with regard to

102 reclassifying all anthelmintics as POM-V. The BVA's argument was underpinned by the
103 assumption that SQP knowledge of parasitology is inferior to that obtained in the course of a
104 full (5 to 6-year) undergraduate veterinary degree (Anon, 2013b). In EU countries such as
105 Denmark and the Netherlands, legislation requires the involvement of a veterinarian and the
106 establishment of a parasitological diagnosis prior to dispensing anthelmintics and prohibits
107 treatment on a prophylactic basis (Nielsen et al., 2006). In a rebuttal from their Secretary
108 General (Anon, 2013a), AMTRA argued that the BVA claims were unsubstantiated and cited
109 facts such as the persistence of anthelmintic resistance in countries employing 'vet-only'
110 prescribing systems, as well as on-going concerns surrounding prescribing practices and
111 resistance with regard to veterinarian-only prescription antimicrobials. With these views in
112 mind, there is little quantitative or qualitative published evidence on which to base the
113 assumptions that either veterinarians or SQPs are better placed to prescribe anthelmintics. For
114 this reason, the aim here was to evaluate levels of knowledge in these channels through
115 execution of a multiple choice question (MCQ) test covering basic helminthology knowledge,
116 prescribing legislation and best practice principles surrounding helminth control.

117

118 **2. Materials and Methods**

119 *2.1. Ethical Statement*

120 Approval for the survey was granted by the UK Department for Environment Food &
121 Rural Affairs (DEFRA) Survey Control Unit. With regards to respondent confidentiality, all
122 information was stored on a secure server at the Moredun Research Institute (MRI). Data on
123 this server is backed up daily at an external site.

124

125 *2.2. Study population*

126 For selection of veterinarians, details of large animal (i.e. livestock and equine) practices in
127 the UK were obtained from the Royal College of Veterinary Surgeons (RCVS) database. The
128 database was cross-checked to group branch practices together, and to omit practices that
129 were not first-opinion practices, such as referral services, or services related to fertility or
130 embryo transfer. The details were also cross-checked with practice websites to establish that
131 the veterinarians contacted currently covered ruminant, pig and/or equine species. This
132 resulted in a list of 755 UK-based veterinarian/practice emails. A further 384
133 veterinarian/practice emails were obtained from a British Equine Veterinary Association
134 (BEVA) list to give a total of 1,139 veterinary surgeons or practices on the mailing list. Note
135 that was not possible to determine the exact number of veterinarians working on each species
136 at each practice. An email inviting the veterinarians to take part in the survey was distributed
137 directly, detailing an introduction to the study and a link to the MCQ test in SurveyMonkey
138 (www.surveymonkey.com, see below). The same link was shared on the pages of the
139 following groups on Twitter (<https://twitter.com/>): the BVA, BEVA, British Cattle Veterinary
140 Association (BCVA), Pig Veterinary Society (PVS), and Sheep Veterinary Society (SVS), as
141 well as the large animal veterinary practice group, XLVets (<http://www.xlvets.co.uk/>). The
142 MCQ test link was also shared via websites or forum pages of the SVS, PVS, BEVA and the
143 BVA. The SQP sample was achieved directly via Mr Stephen Dawson, Secretary General of
144 AMTRA. A total of 2,847 SQPs covering advice provision for the equine, ruminant and pig
145 industries (i.e. E, EA, G, J, K, L and R-SQP license holders (Table 1)) were emailed directly
146 from AMTRA Head Office with the same text and link sent to the veterinarians. The same
147 link was shared on Twitter at <https://twitter.com/SQPWebinars>. Email invitations to take part
148 in the survey were also distributed to SQP members of the Animal Health Distributors'
149 Association (AHDA, <http://www.ahda.co.uk>). This is an organisation comprising UK animal
150 health product distributors and represents 90% of the POM-VPS and Non-Food Animal – Vet,
151 Pharmacist, SQP animal medicines' market.

152

153 2.3. *Study design*

154 The survey comprised several demographic questions to ascertain the profession, age,
155 gender and location of each respondent. These were followed by knowledge-based questions
156 (in MCQ format), all of which were intended to ascertain a respondent's ability to advise on
157 helminth control in line with current UK legislation and best practice guidelines. This was
158 achieved through the assessment of knowledge of basic helminthology (for example, in the
159 identification of common species names of helminths and their host predilection site),
160 epidemiology (for example, the time of year at which acute or chronic helminth-associated
161 disease may be observed), best practice guidelines (for example, advice for quarantine
162 treatments), current UK legislation (for example, meat withdrawal periods of specific
163 products in particular hosts) and the use of parasite diagnostics (for example, the application
164 of faecal worm egg count analysis). The MCQ test (including the correct answers) is
165 available as Supplementary Material (Appendix 1). To maintain consistency between
166 professional groups, a variety of information sources were used to design the test, including
167 veterinary parasitology textbooks, SQP training manuals, and information available from
168 online Continued Professional Development (CPD) guides. Informed consent was sought
169 before answering any questions. There was no time limit imposed upon respondents, who
170 were permitted to return to the form indefinitely (this decision being based on feedback
171 provided in the pilot phase where those working as SQPs stated that this would allow them to
172 be interrupted by customers while completing the survey during working hours). However,
173 respondents were not permitted to return to a page once it had been completed and exited. In
174 the case of all except the demographic questions, respondents were provided with four
175 possible responses to each question, for which only one answer was correct. The respondents
176 were directed to questions based on their particular qualification (in the case of SQPs) or on

177 their client base (in the case of veterinarians). Details of which questions sets were answered
178 by SQPs and veterinarians is presented in Table 1. Of the 78 questions, nine covered general
179 legislation, 38 related to farm animal helminthology (i.e. sheep, cattle and pig associated
180 questions) and 29 related to equine helminthology. In both farm animal and equine sections,
181 the final three questions were “scenario” type questions, based on a real world situation in
182 which a client would be seeking specific advice at the point of dispensing. The survey
183 questions were transferred to and disseminated using the online, cloud-based survey creation
184 software, SurveyMonkey. The test was piloted on a small number of veterinarians and SQPs
185 before being distributed to the participants. The survey was open for 3.5 months (May-
186 September), with monthly reminder emails sent directly to each channel through the
187 distribution lists. Reminders were also posted on Twitter to the relevant group’s sites as
188 detailed above. A flow chart depicting the progression plan for participants of the online
189 survey can be found in Supplementary Materials (Appendix 2). On completion, respondents
190 were redirected to the MRI parasitology homepage (www.moredun.org.uk), last accessed on 8
191 September 2015.

192

193 *2.4. Data Analysis*

194 All responses were recorded using SurveyMonkey software and exported to Microsoft
195 Excel for manipulation (Microsoft Excel for Windows, 2010). Median completion times were
196 calculated for the study sample overall as well as for each channel. As all responses required
197 a single answer, all were coded binomially as 1=correct or 0=incorrect, and the percentages of
198 questions correct overall and in question subsets analysed. Initially, basic descriptive
199 statistics were used. The performance of each channel (‘Vet’; ‘SQP’) and subsequent
200 qualification (SQP) or client base (veterinarian) – ‘farm only’; ‘equine only’; ‘farm and
201 equine’ - were compared in a number of areas including: percentage of questions correct

202 overall, percentage of questions correct for each question type (i.e. helminthology, best
203 practice and legislation type questions) and by host (i.e. questions covering helminthology in
204 relation to cattle (n=12), sheep (n=8), pigs (n=3) and equines (n=16)) and also the average
205 time taken per question, as well as for the entire test. Comparisons were made using two
206 sample t-tests, where data were normally distributed, and a non-parametric alternative, where
207 data were not normally distributed (Mann-Whitney test). Next, univariable and multivariable
208 linear regression analyses were carried out including the demographic variables and channel
209 and host questions to investigate any effect that these may have on the overall percentage of
210 questions answered correctly. A backward elimination was performed whereby variables
211 with p-values >0.05 were removed. All statistical analyses were carried out using Minitab
212 statistical software (Minitab 17; Minitab Inc.). Finally, stage analysis was carried out on the
213 performance on individual questions with the aim of establishing not only differences in
214 knowledge between channels, but to identify gaps in knowledge, within the entire study
215 sample or between channels. This was done by identifying questions where less than 60% of
216 respondents had answered correctly.

217

218 **3. Results**

219 *3.1. Study sample and demographics*

220 A total of 560 respondents were recorded in SurveyMonkey; 342 veterinarians and 218
221 SQPs. Of the respondents that clicked on the link, 227 veterinarians (Table 2) and 57 SQPs
222 (Table 3) completed the test. A total of 244 respondents ceased participation after the
223 demographic questions and a further 32 respondents terminated at the scenario type questions.
224 Further analysis is based on the 284 respondents who completed the test. In the veterinarian
225 sample, the distribution closely matched that of the general veterinary population (taken from

226 those registered with the RCVS) in terms of the proportion of male and female veterinarians
227 registered and the proportionate age range. This was also the case for location distribution,
228 with the majority of veterinarian respondents residing in the southern regions of England. In
229 the SQP sample, the distribution also reflected that of the general SQP population (when
230 compared to those registered with AMTRA), with more female respondents than male and
231 most respondents aged in the range 30-59 years. However, in the SQP sample, slightly more
232 respondents resided in northern England, the Midlands and Scotland, as opposed to southern
233 England, where there is a higher distribution of the general SQP population (as registered
234 with AMTRA).

235

236 *3.2. Completion time*

237 The median completion time overall was 19 min, and ranged from 6 min to 23 h and 45
238 min (Table 4). The median completion time for veterinarians was 18 min (range: 6 min - 22 h
239 and 45 min). The median completion time for SQPs was 31 min (range: 8 min - 23 h and 45
240 min). Further analysis revealed that for veterinarians, 92.5% (n=210) completed the test in
241 less than one hour, 5.7% (n=13) in one to four hours, 0.4% (n=1) in five to twelve hours and
242 1.3% (n=3) in thirteen hours or more. For SQPs, 68.4% (n=39) completed the test in less than
243 one hour, 28.1% (n=16) in one to four hours, 1.8% (n=1) in five to twelve hours and 1.8%
244 (n=1) in thirteen hours or more. Veterinarian completion times were significantly faster than
245 that of the SQPs ($p < 0.01$). When separated into qualification/client base groups (i.e. 'farm
246 only', 'equine only', 'farm and equine'), completion times were significantly different
247 between veterinarian and SQP respondents in all groups, with the exception of the farm only
248 respondents.

249

250 *3.3. Percentage of questions correct*

251 Analysis of the percentage of questions correct overall (Table 5), showed that there was no
252 significant difference between the two channels, although veterinarians (mean 79.7%, range
253 34.0-100%) achieved a higher per cent correct than the SQPs (mean 75.8%, range 38.5-100%)
254 ($p=0.051$). When questions were broken down into respective type, there was a significant
255 difference in performance on helminthology questions ($p=0.001$), with veterinarians
256 achieving a higher percentage correct than SQPs. There was also a significant difference in
257 performance on legislation questions ($p=0.032$), with SQP respondents achieving a higher
258 percentage correct than veterinarians. There was no significant difference in the percentage
259 correct attained by each channel in best practice type questions.

260 In the farm-only question subset, there was no significant difference in percentage of
261 questions correct in both best practice and legislation question types between the veterinarian
262 and SQP respondents. In this subset, there was a significant difference in the percentage
263 correct in helminthology type questions ($p=0.02$), with veterinarians achieving a higher per
264 cent correct than SQPs. In the equine only questions, there was no difference in the
265 percentage correct between the channels in either helminthology or best practice questions;
266 however, there was a significant difference in the percentage correct on legislation type
267 questions ($p=0.01$), with SQPs achieving a higher percentage correct compared to the
268 veterinarian group. There was no significant difference in the percentage of questions correct
269 between the veterinarians and SQPs that answered all questions (i.e. the farm and equine
270 grouping). When the questions were broken down into host species groups (i.e. cattle only
271 questions, sheep only questions, equine only questions, pig only questions and questions
272 relating to more than one host species), analysis of the percentage correct identified that there
273 were no significant differences in performance of the veterinarian and SQP groups (Table 6).

274

275 *3.4. Multivariable linear regression analysis*

276 The final regression model (Table 7) showed that ‘channel’ and ‘question set’ were the
277 only significant variables. The SQP respondents achieved lower scores than the veterinarian
278 respondents and the equine only group performed better than those answering all questions
279 (i.e. farm and equine grouping). Neither age nor gender had an effect on the MCQ test
280 outcomes.

281

282 *3.5. Gaps in knowledge identified through analysis of performance per individual question*

283 The questions for which <60% of the sample achieved correct answers across and within
284 channels can be found in Supplementary Materials (Appendix 3). Of the entire test, there
285 were 11 questions that met this criterion for both channels. There was one additional question
286 for which <60% of respondents were correct in the veterinary channel and a further nine
287 questions with respect to the SQP channel. Of the 11 questions for which <60% of the entire
288 sample gave the correct answer, five were cattle-specific, three were sheep-specific and one
289 each for equine and pig related questions. The majority of these questions covered basic
290 helminthology (i.e. questions relating to helminth species names, predilection sites, times of
291 the year relevant to specific clinical observations and intermediate hosts). There were also a
292 number of legislation questions for which low scores were obtained; these related to the
293 licencing of anthelmintics (with regard to pigs, specifically), and to reporting resistance to the
294 licensing authorities. Of concern, all questions regarding meat withdrawal periods achieved
295 low results. Additionally, with regards to best practice, two scenario questions achieved low
296 results in both channels overall. In the case of the veterinary sample, respondents had lower
297 scores than SQPs on a single equine-related question, relating to the prepatent period of
298 *Parascaris equorum*. The SQPs had lower scores on several questions, particularly those

299 covering Fasciolosis, the predilection site of a helminth in pigs, the migration route of
300 *Strongylus vulgaris*, an equine best practice scenario question and a question relating to the
301 layman's term for *Paramphistomum cervi*.

302

303 **4. Discussion**

304 This study aimed to compare UK-based veterinarian and SQP knowledge with respect to
305 the prescribing and supply of anthelmintic products. The results indicate that, in general, both
306 the veterinarian and SQP groups performed well. The veterinarian group performed better
307 than SQPs on helminthology questions in particular, while the SQP group performed better
308 than veterinarians with regard to legislation knowledge. Equine only respondents performed
309 particularly well in the MCQ test. Stage analysis identified a number of gaps in knowledge
310 relating to the responsible prescribing of anthelmintics in the case of the overall group of
311 respondents, as well as in individual channels. The multivariable linear regression analysis
312 indicated that the veterinary respondents performed better overall than the SQPs after
313 allowing for question set. In reality the difference was small and appeared to be primarily in
314 those questions covering general helminthology knowledge. This observation can perhaps be
315 attributed to differences in the characteristics of veterinarians, particularly, educational ability
316 (Hudson et al., 2009) and a higher level of training as undergraduates. Veterinarians spend
317 time studying helminthology in specific courses within their degree, as well as knowledge
318 garnered in relation to pathogen types, pharmacology, therapeutics and control strategies.
319 Likewise, when the MCQ test results were separated into topics (parasitology, best practice
320 and legislation), the results indicated that veterinarians had a better knowledge of general
321 helminthology than SQPs, while the latter scored a higher percentage correct on legislation
322 questions; in particular, the equine only SQP respondents performed significantly better than
323 the veterinarian group. The results suggested that there was no significant difference in basic

324 knowledge of best practice principles of control between the groups. The generally high
325 results on the best practice questions are encouraging as they indicate that veterinarians and
326 SQPs are informed on the main principles surrounding the responsible prescribing of
327 anthelmintics.

328 The response rate was poor, with relatively low numbers of veterinarians and SQPs
329 proceeding to click the link to the MCQ test. In the case of the SQPs, response rates were
330 particularly disappointing. Only 218 out of the 2,847 emailed AMTRA-registered SQPs
331 covering advice provision for equine, ruminant and pig industries clicked on the MCQ test
332 link. In 2015, according to the AMTRA register database, there were 171 farm only SQPs,
333 1,248 equine only SQPs and 2,242 mixed qualification SQPs – this is equal to 4.7% farm
334 only, 34.1% equine only and 61.2% mixed qualification. These numbers are reflective of the
335 respondents who participated here, as the sample consisted of 7 (12.3%) farm only, 18
336 (31.6%) equine only and 32 (56.1%) mixed qualification SQPs. In the case of the veterinarian
337 respondents, it is not possible to compare the study sample to the entire population of UK
338 registered practicing veterinarians as the RCVS does not hold data regarding the specific
339 practice (i.e. patient species) type of individual veterinarians; however, the general
340 demographic data matched the available RCVS data well. Response rates for web or internet
341 surveys are known to be generally low; it is suggested that this could be due to a lack of
342 information available regarding the most effective ways to motivate individuals to respond in
343 comparison to longer established survey methods such as telephone or mail surveys.

344 The completion rate (i.e. respondents who began the survey and finished the test) was poor
345 in the case of the SQPs, with only 26.1% of respondents completing the test, compared to the
346 veterinarian completion rate which was 66.4%. Most respondents that did not complete the
347 test, did not progress beyond demographic questions. The remainder that did not complete
348 the test terminated when they reached the scenario sections. The length of the test may have

349 discouraged respondents from completing it. It has been suggested that web surveys should
350 adhere to a number of design rules to limit non-response (Dillman et al., 1998). One is to not
351 require that individuals provide an answer before being allowed to proceed to the next
352 question, as this could result in early termination. However, the test here was not a
353 'traditional survey' and this rule would have hindered the ability to calculate an overall score
354 for individuals had they been able to skip questions which they were unsure of. Furthermore,
355 allowing question skipping could result in questions being missed accidentally, adding the
356 issue of ascertaining whether questions had been intentionally skipped or accidentally missed.
357 Self-selection bias is a limitation of online survey research (Wright, 2005) and the findings
358 here must take into account the poor response and completion rate. Individuals who
359 participated and completed the test may have been more likely to do so due to a number of
360 biases, including the likelihood that those who completed it were individuals with greater
361 confidence in their ability and/or knowledge of the subject matter. There is also the
362 possibility that individuals who completed the test were more comfortable with its digital
363 format/question design than those who chose not to. Due to the higher non-completion rates
364 of SQP respondents, these biases could be magnified, leading to a greater possibility of bias
365 amongst this sample. For example, those SQPs who completed the test could be a
366 concentrated sample of the population with high confidence in their knowledge, with more
367 time to dedicate their full attention to the survey or with easier access to the internet. As a
368 result, it would be useful to implement measures in future to increase response and
369 completion rates to try reduce bias; for example, the offering of incentives (this was
370 considered by the authors here; however, was not permitted by the funding body),
371 implementing pre-notifications and/or working closer with stakeholders in encouraging
372 respondents by awarding CPD points to those that complete the test (here, this was offered to
373 AMTRA as an option but was not taken up).

374 Veterinarians completed the test significantly faster than SQPs, though there was only a
375 13-minute difference between the respondent groups in the median completion time. A
376 number of reasons could explain the difference, similar to those considered with regard to
377 differences in test performance. For example, the quicker completion times of veterinarians
378 could be attributed to the fact that those who participated were generally more experienced in
379 MCQ test format or had more time to concentrate on the test without distraction due to the
380 nature of their working environment.

381 The analysis of individual questions highlighted gaps in knowledge that could benefit from
382 having a greater emphasis placed on them during training and CPD programmes. Of the ten
383 questions that did not meet the 60% correct threshold, half covered cattle parasitology. Two
384 questions were within the ‘farm animal scenario’ section, that offered respondents the option
385 to select, ‘More information is required before a recommendation can be given’, which
386 garnered most answers. The issue created as a result of offering this option could be
387 considered similar to that posed by providing respondents with a ‘no-choice option’, whereby
388 respondents, faced with a difficult decision, select this as an easy option, resulting in an
389 absence of the necessary effort needed to come to an appropriate choice resolution (Dhar and
390 Simonson, 2003). However, the basic nature of the remaining three questions on cattle
391 helminths support a recommendation that training needs to be more effectively tailored to the
392 inclusion of helminth species infecting these animals. This might reflect a relative lack of
393 concern regarding anthelmintic resistance within the cattle sector, where, historically, the
394 problem has not been regarded as widespread as in the sheep and equine sectors. It is notable
395 though that anthelmintic resistance in cattle nematodes is being increasingly reported world-
396 wide (Geurden et al., 2015). Another area of concern was in the level of answers correct on
397 questions relating to meat withdrawal period. Low scores here were achieved by
398 veterinarians and SQPs. This is of concern because food safety is an important issue for
399 policy makers and the general public (Cooper et al., 2011). In addition to those questions that

400 achieved low results across both channels, there were a number of questions that achieved
401 low percentage correct totals in each channel. In the SQP sample, 9 questions fell below the
402 60% threshold and the results suggest that SQPs could benefit from adjustments in the
403 syllabus or CPD training regarding information provided on *F. hepatica* epidemiology. This
404 is an important pathogen that has a wide host range (including humans) and has substantial
405 clinical and economic impacts in grazing ruminants (Toet et al., 2014). Other questions
406 achieving low results covered basic parasitology (Latin names or predilection sites), perhaps
407 suggesting that the issues resulting in poor SQP performance on these questions are
408 methodology related (for example, a misunderstanding of the language used in the question
409 or answers provided). Alternatively, this could be the result of differences in learning
410 methods experienced by veterinarians and SQPs; including the shorter time that SQPs spend
411 learning this information compared to veterinarians who are exposed to parasitological-related
412 information at various stages through their full time 5-6 year undergraduate degree. It is
413 important to note that, with the exception of the scenario questions, the questions in this MCQ
414 test demonstrate knowledge/recall that is likely to be at the lower end of the Bloom's
415 Taxonomy cognitive dimension learning pyramid (Krathwohl, 2002). As a result, the
416 outcomes of this study need to be augmented by further work on how this knowledge is
417 applied in 'real life' situations. To do this, the authors are currently exploring (using further
418 questionnaires), the channels' practices, methods and opinions on best practice knowledge
419 dissemination at the point of prescribing by surveying both prescribers and end-users (i.e.
420 farmers and horse owners). Moreover, this MCQ test will be made freely available to those
421 that train UK veterinary undergraduates, SQPs and veterinary pharmacists to be used as a
422 formative revision exercise in large animal helminthology, best practice and anthelmintics
423 prescribing legislation.

424 It is notable that the section of the MCQ test covering equines achieved consistently high
425 results in both channels. It is possible that the greater scores achieved here is due to better

426 knowledge in this specialist host grouping, or methodological reasons, such as the equine
427 question set consisting of easier questions than those of the farm animal question set. The
428 human-horse relationship is one considered to be emotional due to the potential for long term
429 companionship (Schuurman, 2014). It is possible that this type of connection between owner
430 and horse somehow shapes interactions between prescribers and equine clients/customers –
431 resulting in prescribers engaging better in the learning process. Furthermore, horse owners’
432 potential higher awareness of anthelmintic resistance, and therefore expectations for treatment
433 options, could dictate the degree of knowledge required by the prescriber. The increased
434 opportunity to tailor treatment plans to individual animals could also mean that prescribers
435 dealing with equine clients/customers have better knowledge and understanding of
436 information relevant to anthelmintics. It would be advantageous to carry out further
437 investigation to identify factors responsible for this evident better knowledge among equine
438 prescribers so that their principles can be extrapolated and employed in knowledge transfer to
439 other prescribing groups.

440

441 **5. Conclusion**

442 The results show that, despite some apparent gaps in knowledge, both veterinarian and
443 SQP respondents in this study had a sound knowledge of the basic information necessary in
444 the prescription and distribution of anthelmintics. The apparent high level of knowledge in
445 most respondents of best practice principles of control is encouraging with regards to their
446 ability to advise end-users on measures for mitigating the spread of resistance and to preserve
447 efficacy of currently effective anthelmintics. This needs to be balanced, however, against the
448 poor response and completion rates and the unknown quality of knowledge in those who did
449 not perform or complete the test. The identification here of gaps in knowledge in both
450 channels provides the opportunity for improvement in knowledge transfer to further develop

451 the quality of advice at the point of anthelmintics purchase. Furthermore, the evidence
452 gleaned can be put to use in informing policy in terms of those best qualified to prescribe
453 anthelmintics in the UK and abroad.

454

455 **Conflict of Interests**

456 There were no conflicts of interest in the implementation of this project.

457

458 **Acknowledgements**

459 This project was funded by the UK Veterinary Medicines Directorate, an executive agency
460 of the Department for Environment, Food and Rural Affairs. The authors would like to thank
461 the RCVS, AMTRA and AHDA for their assistance in the study and to all of the veterinarians
462 and SQPs who took part in the survey. Additionally, the authors would like to thank the
463 BVA, BEVA, BCVA, PVS, and SVS as well as the large animal veterinary practice group,
464 XLVets, for their assistance in distributing the survey to their members.

465

466 **Reference List**

- 467 Abbott, K.A., Taylor, M.A., Stubbings, L.A., 2012. SCOPS (Sustainable Control of Parasites
468 in Sheep). A Technical Manual for Veterinary Surgeons and Advisers, 4th edition.
469 SCOPS, Malvern.
- 470 AMTRA, 2013. Base Module: The INDUSTRY and its REGULATION. 1-162.
- 471 Anon, 2013a. Anthelmintic prescribing: AMTRA defends the role of SQPs. The Veterinary
472 record 172, 171.
- 473 Anon, 2013b. BVA calls for reclassification of anthelmintics. Veterinary Record 172, 145.
- 474 Bartley, D., 2011. Anthelmintic resistance in cattle nematodes Part 1: A problem for the
475 future? Livestock 16, 19-22.
- 476 Barton, C.H.J., Dale, E.F., Dixon, C., Coles, G.C., 2006. Survey of parasite control on beef
477 farms in south-west England. Veterinary Record 159, 682-684.

- 478 Charlier, J., Van der Voort, M., Kenyon, F., Skuce, P., Vercruysse, J., 2014. Chasing
479 helminths and their economic impact on farmed ruminants. *Trends in Parasitology* 30,
480 361-367.
- 481 Cooper, K.M., Whelan, M., Danaher, M., Kennedy, D.G., 2011. Stability during cooking of
482 anthelmintic veterinary drug residues in beef. *Food Additives and Contaminants* 28.
- 483 Dhar, R., Simonson, I., 2003. The Effect of Forced Choice on Choice. *Journal of Marketing*
484 *Research* 40, 146-160.
- 485 Dillman, D.A., Tortora, R.D., Bowker, D., 1998. Principles for constructing web surveys.
486 Pullman, Washington.
- 487 Fairweather, I., 2011. Liver fluke isolates: a question of provenance. *Veterinary Parasitology*
488 176, 1-8.
- 489 Fairweather, I., Boray, J.C., 1999. Fasciolicides: Efficacy, Actions, Resistance and its
490 Management. *The Veterinary Journal* 158, 81-112.
- 491 Flanagan, A., Edgar, H.W., Gordon, A., Hanna, R.E., Brennan, G.P., Fairweather, I., 2011.
492 Comparison of two assays, a faecal egg count reduction test (FECRT) and a
493 coproantigen reduction test (CRT), for the diagnosis of resistance to triclabendazole in
494 *Fasciola hepatica* in sheep. *Veterinary Parasitology* 176, 170-176.
- 495 Gerwert, S., Failing, K., Bauer, C., 2002. Prevalence of levamisole and benzimidazole
496 resistance in *Oesophagostomum* populations of pig-breeding farms in North Rhine-
497 Westphalia, Germany. *Parasitology Research* 88, 63-68.
- 498 Geurden, T., Chartier, C., Fanke, J., di Regalbono, A.F., Traversa, D., von Samson-
499 Himmelstjerna, G., Demeler, J., Vanimisetti, H.B., Bartram, D.J., Denwood, M.J.,
500 2015. Anthelmintic resistance to ivermectin and moxidectin in gastrointestinal
501 nematodes of cattle in Europe. *International Journal for Parasitology: Drugs and Drug*
502 *Resistance* 5, 163-171.
- 503 Hudson, N.P.H., Rhind, S.M., Moore, L.J., Dawson, S., Kilyon, M., Braithwaite, K., Wason,
504 J., Mellanby, R.J., 2009. Admissions processes at the seven United Kingdom
505 veterinary schools: a review. *Veterinary Record* 164, 583-587.
- 506 Kaminsky, R., Ducray, P., Jung, M., Clover, R., Rufener, L., Bouvier, J., Schorderet Weber,
507 S., Wenger, A., Wieland-Berghausen, S., Goebel, T., Gauvry, N., Pautrat, F.,
508 Skripsky, T., Froelich, O., Komoin-Oka, C., Westlund, B., Sluder, A., Mäser, P.,
509 2008. A new class of anthelmintics effective against drug-resistant nematodes. *Nature*
510 452, 176-180.
- 511 Kaplan, R.M., Vidyashankar, A.N., 2012. An inconvenient truth: Global worming and
512 anthelmintic resistance. *Veterinary Parasitology* 186, 70-78.
- 513 Krathwohl, D.R., 2002. A revision of Bloom's taxonomy: An overview. *Theory into practice*
514 41, 212-218.
- 515 Little, P.R., Hodge, A., Maeder, S.J., Wirtherle, N., Nicholas, D.R., Cox, G.G., Conder, G.A.,
516 2011. Efficacy of combined oral formulation of derquantel-abamectin against the adult
517 and larval stages of nematodes in sheep, including anthelmintic-resistant strains.
518 *Veterinary Parasitology* 181, 180-193.
- 519 Love, S., Murphy, D., Mellor, D., 1999. Pathogenicity of cyathostomin infection. *Veterinary*
520 *Parasitology* 85, 113-121.
- 521 Matthews, J.B., 2014. Anthelmintic resistance in equine nematodes. *International Journal for*
522 *Parasitology: Drugs and Drug Resistance* 4, 310-315.
- 523 McMahon, C., McCoy, M., Ellison, S.E., Barley, J.P., Edgar, H.W., Hanna, R.E., Malone,
524 F.E., Brennan, G.P., Fairweather, I., 2013. Anthelmintic resistance in Northern Ireland
525 (III): uptake of 'SCOPS' (Sustainable Control of Parasites in Sheep) recommendations
526 by sheep farmers. *Veterinary Parasitology* 193, 179-184.
- 527 Mederos, A., Ramos, Z., Banchemo, G., 2014. First report of monepantel *Haemonchus*
528 *contortus* resistance on sheep farms in Uruguay. *Parasites and Vectors* 7, 598-601.

- 529 Morgan, E.R., Coles, G.C., 2010. Nematode control practices on sheep farms following an
530 information campaign aiming to delay anthelmintic resistance. *Veterinary Record* 166,
531 301-303.
- 532 Nielsen, M.K., Monrad, J., Olsen, S.N., 2006. Prescription-only anthelmintics - A
533 questionnaire survey of strategies for surveillance and control of equine strongyles in
534 Denmark. *Veterinary Parasitology* 135, 47-55.
- 535 Papadopoulos, E., Gallidis, E., Ptochos, S., 2012. Anthelmintic resistance in sheep in Europe:
536 a selected review. *Veterinary Parasitology* 189, 85-88.
- 537 Prichard, R.K., 1990. Anthelmintic resistance in nematodes: Extent, recent understanding and
538 future directions for control and research. *International Journal for Parasitology* 20,
539 515-523.
- 540 Relf, V.E., Morgan, E.R., Hodgkinson, J.E., Matthews, J., 2012. A questionnaire study on
541 parasite control practices on breeding Thoroughbred studs. *Equine Veterinary Journal*
542 44, 466-471.
- 543 Schuurman, N., 2014. Blogging situated emotions in human–horse relationships. *Emotion,
544 Space and Society* 13, 1-8.
- 545 Scott, I., Pomroy, W.E., Kenyon, P.R., Smith, G., Adlington, B., Moss, A., 2013. Lack of
546 efficacy of monepantel against *Teladorsagia circumcincta* and *Trichostrongylus*
547 *colubriformis*. *Veterinary Parasitology* 198, 166-171.
- 548 Sutherland, I.A., Leathwick, D.M., 2011. Anthelmintic resistance in nematode parasites of
549 cattle: a global issue? *Trends in Parasitology* 27, 176-181.
- 550 Toet, H., Piedrafita, D.M., Spithill, T.W., 2014. Liver fluke vaccines in ruminants: strategies,
551 progress and future opportunities. *Int Journal for Parasitology* 44, 915-927.
- 552 Wright, K.B., 2005. Researching Internet-Based Populations: Advantages and Disadvantages
553 of Online Survey Research, Online Questionnaire Authoring Software Packages, and
554 Web Survey Services. *Journal of Computer-Mediated Communication* 10.

555

556

557 **Table 1.** Breakdown of criteria that determined the specific question set that respondents were
558 directed to in the MCQ test, based on their qualification (in the case of SQPs) or on their type
559 of clinical practice (in the case of veterinarians)

560

Profession/Qualification	Permissible medicines	MCQ test question set ^c
SQP qualification code ^a		
R-SQP	All VPS medicines	Mixed
G-SQP	VPS: Farm animals and equines only	Mixed
K-SQP	VPS: Farm and companion animals only	Farm only
E-SQP	VPS: Equine and companion animals only	Equine only
L-SQP	VPS: Farm animals only	Farm only
J-SQP	VPS: Equines only	Equine only
JA-SQP	VPS: Equine and avians only	Equine only
EA-SQP	VPS: Companion animals, equines & avians only	Equine only
Veterinarian practice type ^b		
Farm/production animals only	NA	Farm only
Equines only	NA	Equine only
Farm/production animals & equines	NA	Mixed

561

562 ^a SQP qualification codes are based on the specific AMTRA modules which the individual has passed

563 ^b Veterinarian practice types are based on which species individuals work with (i.e. farm/production animals
564 only, equines only or both farm/production animals and equines)

565 ^c Question sets were determined based on animal species that the individual is permitted/likely to prescribe and/or
566 supply to

567 NA - Veterinarians are permitted to prescribe medicines to all species

Demographic category	UK Practising Veterinarian Respondents					Veterinary Sample %	Overall Population % ^b
	Practice Category				Total (n)		
	Farm only	Equine only	Farm and Equine				
	103 (45.4%)	68 (30%)	56 (24.7%)	227 ^a			
Respondent age (years)	18-29	31	15	13	59	26.0	23.1
	30-39	36	31	22	89	39.2	33.9
	40-49	14	13	7	34	15.0	20.9
	50-59	13	6	13	32	14.1	14.9
	60+	8	3	1	12	5.3	7.2
	NA	1	0	0	1	0.4	0.0
Gender	Male	53	23	25	101	44.5	40.7
	Female	50	45	31	126	55.5	59.3
	NA	-	-	-	-	-	0.0
Location	Scotland	7	6	16	29	12.8	10.2
	N. England	14	18	17	49	21.6	19.1
	N. Ireland	1	4	2	7	3.1	3.8
	Wales	2	3	8	13	5.7	5.2
	Midlands	10	15	3	28	12.3	14.2
	S.E. England	22	16	2	40	17.6	34.1
	S.W. England	11	39	6	56	24.7	13.4
	NA	1	2	2	5	2.2	0.0

571 ^a Total respondents that completed the test: i.e. 227/342 (66.4% completion rate)

572 ^b Based on a comprehensive list of UK practising veterinarians - provided by the RCVS on 03/11/2015

573 NA – respondents did not provide demographic data

574 **Table 3.** Survey sample demographics for the SQP group

575

Demographic category	Project sample				SQP Sample %	Overall Population % ^b	
	SQP Category			Total			
	Farm only	Equine only	Farm and Equine				
Total n (%)	7 (12.3%)	18 (31.6%)	32 (56.1%)	57 ^a			
Respondents age (years)	18-29	2	2	6	10	17.5	20.6
	30-39	3	3	5	11	19.3	27.3
	40-49	1	7	4	12	21.1	22.7
	50-59	1	2	11	14	24.6	20.2
	60+	-	4	5	9	15.8	9.0
	NA	-	-	1	1	1.8	0.3
Gender	Male	1	1	16	18	31.6	42.7
	Female	6	17	15	38	66.7	57.2
	NA	-	-	1	1	1.8	0.2
Location	Scotland	3	1	7	11	19.3	9.6
	N. England	6	4	4	14	24.6	18.3
	N. Ireland	-	-	1	1	1.8	6.4
	Wales	-	1	1	2	3.5	13.0
	Midlands	4	1	8	13	22.8	16.3
	S.E. England	4	-	7	11	19.3	19.6
	S.W. England	1	-	4	5	8.8	16.7
	NA	-	-	-	-	0.0	0.0

576

577 ^a Total respondents that completed the test: i.e. 57/218 (26.1% completion rate)

578 ^b Based on a comprehensive list of AMTRA registered SQPs provided by AMTRA on 27/10/2015

579 NA – respondent did not provide demographic data

580

581

582 **Table 4.** Median completion times, ranges for completion times and Mann Whitney results

583 for the MCQ test overall, by channel and by qualification/client base

584

Channel		Completion time (hh:mm:ss)	Range (hh:mm:ss)	Mann Whitney p-value
Overall		00:19:00	00:06:00 – 23:45:00	
Vet		00:18:00	00:06:00 – 22:46:00	<0.001
SQP		00:31:00	00:08:00 – 23:45:00	
Qualification/Client base				Mann Whitney p-value
Farm only	Vet	00:18:00	00:06:00 – 22:46:00	0.3
	SQP	00:21:00	00:12:00 – 01:11:00	
Equine only	Vet	00:13:30	00:06:00 – 04:41:00	0.03
	SQP	00:18:00	00:08:00 – 01:08:00	
Farm and Equine	Vet	00:25:50	00:10:00 – 22:46:00	0.0007
	SQP	00:47:50	00:17:00 – 23:45:00	

585

586

587 **Table 5.** Mean per cent of MCQ test correct by channel (veterinarian, SQP) and question set

588 (farm only, equine only, both farm and equine)

589

Question subsection	Percentage of MCQ test questions correct (95 % CI)		t Test p-value
	Vet (n = 227)	SQP (n = 57)	
Overall	79.7 (78.4-80.9)	75.8 (72.1-79.5)	0.051
Parasitology	80.9 (79.3-82.4)	72.3 (67.4-77.2)	0.001
Best practice	81.7 (80.0-83.4)	78.0 (74.4-81.6)	0.06
Legislation	77.6 (75.9-79.4)	82.1 (78.4-85.9)	0.03
Farm only participants:			
	Vet (n = 103)	SQP (n = 7)	
Parasitology	77.9 (75.7-80.0)	63.8 (53.1-74.5)	0.02
Best practice	83.7 (80.8-86.6)	73.0 (57.5-88.6)	0.1
Legislation	77.4 (74.4-80.4)	78.0 (68.5-87.6)	0.8
Equine only participants:			
	Vet (n = 68)	SQP (n = 18)	
Parasitology	89.7 (87.5-91.9)	84.4 (76.7-92.1)	0.1
Best practice	82.0 (79.4-84.7)	82.1 (76.1-88.1)	0.9
Legislation	78.8 (76-81.6)	87.0 (81.7-92.4)	0.008
Farm and Equine participants:			
	Vet (n= 56)	SQP (n= 32)	
Parasitology	75.6 (72.5-78.6)	69.8 (62.6-76.9)	0.6
Best practice	77.6 (74.4-80.7)	76.7 (71.9-81.6)	0.7
Legislation	76.6 (73.2-79.9)	80.3 (74.6-86.0)	0.2

590

591

592 **Table 6.** Percentage correct across questions covering helminths specific to certain animal
 593 species.

Host species (number of questions per species group)	Mean percentage correct (95% CI)		t Test p –value
	Farm participants ^a :		
	Vet (n=159)	SQP (n=39)	
Cattle (12)	72.7 (70.0-75.4)	62.8 (57.4-68.2)	0.2
Sheep (8)	80.4 (78.1-82.7)	64.7 (59.5-69.9)	0.1
Pigs (3)	78.0 (76.4-79.6)	59.8 (51.1-63.5)	0.1
Mixed (4)	86.6 (84.1-89.2)	76.9 (71.8-82.0)	0.4
	Median percentage correct (range)		Mann Whitney p -value ^c
	Equine participants ^b :		
	Vet (n=124)	SQP (n=50)	
Equine (16)	89.9 (40.0-100.0)	78.0 (56.5-100.0)	0.2

594 ^aAll participants answering farm/production animal questions

595 ^bAll participants answering equine questions

596 ^cAnalyses were carried out using non-parametric methods due to non-normal distribution of the equine
 597 veterinarian data

598

599

600 **Table 7.** Multiple regression model for percentage correct for each respondent on the MCQ
601 test (n =284) including channel (veterinarian and SQP) and question set (All, farm-only and
602 equine-only) variables

	n	Coefficient	Standard error of coefficient	p-value
Intercept		0.76	0.01	<0.001
Channel				
Veterinarians	227	Reference category		
SQPs	57	-0.03	0.02	0.04
Question set				
All	88	Reference category		
Farm	110	0.03	0.02	0.07
Equine	86	0.09	0.02	<0.001

603