



This is a repository copy of *Non-coding RNAs in bone remodelling and bone metastasis : mechanisms of action and translational relevance*.

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/152916/>

Version: Accepted Version

Article:

Puppo, M., Taipaleenmäki, H., Hesse, E. et al. (1 more author) (2019) Non-coding RNAs in bone remodelling and bone metastasis : mechanisms of action and translational relevance. *British Journal of Pharmacology*. ISSN 0007-1188

<https://doi.org/10.1111/bph.14836>

This is the peer reviewed version of the following article: Puppo, M, Taipaleenmäki, H, Hesse, E, Clézardin, P. Non-coding RNAs in Bone Remodelling and Bone Metastasis: Mechanisms of Action and Translational Relevance. *Br J Pharmacol*. 2019., which has been published in final form at <https://doi.org/10.1111/bph.14836>. This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.

Reuse

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

Clézardin Philippe (Orcid ID: 0000-0003-0149-4463)

Non-coding RNAs in Bone Remodelling and Bone Metastasis:

Mechanisms of Action and Translational Relevance

Short title: Non-coding RNAs in Bone Metastasis

Margherita Puppo¹, Hanna Taipaleenmäki², Eric Hesse^{2,3}, Philippe Clézardin^{1,4}.

¹ Department of Oncology and Metabolism, Medical School, University of Sheffield, Sheffield, UK

² Molecular Skeletal Biology Laboratory, University Medical Center Hamburg-Eppendorf, Hamburg, Germany

³ Institute of Molecular Musculoskeletal Research, Faculty of Medicine, LMU Munich, Planegg-Martinsried, Germany

⁴ INSERM, Research Unit UMR_S1033, LyOS, Faculty of Medicine Lyon-Est, University of Lyon 1, Lyon, France

Corresponding author:

Prof Philippe Clézardin

INSERM, UMR_S1033, UFR de Médecine Lyon-Est, 7 Rue Guillaume Paradin,

69372 Lyon cedex 08, France

(Tel.: +33 478785737; e-mail: philippe.clezardin@inserm.fr)

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1111/bph.14836

Department of Oncology and Metabolism, Medical School, University of Sheffield, Beech

Hill Road S10 2RX Sheffield, UK

(Tel.: +44 1142229242; e-mail: p.clezardin@sheffield.ac.uk)

Abbreviations

| | |
|---------------|---|
| AGO | argonaute protein |
| APC | adenomatous polyposis coli |
| BMP | bone morphogenetic protein |
| ceRNA | competing endogenous RNA |
| circRNA | circular RNA |
| COX-2 | cyclooxygenase-2 |
| DKK-1 | dickkopf-related protein-1 |
| ECM | extracellular matrix |
| EV | extracellular vesicle |
| HSC | hematopoietic stem cell |
| IFN | interferon |
| IL | interleukin |
| lncRNA | long non-coding RNA |
| miRNA | microRNA |
| ncRNA | non-coding RNA |
| NSCLC | non-small cell lung cancer |
| NFATc1 | nuclear factor of activated T-cells cytoplasmic 1 |
| OB | osteoblast |
| OC | osteoclast |
| PDGF | platelet-derived growth factor |
| PPAR γ | peroxisome proliferator-activated receptor gamma |
| PRC2 | polycomb complex 2 |
| pre-miRNA | precursor miRNA |
| pre-OB | precursor of osteoblast |
| pre-OC | precursor of osteoclast |
| pri-miRNA | primary miRNA |
| PTHrP | parathyroid hormone-related protein |
| RANKL | receptor activator of nuclear factor kappa-B ligand |
| RBP | RNA-binding protein |
| RISC | RNA-induced silencing complex |
| RNA-seq | RNA-sequencing |
| RUNX2 | runt-related transcription factor 2 |
| SOST-1 | sclerostin-1 |
| TKI | tyrosine kinase inhibitor |
| TNF- α | tumour necrosis factor-alpha |
| Wnt | wingless-INT |

Abstract

Bone metastases are frequent complications in patients with advanced cancer, which can be fatal or may rapidly impede the quality of life of patients. Current treatments for patients with bone metastases are palliative. Therefore, a better understanding of the molecular mechanisms that precede the overt development of skeletal lesions could lead to better therapeutic interventions. In this review, we present evidence that non-coding RNAs (ncRNAs) such as long non-coding RNAs (lncRNAs), microRNAs (miRNAs), and circular RNAs (circRNAs) are emerging as master regulators of bone metastasis formation. We highlight potential opportunities for the therapeutic targeting of ncRNAs. Furthermore, we discuss the possibility that ncRNAs may be used as biomarkers in the context of bone metastases, which might provide insight for improving the response to current bone-targeting therapies.

Keywords

Non-coding RNA, lncRNA, miRNA, circRNA, bone physiology, bone metastasis, biomarkers.

Bullet point summary

What is already known:

- Most studies investigating bone metastasis have focused on the identification and characterization of proteins;
- The role of non-coding RNAs in bone metastasis is only to be unveiled.

What this study adds:

- This study is a comprehensive review of non-coding RNA functions in bone metastasis.

Clinical significance:

- Non-coding RNAs emerge as potential therapeutic targets or biomarkers to predict risk of bone metastasis.

Introduction

Metastasis is a multi-step process by which cancer cells disseminate from primary tumour sites to distant organs (Chaffer et al., 2016). Each type of cancer can metastasize to bone, although some cancer types, such as breast and prostate cancers, are more prone to disseminate to the skeleton (Suva et al., 2011). In bones, metastatic cancer cells tend to colonize vascularized areas, within the red marrow of long bones, sternum, pelvis, ribs and vertebrae (Suva et al., 2011). During bone metastasis formation, cancer cells (i) interfere with normal bone remodelling, (ii) compromise haematopoiesis, and (iii) alter the activity of the immune system, which leads to tumour cell outgrowth and bone degradation (Schmid-Alliana et al., 2018). For these reasons, patients with bone metastases often show severe complications, such as bone pain, pathological fractures, and spinal cord compression, which substantially increase the morbidity and the mortality (Suva et al., 2011).

Approved treatments for bone metastases include bisphosphonates (BPs), an anti-RANKL antibody, and radio-nucleotides, in addition to systemic therapies (Van Poznak et al., 2017). Although bone-targeting agents and systemic therapies increase the overall quality of life of patients with bone metastases, they are only palliative (Van Poznak et al., 2017). Thus, novel approaches to treat patients with bone metastases are needed in the clinics. In order to develop improved therapeutic interventions, it is important to better understand the molecular mechanisms underlying the cellular events preceding the development of overt skeletal lesions.

The majority of studies investigating the mechanisms of bone metastases have focused on the identification and characterization of coding proteins. However, non-coding RNAs (ncRNAs) are emerging as playing a key role in mediating bone metastasis formation (Croset et al., 2015; Hesse & Taipaleenmaki, 2019).

Here we present important aspects of the biogenesis of ncRNAs, and summarise the current knowledge on the role of ncRNAs in bone physiology and bone metastasis formation, highlighting the potential translational impact of ncRNAs as therapeutic targets. Finally, we discuss the possibility of using ncRNAs as potential biomarkers to predict the risk of bone metastases in cancer patients.

Properties of ncRNAs

As suggested by the name, non-coding RNAs (ncRNAs) are a large class of transcripts with no apparent protein-coding potential, but characterized by a functional capacity (**Figure 1**). In general, ncRNAs are classified according to their length as (i) long non-coding RNAs (lncRNAs) (>200 nucleotides), which include long intervening non-coding RNAs (lincRNAs) and long ribosomal RNAs (rRNAs), 18S and 28S, and (ii) small non-coding RNAs (sncRNAs) (<200 nucleotides), which comprises microRNAs (miRNAs), P-element induced wimpy testis (PIWI)-interacting RNAs (piRNAs), short-interfering RNAs (siRNAs), transfer RNAs (tRNAs), 5.8S rRNAs, small nucleolar RNAs (snoRNAs), and small nuclear RNAs (snRNAs) (Bhartiya & Scaria, 2016). Recently, circular RNAs (circRNAs) have been identified as another class of functional and abundant ncRNAs (Bhartiya & Scaria, 2016).

NcRNAs are evolutionary highly conserved and involved in major biological events during embryogenesis, organogenesis, and cell differentiation. In addition, ncRNAs are often implicated in deregulated molecular processes in the context of diseases, including cancer and metastasis (Flynn & Chang, 2014; Hesse & Taipaleenmaki, 2019). Furthermore, ncRNAs (e.g., lncRNAs, miRNAs, and circRNAs) can be secreted by cells, being encapsulated within extracellular vesicles (e.g., microvesicles and exosomes) or in association with high-density lipoproteins or with argonaute (AGO) proteins (Salido-Guadarrama et al., 2014). These ncRNA-containing vesicles or ncRNA-(lipo-)protein complexes can be taken up by recipient cells in which these molecules may exert regulatory functions (Anfossi et al., 2018; Li et al.,

2018b). Secreted ncRNAs can therefore contribute to cell-cell communication and regulate gene expression in recipient cells (Anfossi et al., 2018).

Apart of a small number of lncRNAs such as XIST, HOTAIR, and TERC, the exact function of the vast majority of lncRNAs is unknown (Quinn & Chang, 2016). However, the aberrant expression of lncRNAs appears to play a critical role in tumorigenesis, and 1614 cancer-associated lncRNAs have been reported so far (Lnc2cancer database, <http://www.biogdata.com/lnc2cancer/statistics.html>). The deregulated expression of lncRNAs may potentially contribute to each of the hallmarks of cancer cells (sustained proliferative signalling, evasion from growth suppressors, replicative immortality, invasion and metastasis, induction of angiogenesis, and resistance to cell death) (Quinn & Chang, 2016). This indicates the putative biological importance of these molecules in various cellular processes (Quinn & Chang, 2016). LncRNAs can functionally organize different nuclear sub-compartments, and regulate the chromatin structure accordingly (Akhade et al., 2017). LncRNAs also function as miRNA sponges, thereby acting as competing endogenous RNAs (ceRNAs) in order to regulate gene expression at the post-transcriptional level (Akhade et al., 2017). Furthermore, lncRNAs can trans-activate Staufen1 (STAU1)-mediated mRNA decay (SMD), and serve as a natural antisense transcript for mRNAs and other ncRNAs (Akhade et al., 2017). In addition, lncRNAs can bind to proteins to modulate their functions (Akhade et al., 2017; Pospiech et al., 2018). A further layer of complexity has been provided by recent evidence that some lncRNAs include short open reading frames (ORFs). Thus, these lncRNAs could potentially be translated into peptides and proteins, indicating the current classification of some lncRNAs has some limitations (Niazi & Valadkhan, 2012). Consistently, some annotated lncRNAs have been found to be associated with ribosomes, and to encode for short functional peptides (Anderson et al., 2015; Choi et al., 2018; Huang et al., 2017; Matsumoto et al., 2017; Rossi et al., 2019). However, this ability for some lncRNAs to

be translated into proteins does not preclude their non-coding functions. Moreover, it has been reported that some lncRNAs and their related peptides can cooperate in the same cellular processes (Yu et al., 2017; Rossi et al., 2019).

MiRNAs regulate many cellular events through different molecular mechanisms (**Figure 1**). For instance, miRNAs exert key roles in the post-transcriptional regulation of gene expression by interacting with specific mRNA targets, thereby inhibiting translation and favouring mRNA degradation (Krol et al., 2010). Importantly, one miRNA can modulate the abundance of hundreds of mRNA targets, whereas a single mRNA can be targeted by several miRNAs (Krol et al., 2010). Biologically active miRNAs are the final product of a tightly regulated maturation process. Once a primary miRNA (pri-miRNA) is transcribed by RNA polymerase II, the Drosha-DGCR8 complex processes it as a precursor miRNA (pre-miRNA) (Han et al., 2004). The pre-miRNA is then exported from the nucleus to the cytoplasm where it is cleaved by Dicer, generating a small RNA duplex composed by two RNA strands that are historically referred to 'guide' and 'passenger' strands (Ha & Kim, 2014). The RNA duplex is loaded onto an AGO protein (Salido-Guadarrama et al., 2014) to form an effector complex called RNA-induced silencing Complex (RISC). RISC only retains the guide strand (or mature miRNA) and it is processed into a mature RISC (Nakanishi, 2016). RISC recognises its target(s), usually at the 3' untranslated region (UTR) of mRNAs, based on the base-pairing complementary to miRNA sequence as well as the accessibility of the target site (Jo et al., 2015). Once bound to its target, RISC inhibits protein synthesis either by repressing the translation or by promoting mRNA deadenylation and degradation (Jo et al., 2015). In addition to exerting a repressive function, miRNAs can also increase gene expression in certain cell types and under specific conditions using different mechanisms (Valinezhad Orang et al., 2014). For example, miRNAs can interact with micro-ribonucleoproteins

(miRNPs). These are effector complexes, that can act in trans, thereby promoting the expression of mRNAs (Valinezhad Orang et al., 2014). Furthermore, miRNAs can interact with AU-rich elements (ARE) of mRNAs and thus prevent the ARE-mediated mRNA decay (AMD) (Jing et al., 2005). Changes in miRNA expression or defects in the biogenesis machinery are associated with numerous pathologies, including cancers (Croset et al., 2015; Hesse & Taipaleenmaki, 2019). In particular, deregulated miRNAs can act either as oncogenes or as tumour suppressor genes depending on their target mRNAs (Croset et al., 2015; Hesse & Taipaleenmaki, 2019).

In addition to lncRNAs and miRNAs, circRNAs also act as regulators of gene expression (**Figure 1**). The first circRNA was identified 40 years ago (Hsu & Coca-Prados, 1979). Since then, RNA-sequencing has shown that thousands of circRNAs are generated from protein-coding genes across various eukaryotes (Wilusz, 2018). CircRNAs are generated through a non-canonical splicing mechanism, called back splicing, by which the pre-mRNA splicing machinery covalently links a downstream 3' splice donor to an upstream 5' splice acceptor (which usually belongs to an exon), leading to the formation of a circRNA and a linear RNA (Quan & Li, 2018). CircRNA biogenesis is influenced by transcriptional elongation speed, and it is promoted by some RNA-binding proteins as well as by the presence of inverted repeats in exon extremities or inverted ALU repeat elements in exons (Conn et al., 2015; Quan & Li, 2018). For example, the RNA-binding protein Quaking (QKI) positively regulates circRNA biogenesis during epithelial-to-mesenchymal transition (EMT) (Conn et al., 2015), which is an important process during tumour progression. In general, high circRNA expression levels correlate with low cell proliferation rate and high cell differentiation state (Patop & Kadener, 2018). In tumour cells, the expression of circRNAs is usually reduced, compared to normal cells (Bachmayr-Heyda et al., 2015). In addition to

these general findings, it has been discovered that the expression of specific circRNAs is associated with the expression of oncogenes and/or tumour suppressor genes, suggesting a role in tumour progression (Zhou et al., 2018a). Furthermore, additional mechanisms of action have been proposed, including (i) the regulation of parental gene expression at a transcriptional or post-transcriptional level, (ii) the function as miRNA sponges (or ceRNAs), (iii) the complex formation with other proteins, and (iv) the translation into proteins/peptides (Zhou et al., 2018a).

Functions of ncRNAs in bone physiology

Bone remodelling is a dynamic process that is based on the balanced activities of the bone-forming osteoblasts (OBs) and the bone-resorbing osteoclasts (OCs). The coordinated cellular activities of OBs and OCs are important to ensure bone integrity, repair, and calcium homeostasis (Suva et al., 2011). Any imbalance between OB and OC activities can lead to bone diseases, such as osteoporosis and cancer-associated bone destruction (Suva et al., 2011).

OBs differentiate from bone marrow-derived mesenchymal stem cells (MSCs) in a tightly regulated manner (Almeida et al., 2017). Transforming growth factor- β (TGF- β), bone morphogenetic proteins (BMPs), and Wingless-INT (Wnt) proteins are local factors involved in OB differentiation (Long, 2011). Mature OBs secrete proteins, such as alkaline phosphatase and type I collagen, which are necessary for the formation of the bone extracellular matrix (also known as osteoid), which then becomes mineralized. Afterwards, while most of the OBs die by apoptosis, some reach quiescence as bone lining cells on bone surfaces or become embedded in the bone matrix as osteocytes (Dallas et al., 2013). Osteocytes control calcium and phosphate homeostasis, and serve as mechano-sensors to respond to mechanical loading of the skeleton (Dallas et al., 2013).

OCs are bone-specific multinucleated cells, which derive from the monocyte/macrophage lineage (Long, 2011). Like OBs, OC differentiation and function are regulated by various signals. For instance, the receptor activator of nuclear factor kappa-B ligand (RANKL) and the macrophage-colony stimulating factor (M-CSF) are two of the most important molecules that stimulate OC differentiation and activity (Long, 2011). OCs can also be activated by cytokines, such as interleukin-17 (IL-17), interferon- γ (IFN- γ), and tumour necrosis factor- α (TNF- α) (Long, 2011). OCs remove bone mineral and degrade the collagenous matrix by secreting protons and the proteolytic enzyme cathepsin K, respectively (Long, 2011). As bone is resorbed, growth factors embedded within the bone matrix, such as TGF- β and insulin-like growth factor-1 (IGF-1), are released and stimulate OB function and differentiation (Long, 2011). At the end of their life cycle, OCs undergo apoptosis (Long, 2011). In this respect, ncRNAs regulate OB and OC function, thereby participating in the maintenance of bone integrity, as described below and summarized in **Table 1**.

Role of lncRNAs in bone physiology

Several nuclear and cytoplasmic lncRNAs have been reported to be differentially expressed during osteogenesis (Tye et al., 2018). The first well-characterized lncRNA acting in osteogenesis was Msh Homeobox 1 (Msx1) anti-sense RNA (Msx1-AS RNA) (Blin-Wakkach et al., 2001). Msx1-AS RNA antagonizes the expression of the Msx1 homeodomain protein, which down-regulates the master gene of skeletal differentiation, Runt-related transcription factor 2 (Runx2) (Blin-Wakkach et al., 2001). Thus, the ratio between Msx1 sense and anti-sense RNAs is a very important determinant for the control of skeletal differentiation (Blin-Wakkach et al., 2001).

HOTAIR is another lncRNA involved in skeletal development (Rinn et al., 2007). It acts as a trans-regulator of HoxD gene expression by recruiting the polycomb repressive complex

2 (PRC2) to the HOXD locus (Rinn et al., 2007). In contrast, a recent work suggests that HOTAIR activity in trans is not essential for mouse skeletal development, although significant changes in the expression of the neighbouring genes Hoxc11 and Hoxc12 are detected as a result of HOTAIR deletion in mutant mice, suggesting that HOTAIR can act in cis on the genomic locus (Amandio et al., 2016). Furthermore, HOTAIR regulates the expression of other lncRNAs by targeting their loci, such as the Dlk1-Meg3 locus, the Igf2-H19 locus and the Hottip gene in the Hoxa locus, all of them being involved in bone development (Huynh et al., 2017).

Metastasis-associated lung adenocarcinoma transcript 1 (MALAT1) is an lncRNA that has been originally described in non-small cell lung cancer (NSCLC) (Ji et al., 2003). However, MALAT1 expression is also increased during osteogenic differentiation of adipose-derived mesenchymal stem cells (ADSCs), while miR-30 expression is decreased (Yi et al., 2019). Silencing of MALAT1 or overexpression of miR-30 in ADSCs suppressed the RUNX2-mediated OB differentiation. Overall, the authors showed that MALAT1 sponges miR-30 promote the RUNX2-mediated OB differentiation of ADSCs (Yi et al., 2019).

Maternally expressed gene 3 (MEG3) directly interacts with the transcription factor SOX2, and causes its dissociation from the BMP4 promoter, thereby triggering BMP4 transcription and subsequent differentiation of MSCs into OBs (Zhuang et al., 2015). Interestingly, MSCs from patients with multiple myeloma have frequently low expression of MEG3 in comparison to those obtained from normal donors, suggesting that its down-regulation might explain, at least in part, the lytic nature of skeletal lesions in multiple myeloma (Zhuang et al., 2015).

H19 is involved in developmental biology, and operates through different mechanisms, which are not mutually exclusive (Gabory et al., 2010; Raveh et al., 2015). For example, H19, in concert with miR-675, promotes MSCs differentiation into OBs through inhibition of

the TGF- β 1/Smad3/HDAC signalling pathway (Huang et al., 2015). H19 acts also as a mediator of BMP9-induced osteogenic differentiation of MSCs by modulating Notch signalling-targeting miRNAs (including miR-449b, miR-107, miR-27b, miR-34a, miR-106b, miR-449a, miR-125a, and miR-17) (Liao et al., 2017). Conversely, H19 can act as a ceRNA functioning as a miRNA sponge. For example, H19 activates the Wnt signalling pathway and promotes OB differentiation by binding and thus sequestering miR-141 and miR-22, both of which are negative regulators of osteogenesis (Liang et al., 2016). Similarly, H19 is a positive regulator of mechanical tension-induced osteogenesis of human bone marrow MSCs by acting as a ceRNA for miR-138, which otherwise inhibits MSC differentiation (Wu et al., 2018). Interestingly, H19 also acts as ceRNA for let-7 in primary breast carcinoma (Peng et al., 2017), and let-7 is involved in regulating bone formation (Wei et al., 2014). Although it is not proven yet, it is possible that H19 regulates let-7 function during osteogenesis.

Anti-differentiation ncRNA (ANCR), called also differentiation-antagonizing non-protein coding RNA (DANCR), is required to maintain cells in an undifferentiated state. Its expression is decreased during OB differentiation (Zhu & Xu, 2013). Conversely, its overexpression negatively regulates OB differentiation by inhibiting RUNX2 expression (Zhu & Xu, 2013). Along the same line, ANCR negatively regulates osteogenic differentiation of periodontal ligament stem cells by directly targeting miR-758 (Peng et al., 2018).

RUNX2-AS1 is a lncRNA that has been described as being packed into exosomes secreted by myeloma cells (Li et al., 2018a). These exosomes are transmitted into MSCs and repress OB differentiation of MSCs. More specifically, exosome-derived RUNX2-AS1 forms a duplex with RUNX2 pre-mRNA, thereby repressing RUNX2 expression and subsequent the osteogenic potential of MSCs (Li et al., 2018a).

Linc-ROR is upregulated in MSCs during osteogenesis, and its presence is essential for the correct differentiation of OBs (Feng et al., 2018). Mechanistically, it promotes the osteogenic differentiation of MSCs acting as a ceRNA of two miRNAs, miR-138 and miR-145, known to be negative regulators of osteogenesis by targeting ZEB2 (Feng et al., 2018).

Owing to new bioinformatics techniques, such as next-generation sequencing (NGS) and high-throughput screenings, the list of lncRNAs involved in osteogenesis is expanding very rapidly. An example of a lncRNA identified using these methods is lncRNA-1, whose expression is down-regulated during myoblast differentiation, suggesting a potential role for lncRNA-1 in mesenchymal-to-osteogenic transition (Nardocci et al., 2018). Several other studies used similar bioinformatics strategies, uncovering a potential role for additional lncRNAs in bone physiology and pathology (Gu et al., 2017; Kim et al., 2018; Xie et al., 2016; Zuo et al., 2013).

Function of miRNAs in bone physiology

The first evidence for an important role of miRNAs in skeletal development was obtained by an *in vivo* study that investigated the deletion of the miRNA processing endonuclease Dicer in chondrocytes (Kobayashi et al., 2008). As a result of this deletion, animals had severe skeletal growth defects due to decreased chondrocyte proliferation and accelerated differentiation in post-mitotic hypertrophic chondrocytes (Kobayashi et al., 2008). Dicer deletion in osteoprogenitor cells expressing Osterix also results in severe growth retardation in animals (Raaijmakers et al., 2010). In contrast, conditional deletion of Dicer in mature, osteocalcin-expressing OBs only caused a delay of bone mineralization in mice, which resumed by one month (Gaur et al., 2010). However, a second phenotype of significantly increased bone formation started to develop by two months of age and continued to increase for up to eight months of age, leading to a 2-fold increase in bone mass compared

to control mice (Gaur et al., 2010; Kobayashi et al., 2008; Raaijmakers et al., 2010). Overall, these findings indicate that specific miRNAs must be up- or down-regulated in a timely manner during skeletal development and physiology.

Since the role of miRNAs in bone has already been described in several reviews (Bellavia et al., 2019; Ell & Kang, 2014; Haider & Taipaleenmaki, 2018; Ji et al., 2016), we focus on most recent findings regarding the characterization of miRNA functions. Several miRNAs regulate OB differentiation (Zhang et al., 2011). For example, miR-29b promotes osteogenesis by directly down-regulating known inhibitors of OB differentiation (e.g., HDAC4, TGF- β 3, ACVR2A, CTNNBIP1, DUSP2), through binding to the 3'-UTR seed region of target mRNAs (Li et al., 2009). Along the same line, miR-218 promotes OB differentiation by down-regulating Wnt signalling inhibitors (such as SOST-1, DKK-2, and SFRP2) during the process of osteogenesis (Hassan et al., 2012). Conversely, a panel of miRNAs (including miR-23a, miR-24-2, miR-27a, miR-30c, miR-34c, miR-133a, miR-135a, miR-137, miR-204, miR-205, miR-217, and miR-338) expressed in MSCs was found to negatively control the osteogenic activity of RUNX2 and to impede OB differentiation (Hassan et al., 2010; Zhang et al., 2011). For example, miR-23a, miR-24-2, and miR-27a target SATB2, blocking its synergistic action with the pro-osteogenic transcription factor RUNX2 (Hassan et al., 2010). All of these miRNAs are inversely expressed relative to RUNX2 expression during osteogenesis (Hassan et al., 2010; Zhang et al., 2011). Overall, these findings underline how complex the interactions can be between mRNA targets and miRNAs, and how they can be part of tightly regulated feed-forward mechanisms during osteogenesis. Other miRNAs of particular interest, which have very recently been shown to regulate OB differentiation, are let-7c (Zhou et al., 2018b), the miR-23a cluster (Godfrey et al., 2018), miR-139-3p (Wang et al., 2018b), miR-182-5p (Pan et al., 2018), miR-219a-5p (Aquino-Martinez et al., 2018), miR-221 (Zheng et al., 2018), miR-342-3p (Han et al., 2018),

miR-451a (Karvande et al., 2018), and miR-874 (Lin et al., 2018). In addition, it has recently been reported that miR-21 and miR-218 play a physiologic role in osteocytes, which are differentiated OBs embedded in the bone matrix (Davis et al., 2017). The deletion of miR-21 expression in osteocytes is sufficient to increase apoptosis ex vivo (Davis et al., 2017). Moreover, reduced levels of miR-21 in conjunction with increased miR-218 levels are associated with connexin 43(Cx43)-deficient osteocytes, thereby promoting apoptosis (Davis et al., 2017).

There is ample evidence that miRNAs regulate OC differentiation and activity (Ell & Kang, 2014; Ji et al., 2016). For example, miR-17 mediates the suppression of an osteoclastic protein-tyrosine phosphatase, which in turn promotes the inhibitory effect of EphA4 on OC activity (Lau & Sheng, 2018). MiR-27a is significantly increased during the process of estrogen-mediated inhibition of OC differentiation (Guo et al., 2018b). Mechanistically, miR-27a inhibits peroxisome proliferator-activated receptor gamma (PPAR γ) and adenomatous polyposis coli (APC) expression in OCs through a miR-27a binding site within the 3'-UTR region of PPAR γ and APC. MiR-30a negatively regulates OC differentiation by inhibiting the DC-STAMP-c-Fos-nuclear factor of activated T-cells, cytoplasmic 1 (NFATc1) signalling pathway (Yin et al., 2017). MiR-145 inhibits OC differentiation by decreasing Smad3 expression (Yu et al., 2018). MiR-182 is a negative regulator of protein kinase double-stranded RNA-dependent (PKR), a novel inhibitory factor of OC differentiation, which acts via regulation of the endogenous IFN- β -mediated autocrine feedback-loop (Inoue et al., 2018). MiR-199a-5p expression levels are increased in OCs, compared to their precursors (Guo et al., 2018a). It promotes OC differentiation in vitro by directly targeting transcription factor Mafb (Guo et al., 2018a). Mafb was previously reported to negatively regulate OC differentiation, which suggests that there is a regulatory effect of miR-199a-5p on Mafb mRNA expression (Guo et al., 2018a). MiR-218, beside its pro-osteogenic role (Hassan et

al., 2012), negatively regulates OC differentiation by attenuating the expression of p38 MAPK, c-Fos, and NFATc1 in OCs (Qu et al., 2015). Furthermore, miR-218 directly targets TNF receptor-1, a cell membrane receptor for TNF, leading to the repression of the TNF/nuclear factor- κ B signalling pathway (Wang et al., 2018a). The expression levels of other miRNAs (including miR-33a-5p, miR-133a, miR-141-3p, miR-190, and miR-219-5p) were shown to be decreased after RANKL treatment of OC precursors (Ell et al., 2013). Conversely, miR-16, miR-211, miR-378, and let-7a were significantly up-regulated during OC differentiation (Ell et al., 2013). Additionally, miR-20a and miR-155 regulate autophagy in OCs (Sul et al., 2018; Sun et al., 2015).

As described above, both precursors and mature bone cells express a multitude of miRNAs, which fine-tune OB and OC differentiation. However, exosomal miRNAs derived from bone cells could also participate in bone homeostasis by mediating the crosstalk between OCs and OBs (Li et al., 2016; Xie et al., 2017). For example, OC-derived exosomal miR-214-3p is transferred to OBs to inhibit their activity in vitro and to reduce bone formation in vivo (Li et al., 2016). Conversely, exosomal miR-503-3p from differentiated OBs inhibits RANKL-induced OC differentiation in vitro (Xie et al., 2017).

Relevance of circRNAs in bone physiology

CircRNAs are emerging as important factors that could regulate bone homeostasis. Using microarray analysis, several circRNAs were found to be differentially expressed in MSCs undergoing OB differentiation, compared to their undifferentiated counterparts (Zhang et al., 2018b). Some of these circRNAs possess miRNA response elements enabling their interaction with specific miRNAs, which have osteogenic functions (Zhang et al., 2018b). For example, miR-199b-5p is linked to circIGSF11, and silencing of circIGSF11 promotes OB differentiation and increases miR-199b-5p expression (Zhang et al., 2018b). This study

suggests that circRNA-miRNA interaction actively contributes to the osteogenic differentiation of human MSCs.

Using high-throughput RNA-seq analysis, circRNA expression was measured during the osteogenic differentiation of MC3T3-E1 cells induced by BMP-2 (Qian et al., 2017). Expression of circRNA.5846, circRNA.19142, and circRNA.10042 was significantly increased in BMP2-treated MC3T3E1 cells. However, the signalling pathways controlled by these circRNAs in osteoblastic cells are still unknown (Qian et al., 2017). Similar analyses were conducted in OCs and their precursors following RANKL stimulation (Dou et al., 2016). Over 20 circRNAs were found to be up- or down-regulated at different stages of OC differentiation (Dou et al., 2016). However, no functional analysis was conducted. To further investigate the functions of circRNAs in bone remodelling as well as in cancer, additional investigations are required. The strategy to use high-throughput RNA-seq analyses followed by functional studies (e.g., gene expression modulation of circRNAs in vitro and in vivo) could be useful to address this question.

Dysregulation of ncRNAs in bone metastases

Metastases are formed when cancer cells are released from the primary tumour site, travel through the blood or the lymphatic system, and finally form secondary tumours in distant organs. Bone is the most common metastatic site for primary breast, prostate, and lung tumours, where the bone marrow is a fertile soil for tumour cells to grow (Suva et al., 2011). Tumour cells residing in the bone marrow alter the functions of OCs and OBs (Suva et al., 2011). For example, breast and lung cancer cells cause bone destruction by enhancing OC activity, whereas prostate cancer cells form mixed lesions with osteosclerotic components by increasing OB activity (Suva et al., 2011). Bones weakened due to skeletal metastases are prone to skeletal-related events, such as fractures, hypercalcaemia, cancer-induced bone pain

and spinal cord compression, which substantially increase the morbidity and mortality of patients with advanced cancer (Suva et al., 2011).

At the bone metastatic site, tumour cells induce a distortion of bone remodelling, and bone-derived growth factors released from resorbed bone promote tumour growth (Guisse et al., 1996; Suva et al., 2011). This concept of “vicious cycle” was first proposed by Dr Greg Mundy (University of Texas Health Science Centre, San Antonio, USA). Although it is difficult to identify a starting point for the “vicious cycle”, it is well known that metastatic cancer cells are able to produce a number of factors, such as parathyroid hormone-related protein (PTHrP), interleukin (IL)-11, and prostaglandin E2 that mediate bone resorption by increasing the OB-dependent production of RANKL (Croucher et al., 2016; Dougall, 2012). In turn, RANKL binds to its receptor RANK on OC precursors, leading to the formation of new OCs, as well as promotes the activity of mature OCs, thus favouring OC-mediated bone resorption (Croucher et al., 2016). Under physiological conditions, osteoprotegerin (OPG) binds to RANKL and restricts its activity (Croucher et al., 2016). PTHrP regulates RANKL expression in OBs and stromal cells (Dougall, 2012). Moreover, tumour-derived PTHrP reduces the OB-dependent OPG production, thereby further promoting bone resorption. Tumour cells also secrete IL-8 that has a direct effect on OC differentiation and activation (Suva et al., 2011). Tumour cells not only stimulate OC activity, but also inhibit OB function, thereby worsening the imbalance between bone formation and bone resorption, which promotes bone destruction (Croucher et al., 2016). Different factors produced by tumour cells have been shown to suppress OB differentiation. These include endothelin-1, dickkopf-related protein 1 (DKK-1), sclerostin-1 (SOST-1), activin A, and noggin (NOG) (Croucher et al., 2016). Moreover, bone tissue is an abundant reservoir for growth factors, including TGF- β , IGF-1, BMPs, platelet-derived growth factor (PDGF), and epidermal growth factor (EGF), which are entrapped in the bone matrix and become released in the bone microenvironment

upon matrix degradation by active OCs (Croucher et al., 2016). In turn, these bone-derived growth factors promote tumour growth.

A large body of evidence suggests that ncRNAs regulate the various molecular mechanisms of bone metastasis formation, although only a few studies investigated the role of lncRNAs and circRNAs in bone metastasis so far. By contrast, several articles already reviewed the role of miRNAs in bone metastasis (Baier & Wan, 2016; Browne et al., 2014; Cheung et al., 2017; Croset et al., 2015; Ell & Kang, 2014; Hesse & Taipaleenmaki, 2019). In cancer and metastasis, a loss of tumour-suppressive miRNAs and an increased expression of oncogenic miRNAs are part of the disease mechanism. MiRNA functions as well as those of other ncRNAs in bone metastasis are summarized in **Table 2**. Herein, we will focus on most recent and/or pivotal findings, and show that the dysregulation of ncRNAs in tumour cells or bone cells as well as the shuttling of ncRNAs between tumour and bone cells may lead to the establishment of a cancer niche. This environment is conducive for tumour cells to grow and induce the formation of skeletal lesions.

OC- and OB-derived miRNAs and bone metastases

The role of OC-derived miRNAs in the development of osteolytic bone metastasis has been examined in detail by the group of Dr Yibin Kang (Princeton University, NJ, USA), hypothesizing that miRNAs might be possible therapeutic targets during the differentiation of OC precursors into mature OCs. In this respect, miRNA microarray profiling of mature OCs was performed after exposure of OC precursors to conditioned medium from breast cancer cells (Ell et al., 2013). Four miRNAs (miR-133a, miR-141-3p, miR-190, and miR-219-5p) whose expression was consistently down-regulated were selected for further investigation (Ell et al., 2013). Systemic treatment of healthy mice with each of these miRNAs (whose aim is to replenish the lost miRNA expression during OC differentiation) inhibited bone

resorption *in vivo*. Consistently, treatment of tumour-bearing mice with miR-141 or miR-219 significantly reduced the formation of osteolytic lesions caused by breast cancer cells (Ell et al., 2013). In contrast, systemic treatment of tumour-bearing animals with miR-133a or miR-190 had no effect on the development of bone metastases, compared to control (Ell et al., 2013). Nevertheless, these miRNAs were inhibiting OC differentiation *in vitro* through the functional targeting of essential OC genes, including *Mitf*, *Traf6*, *Calcr*, and *Mmp14*. The observation that miR-133a and miR-190 did not inhibit the formation of lytic bone metastases, whereas they did inhibit normal bone resorption, was of particular interest. Of note, the half-lives of miR-133a and miR-190 in the circulation were shorter than those of miR-141 and miR-219 (Ell et al., 2013). These differences in pharmacokinetics of miRNAs could therefore explain these unexpected results, suggesting that inhibition of osteolytic lesions may require greater miRNA levels to reach the bone than those required for the inhibition of normal bone resorption (Ell et al., 2013).

MiR-34a is another miRNA that is down-regulated during OC differentiation. It inhibits OC differentiation by directly targeting the pro-osteoclastic factor TGIF2 (TGF- β -induced factor 2) (Krzyszinski et al., 2014). The pharmacological administration of a miR-34a mimic delivered in nanoparticles attenuates the formation of breast cancer bone metastases in mice (Krzyszinski et al., 2014). A phase I, open-label, dose-escalation study investigated the safety, pharmacokinetics, and pharmacodynamics of a miR-34a mimic (MRX34), encapsulated in lipid nanoparticles, in patients with advanced or metastatic cancer (Beg et al., 2017). However, the clinical trial was terminated prematurely due to cases of immune-related serious adverse events.

MiR-214-3p expression in OCs is significantly up-regulated in bone specimens from breast cancer patients with bone metastases (Liu et al., 2017). Additionally, the conditioned medium from MDA-MB-231 breast cancer cells stimulates miR-214 expression in mature

OCs during OC differentiation. Mechanistically, miR-214 directly targets the TNF receptor associated factor 3 (TRAF3), a negative regulator of OC differentiation, thereby promoting the formation of mature OCs. The treatment of animals with antagomiR-214-3p encapsulated within an OC-targeting delivery system (D-Asp₈-liposome) attenuates the formation of osteolytic lesions caused by breast cancer cells (Liu et al., 2017). Additionally, OC-derived exosomal miR-214-3p was transferred to OBs, and inhibited OB activity in vitro and reduced bone formation in vivo (Li et al., 2016). Of note, miR-214 is expressed by OBs and inhibits OB differentiation in vitro and bone formation in vivo, through repression of activating transcription factor 4 (Atf4), which is one of the main transcription factors required for OB function (Wang et al., 2013). Taken together, these findings suggest that the therapeutic inhibition of OC- or OB-derived miR-214-3p could be a strategy to impede, respectively, the development of lytic or sclerotic bone metastases (Haider & Taipaleenmaki, 2018; Li et al., 2016; Wang et al., 2013).

Tumour-derived miRNAs and homing of tumour cells to the bone marrow

Searching for regulators of tumour spreading, Tavazoie et al. (2008) identified a series of miRNAs (miR-126, miR-206, and miR-335) that were poorly expressed in lung- or bone-tropic breast cancer cells. Overexpression of these miRNAs in breast cancer cells reduced lung and bone metastasis formation in animals (Tavazoie et al., 2008). MiR-126 suppressed metastases by inhibiting tumorigenesis in vivo and tumour cell proliferation in vitro. In contrast, miR-206 and miR-335 did not affect tumorigenesis, yet suppressed metastasis by inhibiting tumour cell migration and invasion in vitro. MiR-335 directly targets the 3'UTR of mRNAs for transcription factor SOX4 and tenascin C (TNC), both factors being involved in promoting tumour cell migration and invasion (Tavazoie et al., 2008). Elevated expression levels of let-7 in breast cancer cells were also reported to inhibit bone metastasis formation

by directly targeting HMGA2, a chromatin remodelling protein that activates pro-invasive genes (Ell & Kang, 2014). Conversely, miR-10b activated by transcription factor Twist facilitates lung and bone metastasis formation in animals by promoting MDA-MB-231 breast cancer cell invasion through direct targeting of the mRNA encoding homeobox D10 (HOXD10), resulting in an increased expression of the pro-metastatic gene Ras homolog gene family member C (Rhoc) (Croset et al., 2015; Ell & Kang, 2014; Ma et al., 2007). Similarly, miR-17 promotes breast cancer cell migration/invasion in vitro and bone metastases in vivo through repression of the type II TGF- β receptor (T β R2) (Croset et al., 2015). Along the same line, miR-21 promotes the colonization of breast cancer cells in bone by inhibiting the expression of the anti-metastatic genes PTEN, PDCD4, and SPRY2, which in turn leads to an enhanced tumour cell invasion and metastasis formation (Sahay et al., 2015).

In prostate cancer, the ectopic expression of miR-143 and miR-145 in PC-3 tumour cells inhibits bone metastasis formation in vivo by preventing EMT, and by reducing tumour cell migration (Croset et al., 2015; Ell & Kang, 2014). These inhibitory effects of miR-145 on EMT and tumour cell migration occur through repression of the oncogenic protein HEF1 (Croset et al., 2015; Ell & Kang, 2014). MiR-203 is also a tumour suppressor whose expression is inversely correlated with prostate cancer bone metastases (Siu et al., 2018). Reconstitution of metastatic prostate cancer cells with miR-203 suppresses bone metastasis formation in animals by inhibiting EMT, invasion, and motility (Saini et al., 2011). The capacity of miR-203 to inhibit bone metastasis formation has been associated with its ability to directly repress Survivin/BIRC5 expression, an inhibitor of apoptosis (Saini et al., 2011). Additionally, miR-203 represses the activation of the EGFR signalling pathway in prostate cancer cells (Siu et al., 2018). A role for miR-203 in the inhibition of bone metastases caused by non-small cell lung cancer has also been reported (Wei et al., 2017). Several other

miRNAs have been shown to control early stages of bone metastasis formation, whose functions are summarized in **Table 2**.

MSC-derived miRNAs

Entering a foreign environment, such as bone, poses tumour cells with numerous challenges regarding survival and proliferation. It is therefore likely that tumour cells undergo dormancy as a mechanism to facilitate survival until the environmental conditions are sufficiently permissive for tumour outgrowth.

Exosomal miR-23b and miR-222/223 released from MSCs have been reported to stimulate breast cancer dormancy in the bone marrow (Bliss et al., 2016; Ono et al., 2014). For example, miR-23b induces dormancy through the suppression of the myristoylated alanine-rich C-kinase substrate (MARCKS) target gene, which encodes a protein that promotes cell proliferation and motility (Ono et al., 2014).

MiRNAs regulate the adaptation of tumour cells to the bone microenvironment

While residing in the bone marrow, tumour cells exit and re-enter a dormant state. During this phase, cancer cells undergo a selection and acquire the capacity for metastasis formation. Specifically bone metastatic cancer cells express genes that are normally expressed by bone cells (a process called osteomimicry), which endow cancer cells with a full competence for outgrowth in the bone marrow (Croset et al., 2015; Hesse & Taipaleenmaki, 2019). As described below, several miRNAs were reported to regulate cancer cell osteomimicry.

RUNX2 plays a pivotal role in OB differentiation and it is aberrantly expressed in bone-tropic breast- and prostate cancer cells (Haider & Taipaleenmaki, 2018). In bone-tropic tumour cells, RUNX2 promotes the progression of bone metastases and the development of osteolytic lesions (Haider & Taipaleenmaki, 2018). RUNX2 expression is repressed by

several miRNAs, including miR-135, miR-203, and miR-466. In this respect, miR-135 and miR-203 expression is absent in bone-tropic breast cancer cells and in tissue biopsies from breast cancer bone metastases that express RUNX2 (Haider & Taipaleenmaki, 2018). Conversely, transfection of miR-135 and miR-203 mimics in bone-tropic breast cancer cells results in a decreased skeletal tumour growth and decreased formation of osteolytic lesions in tumour-bearing mice (Taipaleenmaki et al., 2015). By repressing RUNX2, miR-135 and miR-203 indirectly inhibit the expression of genes (Il-11, Mmp-13, PthrP) that are associated with bone metastases (Taipaleenmaki et al., 2015). Similarly, miR-203 and miR-466 suppress prostate cancer bone metastases by directly targeting RUNX2 (Colden et al., 2017; Saini et al., 2011). Furthermore, restoring miR-203 in bone-tropic prostate cancer cells represses the expression of other genes associated with osteomimicry, including homeobox protein Dlx-5 (DLX5), osteopontin (SPP1), and osteocalcin (BGLAP) (Saini et al., 2011). Similarly, reconstitution of prostate cancer cells with miR-466 represses SPP1 and BGLAP expression (Colden et al., 2017).

MiRNA members of the miR-30 family (miR-30a, miR-30b, miR-30c, miR-30d, and miR-30e) are suppressors of breast cancer bone metastasis (Croset et al., 2018). Expression of these miRs in bone-tropic triple-negative breast cancer cells was profoundly decreased. Conversely, overexpression of miR-30 family members in triple-negative breast cancer cells results in the reduction of bone metastasis formation in vivo. In vitro, miR-30 family members inhibit tumour cell invasion, and restore bone homeostasis by reversing the effects of tumour cell-conditioned medium on OC and OB formation. Mechanistically, miR-30s directly target a number of genes associated with the stimulation of OC differentiation (IL-8, IL-11), the inhibition of OB differentiation (DKK-1), tumour cell osteomimicry (RUNX2, CDH11), and invasiveness (CTGF, ITGA5, ITGB3) (Croset et al., 2018). Among these genes, silencing of CDH11 or ITGA5 in triple-negative breast cancer cells recapitulates the

inhibitory effects of miR-30 family members on bone metastasis formation *in vivo*. Overall, these findings provide evidence that miR-30 family members employ multiple mechanisms to impede breast cancer bone metastases, including osteomimicry (Croset et al., 2018). Of note, lncRNA MALAT1 acts as a sponge for miR-30, thus promoting OB differentiation of ADSCs (see section on bone physiology) (Yi et al., 2019). MALAT1 is overexpressed in several cancers, including triple-negative breast cancer (Jadaliha et al., 2016). It is therefore tempting to speculate that in triple-negative breast cancer MALAT1 overexpressing cells may be responsible for miR-30 down-regulation, which then facilitates tumour cell osteomimicry.

As previously described, miR-218-5p promotes OB differentiation by down-regulating Wnt signalling inhibitors (e.g., SOST-1, DKK-2, and sFRP2) during the process of osteogenesis (Hassan et al., 2012). MiR-218-5p also enhances Wnt activity in osteotropic breast cancer cells. Moreover, the expression of genes associated with osteomimicry (CXCR-4, BSP, and SSPP1) and OC-mediated bone resorption (PTHrP) is increased in miR-218-expressing breast cancer cells (Haider & Taipaleenmaki, 2018; Hassan et al., 2012; Taipaleenmaki et al., 2016). Transfection of bone-tropic breast cancer cells with anti-miR-218-5p decreases Wnt activity, and reduces skeletal tumour burden and cancer-induced osteolysis *in vivo* (Taipaleenmaki et al., 2016). Furthermore, miR-218 secreted from breast cancer cells can directly down-regulate type I collagen expression by OBs, thereby impairing bone integrity (Liu et al., 2018). Taken together, these results (Haider & Taipaleenmaki, 2018; Liu et al., 2018) indicate that inhibition of miR-218-5p can attenuate the activity of the “vicious cycle” during bone metastasis progression, suggesting that it could be an attractive therapeutic target.

Tumour-derived ncRNAs impair OB and OC function

Several studies have shown that lncRNAs and miRNAs expressed and/or released by tumour cells can be directly involved in the formation of osteolytic or osteoblastic lesions (Table 2).

The lncRNA MAYA mediates a crosstalk between the ROR1-HER3 and Hippo/Yes-associated protein (YAP) pathways in breast cancer cells to promote lytic bone metastases (Li et al., 2017). Mechanistically, neuregulin induces the heterodimerization of receptor tyrosine kinase-like orphan receptor (ROR)-1 and receptor tyrosine kinase erbB-3 (HER3), which leads to the phosphorylation of HER3 in breast cancer cells. In turn, HER3 phosphorylation recruits a MAYA-containing complex that methylates the serine/threonine kinase Hippo/MST1 and activates YAP. This results in the translocation of YAP to the nucleus and the increased expression of connective tissue growth factor (CTGF) (Li et al., 2017). CTGF is a factor that promotes OC differentiation and activity. In vivo, the treatment of animals with an antibody against CTGF prevents the formation of osteolytic metastases caused by human MDA-MB-231 breast cancer cells (Shimo et al., 2006).

Increased MALAT1 expression in NSCLC is associated with a poor clinical outcome, including bone metastases (Liu et al., 2016). MALAT1 is overexpressed in several other cancers, where it promotes tumour cell invasion and proliferation (Liu et al., 2016). It has been reported that MALAT1 expression is increased in PC3 prostate cancer cells if co-cultured with primary mouse OBs (Sebastian et al., 2015). Moreover, MALAT1 expression in PC3 cells is further enhanced upon co-culture with mouse OBs lacking Sost (Sebastian et al., 2015). However, the underlying mechanisms by which SOST regulates MALAT1 expression in tumour cells are unknown.

RUNX2-AS1 is released by myeloma cells and is transferred to MSCs. In MSCs, RUNX2-AS1 represses RUNX2 expression through the formation of a duplex with RUNX2 pre-

mRNA, which then blocks OB differentiation (Li et al., 2018a). MiR-124 is a tumour suppressor whose expression is down-regulated in several cancer types (Cai et al., 2018). Its expression is also substantially reduced in breast cancer bone metastases (Cai et al., 2018). Experimentally, restoration of miR-124 expression in bone-tropic breast cancer cells reduces the progression of bone metastases in animals (Cai et al., 2018). Mechanistically, miR-124 represses the expression of IL-11 by breast cancer cells, which is a tumour-derived factor promoting bone resorption (Cai et al., 2018; Croucher et al., 2016). Of note, IL-11 expression is inversely correlated with the expression of miR-124, and high expression of IL-11 in bone metastases is associated with an overall shorter survival of patients (Cai et al., 2018).

Tumour-derived exosomal miRNAs can also affecting the function of bone cells. For example, exosomal miR-141 and miR-940 produced by prostate cancer cells are taken up by OB precursors and promote OB differentiation and proliferation, thereby facilitating the formation of sclerotic bone metastases (Hashimoto et al., 2018; Ye et al., 2017). Mechanistically, miR-141 activates p38-MAPKinases and increases OPG expression in OBs while miR-940 promotes osteogenic differentiation of MSCs via targeting ARHGAP1 (Rho GTPase Activating Protein 1) and FAM134A (Family with Sequence Similarity 134 Member A) (Hashimoto et al., 2018; Ye et al., 2017). Conversely, miR-218 secreted from breast cancer cells directly inhibits type I collagen deposition by OBs (Liu et al., 2018). Similarly, lung adenocarcinoma cell-derived exosomal miR-21 facilitates OC differentiation in vitro (Hu et al., 2017). MiR-21 exerts its promoting effect on OC differentiation via targeting programme cell death 4 (PDCD4), a repressor of c-Fos, which is an important transcription factor for OC differentiation (Hu et al., 2017).

Circulating ncRNAs and their value as biomarkers

NcRNAs that are released into body fluids (e.g., serum, plasma, urine, saliva) are either encapsulated within extracellular vesicles or are associated with high-density lipoproteins or AGO proteins, which provides protection from nuclease-mediated degradation (Anfossi et al., 2018; Li et al., 2018b). Thus, circulating ncRNAs are rather stable, and a growing body of evidence suggests that they might be useful as biomarkers in cancer (Anfossi et al., 2018; Kang, 2016; Mishra, 2014; Wan-Ibrahim et al., 2016). However, the prognostic value of ncRNAs in predicting the risk of bone metastases in patients with cancer clearly needs to be addressed.

Ell and colleagues identified a series of 4 miRNAs (miR-16, miR-211, miR-378, and let-7a) that were up-regulated during OC differentiation and induced by conditioned medium from breast cancer cells (Ell et al., 2013). The authors thought to investigate this group of miRNAs as potential biomarkers associated with lytic bone metastases in patients. They found that the expression of miR-16 and miR-378 was consistently increased in the serum of breast cancer patients with bone metastases (n = 38), compared to healthy controls (n = 21) (Ell et al., 2013). The amount of miR-10b in the serum was also significantly increased in breast cancer patients with bone metastases (n = 22), compared to breast cancer patients without bone metastases (n = 100) or healthy control subjects (n = 59) (Zhao et al., 2012). In prostate cancer, the abundance of miR-218 was greatly reduced in the serum of patients with bone metastases (n = 38), compared to patients with no metastases (n = 115) (Peng et al., 2019). Conversely, the abundance of miRNAs (miR-409-3p/5p, miR-154*, and miR-379) in the delta-like 1 homolog-deiodinase, iodothyronine 3 (DLK1-DIO3) cluster was increased in the serum of patients with metastatic prostate cancer, and an elevated expression of miR-154 and miR-379 was observed in bone metastatic prostate cancer cells in situ. This suggests that

miRNAs of the DLK1-DIO3 cluster could serve as biomarkers in patients with advanced prostate cancer (Gururajan et al., 2014).

Of note, the identification of circulating bone-derived miRNAs to predict specific stages of the disease has been suggested for bone-related pathologies other than cancer. For example, Seeliger and colleagues identified five miRNAs (miR-21-5p, miR-23a-3p, miR-24-3p, miR-100-5p, and miR-125b-5p) whose expression levels are increased in both the serum and bone tissue of osteoporotic patients with osteoporotic fractures (Seeliger et al., 2014). A correlation with bone mineral density in osteoporotic patients was also observed for miR-21-5p, miR-24-3p, miR-93-5p, miR-100-5p, and miR-125b-5p (Kelch et al., 2017). Additionally, intracellular studies revealed a simultaneous up-regulation of miR-21-5p, miR-93-5p, miR-100-5p and miR125b-5p in OBs and in OCs, whereas miR-148a-3p up-regulation was specific for osteoporotic OCs (Kelch et al., 2017). Overall, these findings (Seeliger et al., 2014; Kelch et al., 2017; Cheng et al., 2019) suggest these miRNAs may be useful biomarkers for early diagnosis of osteoporosis. None of these miRNAs associated with osteoporosis did however overlap with those associated with breast cancer-induced bone destruction (Ell et al., 2013).

Not only miRNAs, but also lncRNAs and circRNAs could potentially serve as biomarkers to predict the risk of bone metastases in patients with advanced cancer. For example, MALAT1 is measurable in the cellular fraction of peripheral human blood, and it has been proposed to be useful as a biomarker for the diagnosis of NSCLC (Weber et al., 2013). Given that MALAT1 is highly expressed in tissues from NSCLC patients with bone metastases compared to tissues from patients without bone metastases (Liu et al., 2016), it is likely that circulating MALAT1 could serve as a biomarker to predict the onset of skeletal-related events in NSCLC.

Interestingly, circRNAs are more stable than miRNAs and lncRNAs (Suzuki & Tsukahara, 2014), and they are enriched in exosomes compared to cells from where they are secreted (Li et al., 2015), suggesting they might be used as biomarkers for human cancers (Kristensen et al., 2018; Meng et al., 2017; Wang et al., 2016).

Concluding remarks

In this review, we describe the properties of ncRNAs, and we show that the dysregulation of ncRNAs in tumour cells and/or bone cells as well as the shuttling of ncRNAs between tumour cells and bone cells lead to the construction of a cancer niche in bone, which is essential for tumour progression (**Figure 2**). Interestingly, the identification of specific ncRNAs involved in bone metastasis formation and progression raises the possibility of developing novel biomarkers, which could provide important insight to predict the risk of relapse in patients with cancer. Likewise, the regulatory activity of ncRNAs in bone homeostasis and their dysregulation in bone metastasis encourage the development of ncRNA-based therapeutics.

Nomenclature of Targets and Ligands:

Key protein targets and ligands in this article are hyperlinked to corresponding entries in <http://www.guidetopharmacology.org>, the common portal for data from the IUPHAR/BPS Guide to PHARMACOLOGY (Harding et al., 2018), and are permanently archived in the Concise Guide to PHARMACOLOGY 2017/18 (Alexander et al., 2017).

Acknowledgements:

P. Clézardin gratefully acknowledge the support of INSERM, “Appel à Projets LIA/LEA 2016” (grant n°: ASC17018CSA), the University of Lyon, the University of Sheffield, and

Weston Park Cancer Charity (grant n°: CA163). H. Taipaleenmäki and E. Hesse acknowledge funding from the Deutsche Forschungsgemeinschaft (TA1154/1-1 and TA1154/2-1 to H.Taipaleenmäki, HE 5208/2-3 to E. Hesse).

Conflict of interest:

The authors declare no conflict of interest.

Declaration of transparency and scientific rigour

This Declaration acknowledges that this paper adheres to the principles for transparent reporting and scientific rigour of preclinical research as stated in the BJP guidelines for Design & Analysis, and as recommended by funding agencies, publishers and other organisations engaged with supporting research.

References

1. Akhade VS, Pal D, & Kanduri C (2017). Long Noncoding RNA: Genome Organization and Mechanism of Action. *Adv Exp Med Biol* 1008: 47-74.
2. Almeida M, Laurent MR, Dubois V, Claessens F, O'Brien CA, Bouillon R, et al. (2017). Estrogens and Androgens in Skeletal Physiology and Pathophysiology. *Physiol Rev* 97: 135-187.
3. Amandio AR, Necsulea A, Joye E, Mascrez B, & Duboule D (2016). Hotair Is Dispensable for Mouse Development. *PLoS Genet* 12: e1006232.
4. Anderson DM, Anderson KM, Chang CL, Makarewich CA, Nelson BR, McAnally JR, et al. (2015). A micropeptide encoded by a putative long noncoding RNA regulates muscle performance. *Cell* 160: 595-606.
5. Anfossi S, Babayan A, Pantel K, & Calin GA (2018). Clinical utility of circulating non-coding RNAs - an update. *Nat Rev Clin Oncol* 15: 541-563.
6. Aquino-Martinez R, Farr JN, Weivoda MM, Negley BA, Onken JL, Thicke BS, et al. (2018). miR-219a-5p Regulates Rorbeta During Osteoblast Differentiation and in Age-related Bone Loss. *J Bone Miner Res*.
7. Bachmayr-Heyda A, Reiner AT, Auer K, Sukhbaatar N, Aust S, Bachleitner-Hofmann T, et al. (2015). Correlation of circular RNA abundance with proliferation--exemplified with colorectal and ovarian cancer, idiopathic lung fibrosis, and normal human tissues. *Sci Rep* 5: 8057.
8. Baier SR, & Wan Y (2016). MicroRNA Exert Macro Effects on Cancer Bone Metastasis. *Curr Osteoporos Rep* 14: 163-169.
9. Beg MS, Brenner AJ, Sachdev J, Borad M, Kang YK, Stoudemire J, et al. (2017). Phase I study of MRX34, a liposomal miR-34a mimic, administered twice weekly in patients with advanced solid tumors. *Invest New Drugs* 35: 180-188.
10. Bellavia D, De Luca A, Carina V, Costa V, Raimondi L, Salamanna F, et al. (2019). Deregulated miRNAs in bone health: Epigenetic roles in osteoporosis. *Bone* 122: 52-75.
11. Bhartiya D, & Scaria V (2016). Genomic variations in non-coding RNAs: Structure, function and regulation. *Genomics* 107: 59-68.
12. Blin-Wakkach C, Lezot F, Ghoul-Mazgar S, Hotton D, Monteiro S, Teillaud C, et al. (2001). Endogenous Msx1 antisense transcript: in vivo and in vitro evidences, structure, and potential involvement in skeleton development in mammals. *Proc Natl Acad Sci U S A* 98: 7336-7341.
13. Bliss SA, Sinha G, Sandiford OA, Williams LM, Engelberth DJ, Guiro K, et al. (2016). Mesenchymal Stem Cell-Derived Exosomes Stimulate Cycling Quiescence and Early Breast Cancer Dormancy in Bone Marrow. *Cancer Res* 76: 5832-5844.

14. Browne G, Taipaleenmaki H, Stein GS, Stein JL, & Lian JB (2014). MicroRNAs in the control of metastatic bone disease. *Trends Endocrinol Metab* 25: 320-327.
15. Cai WL, Huang WD, Li B, Chen TR, Li ZX, Zhao CL, et al. (2018). microRNA-124 inhibits bone metastasis of breast cancer by repressing Interleukin-11. *Mol Cancer* 17: 9.
16. Chaffer CL, San Juan BP, Lim E, & Weinberg RA (2016). EMT, cell plasticity and metastasis. *Cancer Metastasis Rev* 35: 645-654.
17. Cheung DG, Buzzetti M, & Di Leva G (2017). miRNAs in bone metastasis. *Expert Rev Endocrinol Metab* 12: 451-461.
18. Choi SW, Kim HW, & Nam JW (2018). The small peptide world in long noncoding RNAs. *Brief Bioinform*.
19. Colden M, Dar AA, Saini S, Dahiya PV, Shahryari V, Yamamura S, et al. (2017). MicroRNA-466 inhibits tumor growth and bone metastasis in prostate cancer by direct regulation of osteogenic transcription factor RUNX2. *Cell Death Dis* 8: e2572.
20. Conn SJ, Pillman KA, Toubia J, Conn VM, Salmanidis M, Phillips CA, et al. (2015). The RNA binding protein quaking regulates formation of circRNAs. *Cell* 160: 1125-1134.
21. Croset M, Kan C, & Clezardin P (2015). Tumour-derived miRNAs and bone metastasis. *Bonekey Rep* 4: 688.
22. Croset M, Pantano F, Kan CWS, Bonnelye E, Descotes F, Alix-Panabieres C, et al. (2018). miRNA-30 Family Members Inhibit Breast Cancer Invasion, Osteomimicry, and Bone Destruction by Directly Targeting Multiple Bone Metastasis-Associated Genes. *Cancer Res* 78: 5259-5273.
23. Croucher PI, McDonald MM, & Martin TJ (2016). Bone metastasis: the importance of the neighbourhood. *Nat Rev Cancer* 16: 373-386.
24. Dallas SL, Prideaux M, & Bonewald LF (2013). The osteocyte: an endocrine cell ... and more. *Endocr Rev* 34: 658-690.
25. Davis HM, Pacheco-Costa R, Atkinson EG, Brun LR, Gortazar AR, Harris J, et al. (2017). Disruption of the Cx43/miR21 pathway leads to osteocyte apoptosis and increased osteoclastogenesis with aging. *Aging Cell* 16: 551-563.
26. Dou C, Cao Z, Yang B, Ding N, Hou T, Luo F, et al. (2016). Changing expression profiles of lncRNAs, mRNAs, circRNAs and miRNAs during osteoclastogenesis. *Sci Rep* 6: 21499.
27. Dougall WC (2012). Molecular pathways: osteoclast-dependent and osteoclast-independent roles of the RANKL/RANK/OPG pathway in tumorigenesis and metastasis. *Clin Cancer Res* 18: 326-335.

28. Ell B, & Kang Y (2014). MicroRNAs as regulators of bone homeostasis and bone metastasis. *Bonekey Rep* 3: 549.
29. Ell B, Mercatali L, Ibrahim T, Campbell N, Schwarzenbach H, Pantel K, et al. (2013). Tumor-induced osteoclast miRNA changes as regulators and biomarkers of osteolytic bone metastasis. *Cancer Cell* 24: 542-556.
30. Feng L, Shi L, Lu YF, Wang B, Tang T, Fu WM, et al. (2018). Linc-ROR Promotes Osteogenic Differentiation of Mesenchymal Stem Cells by Functioning as a Competing Endogenous RNA for miR-138 and miR-145. *Mol Ther Nucleic Acids* 11: 345-353.
31. Flynn RA, & Chang HY (2014). Long noncoding RNAs in cell-fate programming and reprogramming. *Cell Stem Cell* 14: 752-761.
32. Gabory A, Jammes H, & Dandolo L (2010). The H19 locus: role of an imprinted non-coding RNA in growth and development. *Bioessays* 32: 473-480.
33. Gaur T, Hussain S, Mudhasani R, Parulkar I, Colby JL, Frederick D, et al. (2010). Dicer inactivation in osteoprogenitor cells compromises fetal survival and bone formation, while excision in differentiated osteoblasts increases bone mass in the adult mouse. *Dev Biol* 340: 10-21.
34. Godfrey TC, Wildman BJ, Beloti MM, Kemper AG, Ferraz EP, Roy B, et al. (2018). The microRNA-23a cluster regulates the developmental HoxA cluster function during osteoblast differentiation. *J Biol Chem* 293: 17646-17660.
35. Gong C, & Maquat LE (2011). lncRNAs transactivate STAU1-mediated mRNA decay by duplexing with 3' UTRs via Alu elements. *Nature* 470: 284-288.
36. Gu X, Li M, Jin Y, Liu D, & Wei F (2017). Identification and integrated analysis of differentially expressed lncRNAs and circRNAs reveal the potential ceRNA networks during PDLSC osteogenic differentiation. *BMC Genet* 18: 100.
37. Guise TA, Yin JJ, Taylor SD, Kumagai Y, Dallas M, Boyce BF, et al. (1996). Evidence for a causal role of parathyroid hormone-related protein in the pathogenesis of human breast cancer-mediated osteolysis. *J Clin Invest* 98: 1544-1549.
38. Guo K, Zhang D, Wu H, Zhu Q, Yang C, & Zhu J (2018a). MiRNA-199a-5p positively regulated RANKL-induced osteoclast differentiation by target Mafk protein. *J Cell Biochem*.
39. Guo L, Chen K, Yuan J, Huang P, Xu X, Li C, et al. (2018b). Estrogen inhibits osteoclasts formation and bone resorption via microRNA-27a targeting PPARgamma and APC. *J Cell Physiol* 234: 581-594.
40. Gururajan M, Josson S, Chu GC, Lu CL, Lu YT, Haga CL, et al. (2014). miR-154* and miR-379 in the DLK1-DIO3 microRNA mega-cluster regulate epithelial to

mesenchymal transition and bone metastasis of prostate cancer. *Clin Cancer Res* 20: 6559-6569.

41. Gutschner T, & Diederichs S (2012). The hallmarks of cancer: a long non-coding RNA point of view. *RNA Biol* 9: 703-719.
42. Ha M, & Kim VN (2014). Regulation of microRNA biogenesis. *Nat Rev Mol Cell Biol* 15: 509-524.
43. Haider MT, & Taipaleenmaki H (2018). Targeting the Metastatic Bone Microenvironment by MicroRNAs. *Front Endocrinol (Lausanne)* 9: 202.
44. Han J, Lee Y, Yeom KH, Kim YK, Jin H, & Kim VN (2004). The Drosha-DGCR8 complex in primary microRNA processing. *Genes Dev* 18: 3016-3027.
45. Han Y, Zhang K, Hong Y, Wang J, Liu Q, Zhang Z, et al. (2018). miR-342-3p promotes osteogenic differentiation via targeting ATF3. *FEBS Lett*.
46. Harding SD, Sharman JL, Faccenda E, Southan C, Pawson AJ, Ireland S et al. (2018). The IUPHAR/BPS Guide to PHARMACOLOGY in 2018: updates and expansion to encompass the new guide to IMMUNOPHARMACOLOGY. *Nucl Acids Res* 46: D1091-D1106.
47. Hashimoto K, Ochi H, Sunamura S, Kosaka N, Mabuchi Y, Fukuda T, et al. (2018). Cancer-secreted hsa-miR-940 induces an osteoblastic phenotype in the bone metastatic microenvironment via targeting ARHGAP1 and FAM134A. *Proc Natl Acad Sci U S A* 115: 2204-2209.
48. Hassan MQ, Gordon JA, Beloti MM, Croce CM, van Wijnen AJ, Stein JL, et al. (2010). A network connecting Runx2, SATB2, and the miR-23a~27a~24-2 cluster regulates the osteoblast differentiation program. *Proc Natl Acad Sci U S A* 107: 19879-19884.
49. Hassan MQ, Maeda Y, Taipaleenmaki H, Zhang W, Jafferji M, Gordon JA, et al. (2012). miR-218 directs a Wnt signaling circuit to promote differentiation of osteoblasts and osteomimicry of metastatic cancer cells. *J Biol Chem* 287: 42084-42092.
50. Hesse E, & Taipaleenmaki H (2019). MicroRNAs in Bone Metastasis. *Curr Osteoporos Rep*.
51. Hsu M-T, & Coca-Prados M (1979). Electron microscopic evidence for the circular form of RNA in the cytoplasm of eukaryotic cells. *Nature* 280: 339-340.
52. Hu CH, Sui BD, Du FY, Shuai Y, Zheng CX, Zhao P, et al. (2017). miR-21 deficiency inhibits osteoclast function and prevents bone loss in mice. *Sci Rep* 7: 43191.

53. Huang JZ, Chen M, Chen, Gao XC, Zhu S, Huang H, et al. (2017). A Peptide Encoded by a Putative lncRNA HOXB-AS3 Suppresses Colon Cancer Growth. *Mol Cell* 68: 171-184 e176.
54. Huang Y, Zheng Y, Jia L, & Li W (2015). Long Noncoding RNA H19 Promotes Osteoblast Differentiation Via TGF-beta1/Smad3/HDAC Signaling Pathway by Deriving miR-675. *Stem Cells* 33: 3481-3492.
55. Huynh NP, Anderson BA, Guilak F, & McAlinden A (2017). Emerging roles for long noncoding RNAs in skeletal biology and disease. *Connect Tissue Res* 58: 116-141.
56. Inoue K, Deng Z, Chen Y, Giannopoulou E, Xu R, Gong S, et al. (2018). Bone protection by inhibition of microRNA-182. *Nat Commun* 9: 4108.
57. Jдалиha M, Zong X, Malakar P, Ray T, Singh DK, Freier SM, et al. (2016). Functional and prognostic significance of long non-coding RNA MALAT1 as a metastasis driver in ER negative lymph node negative breast cancer. *Oncotarget* 7: 40418-40436.
58. Ji P, Diederichs S, Wang W, Boing S, Metzger R, Schneider PM, et al. (2003). MALAT-1, a novel noncoding RNA, and thymosin beta4 predict metastasis and survival in early-stage non-small cell lung cancer. *Oncogene* 22: 8031-8041.
59. Ji X, Chen X, & Yu X (2016). MicroRNAs in Osteoclastogenesis and Function: Potential Therapeutic Targets for Osteoporosis. *Int J Mol Sci* 17: 349.
60. Jing Q, Huang S, Guth S, Zarubin T, Motoyama A, Chen J, et al. (2005). Involvement of microRNA in AU-rich element-mediated mRNA instability. *Cell* 120: 623-634.
61. Jo MH, Song JJ, & Hohng S (2015). Single-molecule fluorescence measurements reveal the reaction mechanisms of the core RISC, composed of human Argonaute 2 and a guide RNA. *BMB Rep* 48: 643-644.
62. Kang Y (2016). Dissecting Tumor-Stromal Interactions in Breast Cancer Bone Metastasis. *Endocrinol Metab (Seoul)* 31: 206-212.
63. Karvande A, Kushwaha P, Ahmad N, Adhikary S, Kothari P, Tripathi AK, et al. (2018). Glucose dependent miR-451a expression contributes to parathyroid hormone mediated osteoblast differentiation. *Bone* 117: 98-115.
64. Kim M, Yu Y, Moon JH, Koh I, & Lee JH (2018). Differential Expression Profiling of Long Noncoding RNA and mRNA during Osteoblast Differentiation in Mouse. *Int J Genomics* 2018: 7691794.
65. Kobayashi T, Lu J, Cobb BS, Rodda SJ, McMahon AP, Schipani E, et al. (2008). Dicer-dependent pathways regulate chondrocyte proliferation and differentiation. *Proc Natl Acad Sci U S A* 105: 1949-1954.
66. Kristensen LS, Hansen TB, Venø MT, & Kjems J (2018). Circular RNAs in cancer: opportunities and challenges in the field. *Oncogene* 37: 555-565.

67. Krol J, Loedige I, & Filipowicz W (2010). The widespread regulation of microRNA biogenesis, function and decay. *Nat Rev Genet* 11: 597-610.
68. Krzeszinski JY, Wei W, Huynh H, Jin Z, Wang X, Chang TC, et al. (2014). miR-34a blocks osteoporosis and bone metastasis by inhibiting osteoclastogenesis and Tgfb2. *Nature* 512: 431-435.
69. Lacey DL, Timms E, Tan HL, Kelley MJ, Dunstan CR, Burgess T, et al. (1998). Osteoprotegerin ligand is a cytokine that regulates osteoclast differentiation and activation. *Cell* 93: 165-176.
70. Lau KW, & Sheng MH (2018). A novel miR17/protein tyrosine phosphatase-oc/EphA4 regulatory axis of osteoclast activity. *Arch Biochem Biophys* 650: 30-38.
71. Li B, Xu H, Han H, Song S, Zhang X, Ouyang L, et al. (2018a). Exosome-mediated transfer of lncRUNX2-AS1 from multiple myeloma cells to MSCs contributes to osteogenesis. *Oncogene* 37: 5508-5519.
72. Li C, Wang S, Xing Z, Lin A, Liang K, Song J, et al. (2017). A ROR1-HER3-lncRNA signalling axis modulates the Hippo-YAP pathway to regulate bone metastasis. *Nat Cell Biol* 19: 106-119.
73. Li D, Liu J, Guo B, Liang C, Dang L, Lu C, et al. (2016). Osteoclast-derived exosomal miR-214-3p inhibits osteoblastic bone formation. *Nat Commun* 7: 10872.
74. Li S, Li Y, Chen B, Zhao J, Yu S, Tang Y, et al. (2018b). exoRBase: a database of circRNA, lncRNA and mRNA in human blood exosomes. *Nucleic Acids Res* 46: D106-D112.
75. Li Y, Zheng Q, Bao C, Li S, Guo W, Zhao J, et al. (2015). Circular RNA is enriched and stable in exosomes: a promising biomarker for cancer diagnosis. *Cell Res* 25: 981-984.
76. Li Z, Hassan MQ, Jafferji M, Aqeilan RI, Garzon R, Croce CM, et al. (2009). Biological functions of miR-29b contribute to positive regulation of osteoblast differentiation. *J Biol Chem* 284: 15676-15684.
77. Liang WC, Fu WM, Wang YB, Sun YX, Xu LL, Wong CW, et al. (2016). H19 activates Wnt signaling and promotes osteoblast differentiation by functioning as a competing endogenous RNA. *Sci Rep* 6: 20121.
78. Liao J, Yu X, Hu X, Fan J, Wang J, Zhang Z, et al. (2017). lncRNA H19 mediates BMP9-induced osteogenic differentiation of mesenchymal stem cells (MSCs) through Notch signaling. *Oncotarget* 8: 53581-53601.
79. Lin JC, Liu ZG, Yu B, & Zhang XR (2018). MicroRNA-874 targeting SUFU involves in osteoblast proliferation and differentiation in osteoporosis rats through the Hedgehog signaling pathway. *Biochem Biophys Res Commun* 506: 194-203.

80. Liu J, Li D, Dang L, Liang C, Guo B, Lu C, et al. (2017). Osteoclastic miR-214 targets TRAF3 to contribute to osteolytic bone metastasis of breast cancer. *Sci Rep* 7: 40487.
81. Liu M, Sun W, Liu Y, & Dong X (2016). The role of lncRNA MALAT1 in bone metastasis in patients with non-small cell lung cancer. *Oncol Rep* 36: 1679-1685.
82. Liu X, Cao M, Palomares M, Wu X, Li A, Yan W, et al. (2018). Metastatic breast cancer cells overexpress and secrete miR-218 to regulate type I collagen deposition by osteoblasts. *Breast Cancer Res* 20: 127.
83. Long F (2011). Building strong bones: molecular regulation of the osteoblast lineage. *Nat Rev Mol Cell Biol* 13: 27-38.
84. Ma L, Teruya-Feldstein J, & Weinberg RA (2007). Tumour invasion and metastasis initiated by microRNA-10b in breast cancer. *Nature* 449: 682-688.
85. Matsumoto A, Pasut A, Matsumoto M, Yamashita R, Fung J, Monteleone E, et al. (2017). mTORC1 and muscle regeneration are regulated by the LINC00961-encoded SPAR polypeptide. *Nature* 541: 228-232.
86. Meng S, Zhou H, Feng Z, Xu Z, Tang Y, Li P, et al. (2017). CircRNA: functions and properties of a novel potential biomarker for cancer. *Mol Cancer* 16: 94.
87. Mishra PJ (2014). MicroRNAs as promising biomarkers in cancer diagnostics. *Biomark Res* 2: 19.
88. Nakanishi K (2016). Anatomy of RISC: how do small RNAs and chaperones activate Argonaute proteins? *Wiley Interdiscip Rev RNA* 7: 637-660.
89. Nardocci G, Carrasco ME, Acevedo E, Hodar C, Meneses C, & Montecino M (2018). Identification of a novel long noncoding RNA that promotes osteoblast differentiation. *J Cell Biochem* 119: 7657-7666.
90. Niazi F, & Valadkhan S (2012). Computational analysis of functional long noncoding RNAs reveals lack of peptide-coding capacity and parallels with 3' UTRs. *RNA* 18: 825-843.
91. Ono M, Kosaka N, Tominaga N, Yoshioka Y, Takeshita F, Takahashi RU, et al. (2014). Exosomes from bone marrow mesenchymal stem cells contain a microRNA that promotes dormancy in metastatic breast cancer cells. *Sci Signal* 7: ra63.
92. Pan BL, Tong ZW, Li SD, Wu L, Liao JL, Yang YX, et al. (2018). Decreased microRNA-182-5p helps alendronate promote osteoblast proliferation and differentiation in osteoporosis via the Rap1/MAPK pathway. *Biosci Rep*.
93. Patop IL, & Kadener S (2018). circRNAs in Cancer. *Curr Opin Genet Dev* 48: 121-127.

94. Peng F, Li TT, Wang KL, Xiao GQ, Wang JH, Zhao HD, et al. (2017). H19/let-7/LIN28 reciprocal negative regulatory circuit promotes breast cancer stem cell maintenance. *Cell Death Dis* 8: e2569.
95. Peng P, Chen T, Wang Q, Zhang Y, Zheng F, Huang S, et al. (2019). Decreased miR-218-5p Levels as a Serum Biomarker in Bone Metastasis of Prostate Cancer. *Oncol Res Treat* 42: 165-180.
96. Peng W, Deng W, Zhang J, Pei G, Rong Q, & Zhu S (2018). Long noncoding RNA ANCR suppresses bone formation of periodontal ligament stem cells via sponging miRNA-758. *Biochem Biophys Res Commun* 503: 815-821.
97. Pospiech N, Cibis H, Dietrich L, Muller F, Bange T, & Hennig S (2018). Identification of novel PANDAR protein interaction partners involved in splicing regulation. *Sci Rep* 8: 2798.
98. Qian DY, Yan GB, Bai B, Chen Y, Zhang SJ, Yao YC, et al. (2017). Differential circRNA expression profiles during the BMP2-induced osteogenic differentiation of MC3T3-E1 cells. *Biomed Pharmacother* 90: 492-499.
99. Qu B, Xia X, Yan M, Gong K, Deng S, Huang G, et al. (2015). miR-218 is involved in the negative regulation of osteoclastogenesis and bone resorption by partial suppression of p38MAPK-c-Fos-NFATc1 signaling: Potential role for osteopenic diseases. *Exp Cell Res* 338: 89-96.
100. Quan G, & Li J (2018). Circular RNAs: biogenesis, expression and their potential roles in reproduction. *J Ovarian Res* 11: 9.
101. Quinn JJ, & Chang HY (2016). Unique features of long non-coding RNA biogenesis and function. *Nat Rev Genet* 17: 47-62.
102. Raaijmakers MH, Mukherjee S, Guo S, Zhang S, Kobayashi T, Schoonmaker JA, et al. (2010). Bone progenitor dysfunction induces myelodysplasia and secondary leukaemia. *Nature* 464: 852-857.
103. Raveh E, Matouk IJ, Gilon M, & Hochberg A (2015). The H19 Long non-coding RNA in cancer initiation, progression and metastasis - a proposed unifying theory. *Mol Cancer* 14: 184.
104. Rinn JL, Kertesz M, Wang JK, Squazzo SL, Xu X, Bruggmann SA, et al. (2007). Functional demarcation of active and silent chromatin domains in human HOX loci by noncoding RNAs. *Cell* 129: 1311-1323.
105. Rossi M, Bucci G, Rizzotto D, Bordo D, Marzi MJ, Puppo M, et al. (2019). LncRNA EPR controls epithelial proliferation by coordinating Cdkn1a transcription and mRNA decay response to TGF- β . *Nature Communications* 10: 1969.
106. Sahay D, Leblanc R, Grunewald TG, Ambatipudi S, Ribeiro J, Clezardin P, et al. (2015). The LPA1/ZEB1/miR-21-activation pathway regulates metastasis in basal breast cancer. *Oncotarget* 6: 20604-20620.

107. Saini S, Majid S, Yamamura S, Tabatabai L, Suh SO, Shahryari V, et al. (2011). Regulatory Role of mir-203 in Prostate Cancer Progression and Metastasis. *Clin Cancer Res* 17: 5287-5298.
108. Salido-Guadarrama I, Romero-Cordoba S, Peralta-Zaragoza O, Hidalgo-Miranda A, & Rodriguez-Dorantes M (2014). MicroRNAs transported by exosomes in body fluids as mediators of intercellular communication in cancer. *Onco Targets Ther* 7: 1327-1338.
109. Schmid-Alliana A, Schmid-Antomarchi H, Al-Sahlanee R, Lagadec P, Scimeca JC, & Verron E (2018). Understanding the Progression of Bone Metastases to Identify Novel Therapeutic Targets. *Int J Mol Sci* 19.
110. Sebastian A, Hum NR, Hudson BD, & Loots GG (2015). Cancer-Osteoblast Interaction Reduces Sost Expression in Osteoblasts and Up-Regulates lncRNA MALAT1 in Prostate Cancer. *Microarrays (Basel)* 4: 503-519.
111. Seeliger C, Karpinski K, Haug AT, Vester H, Schmitt A, Bauer JS, et al. (2014). Five freely circulating miRNAs and bone tissue miRNAs are associated with osteoporotic fractures. *J Bone Miner Res* 29: 1718-1728.
112. Shimo T, Kubota S, Yoshioka N, Ibaragi S, Isowa S, Eguchi T, et al. (2006). Pathogenic role of connective tissue growth factor (CTGF/CCN2) in osteolytic metastasis of breast cancer. *J Bone Miner Res* 21: 1045-1059.
113. Siu MK, Abou-Kheir W, Yin JJ, Chang YS, Barrett B, Suau F, et al. (2018). Correction: Loss of EGFR signaling-regulated miR-203 promotes prostate cancer bone metastasis and tyrosine kinase inhibitors resistance. *Oncotarget* 9: 32403.
114. Sul OJ, Sung YB, Rajasekaran M, Ke K, Yu R, Back SH, et al. (2018). MicroRNA-155 induces autophagy in osteoclasts by targeting transforming growth factor beta-activated kinase 1-binding protein 2 upon lipopolysaccharide stimulation. *Bone* 116: 279-289.
115. Sun KT, Chen MY, Tu MG, Wang IK, Chang SS, & Li CY (2015). MicroRNA-20a regulates autophagy related protein-ATG16L1 in hypoxia-induced osteoclast differentiation. *Bone* 73: 145-153.
116. Suva LJ, Washam C, Nicholas RW, & Griffin RJ (2011). Bone metastasis: mechanisms and therapeutic opportunities. *Nat Rev Endocrinol* 7: 208-218.
117. Suzuki H, & Tsukahara T (2014). A view of pre-mRNA splicing from RNase R resistant RNAs. *Int J Mol Sci* 15: 9331-9342.
118. Taipaleenmaki H, Browne G, Akech J, Zustin J, van Wijnen AJ, Stein JL, et al. (2015). Targeting of Runx2 by miR-135 and miR-203 Impairs Progression of Breast Cancer and Metastatic Bone Disease. *Cancer Res* 75: 1433-1444.

119. Taipaleenmaki H, Farina NH, van Wijnen AJ, Stein JL, Hesse E, Stein GS, et al. (2016). Antagonizing miR-218-5p attenuates Wnt signaling and reduces metastatic bone disease of triple negative breast cancer cells. *Oncotarget* 7: 79032-79046.
120. Tavazoie SF, Alarcon C, Oskarsson T, Padua D, Wang Q, Bos PD, et al. (2008). Endogenous human microRNAs that suppress breast cancer metastasis. *Nature* 451: 147-152.
121. Tye CE, Boyd JR, Page NA, Falcone MM, Stein JL, Stein GS, et al. (2018). Regulation of osteogenesis by long noncoding RNAs: An epigenetic mechanism contributing to bone formation. *Connect Tissue Res* 59: 35-41.
122. Ulitsky I, & Bartel DP (2013). lincRNAs: genomics, evolution, and mechanisms. *Cell* 154: 26-46.
123. Valinezhad Orang A, Safaralizadeh R, & Kazemzadeh-Bavili M (2014). Mechanisms of miRNA-Mediated Gene Regulation from Common Downregulation to mRNA-Specific Upregulation. *Int J Genomics* 2014: 970607.
124. Van Poznak C, Somerfield MR, Barlow WE, Biermann JS, Bosserman LD, Clemons MJ, et al. (2017). Role of Bone-Modifying Agents in Metastatic Breast Cancer: An American Society of Clinical Oncology-Cancer Care Ontario Focused Guideline Update. *J Clin Oncol* 35: 3978-3986.
125. Wan-Ibrahim WI, Singh VA, Hashim OH, & Abdul-Rahman PS (2016). Biomarkers for Bone Tumors: Discovery from Genomics and Proteomics Studies and Their Challenges. *Mol Med* 21: 861-872.
126. Wang F, Nazarali AJ, & Ji S (2016). Circular RNAs as potential biomarkers for cancer diagnosis and therapy. *Am J Cancer Res* 6: 1167-1176.
127. Wang W, Yang L, Zhang D, Gao C, Wu J, Zhu Y, et al. (2018a). MicroRNA-218 Negatively Regulates Osteoclastogenic Differentiation by Repressing the Nuclear Factor-kappaB Signaling Pathway and Targeting Tumor Necrosis Factor Receptor 1. *Cell Physiol Biochem* 48: 339-347.
128. Wang X, Guo B, Li Q, Peng J, Yang Z, Wang A, et al. (2013). miR-214 targets ATF4 to inhibit bone formation. *Nat Med* 19: 93-100.
129. Wang Y, Wang K, Hu Z, Zhou H, Zhang L, Wang H, et al. (2018b). MicroRNA-139-3p regulates osteoblast differentiation and apoptosis by targeting ELK1 and interacting with long noncoding RNA ODSM. *Cell Death Dis* 9: 1107.
130. Weber DG, Johnen G, Casjens S, Bryk O, Pesch B, Jockel KH, et al. (2013). Evaluation of long noncoding RNA MALAT1 as a candidate blood-based biomarker for the diagnosis of non-small cell lung cancer. *BMC Res Notes* 6: 518.
131. Wei J, Li H, Wang S, Li T, Fan J, Liang X, et al. (2014). let-7 enhances osteogenesis and bone formation while repressing adipogenesis of human

stromal/mesenchymal stem cells by regulating HMGA2. *Stem Cells Dev* 23: 1452-1463.

132. Wei Y, Li D, Wang D, Qiu T, & Liu K (2017). Evaluation of microRNA-203 in bone metastasis of patients with non-small cell lung cancer through TGF-beta/SMAD2 expression. *Oncol Rep*.

133. Wilusz JE (2018). A 360 degrees view of circular RNAs: From biogenesis to functions. *Wiley Interdiscip Rev RNA* 9: e1478.

134. Wu J, Zhao J, Sun L, Pan Y, Wang H, & Zhang WB (2018). Long non-coding RNA H19 mediates mechanical tension-induced osteogenesis of bone marrow mesenchymal stem cells via FAK by sponging miR-138. *Bone* 108: 62-70.

135. Xie Y, Chen Y, Zhang L, Ge W, & Tang P (2017). The roles of bone-derived exosomes and exosomal microRNAs in regulating bone remodelling. *J Cell Mol Med* 21: 1033-1041.

136. Xie Z, Li J, Wang P, Li Y, Wu X, Wang S, et al. (2016). Differential Expression Profiles of Long Noncoding RNA and mRNA of Osteogenically Differentiated Mesenchymal Stem Cells in Ankylosing Spondylitis. *J Rheumatol* 43: 1523-1531.

137. Ye Y, Li SL, Ma YY, Diao YJ, Yang L, Su MQ, et al. (2017). Exosomal miR-141-3p regulates osteoblast activity to promote the osteoblastic metastasis of prostate cancer. *Oncotarget* 8: 94834-94849.

138. Yi J, Liu D, & Xiao J (2019). LncRNA MALAT1 sponges miR-30 to promote osteoblast differentiation of adipose-derived mesenchymal stem cells by promotion of Runx2 expression. *Cell Tissue Res* 376: 113-121.

139. Yin Y, Tang L, Chen J, & Lu X (2017). MiR-30a attenuates osteoclastogenesis via targeting DC-STAMP-c-Fos-NFATc1 signaling. *Am J Transl Res* 9: 5743-5753.

140. Yu FY, Xie CQ, Sun JT, Peng W, & Huang XW (2018). Overexpressed miR-145 inhibits osteoclastogenesis in RANKL-induced bone marrow-derived macrophages and ovariectomized mice by regulation of Smad3. *Life Sci* 202: 11-20.

141. Yu X, Zhang Y, Li T, Ma Z, Jia H, Chen Q, et al. (2017). Long non-coding RNA Linc-RAM enhances myogenic differentiation by interacting with MyoD. *Nat Commun* 8: 14016.

142. Zhang G, Sun H, Zhang Y, Zhao H, Fan W, Li J, et al. (2018a). Characterization of dysregulated lncRNA-mRNA network based on ceRNA hypothesis to reveal the occurrence and recurrence of myocardial infarction. *Cell Death Discov* 4: 35.

143. Zhang M, Jia L, & Zheng Y (2018b). circRNA Expression Profiles in Human Bone Marrow Stem Cells Undergoing Osteoblast Differentiation. *Stem Cell Rev*.

144. Zhang Y, Xie RL, Croce CM, Stein JL, Lian JB, van Wijnen AJ, et al. (2011). A program of microRNAs controls osteogenic lineage progression by targeting transcription factor Runx2. *Proc Natl Acad Sci U S A* 108: 9863-9868.
145. Zhao FL, Hu GD, Wang XF, Zhang XH, Zhang YK, & Yu ZS (2012). Serum overexpression of microRNA-10b in patients with bone metastatic primary breast cancer. *J Int Med Res* 40: 859-866.
146. Zheng X, Dai J, Zhang H, & Ge Z (2018). MicroRNA-221 promotes cell proliferation, migration, and differentiation by regulation of ZFPM2 in osteoblasts. *Braz J Med Biol Res* 51: e7574.
147. Zhou R, Wu Y, Wang W, Su W, Liu Y, Wang Y, et al. (2018a). Circular RNAs (circRNAs) in cancer. *Cancer Lett* 425: 134-142.
148. Zhou Z, Lu Y, Wang Y, Du L, Zhang Y, & Tao J (2018b). Let-7c regulates proliferation and osteo-differentiation of human adipose-derived mesenchymal stem cells under oxidative stress by targeting SCD-1. *Am J Physiol Cell Physiol*.
149. Zhu L, & Xu PC (2013). Downregulated LncRNA-ANCR promotes osteoblast differentiation by targeting EZH2 and regulating Runx2 expression. *Biochem Biophys Res Commun* 432: 612-617.
150. Zhuang W, Ge X, Yang S, Huang M, Zhuang W, Chen P, et al. (2015). Upregulation of lncRNA MEG3 Promotes Osteogenic Differentiation of Mesenchymal Stem Cells From Multiple Myeloma Patients By Targeting BMP4 Transcription. *Stem Cells* 33: 1985-1997.
151. Zuo C, Wang Z, Lu H, Dai Z, Liu X, & Cui L (2013). Expression profiling of lncRNAs in C3H10T1/2 mesenchymal stem cells undergoing early osteoblast differentiation. *Mol Med Rep* 8: 463-467.

Table 1- Noncoding RNAs (ncRNAs) involved in normal osteoblast (OB) and osteoclast (OC) functions.

| Function | ncRNA name | Target(s) | References |
|--|---------------|--|--|
| Long non-coding RNAs (lncRNAs) | | | |
| Inhibition of OB differentiation | ANCR (DANCR) | Runx2, miR-758 | (Zhu & Xu, 2013; Peng et al., 2018) |
| | lnc-RUNX2-AS1 | Runx2 | (Li et al., 2018a) |
| Promotion of OB differentiation | H19 | TGF- β 1/Smad3/HDAC signalling pathway, miR-449b, miR-107, miR-27b, miR-34a, miR-106b, miR-449a, miR-125a, miR-17, miR-141, miR-22, miR-138, let-7 | (Gabory et al., 2010; Raveh et al., 2015; Huang et al., 2015; Liao et al., 2017; Liang et al., 2016; Wu et al., 2018; Peng et al., 2017; Wei et al., 2014) |
| | HOTAIR | HoxD, Hoxc11, Hoxc12, Dlk1-Meg3, Igf2-H19, Hottip | (Rinn et al., 2007; Amandio et al., 2016; Huynh et al., 2017) |
| | Linc-ROR | miR-138, miR-145 | (Feng et al., 2018) |
| | lncRNA-1 | Runx2/p57, Sp7 | (Nardocci et al., 2018) |
| | MALAT1 | miR-30 | (Yi et al., 2019) |
| Promotion of dental and craniofacial development | MEG3 | Sox2 | (Zhuang et al., 2015) |
| | Msx1-AS | Msx1 | (Blin-Wakkach et al., 2001) |
| MicroRNAs (miRNAs) | | | |
| (1) MiRNAs and osteoblasts (OBs): | | | |
| Inhibition of OB differentiation | let-7c | Scd-1 | (Zhou et al., 2018) |
| | miR-100 | Bmpr2 | (Ell et al., 2014) |
| | miR-106 | Bmp2 | (Ell et al., 2014) |
| | miR-133a | Runx2 | (Zhang et al., 2011) |
| | miR-135a | Runx2 | (Zhang et al., 2011) |
| | miR-137 | Runx2 | (Zhang et al., 2011) |
| | miR-138 | Fak | (Ell et al., 2014) |
| | miR-139-3p | Elk1 | (Wang et al., 2018b) |
| | miR-141 | Dlx5 | (Ell et al., 2014) |
| miR-143 | Osterix | (Ell et al., 2014) | |

| | | |
|-----------------|------------------------------|--------------------------------|
| miR-155 | Socs1, Mitf | (Ell et al., 2014) |
| miR-17-5p | Bmp2 | (Ell et al., 2014) |
| miR-182-5p | Adcy6 | (Pan et al., 2018) |
| miR-204 | Runx2 | (Zhang et al., 2011) |
| miR-205 | Runx2 | (Zhang et al., 2011) |
| miR-200a | Dlx5 | (Ell et al., 2014) |
| miR-206 | Cx43 | (Ell et al., 2014) |
| miR-208 | Ets1 | (Ell et al., 2014) |
| miR-214 | Atf4, Osterix | (Ell et al., 2014) |
| miR-217 | Runx2 | (Zhang et al., 2011) |
| miR-23a | Fak, Runx2, Satb2 | (Ell et al., 2014) |
| miR-23a cluster | HoxA cluster function | (Godfrey et al., 2018) |
| miR-24-2 | Fak, Runx2, Satb2 | (Ell et al., 2014) |
| miR-27a | Fak, Runx2, Satb2 | (Ell et al., 2014) |
| miR-30 | Smad1, Runx2 | (Ell et al., 2014) |
| miR-30c | Runx2 | (Zhang et al., 2011) |
| miR-338 | Runx2 | (Zhang et al., 2011) |
| miR-34 | Satb2, Runx2, Notch pathway | (Ell et al., 2014) |
| miR-34c | Runx2 | (Zhang et al., 2011) |
| miR-542-3p | Bmp7 | (Ell et al., 2014) |
| miR-764-5p | Chip/Stub1 | (Ell et al., 2014) |
| miR-93 | Sp7 | (Ell et al., 2014) |
| miR-15b | Smurf1 | (Ell et al., 2014) |
| miR-17~92 | | (Ell et al., 2014) |
| miR-181a | Tgfb1 | (Ell et al., 2014) |
| miR-20a | PPAR γ , Bambi, Crim1 | (Ell et al., 2014) |
| miR-218 | Sost-1, Dkk-2, Sfrp2 | (Hassan et al., 2012) |
| miR-219a-5p | | (Aquino-Martinez et al., 2018) |
| miR-221 | Zzpm2 | (Zheng et al., 2018) |
| miR-29a | ON, Dkk1, Krm2, sFrp2 | (Ell et al., 2014) |
| miR-332 | Tob2 | (Ell et al., 2014) |
| miR-342-3p | Atf3 | (Han et al., 2018) |
| miR-335-5p | Dkk1 | (Ell et al., 2014) |
| miR-451a | Osr1 | (Karvande et al., 2018) |
| miR-874 | Sufu | (Lin et al., 2018) |

Promotion of OB differentiation

(2) MiRNAs and osteocytes:

| | | | |
|-----------------------------------|---------|---------------------------------------|----------------------|
| Promotion of osteocytes apoptosis | miR-21 | Pten | (Davis et al., 2017) |
| | miR-218 | <i>IκB kinase B</i> | (Davis et al., 2017) |

(3) MiRNAs and osteoclasts (OCs):

| | | | |
|--|-------------|----------------------------------|----------------------|
| Inhibition of OC differentiation | miR-124-3p | NFATc1, RhoA, Rac1 | (Ji et al., 2016) |
| | miR-125a-5p | Traf6 | (Ji et al., 2016) |
| | miR-145 | Sox9, Smad3 | (Yu et al., 2018) |
| | miR-146a-5p | Traf6, Stat1 | (Ji et al., 2016) |
| | miR-155 | Socs1, Mitf | (Ell et al., 2014) |
| | miR-17 | EphA4 | (Lau & Sheng, 2018) |
| | miR-218 | p38 MAPK, NFATc1 | (Qu et al., 2015) |
| | miR-218-5p | p38MAPK-c-Fos-NFATc1 | (Ji et al., 2016) |
| | miR-26a-5p | Ctgf | (Ji et al., 2016) |
| | miR-27a | Runx2 | (Guo et al., 2018b) |
| | miR-30a | | (Yin et al., 2017) |
| | miR-34a-5p | Tgif2 | (Ji et al., 2016) |
| | miR-503-5p | Rank | (Ji et al., 2016) |
| miR-7b-5p | DC-Stamp | (Ji et al., 2016) | |
| Promotion of OC differentiation | miR-148a-3p | MafB | (Ji et al., 2016) |
| | miR-183-5p | HO-1 | (Ji et al., 2016) |
| | miR-199a-5p | Mafb | (Guo et al., 2018a) |
| | miR-29 | Cdc42, Srgap2, Nfia, Cd93, Calcr | (Ji et al., 2016) |
| | miR-31-5p | RhoA | (Ji et al., 2016) |
| | miR-422a | Cbl, Cd226, Igfl, Pag1, Tob2 | (Ji et al., 2016) |
| | miR-9718 | Pias3 | (Ji et al., 2016) |
| Promotion/inhibition of OC differentiation | miR-21-5p | FasL, Pcdcd4 | (Ji et al., 2016) |
| | miR-223-3p | Nfia, Ikk | (Ji et al., 2016) |
| | miR-29b | Cdc42, Srgap2 | (Ell et al., 2014) |
| Promotion of OC activity | miR-133a-3p | Cxcl11, Cxcr3, Slc39A1 | (Ji et al., 2016) |
| | miR-182 | Pkr | (Inoue et al., 2018) |
| Promotion of OC autophagy | miR-155 | Mitf | (Sul et al., 2018) |
| | miR-20a | PPARg, Bambi, Crim1 | (Sun et al., 2015) |
| (4) MiRNAs and chondrocytes: | | | |
| Inhibition of chondrocyte differentiation | miR-145 | Sox9 | (Ell et al., 2014) |
| | miR-34a | | (Ell et al., 2014) |
| | miR-199a-3p | Smad1 | (Ell et al., 2014) |
| Promotion of chondrocyte activity | miR-140 | Hdac4, Dnpep | (Ell et al., 2014) |

| | | |
|--------------------------------------|------------|-----------------------|
| Promotion of chondrocyte development | miR-17~92 | (ElI et al., 2014) |
| CircularRNA (CircRNA) | | |
| Inhibition of OB differentiation | circIGSF11 | (Zhang et al., 2018b) |

Accepted Article

Table 2- Dysregulated non-coding RNAs (ncRNAs) involved in bone metastasis.

| NcRNA (expression level) | Function(s) | Target(s) | References |
|-------------------------------|---|---|---|
| Osteoclasts: | | | |
| miR-133a (up) | Promotes bone resorption in vivo | Mitf, Traf6, Calcr, Mmp14 | (Ell et al., 2013) |
| miR-141-3p (up) | Promotes bone resorption in vivo | Mitf, Traf6, Calcr, Mmp14 | (Ell et al., 2013) |
| miR-190 (up) | Promotes bone resorption in vivo | Mitf, Traf6, Calcr, Mmp14 | (Ell et al., 2013) |
| miR-21 (up) | Promotes bone resorption in vitro | Pdcd4 | (Hu et al., 2017) |
| miR-214-3p (up) | Promotes bone resorption in vivo | Traf3 | (Liu et al., 2017) |
| miR-219-5p (up) | Promotes bone resorption in vivo | Mitf, Traf6, Calcr, Mmp14 | (Ell et al., 2013) |
| miR-34a (up) | Inhibits bone resorption in vivo (MRX34 in clinical trial) | Tgif2 | (Krzyszinski et al., 2014) |
| Osteoblasts: | | | |
| miR-141 (up) | Promotes bone formation in vivo | p38-Mapk, Opg | (Ye et al., 2017) |
| miR-214-3p (up) | Inhibits bone formation in vivo | Atf4 | (Haider & Taipaleenmaki, 2018) |
| miR-218 (up) | Promotes bone disruption in vitro | Collagen-type I | (Liu et al., 2018) |
| Breast cancer cells: | | | |
| let-7g (up) | Inhibits bone metastasis in vivo | Hmga2 | (Ell & Kang, 2014) |
| MALAT1 (up) | Promotes bone metastasis in vitro | miR-30 sponge (?) | (Jadaliha et al., 2016) |
| MAYA (up) | Promotes bone metastasis in vivo | CTGF | (Li et al., 2017a) |
| miR-10b (up) | Promotes bone metastasis in vivo | HoxD10 | (Croset et al., 2015; Ell & Kang, 2014; Ma et al., 2007) |
| miR-124 (down) | Promotes bone metastasis in vivo | Il-11 | (Cai et al., 2018) |
| miR-126 (down) | Promotes bone metastasis in vivo | | (Tavazoie et al., 2008) |
| miR-135 (down) | Promotes bone metastasis in vivo | Runx2 | (Taipaleenmaki et al., 2015) |
| miR-17 (up) | Promotes bone metastasis in vivo | Tbr2 | (Croset et al., 2015; Ell & Kang, 2014) |
| miR-203 (down) | Promotes bone metastasis in vivo | Runx2 | (Taipaleenmaki et al., 2015) |
| miR-206 (down) | Promotes bone metastasis in vivo | | (Tavazoie et al., 2008) |
| miR-21 (up) | Promotes bone metastasis in vitro | Pten, Pdcd4, Spry2 | (Sahay et al., 2015) |
| miR-218 (up) | Promotes bone metastasis in vivo | WNT activity | (Liu et al., 2018; Hassan et al., 2012; Taipaleenmaki et al., 2015) |
| miR-30s (up) | Inhibits bone metastasis in vivo | Il-8, Il-11, Dkk-1 Runx2, Cdh11, Ctgf, Itga5, Itgb3 | (Croset et al., 2018) |
| miR-335 (down) | Promotes bone metastasis in vivo | Sox4, Tnc | (Tavazoie et al., 2008) |
| Prostate cancer cells: | | | |

| | | | |
|--|--|---|---|
| MALAT1 (up) | Promotes bone metastasis in vitro | | (Sebastian et al., 2015) |
| miR-143 (up) | Inhibits bone metastasis in vivo | Hef1 | (Croset et al., 2015; Ell & Kang, 2014) |
| miR-145 (up) | Inhibits bone metastasis in vitro | Hef1 | (Croset et al., 2015; Ell & Kang, 2014) |
| miR-146a (up) | Inhibits proliferation and adhesion in vitro | Rock | (Ell & Kang, 2014) |
| miR-16 (up) | Inhibits bone metastasis in vivo | | (Ell & Kang, 2014) |
| miR-203 (up) | Inhibits bone metastasis in vivo | Survivin/Birc5 EGFR signaling pathway, Runx2, Dlx5, Spp1, Bglap | (Siu et al., 2018; Saini et al., 2011) |
| miR-466 (up) | Inhibits bone metastasis in vivo | Runx2, Spp1, Bglap | (Colden et al., 2017; Saini et al., 2011) |
| Non-small cell lung cancer cells: | | | |
| MALAT1 (up) | Promotes bone metastasis in vivo | | (Liu et al., 2016) |
| miR-203 (down) | Promotes bone metastasis in vivo | Bax, Casp-3, p53, pSmad2, Tgf- β 1 | (Wei et al., 2017) |
| Mesenchymal stem cells: | | | |
| let-7 (up) | Promotion of cancer progression in vivo | Il-6 | (Ell & Kang, 2014) |
| miR-23b (up) | Promotes breast cancer dormancy in vitro | Marcks | (Ono et al., 2014) |
| miR-222/223 (up) | Promotes breast cancer dormancy in vivo | | (Bliss et al., 2016) |
| RUNX2-AS1 (up) | Inhibits osteoblast differentiation in vivo | Runx2 | (Li et al., 2018a) |




Figure 1- Schematic representation of ncRNA mechanisms of action.

NcRNAs regulate gene expression both at transcriptional and post-transcriptional levels. LncRNA (left box), miRNA (middle box), and circRNA (right box) act in a variety of ways promoting or inhibiting the expression of specific targets. Mechanisms of action of ncRNAs are summarized within each box.




Figure 2- Involvement of ncRNAs in ‘vicious cycle’ of bone metastases.

A metastatic cancer cell displays a different gene expression pattern in comparison to a normal cell in part as a result of dysregulated expression of lncRNAs, miRNAs, and circRNAs, which in turn alters the expression of proteins involved in metastasis formation.

Moreover, ncRNAs may serve as inter-cellular messengers for cell-cell communication.

Specifically, ncRNAs can be secreted by cancer cells and bone cells (osteoclast, osteoblast) either encapsulated within extracellular vesicles (EVs), such as microvesicles and exosomes, or as ncRNA-(lipo-)protein complexes, which are taken up by recipient cells. These ncRNAs can then reprogram recipient cells to facilitate bone metastases formation. NcRNAs involved in bone metastasis formation and progression are listed for each cell type (metastatic cancer cell, osteoblast, osteoclast).