

Phylogenomics reveals the evolutionary timing and pattern of butterflies and moths

Akito Y. Kawahara^{a,1}, David Plotkin^{a,b}, Marianne Espeland^{a,c}, Karen Meusemann^{d,e,f}, Emmanuel F. A. Toussaint^{a,g}, Alexander Donath^e, France Gimnich^e, Paul B. Frandsen^{h,i}, Andreas Zwick^f, Mario dos Reis^j, Jesse R. Barber^k, Ralph S. Peters^c, Shanlin Liu^{l,m}, Xin Zhou^m, Christoph Mayer^e, Lars Podsiadlowski^e, Caroline Storer^a, Jayne E. Yackⁿ, Bernhard Misof^e, and Jesse W. Breinholt^{a,o,2}

^aMcGuire Center for Lepidoptera and Biodiversity, Florida Museum of Natural History, University of Florida, Gainesville, FL 32611; ^bEntomology and Nematology Department, University of Florida, Gainesville, FL 32611; ^cArthropoda Department, Alexander Koenig Zoological Research Museum, D-53113 Bonn, Germany; ^dDepartment of Evolutionary Biology and Ecology, Institute for Biology I (Zoology), University of Freiburg, D-79104 Freiburg, Germany; ^eCenter for Molecular Biodiversity Research, Alexander Koenig Zoological Research Museum, D-53113 Bonn, Germany; ^fAustralian National Insect Collection, National Research Collections Australia, Commonwealth Scientific and Industrial Research Organisation, Canberra, ACT 2601, Australia; ^gDepartment of Entomology, Natural History Museum of Geneva, 1208 Geneva, Switzerland; ^hPlant and Wildlife Sciences, Brigham Young University, Provo, UT 84602; ⁱData Science Lab, Smithsonian Institution, Washington, DC 20002; ^jSchool of Biological and Chemical Sciences, Queen Mary University of London, London E1 4NS, United Kingdom; ^kDepartment of Biological Sciences, Boise State University, Boise, ID 83725; ^lChina National GeneBank, BGI-Shenzhen, 518083 Guangdong, China; ^mDepartment of Entomology, College of Plant Protection, China Agricultural University, 100193 Beijing, China; ⁿDepartment of Biology, Carleton University, Ottawa, ON, Canada K1S 5B6; and ^oBioinformatics & Data Analysis Division, RAPiD Genomics, Gainesville, FL 32601

Edited by Douglas Futuyma, Stony Brook University, Stony Brook, NY, and approved September 16, 2019 (received for review May 29, 2019)

Butterflies and moths (Lepidoptera) are one of the major super-radiations of insects, comprising nearly 160,000 described extant species. As herbivores, pollinators, and prey, Lepidoptera play a fundamental role in almost every terrestrial ecosystem. Lepidoptera are also indicators of environmental change and serve as models for research on mimicry and genetics. They have been central to the development of coevolutionary hypotheses, such as butterflies with flowering plants and moths' evolutionary arms race with echolocating bats. However, these hypotheses have not been rigorously tested, because a robust lepidopteran phylogeny and timing of evolutionary novelties are lacking. To address these issues, we inferred a comprehensive phylogeny of Lepidoptera, using the largest dataset assembled for the order (2,098 orthologous protein-coding genes from transcriptomes of 186 species, representing nearly all superfamilies), and dated it with carefully evaluated synapomorphy-based fossils. The oldest members of the Lepidoptera crown group appeared in the Late Carboniferous (~300 Ma) and fed on nonvascular land plants. Lepidoptera evolved the tube-like proboscis in the Middle Triassic (~241 Ma), which allowed them to acquire nectar from flowering plants. This morphological innovation, along with other traits, likely promoted the extraordinary diversification of superfamily-level lepidopteran crown groups. The ancestor of butterflies was likely nocturnal, and our results indicate that butterflies became day-flying in the Late Cretaceous (~98 Ma). Moth hearing organs arose multiple times before the evolutionary arms race between moths and bats, perhaps initially detecting a wide range of sound frequencies before being co-opted to specifically detect bat sonar. Our study provides an essential framework for future comparative studies on butterfly and moth evolution.

Lepidoptera | coevolution | phylogeny | angiosperms | bats

Butterflies and moths (Lepidoptera) are 1 of the 4 major insect superradiations, with close to 160,000 described species (1, 2). They thrive in nearly all terrestrial ecosystems, feature a wide spectrum of ecological adaptations, and are inextricably connected to the natural histories of many plants, predators, and parasitoids. The extraordinary diversity of Lepidoptera is presumed to be tightly linked to the rise of flowering plants (3), and the association between butterflies and angiosperms served as the foundation for Ehrlich and Raven's theory of coevolution (4). Prior studies of lepidopteran evolution postulated that the larvae of the earliest lineages were internal feeders on nonvascular land plants and their adults had mandibulate chewing mouthparts (5, 6). The remaining majority of Lepidoptera are hypothesized to have diversified largely in parallel with angiosperms, with larvae that are

external plant feeders and adults that have tube-like proboscides (7). The proboscis promoted access to flower nectar and may have led to the ability of these ancestors to disperse to greater distances and colonize new host plants.

Nocturnal moths represent >75% of Lepidoptera species diversity (8), and they are thought to have proliferated in the Late

Significance

Lepidoptera play key roles in many biological systems. Butterflies are hypothesized to have evolved contemporaneously with flowering plants, and moths are thought to have gained anti-bat defenses in response to echolocating predatory bats, but these hypotheses have largely gone untested. Using a transcriptomic, dated evolutionary tree of Lepidoptera, we demonstrate that the most recent common ancestor of Lepidoptera is considerably older than previously hypothesized. The oldest moths in crown Lepidoptera were present in the Carboniferous, some 300 million years ago, and began to diversify largely in synchrony with angiosperms. We show that multiple lineages of moths independently evolved hearing organs well before the origin of bats, rejecting the hypothesis that lepidopteran hearing organs arose in response to these predators.

Author contributions: A.Y.K., R.S.P., and B.M. designed research; A.Y.K., D.P., M.E., K.M., E.F.A.T., A.D., F.G., P.B.F., A.Z., M.d.R., S.L., X.Z., C.M., L.P., C.S., J.E.Y., B.M., and J.W.B. performed research; A.Y.K., K.M., P.B.F., A.Z., J.R.B., R.S.P., J.E.Y., B.M., and J.W.B. collected or provided samples; A.Y.K., D.P., M.E., K.M., E.F.A.T., A.D., F.G., P.B.F., A.Z., M.d.R., C.M., L.P., C.S., J.E.Y., B.M., and J.W.B. analyzed data; and A.Y.K., D.P., M.E., K.M., E.F.A.T., A.D., F.G., P.B.F., A.Z., M.d.R., J.R.B., R.S.P., S.L., X.Z., C.M., L.P., C.S., J.E.Y., B.M., and J.W.B. wrote the paper.

The authors declare no competing interest.

This article is a PNAS Direct Submission.

This open access article is distributed under [Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 \(CC BY-NC-ND\)](https://creativecommons.org/licenses/by-nc-nd/4.0/).

Data deposition: Transcriptome assemblies for the newly sequenced transcriptomes used in this study either have been deposited in the National Center for Biotechnology Information Transcriptome Shotgun Assembly archive (all accession nos. provided in [Dataset S1](#)) or in a DRYAD digital repository (<https://doi.org/10.5061/dryad.j477b40>, [Supplementary Archive 1](#)). Other data files have also been archived in the same DRYAD repository. The contents of these archives are listed in the [SI Appendix](#).

¹To whom correspondence may be addressed. Email: kawahara@flmnh.ufl.edu.

²Present address: Division of Bioinformatics, Intermountain Healthcare, Precision Genomics, St. George, UT 84790.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1907847116/-DCSupplemental.

Cretaceous when they evolved bat-detecting ultrasonic hearing organs (9, 10). At least 10 lepidopteran families have these ears (11), which are on different parts of the body in different families (*SI Appendix*, Fig. S1). The timing of when ears arose has remained untested because a stable, dated phylogenetic framework for the Lepidoptera has been unavailable. Previous phylogenetic studies that examined macroevolutionary patterns of Lepidoptera (e.g., refs. 6 and 12–17) lacked sufficient genetic data and carefully evaluated fossil sampling to confidently resolve the evolutionary history of the order.

We constructed the largest dataset of transcriptomes for phylogenetic inference of the order, comprising 2,098 protein-coding genes sampled from 186 extant Lepidoptera species. Using different dating schemes and a set of critically evaluated fossils, we conducted multiple fossil-calibrated phylogenetic analyses. With the dated tree, we examined the evolution and timing of angiosperm feeding, ultrasonic hearing organs, and adult diel activity. Our phylogenetic analyses produced a robust reconstruction of the evolutionary history and timing of these key ecological adaptations in this megadiverse insect order.

Results and Discussion

We assembled de novo characterized transcriptomes and genomes of 186 Lepidoptera species, representing 34 superfamilies, and 17 outgroup species (*Datasets S1–S4*). Phylogenetic analyses were based on nucleotide and amino acid datasets of 2,098 protein coding gene alignments (2,249,363 nucleotides and 749,791 amino acid sites). In addition to maximum likelihood (ML) tree reconstruction, we conducted multispecies coalescence (MSC) analyses and 4-cluster likelihood mapping (FcLM) (18) to alleviate limitations that exist with traditional support metrics, and evaluate the presence of alternative, potentially conflicting signals in our datasets. ML analyses resulted in statistically well-supported trees (*SI Appendix*, Figs. S2–S8), which were largely corroborated by MSC results (*SI Appendix*, Figs. S9 and S10 and *Dataset S5*). The FcLM analyses (*Datasets S6–S8*) indicated that there was little confounding signal. The placement of a few lineages remains uncertain (*SI Appendix*, Fig. S11 and *Dataset S9*), but they do not significantly impact our conclusions. All 8 divergence time analyses resulted in largely congruent age estimates (*SI Appendix*, Figs. S12–S32 and *Datasets S9–S11*). The difference between the oldest and youngest median age estimates for crown Lepidoptera was ~21 Ma, which is ~7% of the median age presented in Fig. 1. When the same calculations are performed on 27 other key lepidopteran crown nodes, only 1 had a median age estimate difference >26 Ma (*Dataset S11*).

Phylogenetic analyses, regardless of dataset or analysis type, recovered the monophyly of Lepidoptera and many other previously hypothesized clades within the order, including Angiospermivora, Glossata, and Ditrysia, with strong branch support (*Dataset S5*). Our results indicate that the most recent common ancestor of crown Lepidoptera appeared in the Late Carboniferous, ~299.5 Ma (95% credibility interval [CI], 312.4 to 276.4 Ma; Fig. 1), significantly predating the oldest known lepidopteran fossil from the Triassic-Jurassic boundary (201 Ma; *Dataset S10*). Among Lepidoptera, the superfamilies Micropterigoidea, Agathiphaogoidea, and Heterobathmioidea have been considered “primitive moths,” because their adults have functional mandibulate mouthparts and other plesiomorphic features, including feeding on nonvascular land plants (19). Prior molecular studies suggested that Agathiphaogoidea, which are seed borers on the ancient conifer family Araucariaceae (19, 20), is the sister group to either Micropterigoidea (12, 21) or Heterobathmioidea (22). Our study contradicts these findings, confidently placing Micropterigoidea, which are detritivores or bryophyte feeders (7), as the sister group to all other Lepidoptera (280.6 Ma; CI, 297.0 to 257.4 Ma), with Agathiphaogoidea as the closest relative to all remaining lineages, corroborating traditional morphology-based studies (19). Our results imply that the ancestral mandibulate moth may have

been feeding on bryophytes ~300 Ma, followed by some lineages transitioning to vascular plants and then to angiosperms, largely following the evolutionary history of plants.

Angiospermivora, which feed predominantly on flowering plants as larvae (21), are composed of the mandibulate Heterobathmioidea and all proboscis-bearing Lepidoptera. The crown age of Angiospermivora is dated at 257.7 Ma (CI, 276.7 to 234.5 Ma), and our analyses unequivocally inferred Heterobathmioidea as a sister group to all other taxa within this clade. Both groups have leaf-mining larvae (19), suggesting that moths were internal plant tissue feeders in the Late Permian. Although the crown age of flowering plants remains controversial (23), there is general agreement that some angiosperm lineages, which are in part now extinct, were present >300 Ma (24), a consensus corroborated by a long stem branch that links angiosperms to gymnosperms in recent studies (24–29). Our results support the hypothesis that Lepidoptera were internal feeders before becoming predominantly external herbivores (1, 3, 7, 19). Large body size, a characteristic of some extant butterflies and moths, may be an ecological consequence of being freed from the constraints of living within plants (7), thereby allowing greater access to nectar sources and the ability to fly far distances.

The adult proboscis is the unifying feature of clade Glossata, which contains >99% of all extant Lepidoptera species (2). The proboscis is considered an important innovation that enabled lepidopteran diversification (19). Adults of Heterobathmioidea, the sister group to Glossata, use their mandibles to eat pollen of the ancient *Nothofagus* tree (19). Adults of Eriocranioidea, part of a 3-family clade sister to all other Glossata, use their proboscis to drink water or sap (19), a behavior that may have been the precursor to nectar feeding. The common ancestor of nectar-feeding Lepidoptera first appeared in the Middle Triassic (241.4 Ma; CI, 261.1 to 218.9 Ma), overlapping with the estimated diversification periods of speciose flowering plant crown groups (Fig. 2).

The clade Ditrysia, which is estimated to have a late-Jurassic crown age (154.7 Ma; CI, 172.1 to 137.5 Ma) is the most taxonomically and ecologically diverse group of Lepidoptera, with >150,000 described species. Ditrysia includes butterflies and many moth superfamilies that predominantly feed on angiosperms as larvae. Monophyly of Ditrysia, recovered in all of our molecular analyses, is supported by a unique apomorphy: females have 2 reproductive openings, 1 for mating and another for egg-laying (17, 19). Relationships among major ditrysonian groups such as Apoditrysia, sensu Regier et al. (12) and Mutanen et al. (22), uniting butterflies and many diverse moth families, are statistically well supported (*Dataset S5*). Most Apoditrysia have additional sensilla on the proboscis (19), implying that further specialization on adult food resources may have arisen in the Early Cretaceous (118.5 Ma; CI, 132.1 to 105.6 Ma).

Butterflies (Papilionoidea) are some of the most popular arthropods, and include nearly 19,000 described species (3). One of the major challenges thus far has been to confidently place butterflies within Lepidoptera (12, 17, 21, 30). We identified Papilionoidea as the sister group to all remaining Obtectomera (including silk moths, owlet moths, grass moths, and others; Fig. 1) with strong nodal support and statistical evidence from topology tests that rule out the impact of confounding signal (*SI Appendix*, Fig. S11). This result confidently shows that butterflies are diurnal moths with a Late Cretaceous crown age (98.3 Ma; CI, 110.3 to 86.9 Ma), slightly younger than other published estimates using butterfly fossils and host-plant maximum age calibrations (13, 31–34). Swallowtails (Papilionoidea) were the sister group to the remaining butterfly families; skippers (Hesperiidae) and neotropical nocturnal moth-like butterflies (Hedylidae) were sister groups (84.4 Ma; CI, 96.6 to 72.2 Ma). Whites and sulfurs (Pieridae; 51.7 Ma; CI, 63.9 to 40.4 Ma) formed a clade that was the sister group to the clade containing the hairstreaks, blues, metalmarks (Lycaenidae + Riodinidae; 66.6 Ma;

hearing organs in Lepidoptera, and animals more broadly, is general auditory surveillance of the environment for broadband sounds produced by animal movement (e.g., predators) (45). Behavioral and neural evidence has shown that moths respond to the low-frequency (<20 kHz) walking and wingbeat sounds of predatory birds (46, 47). Day-flying Lepidoptera and moths endemic to bat-free islands show reduced auditory sensitivity to higher frequencies yet retain reactive thresholds at lower frequencies, comparable to the thresholds of other nocturnal moth species (10). Therefore, hearing organs likely evolved first for auditory surveillance before subsequently being co-opted for bat detection.

In summary, our study reveals that the common ancestor of the butterflies and moths we observe today was likely a small, Late Carboniferous species with mandibulate adults and with larvae that fed internally on nonvascular land plants. Our dating analyses with different fossils and models confirm the hypothesis that the majority of Lepidoptera proliferated with the rise of angiosperms in the Cretaceous. Hearing organs in nocturnal Lepidoptera originated several times separately, before the rise of echolocating bats. Butterflies similarly became diurnal before diversification of the bat crown group, contrary to the hypothesis that butterflies became day-fliers to escape these predators. Instead, butterflies likely became diurnal to capitalize on day-blooming flowers. Our study provides a robust phylogenetic framework and reliable time estimates for future studies on the ecology and evolution of one of the most prominent insect lineages. Data and results from this study will serve as a foundation for future comparative analyses on butterflies and moths.

Materials and Methods

Phylogenomic Analyses. Transcriptome and genome sequences from 186 species of Lepidoptera were used for our study, along with sequences of 17 outgroup species representing other holometabolous insect orders, totaling 203 species (Dataset S1). Transcriptomes from 69 of these species were newly sequenced by either the 1K Insect Transcriptome Evolution (1KITE) consortium or by the Kawahara Lab, Florida Museum of Natural History (FLMNH). RNA extractions, mRNA isolation, fragmentation, cDNA library construction, and transcriptome sequencing were performed using the protocols of Misof et al. (48) and Peters et al. (44) for the 1KITE consortium samples. Protocols of Kawahara and Breinholt (16) were used for the FLMNH samples (SI Appendix, section 2).

Transcriptome assembly and contaminant removal protocols largely followed Peters et al. (44) and Kawahara and Breinholt (16); additional information on these methods is available in SI Appendix, sections 3 and 4. An ortholog set was compiled based on the OrthoDB v7 database (49), using 2 ingroups (*Bombyx mori* and *Danaus plexippus*) and 1 outgroup (*Tribolium castaneum*) as reference species. The assemblies were searched for transcripts of 3,429 single-copy protein coding genes using Orthograph beta4 (50). Orthograph identified a total of 3,427 of the 3,429 orthologs across all included taxa. Amino acid sequences of each ortholog were aligned in MAFFT v7.294 (51) and refined using the protocols of Misof et al. (48). Corresponding nucleotide alignments were generated using a modified version of Pal2Nal v14 (52) with amino acid alignments as blueprints. Supermatrices for the amino acid and nucleotide datasets were generated using FasConCat-G v1.0 (53), and synonymous signal was removed from the nucleotide dataset using Degen v1.4 (54). Additional information on generation of the ortholog set and the final amino acid and “degen1” nucleotide datasets is available in SI Appendix, sections 5 and 6. PartitionFinder2 v2.1.1 (55) and RAxML v8.2.11 (56) were used to obtain the optimal partitioning scheme for the concatenated amino acid dataset; the boundaries of the merged partitions in this scheme were adjusted at the nucleotide level and used for the concatenated degen1 nucleotide dataset. The best-fitting models were reestimated using ModelFinder in IQ-TREE v1.5.5 (57, 58) for both datasets, and maximum likelihood (ML) tree inferences were subsequently conducted. A minimum of 50 ML tree searches were performed for each dataset using IQ-TREE v1.5.5 (57); the tree with the best log-likelihood was selected as the best tree for each analysis. Multiple metrics of statistical support, including nonparametric bootstrap replicates, SH-aLRT (59), and TBE support values (60), were generated to assess the reliability of the best ML tree in each analysis. Multispecies coalescent analyses were also performed on the amino acid dataset in ASTRAL-III v5.6.3 (61), using 2 sets of

individual gene trees that had been inferred with IQ-TREE v1.6.10 (57): a set of the best ML trees for each gene (inferred from 25 ML tree searches per locus) and a set of the consensus trees for each gene partition, with each consensus tree generated using 1,000 ultrafast bootstrap (UFBoot2) replicates (62).

The placement of 3 selected Lepidoptera clades (Alucitidae + Pterophoridae, Bombycoidea, and Papilionoidea) was further assessed using FCLM in IQ-TREE v1.6.7 (57), with predefined groups of taxa, to check for hidden signal and bias due to nonrandomly distributed missing data and (among-lineage) compositional heterogeneity, following the protocol of Misof et al. (48). Additional information on data partitioning, model selection, topology tests, tree inference using the partitioned supermatrix and the multispecies coalescent approach, and branch support is available in SI Appendix, sections 7 and 8.

Divergence Time Estimation. Divergence time estimations were computed on truncated datasets to allow completion of the analyses. AliStat v1.6 (63) was used to generate a subsampled dataset containing 195 Amphimesoptera species (all non-Amphimesoptera removed) and only including sites for which at least 80% of samples had unambiguous amino acids. MCMCTree and codeml (both part of the PAML software package, v4.9g) (64) were used to estimate divergence dates on this subsampled dataset. The best ML tree inferred from the concatenated amino acid dataset was used as the input tree. The input tree was first calibrated using age estimates of 16 carefully selected fossils, following the best-practice recommendations by Parham et al. (65). Although all 16 fossils have diagnostic morphological characteristics that enable reasonably confident placement on the tree, only 3 of these fossils have true synapomorphies. To strictly follow the guidelines of Parham et al. (65), additional analyses were performed using only these 3 fossils. We applied a conservative age constraint on the root of the input tree, with a minimum age of 201 Ma, based on the stem Glossata scale fossils discovered by van Eldijk et al. (66) and a maximum age of 314.4 Ma, based on the absence of Amphimesoptera fossils in the Late Carboniferous. We used 2 well-established approaches to convert fossil ages into calibrations on tree nodes. For the conservative strategy, fossil calibrations were treated with uniform distributions constrained between the corresponding fossil age (the minimum bound) and a hard maximum equal to the maximum root age. For the second strategy, the truncated-Cauchy distribution (67) was used to set calibrations for internal nodes younger than 80 Ma. (Additional information on this approach is provided in SI Appendix, section 10.) We applied both uncorrelated rates and autocorrelated rates to estimate divergence, for a total of 8 analyses: 2 sets of fossil calibrations (16 fossils and 3 fossils) × 2 fossil calibration strategies (uniform and Cauchy priors for nodes <80 Ma) × 2 rate types (independent and autocorrelated). To compare the radiation of Lepidoptera and flowering plants, estimates of the mean age of the ancestral angiosperm were compiled from the literature; these angiosperm ages are presented in Dataset S12 and shown in Fig. 2. Since the estimated timespan between divergence of angiosperms and gymnosperms is large, and it is possible that flowering plants existed long before the crown of angiosperms, we present the interval between the mean age of the crown (node Angiosperm) and the mean age of the stem (node Angiosperm + Gymnosperm) from the above-mentioned studies (Dataset S12). Additional information and justification for our approaches to divergence time estimation is available in SI Appendix, section 10.

Ancestral State Reconstruction. We conducted ancestral state reconstruction on ultrasonic hearing organs and adult diel activity. All ancestral state reconstruction analyses were performed on the best tree from the ML analysis of the concatenated amino acid dataset (SI Appendix, Figs. S33 and S34), using stochastic character mapping, with the “make.simmap” command in the R package Phytools v06-44 (68). Ten thousand stochastic maps were generated for each analysis. The presence of a hearing organ was treated as a binary character in one ASR analysis and as a 7-state character in a separate ASR analysis that accounts for variation in the location of the hearing organ (see matrices A and B in Dataset S13). Diel activity was treated as a 4-state character (Dataset S14). Additional information on ancestral state reconstruction and how characters were coded is provided in SI Appendix, section 11.

Data Availability. Additional data are provided in Supplementary Archives on DRYAD (70).

ACKNOWLEDGMENTS. Data presented here are the result of the collaborative efforts of the 1KITE consortium (<https://www.1kite.org/>). Sequencing and assembly of 1KITE transcriptomes were funded by BGI through support to the

China National GeneBank. This project was also financially supported by multiple grants from the NSF and the German Research Foundation; a list of grant numbers and additional acknowledgements are provided in *SI Appendix*.

1. D. Grimaldi, M. S. Engel, *Evolution of the Insects* (Cambridge Univ Press, Cambridge, UK, 2005), p. 772.
2. E. J. van Nieuwerkerken *et al.*, "Order Lepidoptera Linnaeus, 1758" in *Animal Biodiversity: An Outline of Higher-Level Classification and Survey of Taxonomic Richness*, Z.-Q. Zhang, Ed. (Magnolia Press, Auckland, New Zealand, 2011), vol. 3148, pp. 212–221.
3. C. Mitter, D. R. Davis, M. P. Cummings, Phylogeny and evolution of Lepidoptera. *Annu. Rev. Entomol.* **62**, 265–283 (2017).
4. P. R. Ehrlich, P. H. Raven, Butterflies and plants: A study in coevolution. *Evolution* **18**, 586–608 (1964).
5. B. M. Wiegmann *et al.*, Nuclear genes resolve Mesozoic-aged divergences in the insect order Lepidoptera. *Mol. Phylogenet. Evol.* **15**, 242–259 (2000).
6. A. L. Bazinet *et al.*, Phylotranscriptomics resolves ancient divergences in the Lepidoptera. *Syst. Entomol.* **42**, 305–316 (2017).
7. J. A. Powell, C. Mitter, B. D. Farrell, "Evolution of larval food preferences in Lepidoptera" in *Handbook of Zoology*, N. P. Kristensen, Ed. (Volume IV, Arthropoda: Insecta, Part 35. Lepidoptera, Moths and Butterflies, Volume 1: Evolution, Systematics, and Biogeography, Walter de Gruyter, Berlin, New York, 1998), pp. 403–422.
8. A. Y. Kawahara *et al.*, Diel behavior in moths and butterflies: A synthesis of data illuminates the evolution of temporal activity. *Org. Divers. Evol.* **18**, 13–27 (2018).
9. W. E. Conner, A. J. Corcoran, Sound strategies: The 65-million-year-old battle between bats and insects. *Annu. Rev. Entomol.* **57**, 21–39 (2012).
10. H. M. ter Hofstede, J. M. Ratcliffe, Evolutionary escalation: The bat-moth arms race. *J. Exp. Biol.* **219**, 1589–1602 (2016).
11. N. P. Kristensen, Molecular phylogenies, morphological homologies and the evolution of "moth ears". *Syst. Entomol.* **37**, 237–239 (2012).
12. J. C. Regier *et al.*, A large-scale, higher-level, molecular phylogenetic study of the insect order Lepidoptera (moths and butterflies). *PLoS One* **8**, e58568 (2013).
13. N. Wahlberg, C. W. Wheat, C. Peña, Timing and patterns in the taxonomic diversification of Lepidoptera (butterflies and moths). *PLoS One* **8**, e80875 (2013).
14. M. Heikkilä, M. Mutanen, N. Wahlberg, P. Sihvonen, L. Kaila, Elusive ditrysin phylogeny: An account of combining systematized morphology with molecular data (Lepidoptera). *BMC Evol. Biol.* **15**, 260 (2015).
15. A. L. Bazinet, M. P. Cummings, K. T. Mitter, C. W. Mitter, Can RNA-Seq resolve the rapid radiation of advanced moths and butterflies (Hexapoda: Lepidoptera: Apoditrysia)? An exploratory study. *PLoS One* **8**, e82615 (2013).
16. A. Y. Kawahara, J. W. Breinholt, Phylogenomics provides strong evidence for relationships of butterflies and moths. *Proc. Biol. Sci.* **281**, 20140970 (2014).
17. J. W. Breinholt *et al.*, Resolving relationships among the megadiverse butterflies and moths with a novel pipeline for anchored phylogenomics. *Syst. Biol.* **67**, 78–93 (2018).
18. K. Strimmer, A. von Haeseler, Likelihood-mapping: A simple method to visualize phylogenetic content of a sequence alignment. *Proc. Natl. Acad. Sci. U.S.A.* **94**, 6815–6819 (1997).
19. N. P. Kristensen, *Handbook of Zoology*, N. P. Kristensen, Ed. (Volume IV, Arthropoda: Insecta, Part 35. Lepidoptera, Moths and Butterflies, Volume 1: Evolution, Systematics, and Biogeography, Walter de Gruyter, Berlin, New York, 1998).
20. A. B. Leslie *et al.*, Hemisphere-scale differences in conifer evolutionary dynamics. *Proc. Natl. Acad. Sci. U.S.A.* **109**, 16217–16221 (2012).
21. J. C. Regier *et al.*, A molecular phylogeny and revised classification for the oldest ditrysin moth lineages (Lepidoptera: Tineoidea), with implications for ancestral feeding habits of the mega-diverse Ditrysia. *Syst. Entomol.* **40**, 409–432 (2015).
22. M. Mutanen, N. Wahlberg, L. Kaila, Comprehensive gene and taxon coverage elucidates radiation patterns in moths and butterflies. *Proc. Biol. Sci.* **277**, 2839–2848 (2010).
23. J. M. Beaulieu, B. C. O'Meara, P. Crane, M. J. Donoghue, Heterogeneous rates of molecular evolution and diversification could explain the Triassic age estimate for angiosperms. *Syst. Biol.* **64**, 869–878 (2015).
24. H. Sauquet, S. Magallón, Key questions and challenges in angiosperm macroevolution. *New Phytol.* **219**, 1170–1187 (2018).
25. H.-T. Li *et al.*, Origin of angiosperms and the puzzle of the Jurassic gap. *Nat. Plants* **5**, 461–470 (2019).
26. K. Salomo *et al.*, The emergence of earliest angiosperms may be earlier than fossil evidence indicates. *Syst. Bot.* **42**, 607–619 (2017).
27. C. S. P. Foster *et al.*, Evaluating the impact of genomic data and priors on Bayesian estimates of the angiosperm evolutionary timescale. *Syst. Biol.* **66**, 338–351 (2017).
28. C. S. P. Foster, S. Y. W. Ho, Strategies for partitioning clock models in phylogenomic dating: Application to the angiosperm evolutionary timescale. *Genome Biol. Evol.* **9**, 2752–2763 (2017).
29. J. Barba-Montoya, M. Dos Reis, H. Schneider, P. C. J. Donoghue, Z. Yang, Constraining uncertainty in the timescale of angiosperm evolution and the veracity of a Cretaceous Terrestrial Revolution. *New Phytol.* **218**, 819–834 (2018).
30. J. C. Regier *et al.*, Toward reconstructing the evolution of advanced moths and butterflies (Lepidoptera: Ditrysia): An initial molecular study. *BMC Evol. Biol.* **9**, 280 (2009).
31. M. Espeland *et al.*, A comprehensive and dated phylogenomic analysis of butterflies. *Curr. Biol.* **28**, 770–778.e5 (2018).
32. N. Wahlberg *et al.*, Nymphalid butterflies diversify following near demise at the Cretaceous/Tertiary boundary. *Proc. Biol. Sci.* **276**, 4295–4302 (2009).
33. M. Heikkilä, L. Kaila, M. Mutanen, C. Peña, N. Wahlberg, Cretaceous origin and repeated tertiary diversification of the redefined butterflies. *Proc. Biol. Sci.* **279**, 1093–1099 (2012).
34. N. Chazot *et al.*, Priors and posteriors in Bayesian timing of divergence analyses: The age of butterflies revisited. *Syst. Biol.* **68**, 797–813 (2019).
35. J. E. Yack, J. H. Fullard, Ultrasonic hearing in nocturnal butterflies. *Nature* **403**, 265–266 (2000).
36. J. H. Fullard, "Sensory coevolution of moths and bats" in *Comparative Hearing: Insects*, R. R. Hoy, A. N. Popper, R. R. FayIn, Eds. (Springer, New York, 1998), pp. 279–326.
37. N. Upham, J. A. Esselstyn, W. Jetz, Ecological causes of uneven diversification and richness in the mammal tree of life. bioRxiv:10.1101/504803 (posted March 28, 2019).
38. M. dos Reis *et al.*, Phylogenomic datasets provide both precision and accuracy in estimating the timescale of placental mammal phylogeny. *Proc. Biol. Sci.* **279**, 3491–3500 (2012).
39. J. J. Shi, D. L. Rabosky, Speciation dynamics during the global radiation of extant bats. *Evolution* **69**, 1528–1545 (2015).
40. E. C. Teeling, Hear, hear: The convergent evolution of echolocation in bats? *Trends Ecol. Evol.* **24**, 351–354 (2009).
41. I. Agnarsson, C. M. Zambrana-Torrel, N. P. Flores-Saldana, L. J. May-Collado, A time-calibrated species-level phylogeny of bats (Chiroptera, Mammalia). *PLoS Curr.* **3**, RRN1212 (2011).
42. N. Veselka *et al.*, A bony connection signals laryngeal echolocation in bats. *Nature* **463**, 939–942 (2010).
43. N. B. Simmons, K. L. Seymour, J. Habersetzer, G. F. Gunnell, Primitive Early Eocene bat from Wyoming and the evolution of flight and echolocation. *Nature* **451**, 818–821 (2008).
44. R. S. Peters *et al.*, Evolutionary history of the Hymenoptera. *Curr. Biol.* **27**, 1013–1018 (2017).
45. A. Mikhail, J. E. Lewis, J. E. Yack, What does a butterfly hear? Physiological characterization of auditory afferents in *Morpho peleides* (Nymphalidae). *J. Comp. Physiol. A Neuroethol. Sens. Neural Behav. Physiol.* **204**, 791–799 (2018).
46. D. S. Jacobs, J. M. Ratcliffe, J. H. Fullard, Beware of bats, beware of birds: The auditory responses of eared moths to bat and bird predation. *Behav. Ecol.* **19**, 1333–1342 (2008).
47. J. P. Fournier, J. W. Dawson, A. Mikhail, J. E. Yack, If a bird flies in the forest, does an insect hear it? *Biol. Lett.* **9**, 20130319 (2013).
48. B. Misof *et al.*, Phylogenomics resolves the timing and pattern of insect evolution. *Science* **346**, 763–767 (2014).
49. R. M. Waterhouse, F. Tegenfeldt, J. Li, E. M. Zdobnov, E. V. Kriventseva, OrthoDB: A hierarchical catalog of animal, fungal and bacterial orthologs. *Nucleic Acids Res.* **41**, D358–D365 (2013).
50. M. Petersen *et al.*, Orthograph: A versatile tool for mapping coding nucleotide sequences to clusters of orthologous genes. *BMC Bioinformatics* **18**, 111 (2017).
51. K. Katoh, D. M. Standley, MAFFT multiple sequence alignment software version 7: Improvements in performance and usability. *Mol. Biol. Evol.* **30**, 772–780 (2013).
52. M. Suyama, D. Torrents, P. Bork, PAL2NAL: Robust conversion of protein sequence alignments into the corresponding codon alignments. *Nucleic Acids Res.* **34**, W609–W612 (2006).
53. P. Kück, G. C. Longo, FASconCAT-G: Extensive functions for multiple sequence alignment preparations concerning phylogenetic studies. *Front. Zool.* **11**, 81 (2014).
54. A. Zwick, J. C. Regier, D. J. Zwickl, Resolving discrepancy between nucleotides and amino acids in deep-level arthropod phylogenomics: Differentiating serine codons in 21-amino-acid models. *PLoS One* **7**, e47450 (2012).
55. R. Lanfear, P. B. Frandsen, A. M. Wright, T. Senfeld, B. Calcott, PartitionFinder 2: New methods for selecting partitioned models of evolution for molecular and morphological phylogenetic analyses. *Mol. Biol. Evol.* **34**, 772–773 (2017).
56. A. Stamatakis, RAxML version 8: A tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* **30**, 1312–1313 (2014).
57. L. T. Nguyen, H. A. Schmidt, A. von Haeseler, B. Q. Minh, IQ-TREE: A fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol. Biol. Evol.* **32**, 268–274 (2015).
58. O. Chernomor, A. von Haeseler, B. Q. Minh, Terrace aware data structure for phylogenomic inference from supermatrices. *Syst. Biol.* **65**, 997–1008 (2016).
59. S. Guindon *et al.*, New algorithms and methods to estimate maximum-likelihood phylogenies: Assessing the performance of PhyML 3.0. *Syst. Biol.* **59**, 307–321 (2010).
60. F. Lemoine *et al.*, Renewing Felsenstein's phylogenetic bootstrap in the era of big data. *Nature* **556**, 452–456 (2018).
61. C. Zhang, M. Rabiee, E. Sayyari, S. Mirarab, ASTRAL-III: Polynomial time species tree reconstruction from partially resolved gene trees. *BMC Bioinformatics* **19** (suppl. 6), 153 (2018).
62. D. T. Hoang, O. Chernomor, A. von Haeseler, B. Q. Minh, L. S. Vinh, UFBoot2: Improving the ultrafast bootstrap approximation. *Mol. Biol. Evol.* **35**, 518–522 (2018).
63. T. K. F. Wong *et al.*, CSIRO software collection (AliStat version 1.3, CSIRO, Canberra, Australia, 2014).
64. Z. Yang, PAML 4: Phylogenetic analysis by maximum likelihood. *Mol. Biol. Evol.* **24**, 1586–1591 (2007).
65. J. F. Parham *et al.*, Best practices for justifying fossil calibrations. *Syst. Biol.* **61**, 346–359 (2012).

66. T. J. B. van Eldijk *et al.*, A Triassic-Jurassic window into the evolution of Lepidoptera. *Sci. Adv.* **4**, e1701568 (2018).
67. J. Inoue, P. C. J. Donoghue, Z. Yang, The impact of the representation of fossil calibrations on Bayesian estimation of species divergence times. *Syst. Biol.* **59**, 74–89 (2010).
68. L. J. Revell, Phytools: An R package for phylogenetic comparative biology (and other things). *Methods Ecol. Evol.* **3**, 217–223 (2012).
69. P. A. Hochuli, S. Feist-Burkhardt, Angiosperm-like pollen and Afropollis from the Middle Triassic (Anisian) of the Germanic Basin (Northern Switzerland). *Frontiers Plant Sci.* **4**, 1–14 (2013).
70. A. Y. Kawahara *et al.*, Phylogenomics reveals the evolutionary timing and pattern of butterflies and moths. *Dryad*. <https://doi.org/10.5061/dryad.j477b40>. Deposited 18 September 2019.