

Crenicichla tesay, a new species of cichlid (Perciformes: labroidei) from the río Iguazú basin in Argentina

J Casciotta and A Almirón

Revue Suisse De Zoologie 115:651-660 (2008) <http://biostor.org/reference/115676>



Page images from the Biodiversity Heritage Library, <http://www.biodiversitylibrary.org/>, made available under a Creative Commons Attribution-Noncommercial License <http://creativecommons.org/licenses/by-nc/2.5/>

***Crenicichla tesay*, a new species of cichlid (Perciformes: Labroidei) from the río Iguazu basin in Argentina**

Jorge CASCIOTTA¹ & Adriana ALMIRÓN²

¹ CIC, División Zoología Vertebrados, Facultad de Ciencias Naturales y Museo, UNLP, Paseo del Bosque s/n, 1900 La Plata, Argentina. E-mail: jrcas@fcnym.unlp.edu.ar

² División Zoología Vertebrados, Facultad de Ciencias Naturales y Museo, UNLP, Paseo del Bosque s/n, 1900 La Plata, Argentina.

***Crenicichla tesay*, a new species of cichlid (Perciformes: Labroidei) from the río Iguazu basin in Argentina.** - *Crenicichla tesay* sp. n. differs from its congeners by the following combination of characters: posterior border of preopercle serrated, 48-59 scales on E1 row, snout length 2.6-2.8 times in head length, presence of conspicuous suborbital stripe, flanks with 4 to 6 blotches below the upper lateral line, posteriormost blotch extending on caudal peduncle. Colour pattern with numerous irregularly scattered dots, and absence of vertical stripes on flanks.

Keywords: Neotropical - Cichlidae - *Crenicichla* - Argentina - freshwater fish - new species

INTRODUCTION

The río Iguazu is a tributary of the río Parana that originates at the Serra do Mar in Brazil. This river runs in an east-west direction for 1,320 km of which the final 115 km flow between the Argentina and Brazil border. About 32 km before it joins the río Parana, the Iguazú falls produce a 78 m high drop (Severi & Cordeiro, 1994). These falls have been an effective fish-fauna barrier since their formation in the Oligocene-Miocene period.

The poorness of the fish fauna of the río Iguazu was noted by Haseman (1911) and Godoy (1979); however, very few studies have focused on its ichthyofauna. Out of seventy five species that have been recorded for the río Iuazu basin, 60% are endemic (Agostinho *et al.*, 2005). However this number is probably an underestimation, given the description of several new species in recent years (Garavello, 2005; Casciotta *et al.*, 2006a; 2006b; Garavello & Shibata, 2007).

The genus *Crenicichla* inhabits tropical, subtropical and temperate Cis-Andean environments from Northern South America to río Negro, in Patagonia, Argentina (Casciotta, 1987). *Crenicichla* comprises about 78 species and represents the most speciose genus of cichlid fishes (Kullander & Lucena, 2006). In spite of this great number of species, only two *Crenicichla* species have been recorded in the río Iguazu basin *C. iguassuensis* Haseman, 1911 and *C. yaha* Casciotta *et al.*, 2006. The number of species of *Crenicichla* in the río Iguazu could increase in the future if some *C. iguassuensis* morphs are given species status (Renesto *et al.*, 2001 and Mizoguchi *et al.*, 2007).

The aim of this paper is to describe a new species of the genus *Crenicichla* from tributaries of the río Iguazu upstream from Iguazu falls (Fig. 1).

MATERIAL AND METHODS

Specimens were cleared and counterstained (C&S) following the method of Taylor & Van Dyke (1985). Measurements and counts were taken as described by Kullander (1986). Pharyngeal teeth description and counts of frashed zone concavities follow Casciotta & Arratia (1993). Holotype values are indicated by an asterisk. Body length is expressed as standard length (SL). EI scale counts refer to the scales in the row immediately above that containing the lower lateral line (Lucena & Kullander, 1992).

Institutional abbreviations are as listed in Leviton *et al.* (1985), except for AI (Asociación Ictiológica, La Plata, Argentina) and PNI (Parque Nacional Iguazú, Misiones, Argentina).

Comparative material: A list of comparative material of *Crenicichla niederlemii*, *C. scottii*, *C. vitata* and *C. celidochilus* is available in Casciotta (1987).

In addition, the following material was studied: *Crenicichla hadrostigma*: Argentina AI 220, 1 ex., 72.8 mm, Misiones, Itacaruaré, río Uruguay basin. *Crenicichla iguassuensis*: Brasil. FMNH 54159 (holotype), 137 mm, Porto Uniao da Victoria, Rio Iguassu. *Crenicichla jupiaensis*: Argentina. AI 226, 2 ex., 87.7-93.0 mm, Corrientes, río Parana at Yahape. AI 227, 1 ex., 60.7 mm, Corrientes, río Parana at Yahape. *Crenicichla lepidota*: Argentina. MACN-ict 5067, 4 ex., 67.7-113.4 mm, Misiones, Represa Estación Experimental Cerro Azul. FML 00528, 1 ex., 111.5 mm, Salta, Luna Muerta, Hickman. MACN-ict 3656, 2 ex., 116.0-165.7 mm, Formosa, Riacho de Oro. MACN-ict 7275, 1 ex., 151.6 mm, Chaco, Esteros del Palmar. FML 00312, 1 ex., 138.0 mm, Corrientes, Isla Apipé Grande, Ituzaingó. MACN-ict 4091, 1 ex., 98.4 mm, Entre Ríos, río Uruguay, Concepción del Uruguay. MACN-ict 2314, 6 ex., 59.9-104.2 mm, Buenos Aires, Isla Martín García, Uruguay. MNHN 2087, 1 ex., 72.9 mm, Departamento Colonia, arroyo Limetas. *Crenicichla semifasciata*: Argentina. MACN-ict 3683, 1 ex., 68.8 mm, Formosa, Riacho de Oro. MACN-ict 6239, 1 ex., 176.6 mm, Entre Ríos, arroyo Curupí. *Crenicichla ocellata*: Paraguay. MSNG 33700 (holotype), 257.5 mm, Puerto 14 de Mayo, Bahía Negra, Chaco Boreal. *Crenicichla yaha*: Argentina. Misiones Province. MACN-ict 8924 (holotype), 103.7 mm, arroyo Urugua-í in Isla Palacios. AI 199, 1 ex., 116.6 mm, río Iguazú basin, arroyo Benavente. MTD-F 30606 (paratype), 1 ex., 105.9 mm, arroyo Urugua-í in ruta provincial 19, Parque Provincial Islas Malvinas. AI 200 (paratype), 1 ex., 135.8 mm SL, arroyo Urizú (affluent of A. Urugua-í) in ruta provincial 19, Parque Provincial Islas Malvinas. AI 202 (paratypes), 4 ex., 1 (C&S) 37.4-48.5 mm, arroyo Urugua-í in Isla Palacios.

RESULTS

Crenicichla tesuy sp. n.

Figs 1-4, Table 1

HOLOTYPE: MACN-ict 9016, 115.1 mm, Argentina, Misiones, río Iguazu basin, arroyo Verde (25°40'15.0"S-53°56'00.8"W), coll: F. Nunez, February 2002.

PARATYPES: All from Argentina. MACN-ict 9017, 1 ex., 124.2 mm, Misiones, río Iguazu basin, arroyo Deseado (25°47'08.1"S-54°01'45.0"W), coll: F. Nunez, February 2002. MACN-ict



FIG. 1

Crenicichla tesay sp. n., holotype: MACN-Ict 9016, 115.1 mm SL, Argentina, Misiones, río Iguazu basin, arroyo Verde.

9018, 5 ex., 55.7-252.0 mm. Misiones, río Iguazú basin, arroyo Tateto (25°47'12.8"S-53°58'12.9"W), coll: F. Nunez, February 2001. MHNG, 2708.062. 3 ex., 86.9-119.5 mm, same data as holotype. AI 213, 1 ex., 156.7 mm, Misiones, río Iguazu basin, arroyo Deseado Chico (in road 101) (25°47'19.7"S-54°01'45.0"W), coll: F. Nunez, February 2002. AI 214, 4 ex., 75.8-116.6 mm, Misiones, río Iguazu basin, arroyo Tateto (25°47'12.8"S-53°58'12.9"W), F. Nuñez, February 2001.

NON-TYPES SPECIMENS: Argentina, Misiones Province. AI 215, 1 ex., C&S 69.0 mm, río Iguazu basin, arroyo Tateto (25°47'12.8"S-53°58'12.9"W), coll: F. Nunez, February 2001. MLP 6-III-49-6-7, 2 ex., 97.0-133.0 mm, alto río Iguazu, PNI unreg., 1 ex., 103.2 mm, Parque Nacional Iguazu, arroyo Nandú. PNI unreg., 1 ex., 267.7 mm, Misiones, arroyo Ibicui.

DIAGNOSIS: *Crenicichla tesay* is distinguished from the other species of the genus by the following combination of characters: posterior border of preopercle serrated, 48-59 scales on E1 row, snout length 2.6-2.8 times in head length, presence of conspicuous suborbital stripe, flanks with 4 to 6 blotches below the upper lateral line, posteriormost blotch extending on caudal peduncle. Colour pattern with numerous irregularly scattered dots and absence of vertical stripes on flanks.

DESCRIPTION: Body elongate, depth four to five times in SL (Fig. 1). Head slightly deeper than wider. Snout short, bluntly pointed in lateral view. Jaws isognathous or lower jaw slightly prognathous. Maxilla almost reaching anterior margin of orbit. Lower lip folds widely separated anteriorly. Nostrils dorsolateral, nearer anterior margin of orbit than snout tip. Posterior margin of preopercle serrated (smooth in one specimen). Scales on flank strongly ctenoid. Head scales cycloid. Predorsal scales small, superficially embedded in skin. Prepelvic scales smaller than predorsal ones. Cheek scaled, 6 to 7 scales below eye embedded in skin. Scales in E1 row 48(1), 52(1), 53(3), 55(4*), 56(1), 57(2), 58(1), 59(2). Scales in transverse row 11/14(1), 11/16(3), 12/14(1*), 12/15(2), 12/16(4), 12/17(2), 12/18(1). Three scale rows between lateral lines. Upper lateral line scales slightly larger than the adjacent scales 22(1), 23(1), 24(6*), 25(3), 26(2), 27(1), 28(1). Lower lateral line scales equally in size than adjacent ones 7(1), 10(1), 13(4*), 14(2), 15(5), 16(2). Dorsal, anal, pectoral and pelvic fins naked. Dorsal fin XX1.11(3*), XX1.12(2); XXII.9(1); XXII.11(4); XXII.12(5). Anal fin III.8(4*); III.9(11). Pectoral fin 17(9), 18(6*). Caudal fin squamation




FIG. 2

Crenicichla tesay sp. n., lower pharyngeal tooth plate in occlusal view, AI 215, 69.0 mm SL. Scale bar: 1 mm.

extending almost to middle of fin. Soft-dorsal fin rounded or pointed tip, reaching or scarcely surpassing the caudal-fin base. Soft-anal fin not reaching the caudal-fin base. Caudal fin rounded. Pectoral fin rounded, reaching 75% of pelvic-fin length. Microbranchiospines present on second trough fourth gill arches. Nine gill rakers on ceratobranchial. Four or six patches of unicuspidate teeth on fourth ceratobranchial. Lower pharyngeal tooth plate with unicuspidate recurved and bicuspidate crenulated curved anteriorly teeth, those of posterior row much larger than the remaining ones (Fig. 2). Upper pharyngeal tooth plate with unicuspidate and bicuspidate teeth. Frashed zone bearing one concavity with small unicuspidate teeth. Premaxillary ascending process longer than dentigerous one. Premaxilla with 28 unicuspidate teeth on outer row, larger than the inner ones. Five teeth rows near symphysis. Dentary with 30 unicuspidate teeth on outer row, 4 rows near symphysis. Total vertebrae: 38 (1 C&S ex.). Premaxillary and dentary outer row teeth slightly movable or fixed, inner ones fully depressible.




FIG. 3

Geographical distribution of *Crenicichla tesay* sp. n.: 1. arroyo Verde (type locality), 2. arroyo Deseado; 3. arroyo Deseado Chico; 4. arroyo Taletó; 5. arroyo Ñandú; 6. arroyo Ibicuí.

COLOUR IN ALCOHOL: There is no obvious sexual dichromatism. Lateral line scales lighter than the adjacent ones. Grey preorbital stripe between anterior margin of orbit to snout tip, more conspicuous in smaller specimens (55.7-81.4 mm SL). Wide and greyish, postorbital stripe between posterior margin of orbit to preopercle distal margin, only one specimen with postorbital stripe reaching tip of opercle (75.8 mm SL). Suborbital stripe black, reaching almost the ventral margin of cheek, proximally entire, distally fragmented. Nuchal markings faint in adults, more conspicuous in smaller specimens. Flank with numerous irregularly scattered dark dots; few dots on cheek and opercular region in four specimens. Four to six dark rectangular or sub-circular blotches just below the upper lateral line. Posteriormost blotch extending onto caudal peduncle, except in one specimen. Dorsal, anal, and caudal fins smoky, with numerous dark scattered dots on their surface. Caudal fin with a black subcircular spot well separated from the base of the fin, just above of midline of caudal fin. That spot bears a narrow light ring. Pectoral fin smoky, pelvic fin whitish.

ETYMOLOGY: The specific epithet *tesay*, a noun in apposition, is a Guaraní word that means tears in allusion to the shape of suborbital stripe.

DISTRIBUTION: *Crenicichla tesay* inhabits streams tributaries of the río Iguazu above Iguazu falls in Argentina (Fig. 3).

HABITAT: The depth of the streams was variable, averaging about 80 cm. The bottom was composed of mud, sand and mostly stones, and the stream has falls and pools, with clear, rapidly flowing water. Some areas had scarce submerged vegetation (Fig. 4).

COMPARISON OF SPECIES: Thirteen species of *Crenicichla* have been recorded from freshwater environments in Argentina (Casciotta, 1987; Casciotta *et al.*, 2007; Lucena, 2007; Lucena & Kullander, 1992): *C. celidochilus* Casciotta, 1987; *C. gaucho* Lucena & Kullander, 1992; *C. hadrostigma* Lucena, 2007; *C. jupiaensis* Britski & Luengo, 1968; *C. lepidota* Heckel, 1840; *C. minuano* Lucena & Kullander, 1992; *C. missioneira* Lucena & Kullander, 1992; *C. niederleini* (Holmberg, 1891); *C. scottii* (Eigenmann, 1907); *C. semifasciata* (Heckel, 1840); *C. tendybaguassu* Lucena & Kullander, 1992; *C. vittata* Heckel, 1840; and *Crenicichla yaha* Casciotta *et al.*, 2006.

Crenicichla tesay differs from *C. celidochilus*, *C. tendybaguassu*, *C. minuano*, and *C. missioneira* by having preopercle serrated and a suborbital stripe vs. preopercle smooth and suborbital stripe absent or reduce. *Crenicichla tesay* can be distinguished from *C. vittata* by EI row scales counts, 48-59 vs. 78-85.

The flanks of the new species *C. tesay* have numerous small dots and rectangular or subcircular blotches just below the upper lateral line, a combination of features not present in *C. jupiaensis*, *C. lepidota*, *C. niederleini*, or *C. scottii*. Besides, *C. lepidota* has humeral spot, *C. scottii* has flanks with several regular parallel rows of small dark spots, *C. niederleini* bears narrow vertical bars and a lateral band on the flank, and *C. jupiaensis* bears 14 to 17 narrow vertical bars; all these characters are absent in *C. tesay*.

Crenicichla tesay is easily distinguished from *C. semifasciata* by having half of the caudal fin scaled and the ascending premaxillary process longer than the dentigerous one, vs. an almost completely scaled caudal fin and shorter ascending process. It can be distinguished from *C. gaucho* by the postorbital stripe not reaching the tip of opercle and by the absence of a lateral band (Lucena & Kullander, 1992).

Crenicichla tesay is easily differentiated from *C. hadrostigma* because the latter species bears a conspicuous ocellated posttemporal spot larger than half orbital diameter, which is not found in any other *Crenicichla* species.

Lucena (2007) described *C. empheres* from the río Uruguay drainage in Brasil. The colour pattern of this species resemble that of *C. tesay*, however the new species can be distinguished from *C. empheres* by having flanks with four to six blotches vs. six to eight, and a well developed suborbital stripe that almost reaches the ventral margin of the cheek vs. either reduced to a spot at margin of orbit or two to five small dots below orbits.

The species *C. lacustris* (Castelnau, 1855), *C. maculata* Kullander & Lucena, 2006, and *C. punctata* Hensel, 1870 from the Atlantic coastal rivers of southeastern Brazil share with *C. tesay* the presence of numerous scattered dots on the flanks.



FIG. 4. Habitat of *Crenicichla tesay* sp. n., arroyo Verde, type locality.

TABLE 1. Morphometry of the holotype and 14 paratypes of *Crenicichla tesay* sp. n. expressed as percentage of SL. SD: standard deviation.

	Holotype	Range	Mean	SD
Standard length (mm)	115.1	55.8-252.7		
Head length	32.0	31.9-36.3	33.1	1.29
Snout length	13.2	11.5-14.3	12.3	0.77
Head depth	17.5	14.0-17.9	16.0	1.06
Body depth	24.4	19.2-24.5	21.5	1.52
Orbital diameter	7.2	4.8-8.4	7.2	1.03
Interorbital width	8.5	5.8-9.3	7.2	0.92
Pectoral fin length	22.9	16.1-22.9	20.5	1.83
Caudal peduncle depth	10.6	9.4-11.6	10.2	0.63
Caudal peduncle length	16.2	15.3-16.9	16.2	0.55

However *C. tesay* differs from those species in having a lower EI row scales counts. *Crenicichla tesay* can be distinguished from *C. lacustris* by the absence of a continuous lateral band, dark dots on head and EI row scales counts (48-59 vs. 57-75 [Lucena & Kullander, 1992]). *Crenicichla maculata* has 58-75 scales in the longitudinal series whereas *C. tesay* has 48-59 (Lucena & Kullander, 1992). *Crenicichla punctata* has numerous small dots on the head and 56-70 scales in EI row (Lucena & Kullander, 1992) whereas most specimens of *C. tesay* lack dots on the head (4 ex. with few of them on cheek and opercular region) and 48-59 scales in the EI row.

Three species have been recorded from the upper rio Paraná in Brazil and were not present in the portion of this river that traverse Argentina: *C. britskii* Kullander, 1982; *C. haroldoi* Luengo & Britski, 1974; and *C. jaguarensis* Haseman, 1911. *Crenicichla britskii* has a humeral spot which is absent in *C. tesay*; *C. haroldoi* has brown dots on each lateral line scale which are absent in *C. tesay*. Finally, *C. jaguarensis* has 6 to 7 gill-rakers on the lower anterior arch and eleven vertical bars whereas *C. tesay* has nine gill-rakers and lacks such bars.

Currently only two *Crenicichla* species have been described from the río Iguazú basin: *C. iguassuensis* Haseman, 1911 from upper portion of this river and *Crenicichla yaha* Casciotta *et al.*, 2006, from the arroyo Benavente, just above the Iguazú falls. *Crenicichla tesay* differs from *C. iguassuensis* by having longer snout (2.6-2.8 vs. 3.0-3.5 in head length), lower number of EI row (48-59 vs. 54-64), and having rectangular or subcircular blotches on the flank not extended onto its dorsum. Finally, *C. tesay* can be distinguished from *C. yaha* by having isognathous jaws or slightly prognathous lower jaws vs. isognathous or slightly prognathous upper jaw; head depth 14.0-17.9 vs. 17.9-20.8 % of SL; and (numerous scattered) dots on the flank which are absent in both sexes of *C. yaha*.

ACKNOWLEDGEMENTS

This paper benefited from comments and suggestions of Carlos A. S. de Lucena (PUCRS). We would like to express our gratitude to Carlos Tremouilles (UNLP) for help with some figures. Francisco Firpo helped us with the figure of lower pharyngeal jaws. Simon Reece revised the English version.

REFERENCES

- AGOSTINHO, A. A., THOMAZ, S. M., GOMES L. C. 2005. Conservation of the biodiversity of Brazil's inland waters. *Conservation Biology* 19: 646-652.
- CASCIOTTA, J. R. 1987. *Crenicichla celidochilus* N. Sp. from Uruguay and a multivariate analysis of the *lacustris* group (Perciformes: Cichlidae). *Copeia* 1987: 883-891.
- CASCIOTTA, J. R., ALMIRÓN, A. E. & GÓMEZ, S. E. 2006a. A new species of *Australoheros* (Teleostei: Perciformes: Cichlidae) from the río Iguazu basin, Argentina. *Zoologische Abhandlungen* 55: 77-83.
- CASCIOTTA, J. R., ALMIRÓN, A. E. & GÓMEZ, S. E. 2006b. *Crenicichla yaha* sp. n. (Perciformes: Labroidae: Cichlidae), a new species from the río Iguazu and arroyo Uruguay-í basins, northeastern Argentina. *Zoologische Abhandlungen* 56: 107-112.
- CASCIOTTA, J., ALMIRÓN, A., BECHARA, J., RUIZ DIAZ, F., SANCHEZ, S. & GONZÁLEZ, A. 2007. First record of *Crenicichla jupiaensis* Britski & Luengo, 1968 (Perciformes: Cichlidae) in freshwaters of Argentina. *Ichthyological contributions of Peces Criollos* 4: 1-4.
- CASCIOTTA, J. R. & ARRATIA, G. 1993. Jaws and teeth of American Cichlids (Pisces: Labroidae). *Journal of Morphology* 217: 1-36.
- GARAVELLO, J. C. 2005. Revision of genus *Stemdashneridion* (Siluriformes: Pimelodidae). *Neotropical Ichthyology* 3: 607-623.
- GARAVELLO, J. C. & SHIBATA, O. A. 2007. A new species of the genus *Pimelodus* La Cèpede, 1803 from the rio Iguaçú basin and a reappraisal of *Pimelodus ortmanni* Haseman, 1911 from the rio Paraná system, Brazil (Ostariophysi: Siluriformes: Pimelodidae). *Neotropical Ichthyology* 5: 285-292.
- GODOY, M. P. DE. 1979. Rio Iguaçú, Paraná, Brasil. Reconhecimento da ictiofauna, modificações ambientais e usos múltiplos dos reservatórios. Electrosul, Rio de Janeiro, 33 p.
- HASEMAN, J. D. 1911. An annotated catalog of the cichlid fishes collected by the expedition of the Carnegie Museum to central South America, 1907-10. *Annals of the Carnegie Museum* 7: 329-373.
- KULLANDER, S. O. 1986. Cichlid fishes of the Amazon River drainage of Peru. Swedish Museum of Natural History, Stockholm, 431 pp.
- KULLANDER, S. O. & LUCENA, C. A. S. 2006. A review of the species of *Crenicichla* (Teleostei: Cichlidae) from the Atlantic coastal rivers of southeastern Brazil from Bahia to Rio Grande do Sul State, with description of three new species. *Neotropical Ichthyology* 4: 127-146.
- LEVITON, A. E., GIBBS, Jr., R. H., HEAL, E. & DAWSON, C. E. 1985. Standards in herpetology and ichthyology: Part I. Standard symbolic codes for institutional resource collections in herpetology and ichthyology. *Copeia* 1985: 802-832.
- LUCENA, C. A. S. 2007. Two new species of the genus *Crenicichla* Heckel, 1840 from the upper rio Uruguay drainage (Perciformes: Cichlidae). *Neotropical Ichthyology* 5: 449-456.
- LUCENA, C. A. S. & KULLANDER, S. O. 1992. The *Crenicichla* (Teleostei: Cichlidae) species of the Uruguay River Drainage in Brazil. *Ichthyological Exploration of Freshwaters* 3: 97-160.
- MIZOGUCHI, S. M. H. K., PORTELA-CASTRO, A. L. B. & MARTINS-SANTOS, I. C. 2007. Cytogenetic characterization of *Crenicichla* (Pisces, Perciformes, Cichlidae) of the Iguaçú River. *Genetics and Molecular Research* 6: 650-656.
- RENESTO, E., ZAWADZKI, C. H. & REVALDAVES, E. 2001. Biochemical taxonomy of *Crenicichla* (Pisces: Perciformes: Cichlidae) of the Iguaçú River, Brazil. *Brazilian Archives of Biology and Technology* 44: 15-22.
- SEVERI, W. & CORDEIRO, A. A. M. 1994. Catálogo de peixes da bacia do rio Iguaçú. IAP/GTZ, Curitiba, 128 pp.
- TAYLOR, W. R. & VAN DYKE, G. C. 1985. Revised procedures for staining and clearing small fishes and other vertebrates for bone and cartilage study. *Cybium* 9: 107-119.

