

ON LOCAL DEFINABILITY OF HOLOMORPHIC FUNCTIONS

GARETH JONES, JONATHAN KIRBY, OLIVIER LE GAL, AND TAMARA SERVI

ABSTRACT. Given a collection \mathcal{A} of holomorphic functions, we consider how to describe all the holomorphic functions locally definable from \mathcal{A} . The notion of local definability of holomorphic functions was introduced by Wilkie, who gave a complete description of all functions locally definable from \mathcal{A} in the neighbourhood of a generic point. We prove that this description is no longer complete in the neighbourhood of non-generic points. More precisely, we produce three examples of holomorphic functions which suggest that at least three new operations need to be added to Wilkie's description in order to capture local definability in its entirety. The constructions illustrate the interaction between resolution of singularities and definability in the o-minimal setting.

March 25, 2019

Acknowledgement. This work was partially supported by the Engineering and Physical Sciences Research Council [EP/J01933X/1, EP/N007956/1] to G. J. and [EP/L006375/1] to J. K.; and l'Agence Nationale de la Recherche [ANR-11-BS01-0009] to O. L. G. and [ANR-13-BS01-0006] to T. S..

1. INTRODUCTION

There has long been interaction between the theory of first-order definability and functional transcendence. For instance [Bia97, JKS16, Sfo12, LGR09, LG10, Ran14] use various results on algebraic independence of certain functions to establish results on definability and nondefinability in o-minimal expansions of the real field. The nondefinability results give in turn natural strengthenings of statements from functional transcendence. For instance, van den Dries, Macintyre and Marker, showed, among many other things, that $\int \exp(-x^2) dx$ is not definable in $\mathbb{R}_{\text{an, exp}}$ (Theorem 5.11 in [vdDMM97]). This gives a much stronger form of Liouville's result that this antiderivative is not an elementary function ([Lio33]).

In this spirit, we investigate the holomorphic functions which are locally definable from a given collection \mathcal{A} of holomorphic functions. Roughly speaking, these are the functions whose graph (seen as a subset of \mathbb{R}^{2N} , for a suitable N) can be described, locally, by a first order formula involving finite sums, products and compositions of (the real and imaginary parts of) functions in \mathcal{A} (see Definition 2.1). It is natural, both from the logical point of view and also from the perspective of analytic geometry, to seek a complex analytic characterization of the holomorphic functions locally definable from \mathcal{A} .

Wilkie gave such a characterization around *generic* points in [Wil08]. He showed that if f is a locally definable holomorphic function then around a generic point f is contained in the smallest collection of holomorphic functions containing both the functions in \mathcal{A} and all polynomials, and closed under partial differentiation, Schwarz reflection, composition and extraction of implicit functions. Wilkie conjectured that the same result would hold even without the genericity hypothesis. The aim of this paper is to show that in fact this description is no longer complete around non-generic points. We give three examples of functions which between them suggest that at least three further operations need to be added to those considered by Wilkie in order to capture local definability. We construct functions f_1, f_2, f_3 and g_1, g_2, g_3 with the germ of f_i definable from g_i , but not obtainable from g_i by the operations above. To obtain f_i from g_i requires further operations: f_1 needs monomial division, f_2 needs deramification and f_3 needs blowing-down (Theorems A, B and C, respectively, in Section 2).

2000 *Mathematics Subject Classification.* primary: 03C64; secondary: 14P10.

Key words and phrases. o-minimal structures; first-order definability;

These three new operations arise naturally in resolution of singularities [BM88], which is an essential tool from analytic geometry used in o-minimality (see for example [RSW03, RS15]). This is consistent with Wilkie’s observation that his conjecture is related to resolution of singularities. It is natural to ask whether the operations introduced so far would suffice to give a complete description of the holomorphic functions locally definable from \mathcal{A} (see Question 2.9). Indeed exactly these operations appear in the piecewise description of definable functions given in [RS15]. And based on the latter, it will be shown in the forthcoming [LGSVB17] that, if \mathcal{A} is a collection of *real* analytic functions, then the real analytic functions which are locally definable from \mathcal{A} can be obtained from \mathcal{A} and polynomials by the same list of operations (except Schwarz reflection). Since in the description given in [RS15], the operations involved do not respect any underlying complex structure (for example, exceptional divisors of blow-downs generally have real co-dimension 1), we cannot directly deduce an answer to Question 2.9 from an answer to its real counterpart.

It is also natural to ask if these three new operations are independent, and indeed what the dependencies between them and the other operations are. We introduce the new operations in the order monomial division, deramification, and blowing-down, and our proofs show that each one cannot be obtained from the previous operations. We do expect that none of these three is obtainable from all the other operations, but we do not prove it. However, one can define differentiation using monomial division and the other operations, for example, so giving a complete description of dependencies between all the operations is not a completely trivial matter.

We now briefly discuss the proofs of Theorems A, B and C.

For Theorem A, we use Ax’s functional version of Schanuel’s conjecture to prove that the function $(e^z - 1)/z$ cannot be obtained from e^z without monomial division. There is a connection here with previous work on nondefinability, for example [Bia97, JKS16]. However, these papers concerned arbitrary first-order definability, reduced to existential definability via model completeness. Here, we consider a restricted form of definability characterized by analytic rather than logical considerations.

The proofs of Theorems B and C are less explicit. For Theorem B, we first observe that deramification is independent from the operations used by Wilkie and monomial division (see Proposition 5.2). It is however not easy to witness this independence with a natural function. So instead we build on some of the ideas contained in [LG10]. In particular, we adapt the notion of strongly transcendental function considered there to the setting of several complex variables. A related notion of strongly transcendental function was also independently introduced by Boris Zilber [Zil02] under the name of ‘generic functions with derivatives’, in connection with finding analogues of Schanuel’s conjecture for other complex functions. Our present work then proves the existence of such functions, which was not considered by Zilber.

For Theorem C, the independence of blow-downs from the other operations simply follows from the fact that, unlike the other operators considered here, blow-downs are not local operators: they are applied to all the germs of a function along the exceptional divisor of the corresponding blow-up. As in the case of theorem B, it is not clear how to find an explicit function which witnesses this independence, and we again make use of strongly transcendental functions.

The paper is organized as follows. In Section 2 we introduce some operators and algebras which allow us to give the precise formulations of our results. In Section 3 we define a notion of strong transcendence for holomorphic functions in \mathbb{C}^2 , and prove that the notion is not void and implies an independence property which is the key point for constructing our last two examples. Finally, in Sections 4, 5 and 6 we prove Theorems A, B and C, respectively.

2. MAIN DEFINITIONS AND RESULTS

In this section we introduce some notation, and the various operators and functions we consider. We then give precise statements of our results.

Throughout this paper, we will use the word *definable* in the sense of first-order logic. Unless otherwise specified, sets and functions definable in a given first-order structure will be understood to

be definable with parameters from the domain of the structure. We will use the term \emptyset -definable to denote sets and functions definable without parameters.

Following Wilkie in [Wil08], we say that the restriction $g \upharpoonright \Delta$ of a holomorphic map $g : U \rightarrow \mathbb{C}$ defined on an open set $U \subseteq \mathbb{C}^n$ is a **proper restriction** if $\Delta \subset U$ is an open box, relatively compact in U . Given a family \mathcal{A} of holomorphic functions defined on open subsets of \mathbb{C}^n (for various $n \in \mathbb{N}$), we denote by $\mathcal{A} \upharpoonright$ the collection of all proper restrictions to boxes with corners in the Gaussian rationals, $\mathbb{Q}(i)$, of all functions in \mathcal{A} . We let $\mathbb{R}_{\mathcal{A} \upharpoonright} = \langle \mathbb{R}; 0, 1, +, \cdot, <, \mathcal{A} \upharpoonright \rangle$ be the expansion of the real ordered field by the functions in $\mathcal{A} \upharpoonright$, seen as functions from some (even) power of \mathbb{R} to \mathbb{R}^2 , by identifying \mathbb{C} with \mathbb{R}^2 .

Definition 2.1. A function $g : U \subset \mathbb{C}^n \rightarrow \mathbb{C}$ is **locally definable** in $\mathbb{R}_{\mathcal{A} \upharpoonright}$ if all its proper restrictions, interpreted as real functions, are definable in $\mathbb{R}_{\mathcal{A} \upharpoonright}$. If all the proper restrictions of g to boxes with Gaussian rational corners are \emptyset -definable in \mathcal{R} , we say that g is **locally \emptyset -definable**.

The symbol \mathcal{A} will always denote a collection of holomorphic functions defined on open subsets of \mathbb{C}^n , for various $n \in \mathbb{N}$. More precisely, \mathcal{A} is a family $\{\mathcal{A}_n(U) : n \in \mathbb{N}, U \subseteq \mathbb{C}^n \text{ open}\}$, where $\mathcal{A}_n(U)$ is a collection of holomorphic functions defined on U . It is convenient to suppose that \mathcal{A} is closed under gluing and restrictions, which can always be done, by enriching \mathcal{A} . More precisely, we suppose that, if $(U_\lambda)_{\lambda \in \Lambda}$ is a collection of open subsets of \mathbb{C}^n , then

$$f \in \mathcal{A}_n \left(\bigcup_{\lambda \in \Lambda} U_\lambda \right) \Leftrightarrow \forall \lambda \in \Lambda, f \upharpoonright U_\lambda \in \mathcal{A}_n(U_\lambda).$$

Thus \mathcal{A}_n can be viewed as a sheaf, for each fixed dimension n , whose stalk at $z \in \mathbb{C}^n$, denoted by \mathcal{A}_z , is the collection of all germs at z of the functions which belong to some $\mathcal{A}_n(U)$ with $z \in U$. Note that closing \mathcal{A} under gluing and restrictions does not affect local (\emptyset -)definability in $\mathbb{R}_{\mathcal{A} \upharpoonright}$. By a classical argument of local compactness, writing f_z for the germ of f at z , we have $f \in \mathcal{A}(U)$ if and only if for all $z \in U$, $f_z \in \mathcal{A}_z$.

By abuse of terminology, we will say that \mathcal{A} is a sheaf, and write $\mathcal{A} \subseteq \mathcal{B}$ as a shorthand for $\forall n \in \mathbb{N}, \forall U \subseteq \mathbb{C}^n, \mathcal{A}_n(U) \subseteq \mathcal{B}_n(U)$. In the same way, if \mathcal{A}_n is the sheafification of a family of functions, we will often drop the index n . For instance, we will write $\mathcal{A} = \{\text{exp}\}$ as a shorthand for

$$\forall U \subseteq \mathbb{C}^n, \mathcal{A}_n(U) = \begin{cases} \emptyset & \text{if } n \neq 1 \\ \{\text{exp} \upharpoonright U\} & \text{if } n = 1 \end{cases}.$$

We fix a collection \mathcal{A} of holomorphic functions as in the discussion above. We now construct new sheaves $\mathcal{B}, \mathcal{C}, \mathcal{D}, \mathcal{E}, \mathcal{F}$ by closing \mathcal{A} under certain operations. The operations involved in the definition of $\mathcal{B}, \mathcal{C}, \mathcal{D}$ are local, hence we define them by their action on germs. We denote by $\mathcal{O}_n(U)$ the collection of all holomorphic functions defined on $U \subseteq \mathbb{C}^n$ and by \mathcal{O}_a the collection of all holomorphic germs at $a \in \mathbb{C}^n$. We let \mathcal{P} be the collection of all complex polynomials (in any number of variables) and \mathcal{P}_G be the subcollection of all polynomials with Gaussian rational coefficients. We also use the following notation: $z' = (z_1, \dots, z_{n-1})$ and $z = (z', z_n)$. If $a \in \mathbb{C}^n$ then we write $a = (a', a_n) \in \mathbb{C}^{n-1} \times \mathbb{C}$.

Definition 2.2. We define the following operators:

- (1) **Polynomial** and **Gaussian polynomial** operators. The polynomial operators are operators of arity 0, defined for each $P \in \mathcal{P}$ and $a \in \mathbb{C}^n$ which map the empty set to the germ $P_a \in \mathcal{O}_a$. The Gaussian polynomial operators are similar, but restricted to polynomials $P \in \mathcal{P}_G$.
- (2) The **Schwarz reflection** operator at $a \in \mathbb{C}^n$ maps a germ $f \in \mathcal{O}_a$ to the germ $z \mapsto f(\bar{z}) \in \mathcal{O}_{\bar{a}}$.
- (3) **Composition** operators: if $a \in \mathbb{C}^n$ and $b \in \mathbb{C}^m$, then we consider the operator that maps $(f, g_1, \dots, g_n) \in \mathcal{O}_a \times (\mathcal{O}_b)^n$ to the germ $f \circ (g_1, \dots, g_n) \in \mathcal{O}_b$, whenever $(g_1(b), \dots, g_n(b)) = a$.
- (4) **Partial derivative** operators: for $a \in \mathbb{C}^n$ and $j \in \{1, \dots, n\}$, the j -th partial derivative is the operator that maps $f \in \mathcal{O}_a$ to the germ $\frac{\partial f}{\partial z_j} \in \mathcal{O}_a$.
- (5) The **implicit function** operator at $a \in \mathbb{C}^n$ maps a germ $f \in \mathcal{O}_a$ satisfying $f(a) = 0$ and $\frac{\partial f}{\partial z_n}(a) \neq 0$ to the (unique) germ $\varphi \in \mathcal{O}_{a'}$ satisfying $f(z', \varphi(z')) \equiv 0$ and $\varphi(a') = a_n$.

- (6) **Monomial division** operators: if $a \in \mathbb{C}^n$, then we consider the operator that maps $f \in \mathcal{O}_a$ to the germ at a of the extension by continuity of $z \mapsto \frac{f(z)}{(z_n - a_n)}$, whenever the germ $z' \mapsto f(z', a_n) \in \mathcal{O}_{a'}$ is identically zero.
- (7) **Deramification**: if $a \in \mathbb{C}^n$ and $m \in \mathbb{N}^\times$, then we consider the operator which maps $f \in \mathcal{O}_a$ to the germ $z \mapsto f(z', a_n + \sqrt[m]{z_n - a_n}) \in \mathcal{O}_a$, whenever the germ f satisfies

$$f(z) = f\left(z', a_n + e^{\frac{2i\pi}{m}}(z_n - a_n)\right).$$

Here the condition on f implies that this expression does not depend on the choice of an m^{th} -root for $z_n - a_n$.

Notice that elementary operators can be composed as long as their arities match.

Definition 2.3. We define \mathcal{B}^* to be the set of all operators which can be expressed as finite compositions of polynomial, Schwarz reflection, composition, partial derivative and implicit function operators. The collection $\mathcal{B}^{\theta*}$ is defined similarly, with Gaussian polynomial operators in place of polynomial operators. Analogously, we let \mathcal{C}^* and $\mathcal{C}^{\theta*}$ be the sets of all operators which can be expressed as finite compositions of monomial division operators and operators in \mathcal{B}^* and $\mathcal{B}^{\theta*}$, respectively. And we let \mathcal{D}^* and $\mathcal{D}^{\theta*}$ be the sets of all operators which can be expressed as finite compositions of deramification operators and operators in \mathcal{C}^* $\mathcal{C}^{\theta*}$, respectively.

Once we have defined these operators, we can construct the set of all functions which are “locally obtained from \mathcal{A} ” by the action of such operators. For example, \mathcal{B} will denote the set of all functions which are locally obtained from $\mathcal{A} \cup \mathcal{P}$ by composition, Schwarz reflection, taking partial derivatives and extracting implicit functions.

More formally:

Definition 2.4. We let $\mathcal{B}, \mathcal{B}^\theta, \mathcal{C}, \mathcal{C}^\theta, \mathcal{D}$ and \mathcal{D}^θ be the smallest sheaves containing \mathcal{A} with stalks stable under the action of the operators in $\mathcal{B}^*, \mathcal{B}^{\theta*}, \mathcal{C}^*, \mathcal{C}^{\theta*}, \mathcal{D}^*$ and $\mathcal{D}^{\theta*}$, respectively.

Remark 2.5. If U is an open subset of \mathbb{C}^n and $f \in \mathcal{A}(U)$ satisfies $f(x', 0) = 0$, the germ at $(a', 0)$ of $g(x) = \frac{f(x', x_n)}{x_n}$ belongs to $\mathcal{C}_{(a', 0)}$ by construction. One might wonder if the germ of g at any $a \in U$ belongs to \mathcal{C}_a , to ensure $g \in \mathcal{C}(U)$. Actually, if $a_n \neq 0$, the germ g_a is the implicit function of $x_{n+1}x_n - f(x', x_n)$ at a , so $g_a \in \mathcal{B}_a$. A similar argument holds for functions obtained by taking an m^{th} -root. Along the same lines, if $\varphi : U \rightarrow \mathbb{C}$ is the implicit function of f , its germ at any $a \in U$ is obtained by applying the implicit function operator to f at $(a, \varphi(a))$. In particular, \mathcal{B}^θ coincides with the algebra denoted by $\tilde{\mathcal{F}}$ in [Wil08, Definition 1.7].

It remains to define the functions locally obtained by blow-downs. For this, we say that a sheaf $\mathcal{G} \subseteq \mathcal{O}$ is stable under **blow-downs** if for every $n \in \mathbb{N}$, every open subset $U \subseteq \mathbb{C}^n$ and every blow-up $\pi : V \rightarrow U$ of a smooth analytic manifold $X \subseteq U$, which is locally defined by a system of equations in \mathcal{G} , we have that if $f \circ \pi \in \mathcal{G}(V)$ then $f \in \mathcal{G}(U)$. A more precise (but heavier) definition could be given with no mention of $\mathcal{G}(V)$ (note that V is not an open subset of a power of \mathbb{C}), using local coordinates, but in this paper we will only mention the blow-up of $0 \in \mathbb{C}^2$ and introduce the corresponding local coordinates in Section 6.

Definition 2.6. Let \mathcal{E} and \mathcal{E}^θ be the smallest sheaves which are stable under blow-downs and under the action of the operators in \mathcal{D}^* and $\mathcal{D}^{\theta*}$, respectively.

Finally, we denote by \mathcal{F} and \mathcal{F}^θ the sheaves of all locally definable holomorphic functions and all locally \emptyset -definable holomorphic functions, respectively.

Thanks to the compactness of the fibres of blow-ups, if f is the blow-down of g and g is locally definable in $\mathbb{R}_{\mathcal{A}\uparrow}$, then f is also locally definable in $\mathbb{R}_{\mathcal{A}\uparrow}$.

By construction we have :

$$\begin{array}{ccccccccccc} \mathcal{A} & \subseteq & \mathcal{B}^\theta & \subseteq & \mathcal{C}^\theta & \subseteq & \mathcal{D}^\theta & \subseteq & \mathcal{E}^\theta & \subseteq & \mathcal{F}^\theta \\ & & \cap & & \cap & & \cap & & \cap & & \cap \\ & & \mathcal{B} & \subseteq & \mathcal{C} & \subseteq & \mathcal{D} & \subseteq & \mathcal{E} & \subseteq & \mathcal{F} & \subseteq & \mathcal{O}. \end{array} \quad (*)$$

We can now give a precise formulation of the results announced in the introduction.

Wilkie proposed the following conjecture.

Conjecture 2.7 ([Wil08, Conjecture 1.8]). *Let \mathcal{A} be a family of complex holomorphic functions. Let $z \in \mathbb{C}^n$, U be an open neighbourhood of z and $f : U \rightarrow \mathbb{C}$ be a holomorphic function which is locally \emptyset -definable in $\mathbb{R}_{\mathcal{A}\uparrow}$. Then there is an open box with Gaussian rational corners $\Delta \subseteq U$ such that $z \in \Delta$ and $f \upharpoonright \Delta$ can be obtained from $\mathcal{A} \cup \mathcal{P}_G$ by finitely many applications of composition, Schwarz reflection, taking partial derivatives and extracting implicit functions.*

Conjecture 2.7 can be restated as follows: $\mathcal{B}^0 = \mathcal{F}^0$. By adding all constant functions to \mathcal{A} , the conjecture implies also that $\mathcal{B} = \mathcal{F}$. The conjecture is however not true in general. We show that the six horizontal inclusions between \mathcal{B}^0 and \mathcal{E} in diagram (*) are strict in general:

Theorem A (Monomial division). *Suppose $\mathcal{A} = \{\exp\}$ is the complex exponential function. Then the function $f : z \mapsto (e^z - 1)/z$ (extended by continuity at $z = 0$) is holomorphic and locally \emptyset -definable in $\mathbb{R}_{\mathcal{A}\uparrow}$, but no restriction of f to any neighbourhood of 0 can be obtained from $\mathcal{A} \cup \mathcal{P}_G$ by finitely many applications of composition, Schwarz reflection, taking partial derivatives and extracting implicit functions.*

More precisely, the germ at 0 of f belongs to $\mathcal{C}_0^0 \setminus \mathcal{B}_0$. In particular, $\mathcal{B}^0 \neq \mathcal{C}^0$ and $\mathcal{B} \neq \mathcal{C}$.

Theorem B (Deramification). *There exists a holomorphic function g , with domain a neighbourhood of $0 \in \mathbb{C}$, such that the function $f : z \mapsto g(\sqrt{z})$ is well defined, holomorphic in a neighbourhood of 0 and locally \emptyset -definable in $\mathbb{R}_{\mathcal{A}\uparrow}$ (where $\mathcal{A} = \{g\}$), but no restriction of f to any neighbourhood of 0 can be obtained from $\mathcal{A} \cup \mathcal{P}_G$ by finitely many applications of composition, Schwarz reflection, monomial division, taking partial derivatives and extracting implicit functions.*

More precisely, the germ at 0 of f belongs to $\mathcal{D}_0^0 \setminus \mathcal{C}_0$. In particular, $\mathcal{C}^0 \neq \mathcal{D}^0$ and $\mathcal{C} \neq \mathcal{D}$.

Theorem C (Blow-down). *Let π be the blow-up of the origin in \mathbb{C}^2 . Then there exists a holomorphic function f , with domain a neighbourhood of $0 \in \mathbb{C}^2$, such that, if $\mathcal{A} = \{f \circ \pi\}$, then f is locally \emptyset -definable in $\mathbb{R}_{\mathcal{A}\uparrow}$, but no restriction of f to any neighbourhood of 0 can be obtained from $\mathcal{A} \cup \mathcal{P}_G$ by finitely many applications of composition, Schwarz reflection, monomial division, deramification, taking partial derivatives and extracting implicit functions.*

More precisely, the germ at 0 of f belongs to $\mathcal{E}_0^0 \setminus \mathcal{D}_0$. In particular, $\mathcal{D}^0 \neq \mathcal{E}^0$ and $\mathcal{D} \neq \mathcal{E}$.

Remark 2.8. Wilkie proves in [Wil08, Theorem 1.10] that Conjecture 2.7 holds for all points z which are *generic* with respect to a suitable pregeometry associated to \mathcal{A} . Since the points at which we apply the operators in $\mathcal{E}^* \setminus \mathcal{B}^*$ are not generic, the theorems above are consistent with Wilkie's result. In particular, if z is generic point, then we can restate Wilkie's result [Wil08, Theorem 1.10] in the following form: $\mathcal{B}_z^0 = \mathcal{C}_z^0 = \mathcal{D}_z^0 = \mathcal{E}_z^0 = \mathcal{F}_z^0$.

The purpose of [Wil08, Conjecture 1.8 and Theorem 1.10] was to describe the holomorphic functions locally definable from \mathcal{A} in terms of the functions in \mathcal{A} , using purely complex operations. Since our examples are obtained by means of three natural complex operations, we could consider a modification of conjecture 2.7. But we prefer to formulate this as a question.

Question 2.9. *Let \mathcal{A} be a family of complex holomorphic functions. Let $z \in \mathbb{C}^n$, U be an open neighbourhood of z and $f : U \rightarrow \mathbb{C}$ be a holomorphic function which is locally \emptyset -definable in $\mathbb{R}_{\mathcal{A}\uparrow}$. Is there an open box with Gaussian rational corners $\Delta \subseteq U$ such that $z \in \Delta$ and $f \upharpoonright \Delta$ can be obtained from $\mathcal{A} \cup \mathcal{P}_G$ by finitely many applications of composition, Schwarz reflection, monomial division, deramification, blow-downs, taking partial derivatives and extracting implicit functions?*

Question 2.9 can therefore be restated as follows: do \mathcal{E}^0 and \mathcal{F}^0 coincide?

We conclude this section with some considerations on the local operators defined above, which will be useful to prove our main results.

The operators in \mathcal{D}^* act on germs of holomorphic functions, hence they can also be seen as acting on the Taylor expansion of such germs (we do not make a distinction between an analytic germ and its Taylor expansion). Let us fix some notation.

Notation 2.10. We will often use the following notation for tuples: if $k \in \mathbb{N} \setminus \{0\}$, then $z = z^{(k)}$ means that z is a k -tuple of variables. Similarly, we write $a = a^{(k)}$ for a point in \mathbb{C}^k .

If $\alpha \in \mathbb{N}^k$ and $z = (z_1, \dots, z_k) \in \mathbb{C}^k$ then $z^\alpha = z_1^{\alpha_1} z_2^{\alpha_2} \dots z_k^{\alpha_k}$. We set $\alpha! = \alpha_1! \dots \alpha_k!$, $|\alpha| = \sum_{i=1}^k \alpha_i$ and denote by $\partial_\alpha f$ the partial derivative $\frac{\partial^{|\alpha|} f}{\partial z^\alpha}$ of a function $f : \mathbb{C}^k \rightarrow \mathbb{C}$. If $\alpha = 0$, then $\partial_\alpha f = f$.

Definition 2.11. If $U \subseteq \mathbb{C}^m$, $f \in \mathcal{O}(U)$ and $a = (a_1^{(m)}, \dots, a_n^{(m)}) \in U^n$, the **jet of order k of f at a** is the tuple $j_n^k f(a)$ of all partial derivatives $\partial_\alpha f$ for $|\alpha| \leq k$, evaluated at the points a_i , for $i = 1, \dots, n$:

$$j_n^k f(a) = (\partial_\alpha f(a_1^{(m)}), \dots, \partial_\alpha f(a_n^{(m)}))_{|\alpha| \leq k}.$$

We omit the subscript n if $n = 1$.

For given $a = (a_1, \dots, a_k) \in \mathbb{C}^{m_1} \times \dots \times \mathbb{C}^{m_k}$ and $N \in \mathbb{N}$, we denote by Z_a^N the set

$$Z_a^N = \{(h_1, \dots, h_k) \in \mathcal{O}_{a_1} \times \dots \times \mathcal{O}_{a_k} : \forall i = 1 \dots, k, j^N h_i(a_i) = 0\}.$$

These Z_a^N , when N varies, form a base neighborhood of 0 for a topology on $\mathcal{O}_{a_1} \times \dots \times \mathcal{O}_{a_k}$ called the Krull topology.

The action of our operators on Taylor expansions can be described as follows.

Proposition 2.12. *Let $k, n, m_1, \dots, m_k \in \mathbb{N}$ and $a_1 \in \mathbb{C}^{m_1}, \dots, a_k \in \mathbb{C}^{m_k}, b \in \mathbb{C}^n$. Let $\mathcal{L} : \mathcal{O}_{a_1} \times \dots \times \mathcal{O}_{a_k} \rightarrow \mathcal{O}_b$ be an operator in \mathcal{D}^* , and $(f_1, \dots, f_k) \in \mathcal{O}_{a_1} \times \dots \times \mathcal{O}_{a_k}$ be a k -tuple of germs in the domain of definition of \mathcal{L} . Then, there exist*

- a neighbourhood W of (f_1, \dots, f_k) for the Krull topology,
- a tuple of constants $c = (c_1, \dots, c_\ell) \in \mathbb{C}^\ell$ (for some $\ell \in \mathbb{N}$),
- for each $\alpha \in \mathbb{N}^n$, an integer $n_\alpha \in \mathbb{N}$ and a polynomial $P_\alpha \in \mathbb{Q}[y, \bar{y}]$, where $y = y^{(N_\alpha)}$ and N_α can be computed from $k, m_1, \dots, m_k, \ell, n_\alpha$,

such that, for each $(g_1, \dots, g_k) \in W$, $\mathcal{L}(g_1, \dots, g_k)$ has a Taylor expansion at b of the form

$$\mathcal{L}(g_1, \dots, g_k)(x) = \sum_{\alpha \in \mathbb{N}^n} P_\alpha(c_1, \dots, c_\ell, j^{n_\alpha} g_1(a_1), \dots, j^{n_\alpha} g_k(a_k))(x - b)^\alpha.$$

Proof. We first prove that the proposition is true if \mathcal{L} is an elementary operator.

For all but the implicit function operator, the neighbourhood W will be the whole space. Note that for all but the polynomial and the composition operators, we have $k = 1$ (hence in these cases we will write g instead of g_1).

- (1) If \mathcal{L} is a polynomial operator, then $n_\alpha = 0$ and the constants c_i can be computed from the coefficients of the polynomial and the coordinates of b .
- (2) If \mathcal{L} is the Schwarz reflection operator, then $n_\alpha = |\alpha|$ and $P_\alpha(j^{|\alpha|} g(a)) = \overline{(\alpha!)^{-1} \partial_\alpha g(a)}$.
- (3) If \mathcal{L} is a composition operator, then the proposition follows from the so called Faà Di Bruno formula (see for example [Arb00, p. 92. §115]), with $n_\alpha = |\alpha|$.
- (4) If \mathcal{L} is the partial derivative operator $\frac{\partial}{\partial z_j}$, then $n_\alpha = |\alpha| + 1$ and $P_\alpha(j^{n_\alpha} g(a)) = \frac{\alpha_j}{\alpha!} \frac{\partial}{\partial z_j} (\partial_\alpha g(a))$.
- (5) If \mathcal{L} is the implicit function operator, then set $c_1 = \left(\frac{\partial f}{\partial z_n}(a) \right)^{-1}$, $n_\alpha = |\alpha|$, and

$$W = \left\{ g : g(a) = 0, \frac{\partial g}{\partial z_n}(a) = \frac{\partial f}{\partial z_n}(a) \right\}.$$

It is easy to see that the coefficients of the Taylor expansion up to order $|\alpha|$ of $\mathcal{L}(g)$ at the point $b = (a_1, \dots, a_{n-1})$ can be expressed as polynomials (with coefficients in $\mathbb{Z}[c_1]$) in the derivatives up to order $|\alpha|$ of g at the point a . Hence, $n_\alpha = |\alpha|$ and P_α is a polynomial in the jet of order $|\alpha|$ of g and c_1 .

In order for P_α to be a polynomial and not a rational function, we need $\frac{\partial g}{\partial z_n}(a)$ to be constant and hence to restrict our claim to the neighbourhood W .

- (6) If \mathcal{L} is a monomial division operator, then $n_\alpha = |\alpha| + 1$ and

$$P_\alpha(j^{|\alpha|} g(a)) = ((\alpha_n + 1) \cdot \alpha!)^{-1} \frac{\partial}{\partial z_n} (\partial_\alpha g(a)).$$

(7) If \mathcal{L} is the m th-deramification operator, then $n_\alpha = m|\alpha|$; we leave it to the reader to find the expression for P_α in this case.

Each operator in \mathcal{D}^* is a composition of elementary operators. Suppose that the proposition holds for an operator \mathcal{L} . If $f \in \mathcal{O}_b$ is in the image of \mathcal{L} and $V \subseteq \mathcal{O}_b$ is an open neighbourhood of f in the Krull topology, then $\mathcal{L}^{-1}(V)$ is an open neighbourhood of $\mathcal{L}^{-1}(f)$. This observation and an easy computation show that the conclusion of the proposition is preserved under composition. We illustrate the proof for the composition of two operators $\mathcal{L} : \mathcal{O}_a \rightarrow \mathcal{O}_b$ and $\mathcal{M} : \mathcal{O}_b \rightarrow \mathcal{O}_d$, the general case following as a straightforward but tedious exercise. Suppose that

$$\begin{aligned}\mathcal{L}(f)(y) &= \sum_{\alpha} P_{\alpha}(c_{\mathcal{L}}, j^{n_{\alpha}} f(a))(y-b)^{\alpha}, \\ \mathcal{M}(g)(z) &= \sum_{\beta} Q_{\beta}(c_{\mathcal{M}}, j^{m_{\beta}} g(b))(z-d)^{\beta}.\end{aligned}$$

Then $j^{m_{\beta}} \mathcal{L}(f)(b) = (\partial_{\gamma}(\sum_{\alpha} P_{\alpha}(c_{\mathcal{L}}, j^{n_{\alpha}} f(a))(y-b)^{\alpha})(b))|_{|\gamma| \leq m_{\beta}} = (\gamma! P_{\gamma}(c_{\mathcal{L}}, j^{n_{\gamma}} f(a)))|_{|\gamma| \leq m_{\beta}}$. Hence we can write

$$\mathcal{M}(\mathcal{L}(f))(z) = \sum_{\beta} \tilde{Q}_{\beta}(c_{\mathcal{L}}, c_{\mathcal{M}}, j^{m_{\beta}} f(a))(z-d)^{\beta},$$

for some suitable polynomials \tilde{Q}_{β} . □

3. STRONG TRANSCENDENCE

To prove Theorems B and C, we will need to work with holomorphic functions which satisfy very few functional relations. For this reason, we extend to the complex and multi-dimensional setting the notion of *strong transcendence*, which was introduced in [LG10] in a real and one-dimensional context. Thanks to Proposition 3.4 below, the germs of a strongly transcendental holomorphic function satisfy very few relations. Proposition 3.2 shows that the definition is not empty: strongly transcendental holomorphic functions do exist.

We will use the notation introduced in 2.10 and 2.11.

Definition 3.1. A holomorphic function $f : U \subseteq \mathbb{C}^m \rightarrow \mathbb{C}$ is **strongly transcendental** if, for every $(k, n) \in \mathbb{N}^2$ and for every $z = (z_1^{(m)}, \dots, z_n^{(m)})$ an n -tuple of distinct points of U , we have

$$\text{trdeg}_{\mathbb{Q}} \mathbb{Q}(z, \bar{z}, j_n^k f(z), \overline{j_n^k f(z)}) \geq \text{length}(j_n^k f(z), \overline{j_n^k f(z)}).$$

We first show that such functions exist, then we prove an independence result for their germs at distinct points.

Proposition 3.2. *Let U be an open subset of \mathbb{C}^m . Then the set*

$$ST(U) = \{f \in \mathcal{O}(U); f \text{ is strongly transcendental}\}$$

is residual in $\mathcal{O}(U)$ with respect to the topology induced by uniform convergence on the compact subsets of U . In particular, by the Baire Category Theorem, $ST(U) \neq \emptyset$.

Note that strong transcendence is preserved under restriction but not under gluing, so ST is a presheaf but not a sheaf.

Notation 3.3. If $F : U \rightarrow V$ is a differentiable map between C^1 manifolds, the differential of F at $u \in U$ is denoted by $DF(u)$; it belongs to the space of linear maps $L(T_u U, T_{F(u)} V)$, where $T_u U$ denotes the tangent space of U at u .

Proof. The proof goes as follows. First we express $ST(U)$ as a countable intersection of subsets $B_{n,k,P}$ of $\mathcal{O}(U)$. Each $B_{n,k,P}$ is the complement of the image of a certain map π . We apply a version of the Sard-Smale Theorem for Fréchet manifolds, due to Eftekharinasab in [Eft11], to this π , and obtain that the set of regular values of π is residual. Moreover, we observe that π is everywhere critical, so

the set of its regular values is the complement of its image, then coincides with $B_{n,k,P}$. Hence, $ST(U)$ is a countable intersection of residual sets, therefore also a residual set.

Let us recall that a point $(p, q) \in (\mathbb{C}^{2mn} \times \mathbb{C}^{2N})$ has transcendence degree over \mathbb{Q} at least $2N$ if and only if (p, q) does not satisfy any nonsingular system of $2mn + 1$ polynomial equations over \mathbb{Z} . Hence, given an open subset U of \mathbb{C}^m , a function $f \in \mathcal{O}(U)$ belongs to $ST(U)$ if and only if

$$\begin{aligned} \forall n \in \mathbb{N}^*, \forall k \in \mathbb{N}, \forall P \in (\mathbb{Z}[y^{(2mn+2N)}])^{2mn+1}, \\ \forall z = (z_1^{(m)}, \dots, z_n^{(m)}) \in (\mathbb{C}^m)^n \setminus \Delta_n, \\ \text{if } \text{rk}(DP(z, \bar{z}, j_n^k f(z), \overline{j_n^k f(z)})) = 2mn + 1 \\ \text{then } P(z, \bar{z}, j_n^k f(z), \overline{j_n^k f(z)}) \neq 0, \end{aligned}$$

where $N = n \binom{m+k}{k}$ is the cardinality of the tuple $j_n^k f(z)$ and $\Delta_n = \{(z_1^{(m)}, \dots, z_n^{(m)}) \in (\mathbb{C}^m)^n : \exists i \neq j, z_i^{(m)} = z_j^{(m)}\}$.

In other words, $ST(U)$ can be described as the following countable intersection

$$ST(U) = \bigcap_{\substack{n \in \mathbb{N}^*, k \in \mathbb{N}, \\ P \in \mathbb{Z}[y^{(2mn+2N)}]}} B_{n,k,P},$$

where

$$B_{n,k,P} = \{f \in \mathcal{O}(U) : \forall z = (z_1^{(m)}, \dots, z_n^{(m)}) \in (\mathbb{C}^m)^n \setminus \Delta_n \\ \text{rk}(DP(z, \bar{z}, j_n^k f(z), \overline{j_n^k f(z)})) = 2mn + 1 \Rightarrow P(z, \bar{z}, j_n^k f(z), \overline{j_n^k f(z)}) \neq 0\}.$$

We fix n, k, P as above, define the set V :

$$V := \{(z, f) \in ((\mathbb{C}^m)^n \setminus \Delta_n) \times \mathcal{O}(U) : P(z, \bar{z}, j_n^k f(z), \overline{j_n^k f(z)}) = 0, \\ \text{rk}(DP(z, \bar{z}, j_n^k f(z), \overline{j_n^k f(z)})) = 2mn + 1\},$$

and equip V with the topology induced by the family of seminorms, indexed by the compact subsets K of U :

$$\|\cdot\|_K : (z, f) \mapsto \max \left\{ \|z\|, \sup_{x \in K} |f(x)| \right\}.$$

We can see $B_{n,k,P}$ as $\mathcal{O}(U) \setminus \pi(V)$, where $\pi : V \rightarrow \mathcal{O}(U)$ is the restriction to V of the projection $(\mathbb{C}^m)^n \times \mathcal{O}(U) \rightarrow \mathcal{O}(U)$ onto the second factor.

Since the equations which define V involve conjugations, V does not inherit a complex structure. However, V is a real Fréchet manifold, whose tangent space $T_{(z,f)}V$ at $(z, f) \in V$ is the real vector space given by

$$T_{(z,f)}V = \{(x, g) \in (\mathbb{C}^m)^n \times \mathcal{O}(U) : \\ DP \left(z, \bar{z}, j_n^k f(z), \overline{j_n^k f(z)} \right) \cdot \left[x, \bar{x}, Dj_n^k f(z) \cdot x + j_n^k g(z), \overline{Dj_n^k f(z) \cdot x + j_n^k g(z)} \right] = 0\}.$$

Let us show that the Sard-Smale Theorem for Fréchet manifolds applies to π . We fix $(z, f) \in V$, and denote by \mathcal{M}_z^k the subset of $\mathcal{O}(U)$ of all the functions whose jets of order k at z are equal to zero:

$$\mathcal{M}_z^k = \{g \in \mathcal{O}(U) : j_n^k g(z) = 0\}.$$

First observe that π is a C^1 -Lipschitz-Fredholm function:

- π is a C^1 function, and

$$D\pi(z, f) : (x, g) \in T_{(z,f)}V \mapsto g \in \mathcal{O}(U)$$

is a Lipschitz (with constant 1) operator;

- The kernel of $D\pi(z, f)$ is included in $(\mathbb{C}^m)^n \times \{0\}$, hence it has finite dimension;

- Since $(0, g) \in T_{(z,f)}V$ if $g \in \mathcal{M}_z^k$, we have

$$\mathcal{M}_z^k \subseteq \text{Im}(D\pi(z, f)),$$

so the co-kernel of $D\pi(z, f)$ is a quotient of $\mathcal{O}(U)/\mathcal{M}_z^k$. The space $\mathcal{O}(U)/\mathcal{M}_z^k$ has finite dimension $2N$ over \mathbb{R} , so $\text{co-ker}(D\pi(z, f))$ has finite dimension too.

It remains to compute the index of $D\pi(z, f)$. For this, observe that

- $\ker D\pi(z, f) \subseteq (\mathbb{C}^m)^n \times \{0\}$. In particular,

$$\ker D\pi(z, f) \cap ((\mathbb{C}^m)^n \times \mathcal{M}_z^k) \subseteq (\mathbb{C}^m)^n \times \{0\};$$

- the equations which define $T_{(z,f)}V$ only involve (in terms of g) the jet $j_n^k g(z)$ of order k .

Let $\widetilde{T_{(z,f)}V}$ be the quotient of $T_{(z,f)}V$ by $(\{0\} \times \mathcal{M}_z^k) \cap T_{(z,f)}V$ and $\widetilde{\mathcal{O}(U)} = \mathcal{O}(U)/\mathcal{M}_z^k$. It follows from the previous observations that $D\pi(z, f)$ factors to

$$D\widetilde{\pi}(z, f) : \widetilde{T_{(z,f)}V} \rightarrow \widetilde{\mathcal{O}(U)}$$

and that the index of $D\pi(z, f)$ is equal to the index of $D\widetilde{\pi}(z, f)$. Since $D\widetilde{\pi}(z, f)$ is a linear map between the finite dimensional spaces $\widetilde{T_{(z,f)}V}$ and $\widetilde{\mathcal{O}(U)}$, the index of $D\widetilde{\pi}(z, f)$ is simply $\dim \widetilde{T_{(z,f)}V} - \dim \widetilde{\mathcal{O}(U)}$. Now,

$$\begin{aligned} \widetilde{T_{(z,f)}V} &= \{(x, g) \in (\mathbb{C}^m)^n \times \widetilde{\mathcal{O}(U)} : \\ &\quad DP\left(z, \bar{z}, j_n^k f(z), \overline{j_n^k f(z)}\right) \cdot \left[x, \bar{x}, Dj_n^k f(z) \cdot x + j_n^k g(z), \overline{Dj_n^k f(z) \cdot x + j_n^k g(z)}\right] = 0\}. \end{aligned}$$

is a subspace of a real $(2mn + 2N)$ -dimensional space given by $2(2mn + 1)$ equations. Among this equations, at least $2mn + 1$ are independent, since $\text{rk}\left(DP\left(z, \bar{z}, j_n^k f(z), \overline{j_n^k f(z)}\right)\right) = 2mn + 1$. Hence $\dim \widetilde{T_{(z,f)}V} \leq 2N - 1$. On the other hand, $\dim \widetilde{\mathcal{O}(U)} = 2N$, therefore

$$\text{index}(D\pi(z, f)) \leq (2N - 1) - 2N = -1.$$

We can now conclude. The version of the Sard-Smale Theorem in [Eft11, Theorem 4.3] applies to π since π is C^1 and $1 > 0 = \max\{0, \text{index}(D\pi(z, f))\}$. We deduce that the set of regular values of π is residual in $\mathcal{O}(U)$. On the other hand, since it has negative index, $D\pi(z, f)$ has non trivial co-kernel, so π is nowhere a submersion. In particular the set of its critical values coincides with its image. Hence $B_{n,k,P}$, the complement of this image, is residual. Being a countable intersection of residual sets, $ST(U)$ is residual, which completes the proof. \square

The following proposition expresses strong transcendence in terms of the lack of relations between the germs at distinct points: except for the trivial operator, no operator in \mathcal{D}^* vanishes at any tuple of germs of any strongly transcendental function.

Proposition 3.4. *Let U be an open subset of \mathbb{C}^m , $f \in ST(U)$ be a strongly transcendental holomorphic function on U , $a = (a_1, \dots, a_k) \in (\mathbb{C}^m)^k$ be a k -tuple of distinct points of U , $b \in \mathbb{C}^n$ and $\mathcal{L} : \mathcal{O}_{a_1} \times \dots \times \mathcal{O}_{a_k} \rightarrow \mathcal{O}_b$ be an operator in \mathcal{D}^* . If $\mathcal{L}(f_{a_1}, \dots, f_{a_k}) = 0$, then there exists a neighbourhood W of $(f_{a_1}, \dots, f_{a_k})$ for the Krull topology such that $\mathcal{L}|_W = 0$.*

Proof. We follow the main steps of the proof of Lemma 3.6 in [LG10]. Suppose that \mathcal{L} satisfies the hypotheses of the proposition. Apply Proposition 2.12 to \mathcal{L} and $f_i = f_{a_i}$ to obtain, for all (g_1, \dots, g_k) in some neighbourhood W' of $(f_{a_1}, \dots, f_{a_k})$,

$$\mathcal{L}(g_1, \dots, g_k)(x) = \sum_{\alpha \in \mathbb{N}^n} P_\alpha(c_1, \dots, c_\ell, j^{n_\alpha} g_1(a_1), \dots, j^{n_\alpha} g_k(a_k))(x - b)^\alpha.$$

From the fact that $\mathcal{L}(f_{a_1}, \dots, f_{a_k}) = 0$, we deduce that

$$\forall \alpha \in \mathbb{N}^n, P_\alpha(c_1, \dots, c_\ell, j_k^{n_\alpha} f(a_1, \dots, a_k)) = 0.$$

Denote by $c \in \mathbb{C}^\ell$ the tuple $c = (c_1, \dots, c_\ell)$. Since $f \in ST(U)$, for all $\alpha \in \mathbb{N}^n$ the tuple $(a, \bar{a}, j_k^{n_\alpha} f(a), \overline{j_k^{n_\alpha} f(a)})$ satisfies at most $2mk$ algebraically independent relations. Since the transcendence degree is sub-additive,

$$\begin{aligned} \text{trdeg}_{\mathbb{Q}}(c, \bar{c}, j_k^{n_\alpha} f(a), \overline{j_k^{n_\alpha} f(a)}) &\geq \text{trdeg}_{\mathbb{Q}}(j_k^{n_\alpha} f(a), \overline{j_k^{n_\alpha} f(a)}) \\ &\geq \text{trdeg}_{\mathbb{Q}}(a, \bar{a}, j_k^{n_\alpha} f(a), \overline{j_k^{n_\alpha} f(a)}) \\ &\geq \text{trdeg}_{\mathbb{Q}}(j_k^{n_\alpha} f(a), \overline{j_k^{n_\alpha} f(a)}) - 2\ell, \end{aligned}$$

so $(c, \bar{c}, j_k^{n_\alpha} f(a), \overline{j_k^{n_\alpha} f(a)})$ satisfies at most $2\ell + 2mk$ independent algebraic relations. In particular, all but finitely many of the relations $P_\alpha(c, j_k^{n_\alpha} f(a)) = 0$ are dependent (recall that the P_α are polynomials over \mathbb{Q} in their arguments and their conjugates). This implies that $\{n_\alpha : \alpha \in \mathbb{N}^n\}$ is bounded by some constant $K \in \mathbb{N}$.

Let W be the neighbourhood of $(f_{a_1}, \dots, f_{a_k})$ given by

$$W = W' \cap \{(g_1, \dots, g_k) \in \mathcal{O}_{a_1} \times \dots \times \mathcal{O}_{a_k} : \forall i = 1, \dots, k, j^K g_i(a_i) = j^K f(a_i)\}$$

Then, for every $(g_1, \dots, g_k) \in W$ and every α , we have $j^{n_\alpha} g_i(a_i) = j^{n_\alpha} f(a_i)$. It follows that

$$\begin{aligned} \mathcal{L}(g_1, \dots, g_k)(x) &= \sum_{\alpha \in \mathbb{N}^n} P_\alpha(c_1, \dots, c_\ell, j^{n_\alpha} g_1(a_1), \dots, j^{n_\alpha} g_k(a_k))(x - b)^\alpha \\ &= \sum_{\alpha \in \mathbb{N}^n} P_\alpha(c_1, \dots, c_\ell, j^{n_\alpha} f(a_1), \dots, j^{n_\alpha} f(a_k))(x - b)^\alpha \\ &= \mathcal{L}(f_{a_1}, \dots, f_{a_k})(x) \\ &= 0. \end{aligned}$$

Therefore $\mathcal{L}|_W = 0$, which finishes the proof. \square

4. MONOMIAL DIVISION

In this section we let $\mathcal{A} = \{\exp\}$, and define

$$f(z) = \frac{e^z - 1}{z} \text{ for } z \neq 0 \text{ and } f(0) = 1,$$

so that f is holomorphic and locally \emptyset -definable in $\mathbb{R}_{\mathcal{A}}$.

We shall prove Theorem A. This amounts to proving that the germ of f at zero belongs to $\mathcal{C}^0 \setminus \mathcal{B}$ (so in particular $\mathcal{B}^0 \subsetneq \mathcal{C}^0$ and $\mathcal{B} \subsetneq \mathcal{C}$).

The strategy is the following. Using the differential equation satisfied by the exponential function, it is easy to see that the germs in \mathcal{B}^0 satisfy certain nonsingular systems of exponential polynomial equations (Lemma 4.3). Therefore, if $f \in \mathcal{B}^0$, then there is a tuple Ψ of analytic germs (and some $\frac{1}{d}f(dz)$ is one of them) such that the germs and their exponentials satisfy a nonsingular system of polynomial equations. This gives a certain upper bound M on the transcendence degree of the tuple $(\Psi, \exp(\Psi))$ over \mathbb{C} . On the other hand, by Ax's Theorem [Ax71, Corollary 2], the components of the tuple $(\Psi, \exp(\Psi))$ satisfy few algebraic relations, so that the transcendence degree of the tuple must be at least $M + 1$. This contradicts the fact that $f \in \mathcal{B}^0$ and proves Theorem A.

Notation 4.1. Let $k, l \in \mathbb{N}$, $x = (x_1, \dots, x_k)$ and x' be a sub-tuple of x . If $F(x) = (F_1(x), \dots, F_l(x))$ is an l -tuple of functions and $i \in \{1, \dots, l\}$, then we denote by $\frac{\partial F_i}{\partial x'}(x)$ the column vector whose entries are $\frac{\partial F_i}{\partial x_j}(x)$, for x_j belonging to x' . We denote by $\frac{\partial F}{\partial x'}(x)$ the matrix whose columns are the vectors $\frac{\partial F_i}{\partial x'}(x)$, for $i \in \{1, \dots, l\}$. Finally, we denote by e^x the k -tuple $(e^{x_1}, \dots, e^{x_k})$.

Definition 4.2. Let $n \in \mathbb{N}$, $x = (x_0, x_1, \dots, x_n)$ and $x' = (x_1, \dots, x_n)$. Let $a \in \mathbb{C}$, $g \in \mathcal{O}_a$ and \mathcal{G} a sheaf of holomorphic functions. We say that g is **n -implicitly defined from \mathcal{G}** if there exist $F = (F_1, \dots, F_n) \in (\mathcal{G}_{n+1})^n$ and $\Psi = (\psi_0, \psi_1, \dots, \psi_n) \in (\mathcal{O}_a)^{n+1}$ such that $\psi_0(z) = z$, $\psi_1(z) = g(z)$, $F(\Psi(z)) = 0$ and the matrix $\frac{\partial F}{\partial x'}(\Psi(a))$ is invertible. In this case, F is called an **implicit system** (with coordinates in \mathcal{G}) of **size n** and Ψ is an **implicit solution**.

Lemma 4.3. *Let $a \in \mathbb{C}$ and $g \in \mathcal{O}_a$. Then $g \in \mathcal{B}_a$ if and only if there exists $n \in \mathbb{N}$ such that g is n -implicitly defined from $\mathbb{C}[x, e^x]$.*

The proof of the lemma is routine and is postponed to the end of the section. We now prove Theorem A.

Proof of Theorem A. The germ f_0 of f at zero belongs to \mathcal{C}^\emptyset since it is obtained from the germ $\exp_0 \in \mathcal{A}_0$ by monomial division.

Suppose for a contradiction that $f_0 \in \mathcal{B}_0$.

By Lemma 4.3, there exists $n \in \mathbb{N}$ such that f_0 is n -implicitly defined from $\mathbb{C}[x, e^x]$. Choose n minimal with respect to the following property:

$$\exists d \in \mathbb{N}^\times \text{ s.t. } \frac{1}{d}f_0(dz) \text{ is } n\text{-implicitly defined from } \mathbb{C}[x, e^x].$$

In other words, for all $n', d \in \mathbb{N}^\times$, if $\frac{1}{d}f_0(dz)$ is n' -implicitly defined, then $n' \geq n$.

Let $x = (x_0, x_1, \dots, x_n)$ and $x' = (x_1, \dots, x_n)$. Let $F = (F_1, \dots, F_n) \in (\mathbb{C}[x, e^x])^n$ and $\Psi = (\psi_0, \psi_1, \dots, \psi_n) \in (\mathcal{O}_0)^{n+1}$ be such that $\psi_0(z) = z$, $\psi_1(z) = \frac{1}{d}f_0(dz)$, $F(\Psi(z)) = 0$ and the matrix $\frac{\partial F}{\partial x'}(\Psi(0))$ is invertible. Write $F(x) = P(x, e^x)$, where $P = (P_1, \dots, P_n)$ is an n -tuple of polynomials in $2(n+1)$ variables.

Our first task is to use the minimality of n to prove that the components of the vector

$$(z, \psi_1(z) - \psi_1(0), \psi_2(z) - \psi_2(0), \dots, \psi_n(z) - \psi_n(0))$$

are \mathbb{Q} -linearly independent.

If this is not the case, then there are $\tilde{d}, a_0, \dots, a_{n-1} \in \mathbb{Z}$ where we may suppose without loss of generality that $\tilde{d} \in \mathbb{N}^\times$, and there is $K = \psi_n(0) - \sum_{i=0}^{n-1} a_i \psi_i(0) \in \mathbb{C}$ such that

$$\tilde{d}\psi_n(z) = \sum_{i=0}^{n-1} a_i \psi_i(z) + K.$$

Let

$$\begin{aligned} \tilde{x} &= (x_0, x_1, \dots, x_{n-1}), \quad \tilde{x}' = (x_1, \dots, x_{n-1}) \\ \varphi(\tilde{x}) &= \left(\tilde{d}x_0, \tilde{d}x_1, \dots, \tilde{d}x_{n-1}, \sum_{i=0}^{n-1} a_i x_i + K \right) \\ \tilde{\Psi}(z) &= \left(\frac{\psi_0(z)}{\tilde{d}}, \frac{\psi_1(z)}{\tilde{d}}, \dots, \frac{\psi_{n-1}(z)}{\tilde{d}} \right), \end{aligned}$$

so that $\varphi \circ \tilde{\Psi} = \Psi$. Let $G = F \circ \varphi$. Note that $G \in (\mathbb{C}[\tilde{x}, e^{\tilde{x}}])^n$ and $G(\tilde{\Psi}(z)) = F(\Psi(z)) = 0$. Remark that

$$\frac{\partial G}{\partial \tilde{x}'}(\tilde{\Psi}(0)) = \frac{\partial F}{\partial x'}(\Psi(0)) \cdot \frac{\partial \varphi}{\partial \tilde{x}'}(\tilde{\Psi}(0)).$$

Since $\frac{\partial F}{\partial x'}(\Psi(0))$ is invertible and $\frac{\partial \varphi}{\partial \tilde{x}'}(\tilde{\Psi}(0))$ has rank $n-1$, there exist $n-1$ components of G , say G_1, \dots, G_{n-1} , such that $\frac{\partial(G_1, \dots, G_{n-1})}{\partial \tilde{x}'}(\tilde{\Psi}(0))$ is invertible. It follows that $\tilde{\Psi}(\tilde{d}z)$ satisfies the implicit system $\tilde{G} = 0$. Hence, $\frac{1}{\tilde{d}}f_0(\tilde{d}dz)$ is $(n-1)$ -implicitly defined, which contradicts the minimality of n .

We have thus proved that the components of the vector

$$(z, \psi_1(z) - \psi_1(0), \psi_2(z) - \psi_2(0), \dots, \psi_n(z) - \psi_n(0))$$

are \mathbb{Q} -linearly independent.

By Ax's Theorem [Ax71, Corollary 2], we have that

$$\text{trdeg}_{\mathbb{C}} \mathbb{C}(\Psi(z), e^{\Psi(z)}) \geq (n+1) + 1 = n+2.$$

On the other hand, we claim that the tuple $(\Psi(z), e^{\Psi(z)})$ satisfies a nonsingular system of $(n+1)$ polynomial equations in $2(n+1)$ variables, and hence

$$\text{trdeg}_{\mathbb{C}} \mathbb{C}(\Psi(z), e^{\Psi(z)}) \leq 2(n+1) - (n+1) = n+1.$$

This will provide a contradiction and will prove the theorem.

To prove this second claim, let $y = (y_0, y_1, \dots, y_n) = (y_0, y')$ and $P_0(x, y) = dx_0x_1 + 1 - (y_0)^d$. Notice that, if $F_0(x) = P_0(x, e^x)$, then $F_0(\Psi(z)) = 0$.

We know that

$$\frac{\partial F}{\partial x_j}(x) = \frac{\partial P}{\partial x_j}(x, e^x) + e^{x_j} \frac{\partial P}{\partial y_j}(x, e^x) \quad \forall j = 1, \dots, n$$

and that the matrix $\frac{\partial F}{\partial x'}(\Psi(0))$ is invertible.

Hence there are $u_1, \dots, u_n \in \{x_1, \dots, x_n, y_1, \dots, y_n\}$ such that the vectors

$$\frac{\partial P}{\partial u_1}(\Psi(0), e^{\Psi(0)}), \dots, \frac{\partial P}{\partial u_n}(\Psi(0), e^{\Psi(0)})$$

are \mathbb{C} -linearly independent. Let $\tilde{P} = (P_0, P_1, \dots, P_n)$. Since the first coordinate of the vector $\frac{\partial \tilde{P}}{\partial x_0}(\Psi(0), e^{\Psi(0)})$ is $\frac{\partial P_0}{\partial x_0}(\Psi(0), e^{\Psi(0)}) = f_0(0) = 1$ and the first coordinate of the vectors

$$\frac{\partial \tilde{P}}{\partial u_1}(\Psi(0), e^{\Psi(0)}), \dots, \frac{\partial \tilde{P}}{\partial u_n}(\Psi(0), e^{\Psi(0)})$$

is $\frac{\partial P_0}{\partial u_j}(\Psi(0), e^{\Psi(0)}) = 0$, these $(n+1)$ vectors are \mathbb{C} -linearly independent. This proves the claim and finishes the proof of the theorem. \square

We finish this section by giving a proof of Lemma 4.3.

Proof. We first proceed to prove, by induction on the size n of the system, that \mathcal{B}_a contains the coordinates of all implicit solutions of every implicit system with coordinates in \mathcal{B} . Since $\mathcal{A} \subseteq \mathbb{C}[x, e^x] \subseteq \mathcal{B}$, this will prove the right-to-left implication.

If $n = 1$, then the assertion follows from the fact that \mathcal{B}_a is closed under extracting implicit functions. If $n > 1$, then suppose that \mathcal{B}_a contains the coordinates of all implicit solutions of all implicit systems of size $n-1$ with coordinates in \mathcal{B} .

Let F be an implicit system of size n with coordinates in \mathcal{B} and let Ψ be an implicit solution. After possibly permuting the variables and the coordinates of F , we may suppose that $\frac{\partial F_n}{\partial x_n}(\Psi(a)) \neq 0$.

Let $\tilde{x} = (x_0, x_1, \dots, x_{n-1}) = (x_0, \tilde{x}')$, $\tilde{\Psi}(z) = (\psi_0(z), \psi_1(z), \dots, \psi_{n-1}(z))$ and let $\varphi(\tilde{x}) \in \mathcal{B}_{\tilde{\Psi}(a)}$ be the implicit function of F_n at $\tilde{\Psi}(a)$, so that $\psi_n(a) = \varphi(\tilde{\Psi}(a))$. If we let $\tilde{F}_i(\tilde{x}) = F_i(\tilde{x}, \varphi(\tilde{x}))$ for $i = 1, \dots, n$, then for $j = 1, \dots, n-1$, we have

$$\frac{\partial \tilde{F}_i}{\partial x_j}(\tilde{x}) = \frac{\partial F_i}{\partial x_j}(\tilde{x}, \varphi(\tilde{x})) + \frac{\partial \varphi}{\partial x_j}(\tilde{x}) \frac{\partial F_i}{\partial x_n}(\tilde{x}, \varphi(\tilde{x})).$$

Since the vectors $\frac{\partial F}{\partial x_1}(\Psi(a)), \dots, \frac{\partial F}{\partial x_n}(\Psi(a))$ are linearly independent, there are $i_1, \dots, i_{n-1} \in \{1, \dots, n\}$

such that $\tilde{\Psi}$ is an implicit solution of the implicit system $\tilde{F} = (\tilde{F}_{i_1}, \dots, \tilde{F}_{i_{n-1}})$, which has coordinates in \mathcal{B} , since \mathcal{B} is closed under composition.

By the inductive hypothesis, $\tilde{\Psi} \in (\mathcal{B}_a)^{n-1}$, and, since

$$\psi_n(z) = \varphi(\tilde{\Psi}(z)),$$

we also have $\psi_n \in \mathcal{B}_a$.

It remains to prove the left-to-right implication. Let \mathcal{A}' be the family of all holomorphic functions such that for every $k \in \mathbb{N}$ and $a = a^{(k)} \in \mathbb{C}^k$, \mathcal{A}'_a is the collection of all germs $f \in \mathcal{O}_a$ such that f is n -implicitly defined from $\mathbb{C}[x, e^x]$ (for some $n \in \mathbb{N}$). More precisely, let $z = z^{(k)}$ and $t = t^{(n)}$ be tuples of variables and define

$$\mathcal{A}'_a = \{f \in \mathcal{O}_a : \exists n \in \mathbb{N}, \exists F \in (\mathbb{C}[z, t, e^z, e^t])^{k+n}, \exists \Phi \in (\mathcal{O}_a)^{n-1} \text{ such that,}$$

$$(*) \quad \text{if } \Psi(z) = (z, f(z), \Phi(z)), \text{ then } F(\Psi(z)) = 0 \text{ and } \frac{\partial F}{\partial t}(\Psi(a)) \text{ is invertible}\}.$$

We prove that \mathcal{A}' is stable under the operators in \mathcal{B}^* . Since $\mathcal{A}' \supseteq \mathcal{A}$ and \mathcal{B} is the smallest collection containing \mathcal{A} and stable under the action of \mathcal{B}^* , this will imply that $\mathcal{B} \subseteq \mathcal{A}'$.

(1) *Schwarz reflection.*

It suffices to note that

$$F(\Psi(z)) = 0 \text{ and } \frac{\partial F}{\partial t}(\Psi(a)) \text{ is invertible} \Leftrightarrow \overline{F}(\overline{\Psi(\overline{z})}) = 0 \text{ and } \frac{\partial \overline{F}}{\partial t}(\overline{\Psi(\overline{a})}) \text{ is invertible.}$$

Hence, if $f(z) \in \mathcal{A}'_a$ then $g(z) := \overline{f(\overline{z})} \in \mathcal{A}'_{\overline{a}}$.

(2) *Composition.*

Let $f \in \mathcal{A}'_a$ be n -implicitly defined (following the notation in $(*)$). Let $s \in \mathbb{N}$, $b = b^{(s)} \in \mathbb{C}^s$ and $g = (g_1, \dots, g_k) \in (\mathcal{A}'_b)^k$. Suppose that $g(b) = a$ and let $h := f \circ g \in \mathcal{O}_b$. We aim to prove that $h \in \mathcal{A}'_b$.

It is easy to see that we may suppose that the components of g are implicitly defined by the same implicit system G . In other words, we may suppose that there exists $m \in \mathbb{N}$ such that, if $x = x^{(s)}$, $u = u^{(m)}$ are tuples of variables, then there exist $G \in (\mathbb{C}[x, z, u, e^x, e^z, e^u])^{k+m}$ and $\Gamma \in (\mathcal{O}_b)^m$ such that, if $\Omega(x) = (x, g(x), \Gamma(x))$, then $G(\Omega(x)) = 0$ and $\frac{\partial G}{\partial(z,u)}(\Omega(b))$ is invertible.

Let

$$H(x, z, u, t) := (G(x, z, u), F(z, t)) \in (\mathbb{C}[x, z, u, t, e^x, e^z, e^u, e^t])^{k+m+n}$$

and

$$\Theta(x) := (\Omega(x), h(x), \Phi(g(x))) \in (\mathcal{O}_b)^{s+k+m+n}.$$

Then $H(\Theta(b)) = 0$ and

$$\frac{\partial H}{\partial(z, u, t)}(\Theta(b)) = \begin{pmatrix} \frac{\partial G}{\partial(z, u)}(\Omega(b)) & 0 \\ \frac{\partial F}{\partial(z, u)}(\Psi(a)) & \frac{\partial F}{\partial t}(\Psi(a)) \end{pmatrix},$$

which is invertible. Hence, up to a permutation, h is $(k+m+n)$ -implicitly defined.

(3) *Derivatives.*

Let $f \in \mathcal{A}'_a$ be n -implicitly defined (following the notation in $(*)$) and let $i \in \{1, \dots, k\}$. We aim to prove that $\frac{\partial f}{\partial z_i} \in \mathcal{A}'_a$. Let $w = w^{(n)}$ be a tuple of variables and

$$\tilde{F}(z, t, w) := \frac{\partial F}{\partial z_i}(z, t) + \frac{\partial F}{\partial t}(z, t) \cdot w.$$

Notice that $\frac{\partial \tilde{F}}{\partial w}(z, t, w) = \frac{\partial F}{\partial t}(z, t)$. Let

$$F^*(z, t, w) = (F(z, t), \tilde{F}(z, t, w)) \in (\mathbb{C}[z, t, w, e^z, e^t, e^w])^{2n}$$

and

$$\Psi^*(z) = \left(\Psi(z), \frac{\partial f}{\partial z_i}(z), \frac{\partial \Phi}{\partial z_i}(z) \right) \in (\mathcal{O}_a)^{k+2n}.$$

Then $F^*(\Psi^*(z)) = 0$ and

$$\frac{\partial F^*}{\partial(t, w)}(\Psi^*(z)) = \begin{pmatrix} \frac{\partial F}{\partial t}(\Psi(z)) & 0 \\ \frac{\partial \tilde{F}}{\partial t}(\Psi^*(z)) & \frac{\partial F}{\partial t}(\Psi(z)) \end{pmatrix},$$

which is invertible. Hence, up to a permutation, $\frac{\partial f}{\partial z_i}$ is $2n$ -implicitly defined.

(4) *Implicit function.*

Let $f \in \mathcal{A}'_a$ be n -implicitly defined (following the notation in $(*)$) and suppose that $\frac{\partial f}{\partial z_k}(a) \neq 0$. Write $a = (a', a_k) \in \mathbb{C}^{k-1} \times \mathbb{C}$ and $z = (z', z_k)$, where $z' = (z_1, \dots, z_{k-1})$. Let $\varphi \in \mathcal{O}_{a'}$ be the implicit function of f at a . We aim to prove that $\varphi \in \mathcal{A}'_{a'}$. Write $t = (t_1, t')$, where $t' = (t_2, \dots, t_n)$, and let

$$F^*(z', z_k, t) = (F(z', z_k, t), t_1) \in \left(\mathbb{C} \left[z', z_k, t, e^{z'}, e^{z_k}, e^t \right] \right)^{n+1}$$

and

$$\Psi^*(z') = \Psi(z', \varphi(z')) \in (\mathcal{O}_{a'})^{(k-1)+(n+1)}.$$

Then $F^*(\Psi^*(z)) = 0$ and

$$\frac{\partial F^*}{\partial (z_k, t_1, t')}(\Psi^*(z')) = \begin{pmatrix} \frac{\partial F}{\partial z_k} & \frac{\partial F}{\partial t_1} & \frac{\partial F}{\partial t'} \\ 0 & 1 & 0 \end{pmatrix} (\Psi(z', \varphi(z'))).$$

(Here we mean the derivative of F^* is evaluated at $\Psi^*(z')$).

Differentiating the identity $F(\Psi(z)) = 0$ with respect to the variable z_k we find that

$$\frac{\partial F}{\partial z_k}(\Psi(z)) = -\frac{\partial F}{\partial t_1}(\Psi(z)) \cdot \frac{\partial f}{\partial z_k}(\Psi(z)) - \frac{\partial F}{\partial t'}(\Psi(z)) \cdot \frac{\partial \Phi}{\partial z_k}(z)$$

hence the vector $\frac{\partial F}{\partial z_k}(\Psi^*(a'))$ (which is equal to $\frac{\partial F}{\partial z_k}(\Psi(a))$) is \mathbb{C} -linearly independent from the (\mathbb{C} -linearly independent) vectors

$$\frac{\partial F}{\partial t_2}(\Psi^*(a')), \dots, \frac{\partial F}{\partial t_n}(\Psi^*(a')).$$

so that $\frac{\partial F^*}{\partial (z_k, t_1, t')}(\Psi^*(a'))$ has full rank. □

5. DERAMIFICATION

In this section we prove Theorem B.

In order to do this, we associate to each operator \mathcal{L} in \mathcal{D}^* a function which measures the index shift between a (tuple of) series in the domain of \mathcal{L} and the image under \mathcal{L} of such series.

Definition 5.1. Let $\mathcal{L} : \mathcal{O}_{a_1} \times \dots \times \mathcal{O}_{a_m} \rightarrow \mathcal{O}_b$ be an operator in \mathcal{D}^* . The shift of \mathcal{L} is the function $d_{\mathcal{L}} : \mathbb{N} \rightarrow \mathbb{N}$ given by

$$d_{\mathcal{L}}(n) = \min\{k \in \mathbb{N} : \forall (f_1, \dots, f_m), (g_1, \dots, g_m) \in \mathcal{O}_{a_1} \times \dots \times \mathcal{O}_{a_m}, \\ \text{if } \forall i = 1 \dots, m, j^k f_i(a_i) = j^k g_i(a_i), \text{ then } j^n \mathcal{L}(f_1, \dots, f_m)(b) = j^n \mathcal{L}(g_1, \dots, g_m)(b)\}.$$

The shift function has the following interpretation: to compute the terms of order n of $\mathcal{L}(F)$ it suffices to consider the set of terms of F of order k with $k \leq d_{\mathcal{L}}(n)$. Thanks to Proposition 2.12, the function $d_{\mathcal{L}}(n)$ is well defined for all $\mathcal{L} \in \mathcal{D}^*$. We can be more precise: $d_{\mathcal{L}}(n) = n$ if \mathcal{L} is either the Schwarz reflection, or the composition, or the implicit function operator; $d_{\mathcal{L}}(n) = n + 1$ if \mathcal{L} is a monomial division or a partial derivative operator; finally, $d_{\mathcal{L}}(n) = mn$ if \mathcal{L} is the m th-deramification operator. Moreover, we have:

Proposition 5.2. *Let \mathcal{L} be an operator in \mathcal{C}^* . Then, there exists a constant $N_{\mathcal{L}} \in \mathbb{N}$ such that $d_{\mathcal{L}}(n) \leq n + N_{\mathcal{L}} \forall n \in \mathbb{N}$.*

Proof. The operators in \mathcal{C}^* are finite compositions of operators among polynomials, Schwarz reflections, compositions, implicit functions, for which we set $N_{\mathcal{L}} = 0$, and monomial division and partial derivation, for which we set $N_{\mathcal{L}} = 1$. To obtain a bound on the shift for a general operator $\mathcal{L} \in \mathcal{C}^*$, we first note that if $\mathcal{M}, \mathcal{N}_1, \dots, \mathcal{N}_k$ are operators in \mathcal{D}^* , then the shift of the composition satisfies

$$d_{\mathcal{M} \circ (\mathcal{N}_1, \dots, \mathcal{N}_k)}(n) \leq \max_i d_{\mathcal{N}_i}(d_{\mathcal{M}}(n)).$$

Hence, setting

$$N'_{\mathcal{M}(\mathcal{N}_1, \dots, \mathcal{N}_k)} = N'_{\mathcal{M}} + \max_i N'_{\mathcal{N}_i},$$

we find integers $N'_{\mathcal{L}}$ associated to the representation of \mathcal{L} as a particular composition of elementary operators in \mathcal{C}^* . We can then take $N_{\mathcal{L}}$ to be the minimum of these integers over all representations of \mathcal{L} as a composition of elementary operators in \mathcal{C}^* . \square

This proposition shows in particular that no operator of composition with roots belongs to \mathcal{C}^* . However, there are examples of functions f such that $z \mapsto f(\sqrt{z})$ is obtained from f via operators in \mathcal{C}^* . For instance, if $f(z) = \frac{1}{1-z^2}$, then we have $f(\sqrt{z}) = f(z) + zf(z)$. Hence, to produce our example we need a function which do not satisfy these kind of relations. Strongly transcendental functions have this property.

The following proposition, together with Proposition 3.4, implies Theorem B.

Proposition 5.3. *Let $f \in ST(U)$, where U is an open neighbourhood of $0 \in \mathbb{C}$. Set $g(z) = f(z^2)$ and $\mathcal{A} = \{g\}$. Then, the germ f_0 of f at 0 belongs to $\mathcal{D}_0^\theta \setminus \mathcal{C}_0$.*

Proof. Since $g(z) = f(z^2)$, the germ f_0 is the image of g_0 under the square deramification operator, and hence belongs to \mathcal{D}_0^θ .

Suppose now for a contradiction that $f_0 \in \mathcal{C}_0$. Then there exist distinct points $a_0, \dots, a_k \in \mathbb{C}$ and an operator $\mathcal{L} \in \mathcal{C}^*$ such that $f_0 = \mathcal{L}(g_{a_0}, \dots, g_{a_k})$; we may assume that $a_0 = 0$, so that the germ g_0 does indeed occur in the list. Since $g(z) = f(z^2)$, if we denote by \mathcal{N}_i the operator $\mathcal{N}_i : \mathcal{O}_{a_i^2} \mapsto \mathcal{O}_{a_i}$ of composition with the polynomial $z \mapsto z^2$, we can rewrite this equality as

$$f_0 = \mathcal{L}(\mathcal{N}_0(f_0), \mathcal{N}_1(f_{a_1^2}), \dots, \mathcal{N}_k(f_{a_k^2})).$$

Hence the operator $\mathcal{M} : (\phi_0, \dots, \phi_k) \mapsto \mathcal{L}(\mathcal{N}_0(\phi_0), \mathcal{N}_1(\phi_1), \dots, \mathcal{N}_k(\phi_k)) - \phi_0$, vanishes at the tuple of the germs at distinct points of a strongly transcendental function. By Proposition 3.4, since $\mathcal{M} \in \mathcal{C}^*$, this operator vanishes identically on a neighbourhood of $(f_0, f_{a_1^2}, \dots, f_{a_n^2})$. For a large enough ℓ , we have that \mathcal{M} vanishes at $(f_0 + \lambda z^\ell, f_{a_1^2}, \dots, f_{a_n^2})$ for all $\lambda \in \mathbb{C}$, from which it follows that

$$\forall \lambda \in \mathbb{C}, f_0 + \lambda z_0^\ell = \mathcal{L}(g_0 + \lambda z^{2\ell}, g_{a_1}, \dots, g_{a_k}).$$

From this expression we deduce that $d_{\mathcal{L}}(\ell) \geq 2\ell$ for all sufficiently large ℓ . This, together with Proposition 5.2, contradicts the hypothesis $\mathcal{L} \in \mathcal{C}^*$. \square

6. BLOW-DOWNS

In this section we prove Theorem C.

Notation 6.1. We fix the usual coordinate system for the blow-up of $0 \in \mathbb{C}^2$.

Recall that the blow-up of $0 \in \mathbb{C}^2$ is the map $\pi : V \rightarrow \mathbb{C}^2$ where

$$V = \{(z, p) \in \mathbb{C}^2 \times \mathbb{C}\mathbb{P}^1 : z \in p\}$$

and $\pi(z, p) = z$. Let $D = \pi^{-1}(0) = \{0\} \times \mathbb{C}\mathbb{P}^1$ be the exceptional divisor. On the analytic manifold V we consider the atlas given by the following charts c_λ , for $\lambda \in \overline{\mathbb{C}} := \mathbb{C} \cup \{\infty\}$:

$$\begin{aligned} c_\lambda : \mathbb{C}^2 \ni (z_1, z_2) &\mapsto (z_1, (\lambda + z_2)z_1, [1 : \lambda + z_2]) \text{ if } \lambda \neq \infty \\ c_\infty : \mathbb{C}^2 \ni (z_1, z_2) &\mapsto (z_1 z_2, z_2, [z_1 : 1]), \end{aligned}$$

which, after composition with π , give rise to the following system $\pi_\lambda = \pi \circ c_\lambda$ of local expressions for π :

$$\begin{aligned} \pi_\lambda : \mathbb{C}^2 \ni (z_1, z_2) &\mapsto (z_1, (\lambda + z_2)z_1) \text{ if } \lambda \neq \infty \\ \pi_\infty : \mathbb{C}^2 \ni (z_1, z_2) &\mapsto (z_1 z_2, z_2) \end{aligned}$$

If $f : U \subseteq \mathbb{C}^2 \rightarrow \mathbb{C}$ is a function, then the **blow-up** of f is the function $f \circ \pi : \pi^{-1}(U) \rightarrow \mathbb{C}$, and f is the **blow-down** of $f \circ \pi$. Since we do not want to introduce sheaves on manifolds other than \mathbb{C}^n , we will only use local coordinates. Hence, the **blow-up of $f : U \subseteq \mathbb{C}^2 \rightarrow \mathbb{C}$ centered at λ** is the function $f \circ \pi_\lambda : \pi_\lambda^{-1}(U) \rightarrow \mathbb{C}$; we then say that f is the **blow-down of the family $(f \circ \pi_\lambda)_{\lambda \in \overline{\mathbb{C}}}$** . Note that

the blow-ups of f are obtained from f by composing with polynomials (operators in \mathcal{B}^*). Hence, the fact that \mathcal{E} is stable under blow-downs implies that

$$f_0 \in \mathcal{E}_0 \Leftrightarrow \forall \lambda \in \overline{\mathbb{C}}, (f \circ \pi_\lambda)_0 \in \mathcal{E}_0.$$

The key idea which allows us to construct our counterexample is the observation that blow-downs are not *local* operators. Let $g : V \rightarrow \mathbb{C}$ be a function and suppose that the blow-down f of g exists. Choosing f wisely, we can show that to obtain the germ f_0 , we need the germs of g at *all* points of the exceptional divisor D , whereas if f were obtained from g via operators from \mathcal{D}^* , one would only need the germs of g at finitely many points to construct f_0 . To make this argument work we need to choose an f which satisfies very few relations. The following proposition is useful to establish that strongly transcendental functions have this desired property.

Proposition 6.2. *Let $f \in ST(U)$ be strongly transcendental on an open subset U of \mathbb{C}^2 , let \mathcal{L} be an operator in \mathcal{D}^* of arity n and let $(b_0, b_1, \dots, b_n) \in U^{n+1}$ be distinct points in U . Then $f_{b_0} \neq \mathcal{L}(f_{b_1}, \dots, f_{b_n})$.*

Proof. Suppose $f_{b_0} = \mathcal{L}(f_{b_1}, \dots, f_{b_n})$. Then the operator \mathcal{M} defined by

$$\mathcal{M}(h_0, h_1, \dots, h_n) = \mathcal{L}(h_1, \dots, h_n) - h_0$$

belongs to \mathcal{D}^* as the composition of the operator of composition with a polynomial and the operator \mathcal{L} . Moreover, \mathcal{M} vanishes at $(f_{b_0}, f_{b_1}, \dots, f_{b_n})$, and hence, according to Proposition 3.4, on some neighbourhood W of $(f_{b_0}, f_{b_1}, \dots, f_{b_n})$.

For every $k \in \mathbb{N}$, let

$$B_k = \{(h_0, \dots, h_n) \in \mathcal{O}_{b_0} \times \dots \times \mathcal{O}_{b_n} : j^k h_0(b_0) = 0, \dots, j^k h_n(b_n) = 0\}$$

and choose $k \in \mathbb{N}$ sufficiently large such that $(f_{b_0}, \dots, f_{b_n}) + B_k \subseteq W$. Denote by $b_i = (b_{1,i}, b_{2,i})$ the coordinates of b_i for $i = 0, \dots, n$, and let P be the polynomial

$$P(z_1, z_2) = \prod_{i=0}^n (z_1 - b_{1,i})^k (z_2 - b_{2,i})^k.$$

Since $(P_{b_0}, P_{b_1}, \dots, P_{b_{n+1}})$ and $(0, P_{b_1}, \dots, P_{b_n})$ both belong to B_k , we have

$$\mathcal{L}(f_{b_1} + P_{b_1}, \dots, f_{b_n} + P_{b_n}) - (f_{b_0} + P_{b_0}) = 0$$

$$\text{and } \mathcal{L}(f_{b_1} + P_{b_1}, \dots, f_{b_n} + P_{b_n}) - (f_{b_0}) = 0,$$

from which we deduce $P_{b_0} = 0$, which is absurd. \square

The following proposition, together with Proposition 3.4, implies Theorem C.

Proposition 6.3. *Let U be a neighbourhood of $0 \in \mathbb{C}^2$ and $f : U \rightarrow \mathbb{C}$ be a strongly transcendental function. Let π be the blow-up of $0 \in \mathbb{C}^2$ and $\mathcal{A} = \{f \circ \pi_\lambda : \lambda \in \overline{\mathbb{C}}\}$. Then $f_0 \in \mathcal{E}_0^\emptyset \setminus \mathcal{D}_0$.*

Proof. The germ f_0 is in \mathcal{E}_0^\emptyset since it is the blow-down of the family $\{(f \circ \pi_\lambda)_0 : \lambda \in \overline{\mathbb{C}}\}$, whose elements belong to \mathcal{A}_0 .

Let us suppose for a contradiction that $f_0 \in \mathcal{D}_0$. Then there exist $\mathcal{N} \in \mathcal{D}^*$, $(a_1, \dots, a_k) \in (\mathbb{C}^2)^k$ and $(\lambda_1, \dots, \lambda_k) \in (\overline{\mathbb{C}})^k$ such that

$$(1) \quad f_0 = \mathcal{N}((f \circ \pi_{\lambda_1})_{a_1}, \dots, (f \circ \pi_{\lambda_k})_{a_k}).$$

Let U' be a neighbourhood of $\{c_{\lambda_1}(a_1), \dots, c_{\lambda_k}(a_k)\}$ in V such that $D \not\subseteq \overline{U'}$ (so that $\pi(U')$ is not a neighbourhood of 0). Define $\mathcal{A}' = \{(f \circ \pi_\lambda)|_{c_\lambda^{-1}(U')} : \lambda \in \overline{\mathbb{C}}\}$ and let \mathcal{D}' be the closure of \mathcal{A}' under the action of \mathcal{D}^* .

Equation (1) above shows that $f_0 \in \mathcal{D}'_0$, so by definition there exists a neighbourhood U'' of 0 such that $f|_{U''} \in \mathcal{D}'(U'')$. Since $\pi(U')$ is not a neighbourhood of 0, $U'' \setminus \pi(U') \neq \emptyset$. We fix $b_0 \in U'' \setminus \pi(U')$. Now, since $f|_{U''} \in \mathcal{D}'(U'')$, the germ of f at b_0 can be expressed in terms of some of the germs of the

restriction of $f \circ \pi$ to U' . So there exist $\mathcal{M} \in \mathcal{D}^*$, $(\mu_1, \dots, \mu_n) \in (\overline{\mathbb{C}})^n$ and $(a'_1, \dots, a'_n) \in (\mathbb{C}^2)^n$ with $a'_i \in c_{\mu_i}^{-1}(U')$ for $i = 1, \dots, n$, such that

$$f_{b_0} = \mathcal{M}((f \circ \pi_{\mu_1})_{a'_1}, \dots, (f \circ \pi_{\mu_n})_{a'_n}).$$

This equation can be reformulated as a relation between the germs of f at b_0 and b_1, \dots, b_n , where $b_i = \pi_{\mu_i}(a'_i)$. If $\mathcal{N}_i : \mathcal{O}_{b_i} \rightarrow \mathcal{O}_{a'_i}$ is the operator of composition with the polynomial π_{μ_i} , then $\mathcal{N}_i \in \mathcal{B}^*$ and $(f \circ \pi_{\mu_i})_{a'_i} = \mathcal{N}_i(f_{b_i})$. So we obtain

$$f_{b_0} = \mathcal{M}(\mathcal{N}_1(f_{b_1}), \dots, \mathcal{N}_n(f_{b_n})).$$

Setting $\mathcal{L} = \mathcal{M}(\mathcal{N}_1, \dots, \mathcal{N}_n)$, the operator \mathcal{L} is in \mathcal{D}^* , since it is a composition of operators of \mathcal{D}^* , and we have $f_{b_0} = \mathcal{L}(f_{b_1}, \dots, f_{b_n})$. Up to decreasing n , we may suppose that the points b_1, \dots, b_n are all distinct. Moreover, observe that $b_0 \notin \{b_1, \dots, b_n\}$: for $i = 1, \dots, n$, $a'_i \in c_{\mu_i}^{-1}(U')$, so $b_i \in \pi(U')$ while b_0 has been chosen outside $\pi(U')$. Hence the points b_0, \dots, b_n are all distinct, so Proposition 6.2 applies. But this contradicts the fact that $f_{b_0} = \mathcal{L}(f_{b_1}, \dots, f_{b_n})$. \square

REFERENCES

- [Arb00] L.-F.-A. Arbogast. *Du calcul des dérivations*. À Strasbourg, de l'Imprimerie de Levraut, Frères, An VIII (1800).
- [Ax71] J. Ax. On Schanuel's conjectures. *Ann. of Math. (2)*, 93:252–268, 1971.
- [Bia97] R. Bianconi. Nondefinability results for expansions of the field of real numbers by the exponential function and by the restricted sine function. *J. Symbolic Logic*, 62(4):1173–1178, 1997.
- [BM88] E. Bierstone and P. D. Milman. Semianalytic and subanalytic sets. *Inst. Hautes Études Sci. Publ. Math.*, (67):5–42, 1988.
- [Eft11] K. Eftekharinasab. Sard's theorem for mappings between Fréchet manifolds. *Ukrainian Math. J.*, 62(12):1896–1905, 2011.
- [JKS16] G. Jones, J. Kirby, and T. Servi. Local interdefinability of Weierstrass elliptic functions. *J. Inst. Math. Jussieu*, 15(4):673–691, 2016.
- [LG10] O. Le Gal. A generic condition implying o-minimality for restricted C^∞ -functions. *Ann. Fac. Sci. Toulouse Math. (6)*, 19(3-4):479–492, 2010.
- [LGR09] O. Le Gal and J.-P. Rolin. An o-minimal structure which does not admit C^∞ cellular decomposition. *Ann. Inst. Fourier*, 59(2):543–562, 2009.
- [LGSVB17] O. Le Gal, T. Servi, and E. Vieillard-Baron. Quasianalytic classes: what formal power series have to say about definable sets. *Manuscript in preparation*, 2017.
- [Lio33] J. Liouville. Premier et second mémoire sur la détermination des intégrales dont la valeur est algébrique. *Journal de l'École Royale Polytechnique*, XIV(22):124–193, 1833.
- [Ran14] S. Randriambololona. Two remarks on polynomially bounded reducts of the restricted analytic field with exponentiation. *Nagoya Math. J.*, 215:225–237, 2014.
- [RS15] J.-P. Rolin and T. Servi. Quantifier elimination and rectilinearization theorem for generalized quasianalytic algebras. *Proc. Lond. Math. Soc. (3)*, 110(5):1207–1247, 2015.
- [RSW03] J.-P. Rolin, P. Speissegger, and A. J. Wilkie. Quasianalytic Denjoy-Carleman classes and o-minimality. *J. Amer. Math. Soc.*, 16(4):751–777, 2003.
- [Sfo12] H. Sfouli. On the elementary theory of restricted real and imaginary parts of holomorphic functions. *Notre Dame J. Form. Log.*, 53(1):67–77, 2012.
- [vdDMM97] L. van den Dries, A. Macintyre, and D. Marker. Logarithmic-exponential power series. *J. London Math. Soc. (2)*, 56(3):417–434, 1997.
- [Wil08] A. J. Wilkie. Some local definability theory for holomorphic functions. In *Model theory with applications to algebra and analysis. Vol. 1*, volume 349 of *London Math. Soc. Lecture Note Ser.*, pages 197–213. Cambridge Univ. Press, Cambridge, 2008.
- [Zil02] B. Zilber. A theory of a generic function with derivations. In *Logic and algebra*, volume 302 of *Contemp. Math.*, pages 85–99. Amer. Math. Soc., Providence, RI, 2002.

SCHOOL OF MATHEMATICS, THE UNIVERSITY OF MANCHESTER, OXFORD RD, MANCHESTER, M13 9PL, UK

E-mail address: Gareth.Jones-3@manchester.ac.uk

URL: <http://personalpages.manchester.ac.uk/staff/Gareth.Jones-3/index.php>

SCHOOL OF MATHEMATICS, UNIVERSITY OF EAST ANGLIA, NORWICH RESEARCH PARK, NORWICH, NR4 7TJ, UK

E-mail address: jonathan.kirby@uea.ac.uk

URL: <http://www.uea.ac.uk/mathematics/people/profile/jonathan-kirby>

UNIV. GRENOBLE ALPES, UNIV. SAVOIE MONT BLANC, CNRS, LAMA, 73000 CHAMBÉRY, FRANCE

E-mail address: Olivier.Le-Gal@univ-smb.fr

URL: <http://www.lama.univ-savoie.fr/~LeGal/>

INSTITUT DE MATHÉMATIQUES DE JUSSIEU - PARIS RIVE GAUCHE, UNIVERSITÉ PARIS DIDEROT (PARIS 7), UFR DE MATHÉMATIQUES - BÂTIMENT SOPHIE GERMAIN, CASE 7012, F-75205 PARIS CEDEX 13, FRANCE

E-mail address: tamara.servi@math.univ-paris-diderot.fr

URL: <http://www.logique.jussieu.fr/~servi/index.html>