
Spatial chloroplast-to-nucleus signalling involving plastid-nuclear complexes and stromules

Philip M. Mullineaux^{1*}, Marino Exposito-Rodriguez¹, Pierre Philippe Laissue¹, Nicholas Smirnoff² and Eunsook Park^{3,4}

¹*School of Life Sciences, University of Essex, Wivenhoe Park, Colchester CO4 3SQ, United Kingdom*

²*College of Life and Environmental Sciences, Biosciences, University of Exeter, Exeter EX4 4QD, United Kingdom*

³*Plant Immunity Research Center, College of Agriculture and Life Sciences, Seoul National University, Seoul, 08826, republic of Korea*

⁴*Department of Molecular Biology, College of Agriculture and Natural Resources, University of Wyoming, Laramie, WY 82071, USA*

Keywords: Retrograde signalling, hydrogen peroxide, plastid-nuclear complexes, stromules, gene expression

Summary

Communication between chloroplasts and the nucleus in response to various environmental cues may be mediated by various small molecules. Signalling specificity could be enhanced if the physical contact between these organelles facilitates direct transfer and prevents interference from other subcellular sources of the same molecules. Plant cells have plastid-nuclear complexes, which provide close physical contact between these organelles. plastid-nuclear complexes have been proposed to facilitate transfer of photosynthesis-derived H₂O₂ to the nucleus in high light. Stromules (stroma filled tubular plastid extensions) may provide an additional conduit for transfer of a wider range of signalling molecules, including proteins. However, plastid-nuclear complexes and stromules have been hitherto treated as distinct phenomena. We suggest that plastid-nuclear complexes and stromules work in a co-ordinated manner so that, according to environmental conditions or developmental state the two modes of connection contribute to varying extents. We hypothesise that this association is dynamic and that there may be a link between plastid-nuclear complexes and the development of stromules. Furthermore, the changes in contact could alter signalling specificity by allowing an extended or different range of signalling molecules to be delivered to the nucleus.

*Author for correspondence (mullin@essex.ac.uk).

Introduction

In eukaryotes, the nucleus is the recipient of intracellular signals from every other organelle and compartment (de Souza et al 2017), which strongly suggests that spatial (i.e. 3-dimensional) as well as temporal components in signalling networks are of the utmost importance in terms of signalling specificity and the determination of cell fate. The continual adjustment to cellular metabolism in a fluctuating environment, which every photosynthetically active plant cell in a leaf has to carry out, depends upon the communication from its chloroplasts to the nucleus (hereafter termed retrograde signalling). Conversely, adjustments to primary metabolism involve much communication from the nucleus to plastids (termed anterograde signalling; de Souza et al 2017) and can result in changes to photosynthesis, alter protective mechanisms such as antioxidant capacity and modulate hormone biosynthesis.

In this short paper, we have not considered mitochondrion-nucleus retrograde signalling. Instead, we refer the reader to [other papers in this special issue](#). Rather, we have focussed on two means by which physical contact between plastids and the nucleus have been reported; plastid-nuclear complexes and stromules. We consider what is known about the dynamics of these interactions, the implications of close proximity of these organelles for the specificity of retrograde signalling as raised previously (Selga et al 2010; Higa et al 2014; Exposito-Rodriguez et al 2017) and begin to consider the notion that sub-populations of chloroplasts may have distinct cellular functions.

Plastid-nuclear complexes in higher plants and algae

A close association of plastids, including chloroplasts, and nuclei have been observed in many higher plant species ranging from horsetails (*Equisetum* sp.) to eudicots and monocots (Selga et al 2010; Fig.1A, B). Plastid-nuclear complexes may have a complex but ordered structure because in some images, the peri-nuclear endoplasmic reticulum may be seen to interpose between chloroplasts and their nucleus (Selga et al 2010). Furthermore, an extensive survey of the positioning of plastids around the nuclei of tobacco epidermal

cells strongly suggest a specific positioning between the organelles – the most striking and common being a daisy flower arrangement of plastids associated with the “equator” of the nucleus (Selga et al 2010). This is an arrangement we have also readily observed (Fig.1A; Movie S1). This apparently precise arrangement could mean that the structure of plastid-nuclear complexes is under tight regulation and be amenable to genetic analysis (see below). Algal cells have from one (e.g. *Chlamydomonas*, *Ostreococcus*) to many chloroplasts. In *Chlamydomonas*, the nucleus is enveloped within the cup-shaped chloroplast. In *Ostreococcus tauri*, TEM electron cryotomography shows close association of chloroplast and nucleus with the peroxisome sandwiched between them. At some points during cell division, elongated nuclear envelope processes stream around the chloroplasts (Henderson et al 2007). Since *Ostreococcus* cells are very small, with one copy of each organelle, it is difficult to determine if there are specific physical links. Photosynthetic protists of various kinds have chloroplasts derived from secondary endosymbiosis with algae and therefore have more complex membrane arrangement with 3-4 membranes enclosing the chloroplast and sometimes enclosing a nucleomorph (remnant of the symbiont’s nucleus) (Keeling, 2010). Attachment of the chloroplasts to each other and to the nucleus has been reported in *Euglena*, particularly during cell division (Ehara et al 1990). Chromosomes are prominent near the contact points. In *Ochromonas* the nuclear envelope appears to be continuous with the outer chloroplast membrane, with little or no cytoplasm between them (Gibbs, 1962). Clearly, more extensive data are needed to assess the extent of chloroplast-nuclear attachments in algae and photosynthetic protists.

The study by Selga et al (2010) described plastid-nuclear complexes in 10 plant species that included horsetail, a fern, gymnosperms, eudicots and monocots. This survey suggests that plastid-nuclear complexes in plant cells is the norm, but questions arise about the dynamic nature of plastid-nuclear complexes. For example, is there a turnover of chloroplasts associated with the nucleus? Despite, a range of microscopic methods having been applied to image plastid-nuclear complexes (Selga et al 2010), we have no impression

of their turnover.. In many photosynthetic cell types packed with chloroplasts (e.g. estimates of 50-70 in *Arabidopsis* mesophyll cells; Glynn et al 2009) only a percentage of chloroplasts would be able to engage in direct interactions with chloroplasts (Fig.1B) but it would be difficult to observe turnover. However, in cell types with lower numbers of chloroplasts such as the abaxial epidermal tissue of *Nicotiana benthamiana* (*N. benthamiana*), a single time point sampling revealed 3-12 chloroplasts in contact with the nucleus (Fig.1A, C; Exposito-Rodriguez et al 2017). This could imply a stochastic process but equally could be reflecting turnover in chloroplast numbers in different plastid-nuclear complexes such that at any timepoint different cells display differing number of chloroplasts interacting with their nucleus. Resolving this would require long term observations of the same cell with differentially marked chloroplasts.

Plastid-nuclear complexes and the actin cytoskeleton

In the streptophytes, chloroplasts and nuclei move to anticlinal sides of cells away from high fluence blue light such as in the high light (HL) conditions used by the authors (Exposito-Rodriguez et al 2017). This is called the avoidance response and is controlled by phototropins (Kong and Wada, 2011). The avoidance response of chloroplasts depends upon their interaction with the actin cytoskeleton (Kong and Wada, 2011, Higa et al 2014, Iwabuchi et al 2010). Chloroplasts and the nucleus in each cell are tethered to one another via the actin cytoskeleton and the action of at least three proteins CHLOROPLAST UNUSUAL POSITIONING1 (CHUP1), KINESIN-LIKE PROTEIN FOR ACTIN-BASED CHLOROPLAST MOVEMENT1 (KAC1) and KAC2 that are associated with the plastid outer membrane (Oikawa et al 2003; Caplan et al 2015; Higa et al 2014; Suetsugu et al 2016). These proteins serve primarily to anchor chloroplasts to the plasma membrane but appear also to be crucial for correct tethering of nuclei to chloroplasts. Nuclei have no independent capacity to move along the actin cytoskeleton, instead relying on their physical association with chloroplasts (Higa et al 2014; Suetsugu et al 2016). However, in mutants defective in

one or more of these proteins, nuclei still do move in response to incident light, albeit in an unusual manner. This is because in *chup1* and *kac1kac2* mutants, nuclei retain some connectivity to chloroplasts and therefore some capacity to carry out avoidance. Even a triple mutant (*chup1kac1kac2*) while showing severe attenuation did display some highly aberrant nuclear avoidance responses, implying there was still some nuclear-plastid connectivity (Higa et al 2014; Suetsugu et al 2016). Interestingly and in contrast to these mutants, some plastid division mutants (*plastid division1 (pdv1) / pdv2* double mutant and *paralog of arc6 (parc6)*) are also completely defective in tethering of chloroplasts to the nucleus (Higa et al 2014). In the case of *parc6*, the phenotype shows cell autonomous behaviour with respect to this phenotype (Higa et al 2014) meaning that most cells display a lack of chloroplast-to-nucleus tethering, but a proportion of them do not. Therefore, it may prove possible to compare cells with nuclei attached to chloroplasts alongside cells with separated chloroplasts and nuclei in the same tissue. This may obviate issues around the possibility of pleiotropic effects of such mutants. While there are many questions surrounding the use of *chup1*, *parc6* and other such mutants, they do indicate both the complexity and the likely dynamic nature of these plastid-nuclear complexes. In summary, we conclude that plastid-nuclear complexes are unlikely to be static structures and in considering their interactions with the cytoskeleton and overlap with plastid division they share commonality with stromules (see below).

Recently, it has been proposed that stromules might function to aid the dynamics of plastid-nuclear complexes leading to programmed cell death (PCD) in plant immunity (Kumar et al. 2018). Stromules are tubular protrusions stretched from the plastid body filled with stroma (Kohler et al. 1997; Hanson and Hines 2018; Fig.1C; Fig.2). Recent studies unveiled a potential role of stromules as a path to transfer signalling molecules from plastids to the nucleus (Caplan et al. 2015; Hanson and Hines 2018) and a regulatory factor to maintain the resulting plastid-nuclear complex via actin filaments during PCD (Kumar et al. 2018). Dynamic stromule formation is regulated differentially by actin and microtubule

cytoskeletons (Kumar et al 2018; Hanson and Hines, 2018). Recently and interestingly, the causative defective gene in an *Arabidopsis* mutant displaying enhanced stromule formation in epidermal plastids was shown to be *PARC6* (Itoh et al 2018), which is also implicated in the formation of plastid-nuclear complexes in mesophyll cells (see above). Unlike the chloroplast body that primarily moves along actin filaments (Kong and Wada, 2011, Higa et al 2014, Iwabuchi et al 2010), stromules use microtubules as their guide to undergo directional extension and retraction. Interestingly, actin filaments provide anchor points to regulate stromule length (Kumar et al 2018), showing that movement of the chloroplast body and stromules are not regulated in the same manner. Interestingly, during immune responses, numerous stromules were observed to extend towards the nucleus and often wrap around the nucleus (Caplan et al. 2015). In addition, the tips of stromules can anchor to the periphery of the nucleus followed by a retraction of the stromules resulting in movement of the chloroplast body closer to the nucleus (Fig. 2. Movie S2). This movement might be one of the mechanisms to maintain the plastid-nuclear complex observed in plant immunity (Kumar et al. 2018). However, genetic components to regulate stromules have yet to be identified. Although chloroplast body movement is altered in *chup1* (Oikawa et al 2003; Higa et al 2014; Suetsugu et al 2016), stromules were hyper-induced without pathogen infection in *N. benthamiana* cells showing suppressed *CHUP1* expression by RNA interference (Caplan et al., 2015). These data suggest that CHUP1 is a negative regulator of stromule formation. In these experiments, chloroplast bodies were frequently clustered similar to the plastid-nuclear complexes described above (Fig.1C; Fig. 2), although unfortunately nuclei were not co-visualized (Caplan et al 2015). Nevertheless, these data also suggest that CHUP1, presumably with as yet unidentified components, may also provide stromule-actin connectivity and stromule-mediated chloroplast movement towards the nucleus.

Problems of specificity

In almost all figures illustrating retrograde signalling, a single chloroplast is often depicted as the source of signals transduced to the nucleus (e.g. de Souza et al 2017). The reality in all higher plants' cells is somewhat different; multiple chloroplasts in cells are universal. Furthermore, in response to both internal and external cues it can be expected that not all chloroplasts inside a cell experience the same interaction with the environment. This is especially so for the light environment, where the light avoidance response (see above; Kong and Wada 2011) results in stacking of chloroplasts and ensures that some experience higher light intensity than others. Therefore, from a signalling context, it is feasible that not all chloroplasts in a HL-exposed cell will communicate with the nucleus to the same degree. Thus, how could signalling from multiple chloroplasts be integrated by the nucleus to produce a defined change in gene expression? Likewise, for many small molecules or metabolites that also are signal transducers, more than one source in a cell is possible or likely. The exemplar is hydrogen peroxide (H_2O_2 ; Mullineaux et al 2018) with sources not only from the chloroplasts, but from the peroxisome, mitochondria, plasma membrane and cytosol (Mullineaux et al 2018; Smirnoff and Arnaud, 2019). In which case, how is it possible that an accumulation of H_2O_2 in nuclei but sourced from chloroplasts be distinguished, for example, from H_2O_2 sourced from peroxisomes? Finally, how would a metabolite acting as a signalling molecule avoid being diverted into another pathway *en route* to the nucleus from chloroplasts? The potential advantage of proximity or attachment of chloroplasts and nuclei is that any small molecule signal is directed to the nucleus so that chloroplast conditions are more specifically indicated. However, if metabolites first have to move to the cytosol they will very rapidly equilibrate across the cell. Therefore, for metabolites shared between chloroplasts and the cytosol, this could render them less effective as a chloroplast signal. Alternatively, compounds that are readily metabolised (e.g. H_2O_2) could be removed before entering the nucleus. This is illustrated by the ease of detecting photosynthesis-sourced H_2O_2 in nuclei but not cytosol in response to high light (Exposito-Rodriguez et al 2017). The starting point to answer to all of the above questions could be the spatial context

in which signalling takes place in plastid-nuclear complexes. These complexes would allow direct communication between the origin of the transducing signal (the chloroplasts) and its destination (the nucleus). This is discussed further below, especially in the context of H_2O_2 as a transducing signalling molecule.

Partitioning and direct transfer of H_2O_2 from chloroplasts to the nucleus for signalling – a critical role for plastid-nuclear complexes and stromules?

In HL-exposed photosynthetically active cells, H_2O_2 accumulates in chloroplasts (Nakano and Asada 1980; Fryer et al 2003; Wen et al 2008; Galvez-Valdivieso et al 2009; Mubarakshina et al 2010; Driever and Baker 2011; Exposito-Rodriguez et al 2017). Various biochemical and genetic means of changing ROS levels in plant cells by promoting oxidative stress have been used to study the response of the transcriptome to H_2O_2 as well as other reactive oxygen species (ROS; reviewed in Mignolet-Spruyt et al 2016). The real value of the many independent transcriptomic datasets has been their combined study in meta-analyses using ever more statistically robust methodology (Gadjev et al 2006; Willems et al 2016). This has provided strong evidence that a cohort of H_2O_2 -responsive genes exist which are common to a number of different environmental and cellular cues, including exposure to HL. These meta-analyses do suggest that different subcellular sources of H_2O_2 could provide one element of signal specificity (Mignolet-Spruyt et al 2016; Willems et al 2016). For example, a transcriptomics study of Arabidopsis genotypes with altered H_2O_2 production and scavenging capacities in chloroplasts and peroxisome respectively, which were shifted from non-photorespiratory to photorespiratory conditions, clearly indicated that the source of H_2O_2 may bring about a specificity of response (Sewelam et al 2014). In summary, specificity of H_2O_2 signalling is likely, but how would this be achieved? This is especially the case, if we consider how H_2O_2 could be a retrograde signal transducer. The idea that H_2O_2 could convey a signal out of the chloroplast had been considered to be problematic (Mullineaux et al 2006; Galvez-Valdivieso et al 2009). The problem is that H_2O_2 , in its supposed journey from the chloroplast to the nucleus, would not last long in the

reducing environment of the cytosol. In addition, once exited from the chloroplast, the source specificity of H₂O₂ would surely be lost. Consequently, the view was that H₂O₂ could initiate signalling but not onward transduction out of the chloroplast. Further signal transduction to the nucleus would have to be achieved by some other signalling molecule, which would be stable during its transit of the cytosol. However, subsequent research challenged this view. Isolated chloroplasts secrete H₂O₂ into their medium in a light intensity and photosynthetic electron transport (PET)-dependent manner (Mubarakshina et al 2010) and there was the clear implication that this could also occur *in vivo*. Genetically encoded fluorescent protein biosensors that detect H₂O₂ enabled this question of its mobility and consequent specificity to be addressed (Exposito-Rodriguez et al 2017; Nietzel et al 2019). These biosensors can detect H₂O₂ with a high degree of specificity in real time, non-invasively and quantitatively (Belousov et al 2006; Meyer and Dick 2010; Exposito-Rodriguez et al 2013; Walia et al 2018). Using such a probe (Hyper; Belousov et al 2006) expressed transiently in *N. benthamiana* abaxial epidermal cells and targeted to chloroplast stroma, cytosol and nucleus revealed that under HL, H₂O₂ levels increased in nuclei concomitant with the rates of accumulation in the chloroplast stroma (Exposito-Rodriguez et al 2017). The HL-dependent increase in H₂O₂ (measured as increased HyPer oxidation) in both compartments was dependent upon active photosynthetic electron transport (PET). Furthermore, attenuation of the HL-triggered H₂O₂ accumulation in the chloroplast stroma by over-expressing the H₂O₂-scavenging enzyme ascorbate peroxidase (APX) also crucially inhibited its accumulation in the nucleus. This demonstrated that the H₂O₂ accumulation in the nucleus was directly dependent upon its accumulation in the chloroplast. The simplest, but not only, explanation for these observations is that transfer of H₂O₂ from chloroplasts to the nucleus occurs rapidly upon exposure to HL. Importantly, when a cytosolic isoform of APX was over-expressed it did not significantly attenuate accumulation of H₂O₂ in the nucleus. It was concluded that chloroplast-sourced H₂O₂ does not transit the cytosol and is a direct transfer from chloroplasts to the nucleus. In these abaxial epidermal cells, plastid-

nuclear complexes are readily observed, consisting of a median 7 chloroplasts per nucleus and it was noted that the oxidation of the HyPer probe in PNC chloroplasts was less than in those not associated with the nucleus. It was therefore hypothesised that it is the chloroplasts in plastid-nuclear complexes that transfer H₂O₂ directly to the nucleus.

Interestingly, it has been previously observed (Selga et al 2010) that chloroplasts detached from the nucleus underwent more rapid loss of chlorophyll fluorescence compared with those associated with the nucleus, implying different metabolic states for sub-populations of chloroplasts.

The same treatments that attenuated H₂O₂ in chloroplasts and nuclei also impacted on the expression of a *N. benthamiana* HL-responsive gene, *NbAPX1c*, in the same way, establishing that the H₂O₂ in the nucleus initiates onward signalling leading to the change in expression of at least one HL-responsive gene (Exposito-Rodriguez et al 2017).

While the hypothesis of a direct transfer of H₂O₂ from chloroplasts to the nucleus is the simplest explanation of the data, other, not necessarily mutually exclusive, variations on this retrograde signalling mechanism remain possible. It is clear that chloroplast-sourced H₂O₂ initiates and drives the signalling and that HL-dependent accumulation of H₂O₂ in the nucleus continues that signalling process. Nor is the notion of a spatial dependence of signalling negated. However, it is conceivable that another signalling molecule is transferred to the nucleus which stimulates H₂O₂ synthesis in that compartment or even that chloroplast-sourced H₂O₂ amplifies or activates its nuclear-localised synthesis. For example, nuclear-located cryptochromes (CRYs) when challenged with high fluence blue light can make H₂O₂ (Consentino et al 2015) and *CRY1* has been shown to positively regulate HL-responsive genes that are also responsive to H₂O₂ and require active PET (Kleine et al 2007; Karpinski et al 1999).

H₂O₂ is also known to be generated and have a critical role as a signalling molecule to induce plant immunity (Stael *et al.*, 2015). When PCD occurs, chloroplasts function as a major generator of H₂O₂, which often induce gene expression in the nucleus (Yao and

Greenburg, 2006). Moreover, application of exogenous H₂O₂ to leaves have been shown to increase stromule formation (Gray et al 2012). Recently, H₂O₂ translocation from chloroplast to the nucleus *via* stromules has been raised as a possibility from work using the HyPer H₂O₂ sensor (Caplan et al. 2015). In live cell time-lapsed images, the concentration of H₂O₂ increased in stromules whose tips were anchored to the nucleus. In addition, by using nuclear-localised HyPer, an increase in H₂O₂ in nucleus of plastid-nuclear complexes was monitored. Although these two events were monitored in separate experiments this study does support the hypothesis that H₂O₂ is a retrograde signalling molecule in plant immunity. However, more sophisticated experiments will be required to monitor H₂O₂ translocation from chloroplasts into the nucleus in the same cell, in order to be able to propose that stromules might be a major path for H₂O₂-mediated retrograde signalling in plant immune responses.

Application of exogenous H₂O₂ is sufficient to induce stromule formation vigorously (Gray et al 2012; Caplan et al. 2015; Brunkard and Burch-Smith 2018). Furthermore, evidence has recently been presented that the establishment of pathogen- or effector-triggered immunity or treatment with H₂O₂ also causes the chloroplasts of *N. benthamiana* epidermal pavement cells to cluster around the nucleus (Ding et al 2019). Interestingly though, the authors did not report the presence of stromules during their observations. In summary, evidence may be emerging that H₂O₂ not only produced by chloroplasts, but from other subcellular sources may also promote formation of both plastid-nuclear complexes and stromules. This implies a complex regulatory system, which we are just beginning to perceive. However, all these observations have used agro-infected *N. benthamiana* which might result in an interaction between HL and pathogen-associated molecular pattern (PAMP) responses (Caplan et al 2015; Exposito-Rodriguez et al 2017; Ding et al 2019) and therefore such observations do need to be confirmed in other experimental systems. Furthermore, some plant-pathogen interactions (e.g. that of *Arabidopsis* and *Pseudomonas syringae* DC3000) may suppress photosynthesis and chloroplast-sourced ROS in an effector dependent

manner (de Torres-Zabala et al 2015). In this case, the impact of suppression of chloroplast function and ROS formation on stromule formation is unknown.

Retrograde signalling and H₂O₂ in cytosol microdomains

The close associations between chloroplasts and nuclei do not exclude retrograde signalling involving H₂O₂ also going *via* the cytosol and still achieving signalling specificity. Under HL, *N. benthamiana* abaxial epidermal cells do accumulate H₂O₂ in the cytosol but it is not evenly distributed. It must be assumed that the rate of diffusion of H₂O₂ from chloroplasts that are not part of plastid-nuclear complexes, is sufficient to overcome rates of reducing activity from antioxidant systems in the cytosol for long enough to allow oxidation of the cytosol-located HyPer probe (Exposito-Rodriguez et al 2017). Active transport, *i.e.* secretion of H₂O₂ from chloroplasts, cannot be ruled out but no evidence is available on this point. If the resulting H₂O₂ microdomains are involved in signalling, then there would be temporal and spatial constraints meaning that redox-sensitive signal transducers will have to be in place to meet this localised H₂O₂ exiting from chloroplasts. There are candidate signal transducers that could act in such a role provided their spatial distribution in relation to H₂O₂ microdomains could be confirmed. At least three Arabidopsis heat shock transcription factors, HSFA1D, HSFA8 and HSFA4A, have been shown to be redox-regulated (Jung et al 2013; Giesguth et al 2015; Pérez-Salamó et al 2014). Inter- and intramolecular disulphide bond formation is important in the conversion of inactive cytosol-located monomeric HSF isoforms into active trimeric forms that migrate to the nucleus to carry out their function. The high degree of sequence conservation in extensive plant HSF gene families suggest that such potential redox regulation may extend beyond these three examples (Miller and Mittler 2006) Signal transduction involving H₂O₂ in eukaryotes may involve the transfer of oxidising equivalents by thiol peroxidases (TPXs; Giesguth et al 2015; Mullineaux et al 2018), which again would be required to be located or translocate to where H₂O₂ accumulates in microdomains. A simpler outcome could be that H₂O₂ from such chloroplasts, were it to

continue to accumulate in the cytosol for any length of time, would lead to cellular oxidative stress and trigger PCD (Mullineaux et al 2006).

In summary, regarding the role of H_2O_2 as a signal transducer in retrograde signalling there are clear layers of spatial dependency – plastid-nuclear complexes, stromules and microdomains. The juxtaposition of the players, once identified, in these signal transduction routes with respect to one another and to the accumulation of H_2O_2 will be critical in determining how H_2O_2 -mediated retrograde signalling truly functions.

Spatial considerations of metabolites as retrograde signal transducers

As with H_2O_2 , there are a myriad of small molecules that have single or distinct pools in chloroplasts and are translocated to other part of the cell as part of the normal role in cellular metabolism. Any molecule with a distinct origin or location in plastids has, therefore, the potential to be co-opted as a transducer in retrograde signalling. Recent productive lines of research have established at least 3 such metabolites or metabolic intermediates that fall into this class: 3'-phosphoadenosine 5'-phosphate (PAP; Estavillo et al 2011) with cytosolic and chloroplast pools; methylerythritol phosphate (MEcPP; Xiao et al 2012) a biosynthetic intermediate in plastid isoprenoid production; β -cyclocitral, an oxidation product of carotenoids formed in chloroplasts (Ramel et al 2012). These molecules have all been firmly established in the pantheon of prominent players in retrograde signalling. They have been proposed, and evidence offered, of being able to transduce signals out of the chloroplast and have been shown to strongly influence both whole plant responses to environmental stress and the expression of a distinct cohort of genes (Estavillo et al 2011; Xiao et al 2012; Ramel et al 2012). To our knowledge, no spatial relationship between chloroplasts and the nucleus has been invoked as necessary for their signalling roles to be effective. However, clearly the workings of these signalling pathways could be enhanced if they were functioning in plastid-nuclear complexes or require stromules. For such spatial relationships to be established, specific genetically encoded biosensors would be needed to

allow the necessary investigations to be done. The availability of such probes may still be some way off but would surely be of value to progress this field.

Spatial considerations of proteins as retrograde signal transducers

In contrast to the scores of metabolites and hormones that have been proposed as retrograde signal transducers, only a small number of proteins known to be targeted to the chloroplast have been identified subsequently in the nucleus to function as retrograde signal transducers in response to biotic and abiotic stresses. WHIRLY1 has been proposed to convey the redox status in chloroplasts to the nucleus in a salicylic acid-dependent manner (Foyer et al., 2014). WHIRLY1 proteins localize to both chloroplast and nucleus (Grabowski et al., 2008; Isemer et al., 2012). Expression of WHIRLY1 protein without its N-terminal plastid transit peptide sequence resulted in localization in the nucleus and successfully rescued the *whirly1* mutant phenotype (Isemer et al., 2012). Although dual localization of WHIRLY1 has been shown by several different approaches, how the translocation of WHIRLY1 from chloroplasts to the nucleus might occur is still not shown. An interesting chloroplast outer envelope protein, PTM (a PHD type transcription factor with transmembrane domains) was proposed to translocate to the nucleus to regulate high light-responsive gene expression (Sun et al 2011). This translocation of PTM was proposed to allow its binding to the promoter of *ABSCISIC ACID INSENSITIVE4* to induce expression of light-responsive genes (Sun et al., 2011) and to the promoter of *FLOWERING LOCUS C* to control flowering under HL (Feng et al 2016). However, the identity of the signal from the chloroplast to induce an intramembrane proteolytic cleavage of the PTM is unknown and how the N-terminal moiety of the PTM is released from the chloroplast and finally ends up in the nucleus remains to be investigated. Subsequently, doubt about this proposed role of PTM was raised by the lack of impairment of a genomes-uncoupled phenotype in *ptm* mutants treated with norflurazon and lincomycin (Page et al 2017).

Several GFP-tagged proteins have shown to be present in stromules (e.g. carbonic anhydrase) and GFP photoconversion and photobleaching experiments suggest this is a

dynamic process and that transfer of proteins between plastids can occur *via* stromules (reviewed by Hanson and Hines 2018). Recent studies of NRIP1 translocation from chloroplasts to nuclei *via* stromules might aid an understanding of the mechanism of translocation (Caplan et al 2015). NRIP1 is a helper of N protein, which recognizes the p50 protein of TMV (tobacco mosaic virus) and, in turn, rapidly triggers plant immunity (Caplan et al. 2008). NRIP1 protein is localised in the stroma of chloroplasts of tobacco plant cells in normal conditions. However, upon TMV infection, NRIP1 proteins can translocate into the nucleus through stromules anchored to the nucleus (Caplan et al. 2015).

Without further experimental support, it is hard to propose whether any of these above exemplar proteins translocate through stromules or directly by the plastid-nuclear complexes. However, given the proposed role of the stromules and the plastid-nuclear complexes to provide a path to transfer signalling molecules from chloroplasts to the nucleus in response to rapid changes of environmental status, it would be worth examining levels of stromules and the frequency of plastid-nuclear complexes in the WHIRLY1, PTM and NRIP1 activation conditions and their translocation *via* stromules and/or the plastid-nuclear complexes.

A time for stromules and a time for plastid-nuclear complexes: Is the link photosynthesis?

Both plastid-nuclear complexes and stromules are now proposed to provide a spatial element to retrograde signalling. Especially in the case of H₂O₂-mediated retrograde signalling, such direct contacts between chloroplasts and their nucleus provide signalling specificity and permit this ROS to be a direct carrier of a signal from chloroplasts to their associated nucleus. Exactly the same argument and evidence is provided for stromules regarding H₂O₂-mediated retrograde signalling. The difference in signalling roles between plastid-nuclear complexes and stromules may be that the latter are able to provide a specific conduit for a much wider range of signalling molecules from the chloroplasts,

including proteins (Hanson and Hines 2018). However, to our knowledge, no evidence is available that protein-mediated retrograde signalling is definitively limited to stromules. We have considered that plastid-nuclear complexes potentially provide a spatial component for signalling without stromules, but some studies show a high degree of stromule-producing chloroplasts present in such structures and stromules apparently facilitating the entry of their chloroplast into close contact with the nucleus (Fig. 2; Movie S2). All of this points to a distinct function for stromules over and above any signalling role that is also achieved by direct contact between chloroplasts and the nucleus.

We have commented above that that some researchers observed stromules in their experimental systems and others do not. This suggests that specific physiological states of chloroplasts and cells are important in determining the circumstances which give rise to stromule formation. While the predominance of observations has been made in cells undergoing PCD, either as senescence or in the induction of pathogen- or elicitor-induced immunity (see above), it is premature to assume that stromule formation is a phenomenon linked to this process. This is because drought, salinity, phosphate limitation and ABA (possibly *via* strigolactone signalling) can also induce stromule formation (Gray et al 2012; Vismans et al 2016) and these treatments, to our knowledge, do not induce PCD.

Furthermore, isolated chloroplasts have been reported to be able to form stromules (Brunkard et al 2015; Ho and Theg 2016). We propose instead that all these situations have in common a diminution in photosynthesis and primary metabolism. Induction of drought stress or phosphate limitation, and more controversially, exogenous ABA often disrupt photosynthesis (Spencer & Foyer 1986; Pinheiro and Chaves 2010; Bechtold et al 2016). The impact of immunity and senescence on photosynthesis is always associated with a decline in this function (Balazadeh et al 2008; de Torres-Zabala et al 2015). Furthermore, any restriction of photosynthesis and consequent rise in the oxidation state of the stroma is a likely pre-requisite for stromule formation (Brunkard et al 2015).

Acknowledgments

The authors are grateful to Dr. S. P. Dinesh-Kumar (Department of Biological Sciences,

University of California, Davis, USA) for allowing us to present image sequences that EP generated when she worked in his laboratory. EP acknowledges the support of the National Research Foundation of Korea (NRF) grant by the Korea government (MSIT) (# 2018R1A5A1023599, SRC), the Next Generation Biogreen 21 program of RDA (# PJ013201) and a NSF start-up award (# EPS-1655726). PMM, PPL, MER and NS acknowledge the support of the UK Biotechnology and Biological Sciences Research Council (BB/P026656/1, BB/I020071/1, BB/I020004/1 and BB/N001311/1) for the research underpinning this article.

References

Adam Z (2015). Plastid intramembrane proteolysis. *Biochim. Biophys. Acta* 1847: 910–914. DOI: 10.1016/j.bbabi.2014.12.006.

Balazadeh S, Parlitz S, Mueller-Roeber B & Meyer RC (2008), Natural developmental variations in leaf and plant senescence in *Arabidopsis thaliana*. *Plant Biology*, 10: 136-147. DOI:10.1111/j.1438-8677.2008.00108.x.

Bechtold U, Penfold CA, Jenkins DJ, Legaie R, Moore JD, Lawson T, Matthews JSA, Violet-Chabrand SRM, Baxter L, Subramaniam S, Hickman R, Florence H, Sambles C, Salmon DL, Feil R, Bowden L, Hill C, Baker NR, Lunn JE, Finkenstädt B, Mead A, Buchanan-Wollaston V, Beynon J, Rand DA, Wild DL, Denby KJ, Ott S, Smirnov N & Mullineaux PM (2016) Time-series transcriptomics reveals that *AGAMOUS-LIKE22* affects primary metabolism and developmental processes in drought-stressed *Arabidopsis*. *Plant Cell* 28: 345-366. DOI: 10.1105/tpc.15.00910.

Belousov VV, Fradko AF, Lukyanov KA, Staroverov DB, Shakhbazov KS, Terskikh AV & Lukyanov S (2006) Genetically encoded fluorescent indicator for intracellular hydrogen peroxide. *Nature Methods* 3: 281-286. DOI: 10.1038/nmeth866.

Brunkard JO, Runkel AM & Zambryski PC (2015) Chloroplasts extend stromules independently and in response to internal redox signals. *Proc Natl. Acad. Sci. USA* 112: 10044-10049. DOI: 10.1073/pnas.1511570112.

Brunkard JO & Burch-Smith TM (2018) Ties that bind: the integration of plastid signalling pathways in plant cell metabolism. *Essays Biochem* 62: 95-107 DOI: 10.1042/EBC20170011.

Caplan JL, Mamillapalli P, Burch-Smith TM, Czymmek K & Dinesh-Kumar SP (2008). Chloroplastic protein NRIP1 mediates innate immune receptor recognition of a viral effector. *Cell* 132: 449–462. DOI: 10.1016/j.cell.2007.12.031.

Caplan JL, Kumar AS, Park, E, Padmanabhan MS, Hoban K, Modla S, Czymmek K, Dinesh-Kumar SP (2015) Chloroplast stromules function during innate immunity. *Dev. Cell* 34, 45–57. DOI: 10.1016/j.devcel.2015.05.011.

Consentino L, Lambert S, Martino C, Jourdan N, Bouchet P-E, Witczak J, Castello P, El-Esawi M, Corbineau F, d'Harlingue A & Ahmad M (2015). Blue-light dependent reactive oxygen species formation by *Arabidopsis* cryptochrome may define a novel evolutionarily conserved signaling mechanism. *New Phytol.* 206: 1450–1462. DOI: 10.1111/nph.13341.

de Souza A, Wang J, Dehesh K 2017 Retrograde signals: Integrators of Interorganellar communication and orchestrators of plant development. *Annu Rev Plant Biol* 68; 85-108. DOI: 10.1146/annurev-arplant-042916-041007

de Torres-Zabala M, Littlejohn G, Jayaraman S, Studholme D, Bailey T, Lawson T, Tillich M, Licht D, Bolter B, Delfino L, Truman W, Mansfield J, Smirnov N, Grant M. 2015. Chloroplasts play a central role in plant defence and are targeted by pathogen effectors. *Nature Plants*, 1: 15074. DOI: 10.1038/nplants.2015.74.

Ding X, Jimenez-Gongora T, Krenz B, Lozano-Duran R (2019) Chloroplast clustering around the nucleus is a general response to pathogen perception in *Nicotiana benthamiana*. *Mol Plant Pathol*. DOI: 10.1111/mpp.12840.

Driever S & Baker NR (2011) The water-water cycle in leaves is not a major alternative electron sink for dissipation of excess excitation energy when CO₂ assimilation is restricted. *Plant Cell Environ*. 34: 837-846. DOI: 10.1111/j.1365-3040.2011.02288.x.

Ehara T, Osafune T & Hase E. (1990) Interactions between the nucleus and cytoplasmic organelles during the cell cycle of *Euglena gracilis* in synchronized cultures IV. An aggregate form of chloroplasts in association with the nucleus appearing prior to chloroplast division. *Exp Cell Res* 190, 104-112. DOI: 10.1016/0014-4827(90)90150-9.

Estavillo GM, Crisp PA, Pornsiriwong W, Wirtz M, Collinge D, Carrie C, Giraud, E, Whelan, J, David, P, Javot H, Brearley C, Hell R, Marin E & Pogson BJ (2011) Evidence for a SAL1-PAP Chloroplast Retrograde Pathway That Functions in Drought and High Light Signaling in *Arabidopsis*. *Plant Cell* 23: 3992-4012 DOI: 10.1105/tpc.111.091033.

Exposito-Rodriguez M, Laissue PP, Yvon-Durocher G, Smirnov N & Mullineaux PM (2017) Photosynthesis-dependent H₂O₂ transfer from chloroplasts to nuclei provides a high-light signalling mechanism. *Nature Comm*. 8: art. 49. DOI: 10.1038/s41467-017-00074-w.

Feng P, Guo H, Chi W, Chai X, Sun X, Xu X, Ma J, Rochaix J-D, Leister D, Wang H, Lu C, Zhang L (2016) Chloroplast retrograde signal regulates flowering *Proc. Natl. Acad. Sci. USA* 113: 10708-10713. DOI: 10.1073/pnas.1521599113

Foyer, C. & Spencer, C. *Planta* (1986) The relationship between phosphate status and photosynthesis in leaves. *Planta* 167: 369-375. DOI: 10.1007/BF00391341

Foyer CH, Karpinska B & Krupinska K (2014) The functions of WHIRLY1 and REDOX-RESPONSIVE TRANSCRIPTION FACTOR 1 in cross tolerance responses in plants: a hypothesis. *Phil. Trans. R. Soc. Lond. B Biol. Sci.* 369: 20130226. DOI: 10.1098/rstb.2013.0226.

Fryer MJ, Ball L, Oxborough K, Karpinski S, Mullineaux PM & Baker NR (2003) Control of Ascorbate Peroxidase 2 expression by hydrogen peroxide and leaf water status during excess light stress reveals a functional organisation of *Arabidopsis* leaves. *Plant J* 33: 691-705. DOI: 10.1046/j.1365-313X.2003.01656.x.

Gadjev I, Vanderauwera S, Gechev TS, Laloi C, Minkov IN, Shulaev V, Apel K, Inzé D, Mittler R, Van Breusegem F (2006). Transcriptomic footprints disclose specificity of reactive oxygen species signaling in *Arabidopsis*. *Plant Physiol*. 141: 436-445 DOI: 10.1104/pp.106.078717.

Galvez-Valdivieso G, Fryer MJ, Lawson T, Slattery K, Truman W, Smirnov N, Asami T, Davies WJ, Jones AM, Baker NR & Mullineaux PM (2009). The High Light Response in *Arabidopsis* Involves ABA Signaling between Vascular and Bundle Sheath Cells. *Plant Cell* 21: 2143-2162. DOI: 10.1105/tpc.108.061507.

Gibbs SP (1962) Nuclear envelope-chloroplast relationships in algae. *J. Cell Biol.* 14: 433-444. DOI:10.1083/jcb.14.3.433.

Giesguth M, Sahm A, Simon S & Dietz K-J (2015) Redox-dependent translocation of the heat shock transcription factor AtHSFA8 from the cytosol to the nucleus in *Arabidopsis thaliana*. *FEBS Lett.* 589: 718-725. DOI: 10.1016/j.febslet.2015.01.039.

Glynn JM, Yang Y, Vitha S, Schmitz AJ, Hemmes M, Miyagishima S & Osteryoung K (2009). PARC6, a novel chloroplast division factor, influences FtsZ assembly and is required for recruitment of PDV1 during chloroplast division in *Arabidopsis*. *Plant J.*, 59: 700-711. DOI:10.1111/j.1365-313X.2009.03905.x.

Grabowski E, Miao Y, Mulisch M & Krupinska K (2008). Single-stranded DNA-binding protein Whirly1 in barley leaves is located in plastids and the nucleus of the same cell *Plant Physiol.*, 147: 1800-1804. DOI: 10.1104/pp.108.122796.

Gray JC, Hansen MR, Shaw DJ, Graham, K, Dale R, Smallman P, Natesan SKA, Newell, CA (2012) Plastid stromules are induced by stress treatments acting through abscisic acid. *Plant J* 69: 387-398. DOI: 10.1111/j.1365-313X.2011.04800.x.

Hanson MR & Hines KM (2018) Stromules: Probing form and function. *Plant Physiol.* 176: 128-137. DOI: 10.1104/pp.17.01287

Henderson GP, Gan L & Jensen GJ (2007) 3-D ultrastructure of *O. tauri*: electron cryotomography of an entire eukaryotic cell. *PLoS One* 2: e749. DOI:10.1371/journal.pone.0000749.

Higa T, Suetsugu N, Kong S and Wada M (2014) Actin-dependent plastid movement is required for motive force generation in directional nuclear movement in plants. *Proc Natl. Acad. Sci. USA* 111: 4327-4331. DOI: 10.1073/pnas.1317902111.

Ho J, Theg SM (2016) The formation of stromules *in vitro* from chloroplasts isolated from *Nicotiana benthamiana*. *PLoS One* 11: e0146489. DOI: 10.1371/journal.pone.0146489

Isemer R, Krause K, Grabe N, Kitahata N, Asami T & Krupinska K (2012) Plastid located WHIRLY1 enhances the responsiveness of *Arabidopsis* Seedlings toward abscisic acid. *Front. Plant Sci.* 3: 283. DOI: 10.3389/fpls.2012.00283.

Itoh RD, Ishikawa H, Nakajima KP, Moriyama S & Fujiwara MT (2018) Isolation and analysis of a stromule-overproducing *Arabidopsis* mutant suggest the role of *PARC6* in plastid morphology maintenance in the leaf epidermis. *Physiol. Plant.* 162: 474-494. DOI: 10.1111/ppl.12648

Iwabuchi K, Minamino R & Takagi S (2010) Actin Reorganization Underlies Phototropin-Dependent Positioning of Nuclei in *Arabidopsis* Leaf Cells. *Plant Physiol.* 152: 1309-1319. DOI: 10.1104/pp.109.149526.

Jung H-S, Crisp PA, Estavillo GM, Coleb B, Hong F, Mockler TC, Pogson BJ & Chory J (2013) Subset of heat-shock transcription factors required for the early response of *Arabidopsis* to excess light. *Proc. Natl. Acad. Science USA* 110: 14474-14479. DOI: 10.1073/pnas.1311632110.

- Keeling PJ (2010) The endosymbiotic origin, diversification and fate of plastids. *Phil. Trans. Royal Soc. B Biol. Sci.* 365: 729-748. DOI:10.1098/rstb.2009.0103.
- Köhler RH, Cao J, Zipfel WR, Webb WW & Hanson MR (1997) Exchange of protein molecules through connections between higher plant plastids. *Science* 276: 2039-2042. DOI: 10.1126/science.276.5321.2039.
- Kong SG and Wada M 2011 New insights into dynamic actin-based chloroplast photo-relocation movement. *Mol. Plant* 4: 771-781. DOI: 10.1093/mp/ssr061.
- Kumar AS, Park E, Nedo A, Alqarni A, Ren L, Hoban K, Modla S, McDonald JH, Kambhamettu C, Dinesh-Kumar SP & Caplan JL (2018) Stromule extension along microtubules coordinated with actin-mediated anchoring guides perinuclear chloroplast movement during innate immunity. *eLife* e23625. DOI: 10.7554/eLife.23625.
- Meyer AJ & Dick TP (2010) Fluorescent Protein-Based Redox Probes. *Antiox. Redox Signaling* 13: 621-650. DOI: 10.1089/ars.2009.2948.
- Mignolet-Spruyt L, Xu E, Idänheimo N, Hoerberichts FA, Mühlenbock P, Brosché M, Van Breusegem F, Kangasjärvi J (2016) Spreading the news: subcellular and organellar reactive oxygen species production and signalling. *J. Exp. Bot.* 67: 3831-3844. DOI:10.1093/jxb/erw080.
- Miller G & Mittler R (2006) Could Heat Shock Transcription Factors Function as Hydrogen Peroxide Sensors in Plants? *Ann. Bot.* 98: 279-288. DOI: 10.1093/aob/mcl107.
- Mubarakshina MM, Ivanov BN, Naydov IA, Hillier W, Badger MR, Krieger-Liszkay (2010) Production and diffusion of chloroplastic H₂O₂ and its implication to signalling. *J. Exp. Bot.* 61: 3577-3587. DOI: 10.1093/jxb/erq171.
- Mullineaux PM, Karpinski S and Baker NR (2006) Spatial dependence for hydrogen peroxide-directed signaling in light-stressed plants. *Plant Physiol* 141: 346-350. DOI: 10.1104/pp.106.078162
- Mullineaux PM, Exposito-Rodriguez M, Laissue PP & Smirnoff N (2018). ROS-dependent signalling pathways in plants and algae exposed to high light: Comparisons with other eukaryotes. *Free Rad. Biol. Med.* 122: 52-64. DOI: 10.1016/j.freeradbiomed.2018.01.033.
- Nakano Y & Asada K (1980) Spinach chloroplasts scavenge hydrogen peroxide on illumination. *Plant Cell Physiol.* 21: 1295-1307. DOI: 10.1093/oxfordjournals.pcp.a07612.
- Nietzel T, Elsässer M, Ruberti C, Steinbeck J, Ugalde JM, Fuchs P, Wagner S, Ostermann L, Moseler A, Lemke P, Fricker MD, Müller-Schüssele SJ, Moerschbacher BM, Costa A, Meyer AJ & Schwarzländer M (2019). The fluorescent protein sensor roGFP2-Orp1 monitors *in vivo* H₂O₂ and thiol redox integration and elucidates intracellular H₂O₂ dynamics during elicitor-induced oxidative burst in Arabidopsis. *New Phytol.* 221: 1649-1664. DOI:10.1111/nph.15550.
- Oikawa K, Kasahara M, Kiyosue T, Kagawa T, Suetsugu N, Takahashi F, Kanegae T, Niwa Y, Kadota A, Wada M (2003). Chloroplast unusual positioning1 is essential for proper chloroplast positioning. *Plant Cell* 15: 2805-2815. DOI: 10.1105/tpc.016428.

- Page MT, Kacprzak SW, Mochizuki N, Okamoto H, Smith AG, Terry MJ (2017) Seedlings lacking the PTM protein do not show a *genomes uncoupled* (*gun*) mutant phenotype Plant Physiol. 174, 21-26. doi: 10.1104/pp.16.01930.
- Pérez-Salamó I, Papdi C, Rigó G, Zsigmond L, Vilela B, Lumbreras V, Nagy I, Horváth B, Domoki M, Darula Z, Medzihradzky K, Bögre L, Koncz C & Szabados L (2014) The Heat Shock Factor A4A Confers Salt Tolerance and Is Regulated by Oxidative Stress and the Mitogen-Activated Protein. Plant Physiol. 165: 319-334. DOI: 10.1104/pp.114.237891.
- Pinheiro C & Chaves MM (2011) Photosynthesis and drought: can we make metabolic connections from available data? J. Exp. Bot. 62: 869-892. DOI: 10.1093/jxb/erq340
- Ramel F, Birtic S, Ginies C, Soubigou-Taconnat L, Triantaphylides C, Havaux M (2012). Carotenoid oxidation products are stress signals that mediate gene responses to singlet oxygen in plants. Proc. Natl. Acad. Sci. USA 109: 5535-5540. DOI: 10.1073/pnas.1115982109.
- Selga T, Selga M, Gobins V, Ozolina A (2010) Plastid-nuclear complexes: permanent structures in photosynthesizing of vascular plants. Environ. Exp. Biol. 8: 85-92. DOI: doi.org/10.22364/eeb.
- Sewelam N, Jaspert N, Van Der Kelen K, Tognetti VB, Schmitz J, Frerigmann H, Stahl E, Zeier J, Van Breusegem F & Maurino VG (2014). Spatial H₂O₂ Signaling Specificity: H₂O₂ from Chloroplasts and Peroxisomes Modulates the Plant Transcriptome Differentially. Mol. Plant 7: 1191-1210. DOI: 10.1093/mp/ssu070.
- Smirnoff N & Arnaud D (2019) Hydrogen peroxide metabolism and functions in plants. New Phytol. 221: 1197-1214. DOI: 10.1111/nph.15488.
- Spencer C & Foyer C (1986) The relationship between phosphate status and photosynthesis in leaves. Planta 167: 369-375. DOI: 10.1007/BF00391341
- Stael S, Kmiecik P, Willems P, Van Der Kelen K, Coll NS, Teige M & Van Breusegem F (2015). Plant innate immunity--sunny side up? Trends Plant_Sci. 20: 3-11. DOI: 10.1016/j.tplants.2014.10.002.
- Suetsugu N, Higa T, Gotoh E & Wada M (2016). Light-induced movements of chloroplasts and nuclei are regulated in both cp-actin-filament- dependent and -independent manners in Arabidopsis thaliana. PLoS ONE 11(6): e0157429. DOI: 10.1371/ journal.pone.0157429.
- Sun X, Feng P, Xu X, Guo, H, Ma J, Chi, W, Lin R, Lu C & Zhang L (2011). A chloroplast envelope-bound PHD transcription factor mediates chloroplast signals to the nucleus. Nat. Comm. 2, 477. DOI: 10.1038/ncomms1486.
- Vismans G, van der Meer T, Langevoort O, Schreuder M, Bouwmeester H, Peisker H, Dörman P, Ketelaar T & van der Krol A (2016) Low- phosphate induction of plastidal stromules is dependent on strigolactones but not on the canonical strigolactone signaling component MAX2. Plant Physiol 172: 2235–2244 DOI: 10.1104/pp.16.01146
- Walia A, Rainer W & Jones A (2018). Annu. Rev. Plant Biol. 69: 497-524. DOI: 10.1146/annurev-arplant-042817-040104.

Wen F, Xing D, Zhang L (2008). Hydrogen peroxide is involved in high blue light-induced chloroplast avoidance movements in *Arabidopsis*. *J. Exp. Bot.* 59: 2891-2901. DOI: 10.1093/jxb/ern147.

Willems P, Mhamdi A, Stael S, Storme V, Kerchev P, Noctor G, Gevaert K, Van Breusegem Frank (2016). The ROS Wheel: Refining ROS Transcriptional Footprints. *Plant Physiol.* 171: 1720-1733. DOI: 10.1104/pp.16.00420.

Xiao Y, Savchenko T, Baidoo EEK, Chehab WE, Hayden DM, Tolstikov V, Corwin JA, Kliebenstein DJ, Keasling JD & Dehesh K (2012). Retrograde signaling by the plastidial metabolite MEcPP regulates expression of nuclear stress-response genes. *Cell* 149: 1525–1535. DOI: 10.1016/j.cell.2012.04.038.

Yao N & Greenberg JT (2006). *Arabidopsis* ACCELERATED CELL DEATH2 modulates programmed cell death. *Plant Cell*.18: 397-411. DOI:10.1105/tpc.105.036251.

Figure captions

Fig. 1. Nucleo-plastid association in *Arabidopsis thaliana* and *N. benthamiana*. All chloroplasts are magenta, all nuclei green. All scale bars 10 μ m. A) Nuclei are surrounded by chloroplasts in the typical 'daisy flower' arrangements in *N.benthamiana* abaxial epidermal cells. B) In the spongy mesophyll of *Arabidopsis*, nuclei are in contact with but not surrounded by chloroplasts. C) A nucleus with surrounding chloroplasts from *N. benthamiana* abaxial epidermal cells, displaying occasionally observed stromules under low light conditions.

Fig. 2. Stromule-mediated perinuclear clustering (PNC) of the chloroplast during plant immune responses.

NRIP1(cTP)-TagRFP (magenta) were transiently expressed to visualize chloroplasts and stromules in GFP-TUA6 (green) transgenic *N. benthamiana* leaf epidermal cells. Images are 6 representative images in indicated time points from the Movie S2. When *N. benthamiana* leaf epidermal cells are infected by *Pseudomonas syringae*, stromules are vigorously induced and attached to the nucleus. Dynamic stromule retractions bring about chloroplast body movement toward the nucleus (yellow arrow) and extension of stromules also occurs to withdraw the chloroplast body from the nucleus (white arrowhead), controlling the extent of the PNC during plant immunity.

Supplementary material

Movie S1. Changing views of plastid-nuclear complexes in *N. benthamiana* abaxial epidermal cells. All chloroplasts are false-coloured magenta, all nuclei green. Nuclei and the cytosol are expressing HyPer as described previously (Exposito-Rodriguez et al 2017).

Movie S2. Stromules mediate chloroplast movement towards and away from the nucleus. Chloroplasts and stromules were visualized by transient expression of NRIP1(cTP)-TagRFP (magenta) in GFP-TUA6 (green) transgenic *N. benthamiana* leaf epidermal cells. 8 hours after *Pseudomonas syringae* infection, time-lapsed images were acquired every 5 min and compiled in AVI format to run in 6 frames per second. Representative images are shown in Figure 2.