

Complete Genome Sequence of *Campylobacter fetus* subsp. *venerealis* Biovar Intermedius, Isolated from the Prepuce of a Bull

Gregorio Iraola,^{a,b} Ruben Pérez,^a Hugo Naya,^{b,c} Fernando Paolicchi,^d David Harris,^e Trevor D. Lawley,^e Natalia Rego,^b Martín Hernández,^a Lucía Calleros,^a Luis Carretto,^a Alejandra Velilla,^d Claudia Morsella,^d Alejandra Méndez,^d Andrea Gioffre^f

Sección Genética Evolutiva, Facultad de Ciencias, Montevideo, Uruguay^a; Unidad de Bioinformática, Institut Pasteur de Montevideo, Montevideo, Uruguay^b; Departamento de Producción Animal y Pasturas, Facultad de Agronomía, Universidad de la República, Montevideo, Uruguay^c; Laboratorio de Bacteriología, Unidad Integrada INTA-Universidad Nacional de Mar del Plata, Balcarce, Argentina^d; Wellcome Trust, Sanger Institute, Hinxton, Cambridgeshire, United Kingdom^e; Instituto de Biotecnología, Instituto Nacional de Tecnología Agropecuaria, Buenos Aires, Argentina^f

***Campylobacter fetus* subsp. *venerealis* is the causative agent of bovine genital campylobacteriosis, a sexually transmitted disease distributed worldwide. *Campylobacter fetus* subsp. *venerealis* biovar Intermedius strains differ in their biochemical behavior and are prevalent in some countries. We report the first genome sequence for this biovar, isolated from bull prepuce.**

Received 12 June 2013 Accepted 24 June 2013 Published 1 August 2013

Citation Iraola G, Pérez R, Naya H, Paolicchi F, Harris D, Lawley TD, Rego N, Hernández M, Calleros L, Carretto L, Velilla A, Morsella C, Méndez A, Gioffre A. 2013. Complete genome sequence of *Campylobacter fetus* subsp. *venerealis* biovar Intermedius, isolated from the prepuce of a bull. *Genome Announc.* 1(4):e00526-13. doi:10.1128/genomeA.00526-13.

Copyright © 2013 Iraola et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](http://creativecommons.org/licenses/by/3.0/).

Address correspondence to Gregorio Iraola, goyo1987@gmail.com.

Campylobacter fetus is an important veterinary pathogen. This species is currently divided into two subspecies, *Campylobacter fetus* subsp. *fetus*, causative of abortion in sheep, and *Campylobacter fetus* subsp. *venerealis*, the etiologic agent of bovine genital campylobacteriosis (1), a disease that has spread worldwide and causes economic losses mainly in countries where natural breeding is frequent (2). A distinct group of *C. fetus* strains known as *C. fetus* subsp. *venerealis* biovar Intermedius has also been determined; these strains phenotypically resemble *C. fetus* subsp. *venerealis*, but they react positively to the H₂S test (typically positive for *C. fetus* subsp. *fetus*) (3). In recent years, an increase in the prevalence of this biovar has been noticed in some countries (e.g., South Africa) (4). However, the lack of genomic information for these atypical strains has hindered the development of molecular diagnostic tools and the study of their genomic evolution. Here we present the first complete genome sequence for *Campylobacter fetus* subsp. *venerealis* biovar Intermedius INTA 99/541, isolated from the prepuce of a naturally infected bull (5).

Sequencing was performed on an Illumina Hi-Seq 2000 platform and generated 13,953,630 paired-end reads (2 × 100 cycles). The resulting library was first corrected using ALLPATHS-LG (6) and then assembled with Velvet (7) software, producing 111 contigs with an average coverage of 130-fold. The assembly quality was improved using the PAGIT toolkit (8), based on the genome sequence of *C. fetus* subsp. *fetus* 82-40 (accession no. NC_008599) as the reference. The final assembly quality was evaluated with an assembly likelihood estimator (ALE) (9). The resulting pseudomolecule produced by contig scaffolding was automatically annotated with RAST (<http://rast.nmpdr.org/>).

Campylobacter fetus subsp. *venerealis* biovar Intermedius INTA 99/541 has a circular chromosome of 1,774,509 bp with an average GC content of 33%, including 2,421 putative protein-coding open reading frames (1.36 genes per kb), 3 rRNA operons,

and 40 tRNA genes. BLAST analysis between those contigs that were not used in chromosome scaffolding and the GenBank plasmids database revealed high sequence homology with the pTet *Campylobacter jejuni* plasmid (accession no. NC_008790) and strong synteny conservation of Cpp protein-coding genes, important for plasmid mobilization (10). However, more extensive analyses are needed to confirm that this strain is carrier of extra-chromosomal replicons.

Comparison between *C. fetus* subsp. *venerealis* biovar Intermedius INTA 99/541, *C. fetus* subsp. *venerealis* Azul-94 (10), *C. fetus* subsp. *venerealis* NTCT 10354^T (11), and *C. fetus* subsp. *fetus* 82-40 genomes using the Artemis comparison tool (12) revealed that sequence identity and synteny are conserved along genomes. Further analysis of these genomes will provide information regarding the basis of the unique physiological and biochemical features of *C. fetus* subsp. *venerealis* biovar Intermedius. Moreover, the availability of the first genome from this organism is an important achievement in the development of specific molecular tools for diagnosis and will shed light on the genomic evolution of *Campylobacter* species, although a representative number of genomes for this biovar will be needed to conduct more robust comparisons.

Nucleotide sequence accession numbers. This whole-genome shotgun project has been deposited at DDBJ/EMBL/GenBank under the accession number [ASTK00000000](https://www.ncbi.nlm.nih.gov/nuccore/ASTK00000000). The version described in this paper is version [ASTK01000000](https://www.ncbi.nlm.nih.gov/nuccore/ASTK01000000).

ACKNOWLEDGMENTS

This work was funded by Wellcome Trust grant 098051, Comisión Sectorial de Investigación Científica (C. S. I. C.), and INTA Argentina Projects PE 242121 and PNSA-1115053.

We thank Gordon Dougan and Julian Parkhill for their useful advice during this work.

REFERENCES

1. Debruyne L, Gevers D, Vandamme P. 2008. Taxonomy of the family *Campylobacteraceae*, p 3–25. In Nachamkin I, Szymanski CM, Blaser MJ (ed), *Campylobacter*, 3rd ed. ASM Press, Washington, DC.
2. Mshelia GD, Amin JD, Woldehiwet Z, Murray RD, Egwu GO. 2010. Epidemiology of bovine venereal campylobacteriosis: geographic distribution and recent advances in molecular diagnostic techniques. *Reprod. Domest. Anim.* 45:e221–e230. doi:10.1111/j.1439-0531.2009.01546.x.
3. Véron M, Chatelain R. 1973. Taxonomic study of the genus *Campylobacter* Sebald and Véron and designation of the neotype strain for the type species, *Campylobacter fetus* (Smith and Taylor) Sebald and Véron. *Int. J. Syst. Bacteriol.* 23:122–134.
4. Schmidt T, Venter EH, Picard JA. 2010. Evaluation of PCR assays for the detection of *Campylobacter fetus* in bovine preputial scrapings and the identification of subspecies in South African field isolates. *J. S. Afr. Vet. Assoc.* 81:81–92.
5. van Bergen MAP. 2005. Ph.D. thesis. Utrecht University, Utrecht, Netherlands.
6. Butler J, MacCallum I, Kleber M, Shlyakhter IA, Belmonte MK, Lander ES, Nusbaum C, Jaffe DB. 2008. ALLPATHS: de novo assembly of whole-genome shotgun microreads. *Genome Res.* 18:810–820.
7. Zerbino DR, Birney E. 2008. Velvet: algorithms for de novo short read assembly using de Bruijn graphs. *Genome Res.* 18:821–829.
8. Swain MT, Tsai IJ, Assefa SA, Newbold C, Berriman M, Otto TD. 2012. A post-assembly genome-improvement toolkit (PAGIT) to obtain annotated genomes from contigs. *Nat. Protoc.* 7:1260–1284.
9. Clark SC, Egan R, Frazier PI, Wang Z. 2013. ALE: a generic assembly likelihood evaluation framework for assessing the accuracy of genome and metagenome assemblies. *Bioinformatics* 29:435.
10. Moolhuijzen PM, Lew-Tabor AE, Wlodek BM, Agüero FG, Comerci DJ, Ugalde RA, Sanchez DO, Appels R, Bellgard M. 2009. Genomic analysis of *Campylobacter fetus* subspecies: identification of candidate virulence determinants and diagnostic assay targets. *BMC Microbiol.* 9:86.
11. Stynen AP, Lage AP, Moore RJ, Rezende AM, de Resende VD, Ruy Pde C, Daher N, Resende Dde M, de Almeida SS, Soares Sde C, de Abreu VA, Rocha AA, dos Santos AR, Barbosa EG, Costa DF, Dorella FA, Miyoshi A, de Lima AR, Campos FD, de Sá PG, Lopes TS, Rodrigues RM, Carneiro AR, Leao T, Cerderia LT, Ramos RT, Sliva A, Azevedo V, Ruiz JC. 2011. Complete genome sequence of type strain *Campylobacter fetus* subsp. *venerealis* NCTC 10354T. *J. Bacteriol.* 193:5871–5872.
12. Carver TJ, Rutherford KM, Berriman M, Rajandream MA, Barrell BG, Parkhill J. 2005. ACT: the Artemis comparison tool. *Bioinformatics* 21:3422–3423.