

[54] **RESTORATION OF IMPAIRED CARDIAC FUNCTION OF PATIENTS WITH DIVERSE MUSCULAR DYSTROPHIES BY THERAPY WITH COENZYME Q10**

[75] **Inventors:** Karl Folkers, Austin, Tex.; Janusz Wolaniuk, Indianapolis, Ind.

[73] **Assignee:** Board of Regents, The University of Texas System, Austin, Tex.

[21] **Appl. No.:** 155,891

[22] **Filed:** Feb. 16, 1988

Related U.S. Application Data

[63] Continuation-in-part of Ser. No. 725,059, Apr. 19, 1985, abandoned.

[51] **Int. Cl.⁴** **A61K 37/48**

[52] **U.S. Cl.** **424/94.1; 514/907**

[58] **Field of Search** **424/94.1; 514/907**

[56] **References Cited**

U.S. PATENT DOCUMENTS

4,068,003	1/1978	Miyata	424/94.1
4,156,718	5/1979	Blizakov	424/94.1
4,491,594	1/1985	Ogawa et al.	424/94.1

OTHER PUBLICATIONS

Folkers et al., "Evidence for a Deficiency of Coenzyme Q₁₀ in Human Heart Disease", *International Journal for Vitamin Research*, 1970, pp. 380-390.

Ganong, W. F., "The Heart as a Pump", In: *Review of Medical Physiology*, Los Altos, Lange Medical Publication, pp. 441-451 (1981).

Baker et al., "Use of Electrical Impedance to Measure Cardiac Function In Studies of CoQ₁₀", Elsevier/North-Holland Biomedical Press, 1981, pp. 193-206.

Vadhanavikit et al., "Micro-Analysis for Coenzyme Q₁₀ in Endomyocardial Biopsies of Cardiac Patients and Data on Bovine and Canine Hearts", *Biochemical and Biophysical Resrch. Comm. Vo. 123, No. 3, 1984* pp. 1165-1169.

Mortensen et al., "Deficiency of Coenzyme Q₁₀ in Myocardial Failure," *Drugs Exptl. Clin. Res. (7)*, 1984, pp. 497-502.

Folkers et al., "Biochemical Rationale and The Cardiac Response of Patients With Muscle Disease To Therapy With Coenzyme Q₁₀", *Proc. Natl. Acad. Sci. USA*, vol. 82, Jul. 1985, pp. 4513-4516.

DePierre et al., "Enzyme Topology of Intracellular Membranes", *Ann. Rev. Biochem.*, 1977, pp. 201-262.

Crane et al., "Isolation of a Quinone From Beef Heart Mitochondria", *Biochem. Biophys. Acta.*, vol. 25, 1957, pp. 220-221.

Saviotti et al., "Electrocardiography in Duchenne Muscular Dystrophy", *G. Neuropsychiatr Eta Evol.*, 1984, 4/Suppl. 1 (67-69), p. 337.

Venco et al., "The Heart in Progressive Muscular Dystrophy", *G. Neuropsychiatr Eta Evol.*, 1984, 4/Suppl. 1 (61-66), p. 341.

Carafoli et al., "Molecular Aspects of Medicine" (1980) *Molec. Aspects of Med.*, V3, pp. 295-429.

Sunamori et al., "Clinical Application of Coenzym Q

To Coronary Bypass Graft Surgery" (1984) *Biomed. and Clin, Aspects of Coenzyme Q*, ed Folkers et al., V4, pp. 333-342.

Michelson et al., "Dystrophia Muscularis: A Hereditary Primary Myopathy In the House Mouse", 1955) *Proc. Nat'l. Acad. Sci.*, V 41, pp. 1079-1084.

West, "Hereditary Mouse Muscular Dystrophy With Particular Emphasis on Pathogenesis and Attempts At Therapy", (1966) *Annals. N.Y. Accas. Of Sciences*, V 138, pp. 4-13.

Welsh et al., "Cardiac Findings in 73 Patients with Muscular Dystrophy", (1963) *Arch. Int. Med.*, V 112, pp. 97-104.

Durnin et al., "The Electrocardiogram in Pseudohypertrophic or Duchenne's Muscular Dystrophy (Type IIIa)", (1960) *J. Iowa Med. Soc.*, V 59, pp. 113-115.

Morgan-Hughes, "Disorders of Mitochondrial Metabolism Some Clinical and Biochemical Mechanisms", (1983), *Advanced Med.*, V 19 ed by Saunders, pp. 243-260.

Farley et al., "Response Of Genetically Dystrophic Mice to Therapy With Hexahydrocoenzyme Q₄", (196) *Biochem. and Biophys. Res. Comm. V 24*, pp. 299-303.

Nilsson et al., "Biosynthesis and Levels of Coenzyme Q in Genetically Dystrophic Mice", (1968) *Arch. Biochem. and Biophys.* V 133, pp. 422-426.

Lenaz et al., "Organic Structural Specificity and Sites of Coenzyme Q in Succinoxidase and DPNH-Oxidase Systems", (1968) *Arch. Biochem. and Biophys.*, V 123 pp. 539-550.

(List continued on next page.)

Primary Examiner—John W. Rollins

Attorney, Agent, or Firm—Arnold, White & Durkee

[57] **ABSTRACT**

The present invention relates to the use of Coenzyme Q in the treatment of slow muscle degeneration, commonly known to those of skill in the art so a dystrophy or atrophy, and the accompanying cardiac complications typically identified in such patients. Administration of Coenzyme Q, and particularly the analog Coenzyme Q₁₀ (CoQ₁₀) to humans increases the pumping of blood by the heart, and thereby increases tissue oxygenation throughout the body. The net physiological effect halts the progression of muscle deterioration and improves cardiac function. An overall improvement in the quality of life for these human subjects is also observed, said patients reportedly experiencing less fatigue.

A method for treating human patients with progressive muscular dystrophies or the neurogenic atrophies with Coenzyme Q₁₀ (CoQ₁₀) specifically disclosed. The method is similarly effective for the treatment of any form of muscle degeneration or cardiac muscular dysfunction independently.

1 Claim, No Drawings

OTHER PUBLICATIONS

- Scholler et al., "Therapeutic Activity of Coenzyme Q for Reproduction", (1968) *International J. for Vit. Res.*, V 38, pp. 362-368.
- Littarru et al., "Deficiency of Coenzyme Q₉ in Mice Having Hereditary Muscular Dystrophy", (1970) *Biochem. and Biophys. Res. Comm.*, V 41, pp. 1306-1313.
- Danowski et al., "Tabulation of Findings in the Muscular Dystrophies and in Myotonia Dystrophica", (1971) (I) *Arch. of Physical Med. and Rehab.*, V 52, pp. 193-200.
- Sovik et al., "Coenzyme Q in Duchenne Muscular Dystrophy", (1971) *Acta Pediat. Scand.*, V 60, pp. 428-432.
- Danowski et al., "Hexahydrocoenzyme Q₄ in Pseudohypertrophic Muscular Dystrophy", (1971) II *J. Pharmacent. Sci.*, V 60, pp. 640-642.
- Zellweger et al., "The Diagnostic Significance of Serum Enzymes and Electrocardiogram in Various Muscular Dystrophies", (1972) *Internat. J. Vit. Nutr. res.*, V 42, pp. 139-163.
- Folkers et al., "Survey and New Clinical Studies on Coenzyme Q in Human Muscular Dystrophy", (1972) *Internat. J. Vit. Nutr. Res.*, V 42, pp. 139-163.
- Griggs (1974) *Circul. Res.*, V 34 and 35, pp. II-145-II-150.
- Folkers et al., "Effect of Coenzyme Q on Serum Levels of Creatine Phosphokinase in Preclinical Muscular Dystrophy", (1974) *Circul. Res.*, V 34 and 35, pp. II-145-II-150.
- Kuhn et al., "Early Myocardial Disease and Cramping Myalgia in Becker-Type Muscular Dystrophy: A Kindred", (1979) *Neurology*, V 29, pp. 1144-1149.
- Reeves et al., "Echocardiographic Evaluation of Cardiac Abnormalities in Duchene's Dystrophy and Myotonic Muscular Dystrophy", (1980) *Arch. Neurol.*, V 37, pp. 273-277.
- Hopkins et al., "Emery-Dreifuss Humeroperoneal Muscular Dystrophy: An X-linked Myopathy with Unusual Contractures and Bradycardia", *Annals of Neurol.* (1981), V 10, pp. 230-237.
- Kishi et al., "Estimation of Plasma Levels of Coenzyme Q₁₀ and Relationship to Oral Dosage", (1981) *Biomed. and Clin. Aspects of Coenzyme Q*, V 3, eds Folders et al., pp. 67-78.
- Couch et al., "Coenzyme Q In Therapy of Limb-Girdle Muscular Dystrophy", (1981) *ibid.*, pp. 257-266.
- Folkers et al., "New Progress on the Biomedical and Clinical Research on Coenzyme Q", (1981) *ibid.*, pp. 399-412.
- Goldberg et al., "Serial Two-Dimensional Echocardiography in Duchenne Muscular Dystrophy", (1982) *Neurology*, V 32 pp. 1101-1105.
- Hawley et al., "Families With Myotonic Dystrophy With and Without Cardiac Involvement", (1983) *Arch. Int. Med.*, V 143, pp. 2134-2136.
- Bender, "Diagnosis: Muscular Dystrophy", (1984) *Hospital Med. Hanuary*, pp. 95-125.
- Morgan-Hughes et al., "Mitochondrial Myopathies Results of Exploratory Therapeutic Trials", (1984) *Biomed and Clin. Aspects of Coenzyme Q*, Elsevier Science Pub., V 4, pp. 417-424.
- Ho et al., "Genetic Dystrophy in Mice, A Possible Coenzyme Q Dependency State", (1975) *Abstract of Presented Paper*.
- Kubicek et al., "Development and Evaluation of an Impedance Cardiac Output System", (1966) *Aerospace Med.* (Dec. 1966) pp. 1208-1212.

**RESTORATION OF IMPAIRED CARDIAC
FUNCTION OF PATIENTS WITH DIVERSE
MUSCULAR DYSTROPHIES BY THERAPY WITH
COENZYME Q₁₀**

This application is a continuation-in-part of application Ser. No. 06/725,059 filed 04/19/85 now abandoned.

THE INVENTION

In January, 1984, (Bender, A. N., *Hospital Medicine*, p. 95, January, 1984), stated:

—“The term muscular dystrophy applies to a variable group of disorders—some well defined, and others less well defined—categorized by general features—. All are primarily diagnosed on clinical and genetic grounds, and none of them is treatable at the present time. Accurate diagnosis allows realistic prognosis, effective rehabilitation, and prevention of further cases through genetic counseling.”

Griggs, M. D., (Supplement II to Circulation Research, vols. 34 and 35) has stated that:

—“Cardiac disease appears to be the rule rather than the exception in virtually every form of hereditary and acquired skeletal myopathy.”

We have made three related discoveries on the administration of coenzyme Q₁₀ (vitamin Q₁₀) to patients with diverse muscular dystrophies and diverse neurogenic atrophies.

The first discovery was the recognition that, *in spite of the unknown and variable genetic defects, a patient with a form of muscular dystrophy or a patient with a form of neurogenic atrophy, could have the impaired cardiac function be significantly improved by treatment with coenzyme Q₁₀ and not improved by treatment with a matching placebo.*

The second discovery was the recognition that *all patients with eight diverse forms of muscular dystrophies and neurogenic atrophies had an increase in their impaired cardiac function by treatment with coenzyme Q₁₀. It was almost unbelievable that all patients with such diverse genetic diseases should all respond by this therapy and show a significant increase in the pumping of the blood by their hearts.*

The third discovery was the recognition that the *impaired cardiac function of many such patients with such diverse disease was not only improved, but was restored to a normal level of cardiac function. Such restoration was not even considered. It was thought that a significant increase of the impaired cardiac function to a level below that of the normal range might be possible.*

Reduction to practice may be considered as achieved only by the two double blind trials as required by the Food and Drug Administration. The two double blind trials constitute the Reduction to Practice.

These three related discoveries constitute a basis for the Food and Drug Administration to grant an NDA for the clinical use of CoQ₁₀ as an orphan drug for months and years of administration and on the basis of complete safety. Since such patients have their genetic defects for a life-time, they may be safely treated with CoQ₁₀ for their life-time. A pharmaceutical company may now proceed to market CoQ₁₀ as an orphan drug

for such patients with an NDA from the F&DA, and with the protection of a use patent.

SUMMARY OF THE APPLICATION

5 It is known that both cardiac and skeletal muscle contract.

It is known that the contraction of both cardiac and skeletal muscle require energy or “bioenergetics.”

10 Skeletal muscle comprises the great mass of the somatic musculature and has well-developed cross striations, and does not normally contract in the absence of nervous stimulation. Skeletal muscle is generally under voluntary control.

Cardiac muscle does have cross-striations, but is functionally syncytial in character and contracts rhythmically in the absence of external innervation owing to the presence in the myocardium of pacemaker cells that discharge spontaneously. Cardiac muscle contains far larger numbers of elongated mitochondria in close contact with the fibrils than does skeletal muscle.

20 All forms of dystrophy and atrophy involve genetic defects and these defects vary widely as evidenced by the widely differing symptoms of all of the forms of dystrophy and atrophy or myopathy. These genetic defects have been studied on skeletal muscle because skeletal muscle is available by biopsy in contrast to cardiac muscle which is very difficult to obtain from living patients for ethical reasons. Consequently, little or nothing is known about the differences in genetic defects between cardiac and skeletal muscle although such differences very probably must exist.

30 Coenzyme Q₁₀ is known to provide clinical benefit to patients with congestive heart failure but who have no muscular dystrophy or atrophy or myopathy. There was no basis to predict whether coenzyme Q₁₀ would or would not be beneficial to the cardiac dysfunction of patients with these muscular diseases and because of the great diversity of the genetic defects and the great diversity of the clinical symptoms of all these diseases. It appeared probable that any clinical usefulness of administering coenzyme Q₁₀ to such patients would be very limited or negative entirely, because of the apparent and diverse genetic defects.

There was no assurance that CoQ₁₀ would benefit the cardiac function of dystrophic patients. The first double-blind and open cross-over trial on the oral administration of coenzyme Q₁₀ (CoQ₁₀) to 12 patients with progressive muscular dystrophies and neurogenic atrophies was conducted. These diseases included the Duchenne, Becker and, limb-girdle dystrophies, myotonic dystrophy, Charcot-Marie tooth disease and Welander disease. Cardiac function was noninvasively and extensively monitored by impedance cardiography. Solely by significant change or no change in stroke volume and cardiac output, all eight patients on blind CoQ₁₀ and all four on blind placebo were correctly assigned, $p < 0.003$. The monitoring of cardiac function of these 12 patients before the trial revealed degrees of impaired cardiac function ranging from relatively severe to definite to low impairment.

65 It was discovered that the oral administration of CoQ₁₀ gave a substantial and surprising improvement to the impaired cardiac function. This initial double blind trial was only on 12 patients, 8 on blind CoQ₁₀ and 4 on blind placebo. It was apparent that even though the cardiac response was significant, $p < 0.003$, the two groups of 8 patients and 4 patients, respectively, had too few patients to support biologically the discovery of a

positive clinical benefit to cardiac function, when the possibility of such a discovery was so questionable. In a second double blind trial with a total of 15 similar patients, 8 on blind CoQ₁₀ and 7 on blind placebo, the discovery was confirmed. All 8 patients on blind CoQ₁₀ showed significant cardiac improvement and all 7 on blind placebo were negative. The results of the two trials when combined involved 16 patients on blind CoQ₁₀ and 11 patients on blind placebo which is sufficient to document the discovery that CoQ₁₀ benefits the cardiac function of dystrophic patients.

Two more discoveries resulted from this research. The second discovery was that CoQ₁₀ therapy benefited eight diverse forms of dystrophy and myopathy which is remarkable since these eight dystrophies and myopathies are genetically different and reveal very different clinical symptomatology including early death for the Duchenne dystrophy (up to 20 years of age) and decades of life for other dystrophies such as the Charcot-Marie tooth disease.

The third discovery from the second double blind trial was that the cardiac functions of some of the patients were restored to a normal range. A higher dosage and longer treatment could likely increase the cardiac function of even more of the patients.

Consequently, the oral administration of coenzyme Q₁₀ to patients with very diverse forms of muscular dystrophy and neurogenic atrophy is very useful, because the resulting improved cardiac function definitely benefits the health through more blood providing more oxygenation throughout the body. Since CoQ₁₀ is exceptionally safe, this therapy for patients with such disease may be used as long as required or even for a life-time.

BACKGROUND ON INVENTION STUDIES

Coenzyme Q in Mice with Genetic Dystrophy

Lack of Significance of Animal Dystrophy. There have been many studies involving mice based on the biochemistry of CoQ, but there is no absolute certainty of the relationship of a genetic disease in mice in comparison with that of man. It hardly seems that the genetic defects of mice and man are truly identical, although there can be superficial similarities. One fact proving a difference rather than identity is the existence of CoQ₉ in mice and CoQ₁₀ in man. Obviously, a promising benefit for mice with a genetic disease does not guarantee a response in man by the same treatment.

An account of studies with CoQ in mice with genetic dystrophy are as follows.

Michelson, A. M., Russel, E. S., and Pinckey, K. H., (*Proc. Natl. Acad. Sci. USA* 41, 1079 (1955)), described a mutation of mice that have a muscular degeneration which resembles progressive muscular dystrophy in humans. West, W. T., Meier, H., and Hoag, W. G., (*Ann. N.Y. Acad. Sci.*, 138, 4 (1966)) considered that dystrophy occurred as a spontaneous autosomal mutation in mice 129/Re which causes muscular weakness, atrophy and a reduced life-span. These mice showed clinical, histological, and physiological similarities to juvenile progressive pseudohypertrophic and myotonic dystrophy of man, Erb's dystrophy and, except for the difference in inheritance, the Duchenne dystrophy.

Dystrophic mice which were treated with hexahydrocoenzyme Q₄ (H₆CoQ₄) improved so that severely dystrophic animals responded and were able to walk using all their legs; (Farley, T. M., Scholler, J., and Folkers, K., *Biochem. Biophys. Res. Commun.* 24, 299

(1966)). A study of the biosynthesis and levels of CoQ in these mice indicated a possible genetic defect in the biosynthesis of p-hydroxybenzoic acid (HBA) from tyrosine rather than a defect between CoQ and HBA or some other defect involving CoQ; (Nilsson, L. G., Farley, T. M., Scholler, J., and Folkers, K., *Arch. Biochem. Biophys.* 123, 422 (1968)). In later studies, these dystrophic mice were treated with H₆CoQ₄ when they became unable to use their hind limbs. After two weeks of therapy, their improved condition allowed the use of one or both hind legs, and survival was four times that of control mice; (Scholler, J., Farley, T. M., Folkers, K., *Intern. J. Vitamin Res.* 38, 369 (1968)). This dystrophy was studied as a possible coenzyme Q dependency state; (Ho, L., Folkers, K., *Abstract for 27th Meeting of Southwest Region of American Chemical Society* 88, 38, San Antonio, Tex., December, 1971). Determination of the specific activities of the succinate dehydrogenase-coenzyme Q₉ reductase of hearts and hind leg muscles of dystrophic mice revealed a deficiency, $p < 0.01$, of the intrinsic CoQ₉; (Littarru, G. P., Jones, D., Scholler, J., and Folkers, K., *Biochem. Biophys. Res. Commun.* 41(5), 1306 (1970)). When these dystrophic mice were curatively treated with CoQ₇, instead of the intrinsic CoQ₉, in the fifth month of their life span of eight months, survival increased to twice that of the control group. CoQ₇ was isolated from the mitochondria of the hearts and hind legs of the orally treated mice, substantiating that the CoQ₇ was substituting for and correcting the deficiency of CoQ₉ in the dystrophy.

Studies with Coenzyme Q on Patients with Genetic Muscle Disease.

Significance of Open Very Limited Trials. The exploratory trials of administering H₆CoQ₄ to humans with dystrophy were negative. Couch, J. R. and Folkers, K. (*Biomedical and Clinical Aspects of Coenzyme Q*, Vol 3, K. Folkers and Y. Yamamura, eds., Elsevier/North Holland Biomedical Press, Amsterdam, 257-266, (1981)) administered CoQ₁₀ to 2 adults with limb-girdle dystrophy and recorded a transient small improvement in strength, and which appeared to slow deterioration. Such observation is clinically meaningless because the transient small improvement could have been the variation in the symptoms of the disease which are known to occur with the limb-girdle dystrophy and which is not as fatal as Duchenne. It is well known that the symptoms of this dystrophy decrease and improve with time and independent of any treatment.

Folkers, K., Baker, L., Richardson, P. C., Shizukui-shi, S., Takemura, K., Drzewoski, J., Lewandowski, J., Ellis, J., (*Biomedical and Clinical Aspects of Coenzyme Q*, Vol. 3, K. Folkers and Y. Yamamura, eds., Elsevier/North Holland Biomedical Press, Amsterdam, 399-412 (1981)) reported the administration of CoQ₁₀ to a single adult with a late onset form of muscle disease and observed that the expected deterioration appeared to be retarded. Again, such observation is clinically meaningless, because one does not know what the rate of deterioration would have been in the absence of administration of CoQ₁₀.

Again, the patient with this mild dystrophy which is not fatal until late decades is known to have decreased symptoms and increased symptoms over time without any treatment.

Danowski, T. S., Wissinger, H. A., Hohmann, T. C., Gerneth, J. A., Folkers, K., Vester, J. W., and Fisher,

E. R., (*Archives of Physical Medicine and Rehabilitation* 52, 193 (1971)), clinically studied the administration of H_6CoQ_4 in the muscular dystrophies and in myotonia dystrophica, but no clinical benefit was observed in the human pseudohypertrophic muscular dystrophy. Sovik, O., Stromme, J., Folkers, K., (*Acta Paediat. Scand.* 60, 428 (1971)), administered H_6CoQ_4 to four cases of Duchenne muscular dystrophy. No improvement in muscular strength was observed, but there was a statistically significant decrease of CPK and aldolase for one boy. Danowski, T. S., Folkers, K., Wissinger, H. A., Hohmann, T. C., Gerneth, J. A., and Vester, J. W., (*Journal of Pharm. Sciences* 60, 640 (1971)) investigated the administration of H_6CoQ_4 to 19 boys with pseudohypertrophic muscular dystrophy of the Duchenne type. A battery of clinical and laboratory indices were monitored. No beneficial change was observed. The failure of H_6CoQ_4 to elicit a clinical effect in humans, but to do so in mice, might be partly explained by biochemical data that showed H_6CoQ_4 has only about 10-15% of the activity of CoQ_{10} for DPNH-oxidase; (Lenaz, G., Daves, G. D., Folkers, K., *Arch. Biochem. Biophys.* 123, 539 (1968)), by dosage, species difference and protocol. Folkers, K., Littarru, G. P., Nakamura, R., and Scholler, J., (*Internat. J. Vit. Nutr. Res.* 42, 139 (1972)) used new enzyme methodology on muscle biopsies of dystrophic patients, and found that the succinate dehydrogenase-coenzyme Q_{10} reductase was inactive for some patients and poorly active for others, which might result in no or poor response to treatment with CoQ . However, administration of CoQ and placebo appropriately reduced CPK. They emphasized the desirability of treatment with CoQ of pre-clinical stages of dystrophy, and Folkers, K., Nakamura, Littarru, G. P., Zellweger, H., Brunkhorst, J. B., Williams, C. Y., Langston, J. H., (*Proc. Nat'l. Acad. Sci.* 71(5), 2098-2102 (1974)) did administer H_6CoQ_4 and CoQ_7 to children with pre-clinical dystrophy. Extensive monitoring of enzyme levels revealed significant reductions of CPK. Couch, J. R., and Folkers, K., (*Biomedical and Clinical Aspects of Coenzyme Q*, Vol. 3, K. Folkers and Y. Yamamura, eds., Elsevier/North Holland Biomedical Press, Amsterdam, 257-266, (1981)) administered CoQ_{10} to two adults with limb-girdle dystrophy, and recorded a transient small improvement in strength, which appeared to slow deterioration. Folkers, K., Baker, L., Richardson, P. C., Shizukuishi, S., Takemura, K., Drzewoski, J., Lewandowski, J., Ellis, J., (*Biomedical and Clinical Aspects of Coenzyme Q*, Vol. 3, K. Folkers and Y. Yamamura, eds., Elsevier/North Holland Biomedical Press, Amsterdam, 399-412 (1981)) reported the administration of CoQ_{10} to an adult with a late onset form of muscle disease for six years, and the expected deterioration of physical competence was markedly retarded.

Association of Muscular Disease with Cardiac Disease.

Welsh, J. D., Lynn, T. N., Haase, G. R., (*Archives of Internal Medicine* 112, 199-206 (1963)) reported upon the cardiac findings in 73 patients with muscular dystrophy, including the Duchenne, the limb-girdle, and the facioscapulohumeral forms, and patients with dystrophia myotonica. Zellweger, H., Durnin, R., Simpson, J., (*Acta Neurol. Scandinav.* 48, 87 (1972)) evaluated the electrocardiograms for various muscular dystrophies. Kuhn, E., Fiehn, W., Schroder, J. M., Assmus, H., and Wagner, A., (*Neurology* 29, 1144-1149 (1979)) described the early myocardial disease of the Becker muscular

dystrophy. Reeves, W. C., Griggs, R., Nanda, N. C., Thomson, K., Gramiak, R., (*Arch. Neurol.* 37, 273-277 (1980)) evaluated echocardiographic abnormalities in 24 subjects with Duchenne's dystrophy and 29 with myotonic muscular dystrophy. A significant incidence of cardiac abnormalities were demonstrable by echocardiography. Hopkins, L. C., Jackson, J. A., Elsas, L. J., (*Ann. Neurol.* 10, 230-237 (1981)) found that Emery-/Dreifuss humeroperoneal muscular dystrophy may be associated with a potentially lethal cardiac arrhythmia. Goldberg, S. J., Stern, L. Z., Feldman, L., Allen, H. D., Sahn, D. J., Valdes-Cruz, L. M., (*Neurology* 32, 1101-1105 (1982)), by a computerized study, found ventricular abnormalities as Duchenne patients develop severe cardiac disease. Hawley, R. J., Gottdiener, J. S., Gay, J. A., Engel, W. K., (*Arch. Intern. Med.* 143, 2134-2136 (1983)) evaluated 30 patients with myotonic dystrophy and the incidence of substantial cardiac involvement. Durnin, R. E., Ziska, J., Chandramouli, B., (*Journal of Iowa Medical Society*, (1969)), found that 80% of 52 patients with progressive muscular dystrophy had abnormal electrocardiograms.

The association of cardiac disease with muscular disease is established.

Mitochondrial Myopathies.

Carafoli, E., Roman, I., (*Molecular Aspects of Medicine* 3, 295 (1980)) and Morgan-Hughes, J. A., (*Advanced Medicine* 19, K. B. Saunders, ed., Pitman Publishers Ltd., London, 243-260 (1983)), identified deficiencies of the mitochondrial respiratory chain or its associated phosphorylation system in muscle mitochondria from myopathies in man. Since CoQ_{10} is an established constituent of the mitochondrial respiratory chain, these studies are basic to elucidation of human muscle disease.

Morgan-Hughes, J. A., Hayes, D. J., Clark, J. B., Cooper, J. M., (*Biomedical and Clinical Aspects of Coenzyme Q*, Vol. 4, K. Folkers and Y. Yamamura, eds., Elsevier/North Holland Biomedical Press, Amsterdam, 417-424 (1984)) conducted a most important study on isolated muscle mitochondria from 18 patients with myopathies. The respiratory activity, cytochrome content, and activities of several citric acid cycle enzymes were measured. The purity and integrity of the mitochondrial samples were documented. In ten cases of myopathy, the defect was in the first respiratory complex. In five cases, the deficiency was localized to the coenzyme Q-cytochrome bc₁ reductase complex. Respiratory rates were either low or not demonstrable with NAD-linked substrates or with succinate. Complex III deficiency was also monitored.

Measurement of Cardiac Function of Patients with Muscle Disease

All of the background research on the detection and some measurement of impaired cardiac function in patients with muscular disease only demonstrated an association of cardiac disease with muscular disease. Not one of those past studies involved the testing of anything to improve an impaired cardiac function. The conventional methods (echo cardiography, etc.) in cardiology of measuring cardiac function are essentially qualitative procedures useful for confirmation of diagnosis. These procedures are not sufficiently reliable and quantitative to monitor accurately over months of time cardiac function to determine whether the administra-

tion of any therapeutic agent is or is not effectively benefiting cardiac function.

Impedance cardiography offers a noninvasive, reproducible and acceptably quantitative measurement to monitor cardiac function over months of time. This technique has been described by Kubicek, W. G., Karnegis, J. N., Patterson, R. P., Witsoe, D. A. and Mattson, K. H., (*Aerospace Med.* 37, 1208-1212, 1966). Therefore, impedance cardiography is the instrumentation of choice to monitor cardiac function of patients with muscle disease over months of time as coenzyme Q₁₀ is orally administered.

General Method (The First Double Blind Trial)

The selected patients, as volunteers, were monitored by four to six impedance cardiographic measurements during a 30 to 75-day control period. The instrument consisted of a cardiograph, a computer and a printer purchased from Surcom, Inc., 4180 Edmond Blvd., Minneapolis, Minn. 55406. The double-blind treatment was with daily capsules of CoQ₁₀ (33 mg CoQ₁₀/capsule, t.i.d.) and a matching placebo. Stroke volume, cardiac output and heart rate were monitored for a minimum of three measurements/month during of each month of a three-month period. Multiple measurements allowed statistical calculations. When there was a sustained clinical response judged by the cardiac output and or in the stroke volume, with statistical significance of $p < 0.01$ – $p < 0.001$ during a three-month period, a blind assignment of CoQ₁₀ was made. When cardiac output and stroke volume had not significantly changed, not even by $p < 0.05$ during a blind treatment period of three months, placebo was assigned, and the patients were then provided with open CoQ₁₀ on a crossover basis.

The blood levels of CoQ₁₀ were determined essentially by the method of Kishi, T., Okamoto, T., Kanamori, N., Yamagami, T., Kishi, H., Okada, A., and Folkers, K., (*Biomedical and Clinical Aspects of Coenzyme Q*, Vol. 3, K. Folkers and Y. Yamamura, eds., Elsevier/North Holland Biomedical Press, Amsterdam, 67-78 (1981)), during the control and the blind periods of treatment. These data from the blind treatment period were not made available until after assignments of the patients to treatment with CoQ₁₀ or placebo were made, because knowledge of blood data prior to the blind assignments would have allowed assignments on the basis of blood data alone.

Results of the First Double Blind Trial

Blood Levels of CoQ₁₀. The control blood levels of CoQ₁₀ ranged from 0.50–0.84, which are lower than a mean plasma (only) level of 0.79 ± 0.23 $\mu\text{g/ml}$ of presumably normal subjects; Kishi, et al., 1981 (31). These differences between levels in muscle disease and normal may be numerically small, but bioenergetically high. After oral CoQ₁₀, the increased blood levels ranged from 1.11–2.93 $\mu\text{g/ml}$.

Cardiac Function (Objective). Table I contains the data on the cardiac function of the patients during the double-blind trial with capsules of CoQ₁₀ and a matching placebo, and the crossover from blind placebo to open CoQ₁₀ at the time of assignment and decoding. Table II contains the data on assignments, code and CoQ₁₀-blood levels.

For a total of 12 patients, seven had progressive muscular dystrophies including the Duchenne, Becker, and limb-girdle dystrophies and five had neurogenic atro-

phies including the Charcot-Marie Tooth disease, and the Wohlfart-Kugelberg Welander disease.

Before decoding and the availability of the blood data, eight of the twelve patients were assigned CoQ₁₀ and the remaining four were assigned placebo. Decoding revealed that not one mistake of assignment had been made, $p < 0.003$.

The patients who initially received blind CoQ₁₀ had the Duchenne and Becker dystrophies and Welander's disease and the patients initially receiving blind placebo had the limb-girdle and myotonic dystrophies and Charcot-Marie Tooth disease. All four patients initially receiving blind placebo were then treated for 3 months with open CoQ₁₀, and $\frac{3}{4}$ showed statistically significant increases in cardiac function.

Of the eight patients who were first treated with blind CoQ₁₀, two resigned from the study before the crossover period. Five of the six who were first treated with blind CoQ₁₀ maintained improved cardiac function during crossover with open CoQ₁₀, and one of these five, L. R., with Duchenne dystrophy, age 7, responded better ($p < 0.001$) in the 4th–6th month on CoQ₁₀ than in months 1–3 ($p < 0.054$ for S. V.). The one patient, H. M. K. with Charcot-Marie Tooth disease, 63 years, relapsed to the impaired control level of cardiac function on placebo, after significant response, $p < 0.001$, to blind CoQ₁₀. The control stroke volume of all 12 patients ranged from 29 to 100 ml./beat, but even the one patient (Becker's dystrophy) with a stroke volume of 100 ± 16 increased to 130 ± 9 ($p < 0.001$) during CoQ₁₀ treatment. Stroke volume, in ml. of blood pumped/heart beat, is appraised as a prime criterion because it is independent of the heart rate, and impedance cardiography is superior in allowing beat-by-beat data. The cardiac outputs for all 12 subjects ranged from 3.1 to 7.2 L/min. The subject with CO of 7.2 ± 1.6 (Becker's dystrophy) increased to 9.0 ± 0.4 ($p < 0.001$) on CoQ₁₀.

Introduction to Second Double Blind Trial

Coenzyme Q₁₀ (CoQ₁₀) is essential in the bioenergetics of the human cell and is one of the factors that sustains muscle activity. CoQ₁₀ is lipid-soluble and is present in the mitochondrial membrane and is an intermediate between NADH and succinate dehydrogenase and cytochrome b in the respiratory chain of human muscle tissue. (1,2). Folkers et al. (3) and Kamikawa et al. (4) have documented that coenzyme Q₁₀ is essential in the maintenance of normal heart function by regulating the energy metabolism in the myocardium. Diseases of the human heart are often correlated with a decrease in the levels of myocardial CoQ₁₀ (5,6). Many investigators have reported the association of cardiac disease with muscular disease (7, 8, 9, 10, 11, 12, 13, 14). Our first double blind study (15) showed the impaired myocardial function of patients with muscular dystrophy improved on CoQ₁₀ therapy, and there was also observed improved physical performance. The aim of this present study was to extend and to confirm our initial double blind trial on the administration of coenzyme Q₁₀ to patients with muscular dystrophy.

General Method (Second Double Blind Trial)

Fifteen muscular dystrophy patients (Table III), as volunteers, were monitored by four to six impedance cardiographic measurements during 14 to 96-day control periods. Cardiac performance was determined as stroke volume (SV) and cardiac output (CO) and was recorded at rest (16). The impedance cardiography

provides a noninvasive, reproducible and acceptably quantitative measurement and is known to correlate with angiographic, dilution or ultrasound procedures to measure cardiac function (17, 18). The double blind treatment was with daily capsules of CoQ₁₀ (33 mg of CoQ₁₀ per capsules, t.i.d.) and a matching placebo. Heart rate, SV and CO were monitored for a minimum of three measurements per month during each month of the investigation period. Multiple measurements allowed statistical calculations. A sustained clinical response, judged by the cardiac output and/or the stroke volume with statistical significance of $p < 0.01$ to $p < 0.001$ during observation in the blind period allowed an assignment of CoQ₁₀ to be made. When cardiac output and stroke volume had not significantly changed during the blind treatment period, placebo was assigned, and the patients were then provided with open CoQ₁₀ on a crossover basis. Patient 8 with Becker's M. D. was a male, 14 years, who was obviously growing, and showed a modest increase in cardiac function on placebo and a greater increase on CoQ₁₀.

The blood levels of CoQ₁₀ were determined (19) during the control and blind periods of treatment. Blood data during treatment were not made available until after assignments of the patients to treatment with CoQ₁₀ or to placebo were made, because knowledge of blood data prior to the blind assignments would have allowed assignments on the basis of blood data alone. Statistic comparison of results were obtained using the Student-t test.

Results of the Second Double Blind Trial

The cardiac function of all 8 patients treated with CoQ₁₀ during double blind and all 7 placebo patients during crossover improved significantly (Table III). The cardiac performance of 6 patients on blind placebo did not significantly improve.

Of the 15 patients, 8 had cardiac functions at rest which were lower than the low value of a normal range (20). Four (having limb-girdle, Becker's and Charcot-Marie Tooth diseases) of these 8 had cardiac functions on therapy which were restored to a normal range. However, each of the 15 patients had a significantly improved cardiac function on therapy.

Patient #8, a fourteen year-old boy with Becker's muscular dystrophy, improved somewhat by SV and CO in the blind placebo period, but improved far more during crossover on CoQ₁₀. His improved cardiac performance in the placebo period was apparently caused by his markedly increased growth. His degree of growth was the basis of assignment of placebo. Two children (patients #1 and #15) did not show any marked body growth during the time of the study.

All assignments of the patients to treatment with CoQ₁₀ or placebo were made correctly (Table IV). The control blood levels of CoQ₁₀ ranged from 0.58 to 1.17 $\mu\text{g/ml}$ and increased more than two-fold during the CoQ₁₀ treatment period (Table II). Compliance is difficult to know, and it is doubtful that all patients had perfect compliance, which would relate to the blood levels of CoQ₁₀.

Our first double blind trial (15) was conducted with 12 patients, and this second trial was with 15 patients. Not a single mistake was made in assignment of treatment with CoQ₁₀ or placebo during the two double blind trials with a total of 27 patients with diverse muscular dystrophies.

Cardiac Performance. Muscular dystrophies are generally regarded as involving genetic defects and as incurable diseases. However, therapeutic benefit is possible for the sequelae of the genetic defects.

Cardiac involvement in the form of cardiomyopathy is a well known essential feature of muscular dystrophies. Ultrastructural studies have demonstrated that cardiac changes are similar to those detected in the "dystrophic" skeletal muscle (8, 9). The diagnosis of cardiac involvement was also based on the demonstration of typical electrocardiographic changes, namely tall R waves in V₁, with R/S ratio > 1 , and deep and narrow Q-waves in limb leads and over the left lateral precordium. Physical and radiographic examination of the heart appears less specific, because of the frequent presence of thoracic deformities and high diaphragms. In recent years, echocardiography has gained increasing recognition, also in the preclinical state, in the noninvasive assessment of functional and anatomic cardiac changes. In the present study, we have used impedance cardiography to evaluate cardiac performance. Impedance cardiography is a noninvasive, reproducible and is a quantitative measurement to monitor cardiac function over months of time. This technique has superiorities, as evaluated by Baker and Mistry (21), in comparison with other techniques to measure cardiac function in the study of CoQ₁₀ and dystrophy.

Mitochondrial Myopathies. Over the past decade, there have been many studies on the treatment of different cardiac diseases with CoQ₁₀ (22, 23, 24, 25, 26, 27). Carafoli et al. (28) and Morgan-Hughes et al. (29, 30) identified deficiencies in the mitochondrial respiratory chain or its associated phosphorylation system, in muscle mitochondria from myopathies in man. In 1986, Sholte et al. (31) found that the most frequently discovered defects in human mitochondria of patients with muscular disease are multiple defects in oxidative phosphorylation, NADH-CoQ reductase, CoQ₁₀-cytochrome bc₁, cytochrome aa₃, multiple respiratory chain defects, loose coupling and adenine nucleotide translocase. The rationale of this trial was based on known myocardial myopathies which involved respiratory enzymes, the known presence of CoQ₁₀ in respiration, and prior clinical data on CoQ₁₀ and dystrophy. The results show improved cardiac function by double blind CoQ₁₀ therapy for these dystrophic patients and are identical to the results of the prior study (15).

Conclusion. Therapy with CoQ₁₀ is oral, without any side effect, and may be administered for the lifetime of such patients with muscular dystrophy. At present, there is no therapy for such patients. The improved cardiac performance of such patients is apparently established.

REFERENCES FOR THE SECOND DOUBLE BLIND TRIAL WHICH ARE INCORPORATED BY REFERENCES HEREIN FOR REASONS STATED EARLIER

1. Crane, F. S., Hatefi, Y., Lester, R. L., Widmer, C., *Biochem. Biophys. Acta.* 25, 220-221 (1957).
2. De Pierre, J. W., Ernster, L., *Ann. Rev. Biochem.* 46, 201-262 (1977).
3. Folkers, K., Watanabe, T., Kaji, M., *J. Mol. Med.* 2, 431-460 (1977).
4. Kamikawa, T., Kobayashi, A., Yamashita, T., Hayashi, H., Yamazaki, N., *Am. J. Cardiol.* 56, 247-251 (1985).

5. Folkers, K., Littarru, G. P., Ho, L., Runge, T. H. M., *Inter. J. Vitamin. Nutr. Res.* 40, 380-390 (1970).
6. Mortensen, S. A., Vadhanavikit, S., Folkers, K., *Drugs Exptl. Res.* 10, 497-502 (1984).
7. Welsh, J. D., Lynn, T. N., Haase, G. F., *Arch. Intern. Med.* 112, 199-206 (1963).
8. Venco, A., Grandi, A., Barzizza F., Malamani, G., Ghisoni, A., Basano, D., *Neuropsychiatr. Eta. Evol.* 4(1), 61-66 (1984).
9. Kuhn, E., Fiehn, W., Schroder, J. M., Assmus, H., Wagner, A., *Neurology* 29, 1144-1149 (1979).
10. Carstens, V., Behrenbeck, D. S., *Herz Kreisl* 17, 147-153 (1985).
11. Hopkins, L. C., Jackson, J. A., Elsas, L. J., *Ann. Neurol.* 10, 230-237 (1981).
12. Goldberg, S. J., Stern, L. Z., Feldman, L., Allen, H. D., Sahn, D. J., Valdes-Cruz, L. M., *Neurology* 32, 1101-1105 (1982).
13. Hawley, R. J., Gottdiener, J. S., Gay, J. A., Engel, W. K., *Arch. Intern. Med.* 143, 2134-2136 (1983).
14. Saviotti, M., Casazza, F., Fiorista, F., Murpurgo, M., *Neuropsychiatr. Eta. Evol.* 4(1), 67-69 (1984).
15. Folkers, K., Wolaniuk, J., Simonsen, R., Morishita, M., Vadhanavikit, S., *Proc. Natl. Acad. Sci. USA* 82, 4513-4516 (1985).
16. Kubicek, W. G., Karnegis, J. N., Patterson, P. R., Witsoe, D. A., Mattson, R. H., *Aerospace Med.* 37, 1208-1212 (1966).
17. Edmunds, A. T., Godfrey, S., Tooley, M., *Clin. Sci.* 63, 107-113 (1982).
18. Veigl, V. L., Judy, W. V., *Cardiovas. Res.* 17, 728-734 (1983).
19. Vadhanavikit, S., Morishita, M., Duff, G. A., Folkers, K., *Biochem. Biophys. Res. Comm.* 123, 1165-1169 (1984).
20. Ganong, W. F., In: Review of Medical Physiology. 35 Los Altos: Lange Medical Publication, 441-451 (1981).
21. Baker, L., Mistry, G. D., In: Biomedical and Clinical Aspects of Coenzyme Q, Vol. 3, K. Folkers and Y. Yamamura, eds., Amsterdam: Elsevier/North Holland Biomedical Press, 193-206 (1981).
22. Biomedical and Clinical Aspects of Coenzyme Q, Vol. 1. K. Folkers and Y. Yamamura, eds. Amsterdam: Elsevier/North Holland Biomedical Press (1977).
23. Biomedical and Clinical Aspects of Coenzyme Q, Vol. 2. K. Folkers, Y. Yamamura, and Y. Ito, eds. Amsterdam: Elsevier/North Holland Biomedical Press (1979).
24. Biomedical and Clinical Aspects of Coenzyme Q, Vol. 3. K. Folkers and Y. Yamamura, eds. Amsterdam: Elsevier/North Holland Biomedical Press (1981).
25. Biomedical and Clinical Aspects of Coenzyme Q, Vol. 4. K. Folkers and Y. Yamamura, eds. Amsterdam: Elsevier/North Holland Biomedical Press (1984).
26. Biomedical and Clinical Aspects of Coenzyme Q, Vol. 5. K. Folkers and Y. Yamamura, eds. Amsterdam: Elsevier/North Holland Biomedical Press (1986).
27. Coenzyme Q. G. Lenaz, ed. New York: John Wiley and Sons Ltd. (1985).
28. Carafoli, E., Roman, I., *Molecular Aspects of Medicine* 3, 295 (1980).
29. Morgan-Hughes, J. A., In: Advanced Medicine, Vol. 19, K. B. Saunders, ed., London: Pitman Publishers Ltd., 243-260 (1983).
30. Morgan-Hughes, J. A., Hayes, D. J., Clark, J. B., Cooper, J. M., Biomedical and Clinical Aspects of Coenzyme Q, K. Folkers and Y. Yamamura, eds., Elsevier/North Holland Biomedical Press, Amsterdam, 417-424 (1984).
31. Sholte, H. R., Luyt-Houwen, I. E. M., Busch, H. F. M., Vaandrigen-Verduin. New York Academy of Sciences, Como meeting (May, 1986).

TABLE I

Patient		Treatment* Duration, (days)			Functional Data; Mean \pm S.D. a, Cardiac Output (CO), L/min. b, Stroke Volume (SV), ml. c, Heart Rate (HR), beats/min.				
Sex, Age	Diagnosis	C*	B*	O*	C, Control		B, Double Blind		O, Open
1. M. B. F 45	Limb-Girdle Dystrophy	48	60	300	a, 5.2 \pm 0.3	5.0 \pm 0.6	n.s.	6.4 \pm 0.4	p < 0.001
					b, 57 \pm 7	60 \pm 1	n.s.	69 \pm 4	p < 0.001
					c, 96 \pm 8	84 \pm 11	—	98 \pm 3	—
2. L. R. M 7	Duchenne Dystrophy	75	70	270	a, 3.1 \pm 0.4	3.4 \pm 0.2	n.s.	4.0 \pm 0.1	p < 0.001
					b, 30 \pm 5	34 \pm 7	p < 0.5	37 \pm 8	p < 0.001
					c, 102 \pm 8	101 \pm 9	—	107 \pm 7	—
3. D. M. M 16	Myotonic Dystrophy	50	61	180	a, 3.3 \pm 0.4	3.4 \pm 0.2	n.s.	4.2 \pm 0.5	p < 0.001
					b, 29 \pm 4	30 \pm 1	n.s.	42 \pm 8	p < 0.001
					c, 118 \pm 6	110 \pm 2	—	101 \pm 10	—
4. K. A. C. M 7	Duchenne Dystrophy	29	113	172	a, 4.2 \pm 0.6	5.2 \pm 0.2	p < 0.001	4.5 \pm 0.7	p < 0.001
					b, 38 \pm 6	48 \pm 3	p < 0.001	42 \pm 8	p < 0.001
					c, 108 \pm 3	107 \pm 8	—	107 \pm 4	—
5. H. L. D. F 39	Limb-Girdle Dystrophy	37	122	110	a, 6.0 \pm 0.5	5.7 \pm 0.4	n.s.	5.5 \pm 0.3	n.s.
					b, 79 \pm 7	72 \pm 2	n.s.	73 \pm —	n.s.
					c, 77 \pm 3	76 \pm 6	—	75 \pm —	—
6. J. L. J. M 25	Becker's Dystrophy	29	180	180	a, 7.2 \pm 1.6	9.0 \pm 0.4	p < 0.001	D	—
					b, 100 \pm 16	130 \pm 9	p < 0.001	D	—
					c, 69 \pm 4	71 \pm 4	—	—	—
7. D. L. K. M 37	Becker's Dystrophy	60	150	150	a, 5.9 \pm 1.1	6.9 \pm 1.0	p < 0.001	7.42 \pm 1.36	p < 0.001
					b, 80 \pm 12	99 \pm 16	p < 0.001	106 \pm 26.63	p < 0.01
					c, 76 \pm 6	68 \pm 4	—	70	—
8. W. P. F 69	Charcot-Marie Tooth Disease	28	121	90	a, 5.2 \pm 0.2	5.7 \pm 0.2	n.s.	6.3 \pm 0.8	p < 0.001
					b, 67 \pm 6	69 \pm 2	n.s.	85 \pm 12	p < 0.001
					c, 77 \pm 3	76 \pm 3	—	74 \pm 2	—
9. H. M. K. F 63	Charcot-Marie Tooth Disease	32	91	96	a, 5.3 \pm 0.8	6.5 \pm 1.1	p < 0.001	4.6 \pm 0.4	n.s. E
					b, 71 \pm 9	89 \pm 13	p < 0.001	67 \pm 2	n.s. E
					c, 73 \pm 4	73 \pm 2	—	71 \pm 6	—

TABLE I-continued

		Data on Cardiac Function of Patients			Functional Data; Mean \pm S.D.					
Patient	Sex, Age	Diagnosis	Treatment* Duration, (days)			a, Cardiac Output (CO), L/min. b, Stroke Volume (SV), ml. c, Heart Rate (HR), beats/min.				
			C*	B*	O*	C, Control	B, Double Blind	O, Open		
10. B. H. M 31	Charcot-Marie Tooth Disease	30 105			a,	4.0 \pm 0.4	4.2 \pm 0.6	n.s.	D	
					b,	61 \pm 8	70 \pm 11	p < 0.001	D	
					c,	67 \pm 4	60 \pm 6	—	—	
11. M. D. F 63	Charcot-Marie Tooth Disease	45 90 300			a,	5.0 \pm 0.3	5.3 \pm 0.2	n.s.	6.3 \pm 0.6	p < 0.001
					b,	68 \pm 8	76 \pm 5	p < 0.001	83 \pm 5	p < 0.001
					c,	74 \pm 6	70 \pm 3	n.s.	75 \pm 4	—
12. C.S. F 49	Welander Disease	43 120 180			a,	3.9 \pm 0.5	4.5 \pm 0.5	p < 0.001	4.6 \pm 0.5	p < 0.001
					b,	68 \pm 6	84 \pm 9	p < 0.001	74 \pm 11	p < 0.001
					c,	57 \pm 6	54 \pm 5	—	62 \pm 5	—

*C = Control,
B = Blind, coded period of study,
O = Open period of study
D = Resigned from study
E = Crossover placebo period

TABLE II

		Double Blind and Open Trial Data and Blood Data				
Patient	Sex, Age	Double Blind		CoQ ₁₀ Blood Levels Average μ g/ml		
		Assignment	Code	Control	Blind	Open
1. M. B.; F 45		Placebo	Placebo	0.63	0.61	1.40; CoQ ₁₀
2. L. R.; M 7		CoQ ₁₀	CoQ ₁₀	0.62	1.11	1.24; CoQ ₁₀
3. D. M.; M 16		Placebo	Placebo	0.58	0.61	1.32; CoQ ₁₀
4. K. A. C.; M 7		CoQ ₁₀	CoQ ₁₀	0.50	1.94	1.47; CoQ ₁₀
5. H. L. D.; F 39		Placebo	Placebo	0.68	0.63	1.76; CoQ ₁₀
6. J. L. J.; M 25		CoQ ₁₀	CoQ ₁₀	0.71	1.22	—
7. D. L. K.; M 37		CoQ ₁₀	CoQ ₁₀	0.55	1.16	1.13; CoQ ₁₀
8. W. P.; M 69		Placebo	Placebo	0.84	0.77	2.13; CoQ ₁₀
9. H. M. K.; F 63		CoQ ₁₀	CoQ ₁₀	0.82	2.93	1.12; P
10. B. H.; M 31		CoQ ₁₀	CoQ ₁₀	0.59	2.09	—
11. M. D.; F 63		CoQ ₁₀	CoQ ₁₀	0.68	1.96	1.43; CoQ ₁₀
12. C. S.; F 49		CoQ ₁₀	CoQ ₁₀	0.78	1.58	2.03; CoQ ₁₀

TABLE III

MUSCULAR DYSTROPHY PATIENTS
DETAILS OF DATA ON CARDIAC FUNCTION

		Treatment (Days)			Functional Data; Mean \pm S.D. A. Cardiac Output (CO), L/min B. Stroke Volume (SV) ml/beat C. Heart Rate (HR) beats/min								
No.	Patient, Sex, Age	Diagnosis	Ctrl ^a	DB ^b	O ^c	Control ^a	Double Blind ^b	Open ^c Q ₁₀ , crossover.					
1.	J. A. (M,8)	Duchenne M.D.	96	105	182	A.	5.3 \pm 0.7	A.	5.6 \pm 0.7	n.s.	A.	6.6 \pm 0.8	p < 0.001
						B.	65 \pm 9	B.	66 \pm 16	n.s.	B.	80 \pm 7	p < 0.001
						C.	80 \pm 9	C.	86 \pm 10	—	C.	87 \pm 8	—
2.	J. P. B. (M,43)	Limb-girdle M.D.	78	102	862	A.	5.3 \pm 0.3	A.	5.5 \pm 0.3	n.s.	A.	5.8 \pm 0.7	p < 0.001
						B.	73 \pm 5	B.	75 \pm 5	n.s.	B.	73 \pm 10	n.s.
						C.	77 \pm 2	C.	73 \pm 4	—	C.	79 \pm 5	—
3.	D. C. (M,33)	Fascioscapulo- humeral M.D.	30	186	69	A.	7.9 \pm 0.7	A.	8.1 \pm 1.4	n.s.	A.	11.1 \pm 1.4	< 0.001
						B.	129 \pm 1	B.	132 \pm 8	n.s.	B.	168 \pm 21	p < 0.001
						C.	64 \pm 4	C.	63 \pm 10	—	C.	65 \pm 7	—
4.	D. H. (M,54)	Charcot-Marie- Tooth M.D.	20	126	34	A.	7.9 \pm 0.8	A.	8.1 \pm 0.7	n.s.	A.	11.8 \pm 2.3	p < 0.001
						B.	126 \pm 14	B.	129 \pm 10	n.s.	B.	186 \pm 31	p < 0.001
						C.	62 \pm 1	C.	61 \pm 2	—	C.	78 \pm 4	—
5.	P. K. (F,31)	Limb-girdle M.D.	34	481	890	A.	4.5 \pm 0.4	A.	5.4 \pm 0.4	p < 0.001	A.	6.2 \pm 0.5	p < 0.001
						B.	46 \pm 9	B.	59 \pm 7	p < 0.001	B.	69 \pm 4	p < 0.001
						C.	99 \pm 9	C.	91 \pm 5	—	C.	89 \pm 5	—
6.	M. R. (M,27)	Charcot-Marie- Tooth M.D.	14	158	68	A.	9.1 \pm 0.8	A.	12.2 \pm 1.7	p < 0.001	A.	11.9 \pm 1.4	p < 0.001
						B.	135 \pm 17	B.	188 \pm 14	p < 0.001	B.	195 \pm 11	p < 0.001
						C.	68 \pm 8	C.	64 \pm 5	—	C.	62 \pm 6	—
7.	M. R. (F,33)	Hypotonic congenitale	24	118	260	A.	6.9 \pm 0.6	A.	7.5 \pm 0.1	p < 0.001	A.	10.4 \pm 2.1	p < 0.001
						B.	82 \pm 6	B.	95 \pm 5	p < 0.001	B.	124 \pm 19	p < 0.001
						C.	84 \pm 7	C.	79 \pm 5	—	C.	83 \pm 6	—
8.	T. A. W. (M,14)	Becker's M.D.	16	131	124	A.	6.1 \pm 0.5	A.	7.3 \pm 0.7	p < 0.001	A.	9.7 \pm 1.7	p < 0.001
						B.	57.2 \pm 11	B.	74 \pm 7	p < 0.001	B.	97 \pm 15	p < 0.001
						C.	104 \pm 2	C.	99 \pm 4	—	C.	99 \pm 3	—
9.	J. M. (F,58)	Charcot-Marie- Tooth M.D.	30	120	359	A.	5.0 \pm 0.9	A.	5.5 \pm 0.9	n.s.	A.	6.6 \pm 1.7	p < 0.001
						B.	60 \pm 9	B.	57 \pm 9	n.s.	B.	70 \pm 16	p < 0.001
						C.	84 \pm 2	C.	98 \pm 6	—	C.	93 \pm 5	—
10.	T. D. (M,31)	Limb-girdle M.D.	21	66	667	A.	3.8 \pm 0.3	A.	4.8 \pm 0.7	p < 0.001	A.	5.6 \pm 0.5	p < 0.001
						B.	46 \pm 4	B.	68 \pm 5	p < 0.001	B.	72 \pm 6	p < 0.001

TABLE III-continued
MUSCULAR DYSTROPHY PATIENTS
DETAILS OF DATA ON CARDIAC FUNCTION

No.	Patient, Sex, Age	Diagnosis	Treatment (Days)			Functional Data; Mean \pm S.D.							
			Ctrl ^a	DB ^b	O ^c	Control ^a		Double Blind ^b		Open ^c Q ₁₀ crossover.			
						A. Cardiac Output (CO), L/min		B. Stroke Volume (SV) ml/beat		C. Heart Rate (HR) beats/min			
11.	L. L. (M,38)	Myotonic M.D.	27	172	539	C.	83 \pm 4	C.	72 \pm 2	—	C.	77 \pm 2	—
						A.	3.0 \pm 0.4	A.	3.4 \pm 0.3	n.s.	A.	3.7 \pm 0.4	p < 0.01
						B.	50 \pm 7	B.	60 \pm 4	p < 0.01	B.	65 \pm 5	p < 0.001
12.	T. E. P. (M,74)	Limb-girdle M.D.	19	120	721	C.	61 \pm 8	C.	58 \pm 6	—	C.	57 \pm 7	—
						A.	2.7 \pm 0.6	A.	3.4 \pm 0.3	p < 0.01	A.	3.3 \pm 0.4	p < 0.001
						B.	41 \pm 8	B.	52 \pm 4	p < 0.01	B.	51 \pm 4	p < 0.001
13.	M. T. (F,69)	Limb-girdle M.D.	20	150	560	C.	65 \pm 2	C.	65 \pm 5	—	C.	69 \pm 5	—
						A.	3.2 \pm 0.5	A.	3.5 \pm 0.3	n.s.	A.	4.9 \pm 0.9	p < 0.001
						B.	39 \pm 6	B.	41 \pm 4	n.s.	B.	56 \pm 8	p < 0.001
14.	B. H. (F,53)	Welander M.D.	22	75	695	C.	81 \pm 3	C.	85 \pm 6	—	C.	87 \pm 6	—
						A.	3.2 \pm 0.5	A.	4.6 \pm 0.4	p < 0.001	A.	4.2 \pm 0.3	p < 0.001
						B.	43 \pm 4	B.	62 \pm 3	p < 0.001	B.	55 \pm 8	p < 0.001
15.	B. S. (M,15)	Becker's M.D.	31	90	348	C.	75 \pm 6	C.	74 \pm 7	—	C.	77 \pm 7	—
						A.	8.1 \pm 1.2	A.	9.1 \pm 1.5	n.s.	A.	11.1 \pm 2.5	p < 0.001
						B.	96 \pm 11	B.	115 \pm 8	p < 0.001	B.	130 \pm 27	p < 0.001
					C.	84 \pm 5	C.	79 \pm 9	—	C.	84 \pm 6	—	

^aControl^bDouble Blind, Coded Period of Study^cOpen Period of Study

TABLE IV

RESULTS OF DOUBLE BLIND AND OPEN TRIAL
DATA AND BLOOD DATA

No.	Patient; Sex, Age	Double Blind		CoQ ₁₀ Blood Levels Average μ g/ml		
		Assignment	Code	Control	Blind	Crossover
1.	J. A.; M, 8	Placebo	Placebo	0.61	0.53	1.62
2.	J. P. B.; M, 43	Placebo	Placebo	0.72	0.84	2.48
3.	D. C.; M, 33	Placebo	Placebo	0.94	0.75	1.68
4.	D. H.; M, 54	Placebo	Placebo	0.89	0.83	0.94
5.	P. K.; F, 31	CoQ ₁₀	CoQ ₁₀	0.66	2.24	2.84
6.	M. R.; M, 27	CoQ ₁₀	CoQ ₁₀	0.64	1.05	1.17
7.	M. R.; F, 33	CoQ ₁₀	CoQ ₁₀	0.74	1.81	1.27
8.	T. A. W.; M, 14	Placebo	Placebo	0.81	0.82	1.42
9.	J. M.; F, 58	Placebo	Placebo	0.93	1.00	2.05
10.	T. D.; M, 31	CoQ ₁₀	CoQ ₁₀	1.17	2.74	2.48
11.	L. L.; M, 38	CoQ ₁₀	CoQ ₁₀	0.65	1.85	1.72
12.	T. E. P.; M, 74	CoQ ₁₀	CoQ ₁₀	1.04	2.29	2.45
13.	M. T.; F, 69	Placebo	Placebo	0.90	0.80	1.70
14.	B. H.; F, 53	CoQ ₁₀	CoQ ₁₀	0.95	2.20	2.50
15.	B. S.; M, 15	CoQ ₁₀	CoQ ₁₀	0.58	1.38	1.44
	Mean \pm S.D.			0.81 \pm 0.17	p < 0.001	1.85 \pm 0.56

What is claimed:

1. A method for treating impaired cardiac function of human patients having the Duchenne dystrophy, the Becker dystrophy, the limb-girdle dystrophies, myotonic dystrophy, Charcot-Marie-Tooth disease fascioscapulothoracic muscular disease, hypotonia congenitale

or Welander disease, or a closely related disease, said method comprising administering to the patients having any one of these dystrophies or myopathies a therapeutically effective amount of a formulation consisting essentially of coenzyme Q₁₀.

* * * * *

55

60

65