

Improving Clinical Performance of an Interprofessional Emergency Medical Team through a One-day Crisis Resource Management Training

Teodora Sorana Truta¹, Cristian Marius Boeriu^{1*}, Marc Lazarovici², Irina Ban³, Marius Petrișor¹, Sanda-Maria Copotoiu¹

¹ University of Medicine and Pharmacy Tirgu-Mures, Street Gheorghe Marinescu 38, 540139 Tirgu-Mures, Mures, Romania

² Institut für Notfallmedizin und Medizinmanagement- INM, Klinikum der Universität, Ludwig-Maximilians-Universität München, Schillestraße 53, 80336 München, Germany

³ Department of Medicine University of Padova, Via V. Galluci, 13-35121, Padova, Italy

ABSTRACT

Introduction: Errors are frequent in health care and Emergency Departments are one of the riskiest areas due to frequent changes of team composition, complexity and variety of the cases and difficulties encountered in managing multiple patients. As the majority of clinical errors are the results of human factors and not technical in nature or due to the lack of knowledge, a training focused on these factors appears to be necessary. Crisis resource management (CRM), a tool that was developed initially by the aviation industry and then adopted by different medical specialties as anesthesia and emergency medicine, has been associated with decreased error rates.

The aim of the study: To assess whether a single day CRM training, combining didactic and simulation sessions, improves the clinical performance of an interprofessional emergency medical team.

Material and Methods: Seventy health professionals with different qualifications, working in an emergency department, were enrolled in the study. Twenty individual interprofessional teams were created. Each team was assessed before and after the training, through two in situ simulated exercises. The exercises were videotaped and were evaluated by two assessors who were blinded as to whether it was the initial or the final exercise. Objective measurement of clinical team performance was performed using a checklist that was designed for each scenario and included essential assessment items for the diagnosis and treatment of a critical patient, with the focus on key actions and decisions. The intervention consisted of a one-day training, combining didactic and simulation sessions, followed by instructor facilitated debriefing. All participants went through this training after the initial assessment exercises.

Results: An improvement was seen in most of the measured clinical parameters.

Conclusion: Our study supports the use of combined CRM training for improving the clinical performance of an interprofessional emergency team. Empirically this may improve the patient outcome.

Keywords: interprofessional, clinical performance, training, crisis resource management, emergency department

Received: 18 August 2018 / Accepted: 08 September 2018

INTRODUCTION

In the recent years, the need to enhance patient safety has become a priority as the majority of clinical errors (approximately 70%) are the results of human factors [1] and not technical in nature or due to the lack of knowledge (failure in team communication, situational awareness, resource utilization and leadership) [2]. Ef-

forts are necessary to improve the quality of the medical care provided and the patient outcome, especially in high risk areas as emergency medicine, anesthesia, surgery and intensive care.

How can we achieve this if just improving the knowledge is not enough? Steps were taken by adopting and adapting the Crises Resource Management (CRM) training, a tool that is oriented towards human

* Correspondence to: Boeriu Cristian, University of Medicine and Pharmacy Tirgu Mures, Gheorghe Marinescu Street 38, 540139 Tirgu-Mures, Romania. Email: cboeriu@gmail.com

factors and was initially developed by the aviation industry. CRM training is considered to be responsible for the decrease in aircraft accidents over the last four decades [3].

The primary objectives of CRM training are to improve team dynamics, to identify and help change mental models that create barriers in adopting effective communication, effective task management, healthy leadership and fellowship behaviors and an increasing awareness [4].

Anesthesia was one of the first specialties to adopt the CRM training and an improvement of team performance and a reduced risk of errors during medical and surgical crisis in the operating room has been shown [5].

Early interest in this type of training was demonstrated also by the emergency medicine specialty.

Morey et al. (2002), in their evaluation of MedTeams Project's results found that formal teamwork training was effective in improving team behaviors and in reducing errors in Emergency Departments (ED) [6]. Despite this, human factors training is not widely disseminated through professionals working in ED and most studies focused only on emergency medicine residents and interdisciplinary trauma teams.

According to Chiniara (2003), simulation training is most suitable for low frequency high-risk situations that are potentially harmful for the patient [7]. These types of situations are those where crises arise, and CRM training may play an important role. As crises events are rare in real practice, the simulation room is an ideal setting for teaching CRM principles [7] and simulation-based CRM training was shown to be effective and relevant for the emergency medicine residents' practice [2].

Different scales were developed to assess the non-technical skills/behavioral performance and many studies reported on the reaction of participants toward the training, the acquisition of knowledge and the changes in attitude in non-clinical settings. Only a limited number of studies focused on transfer of the learning to the working place and even less measured the effect on patient outcomes [8].

As morbidity and mortality are indicators difficult to track in prospective studies, surrogates such as time to disposition, delays in achieving tasks in a timeframe, number of errors [9] have been used as well as checklists, to assess the clinical performance.

This study is aimed at assessing whether a single day CRM oriented team training combining didactic and simulation sessions improves the clinical performance of interprofessional emergency medicine teams.

■ METHODS

Study design

The study took place in the emergency department of the Tirgu-Mures Emergency Clinical County Hospital, Romania between March and July 2016. The hospital has an emergency medicine residency training program and is the site of an affiliated emergency and disaster medicine simulation center that focuses on multidisciplinary and inter-professional training. The emergency department's annual census is around 77,000 patients, approximately 10% of them being "critically ill".

Emergency department clinical staff without prior CRM training were invited to participate. Residents with less than three months clinical experience in the emergency department were excluded. Twenty board-certified emergency medicine (EM) physicians, ten emergency medicine residents, and forty nurses volunteered for the study.

The study protocol was approved by the Ethics Committee of the Tirgu-Mures Emergency Clinical County Hospital. Written informed consent was obtained from all study participants. Twenty mixed teams were constructed according to participants' work schedules.

Each team utilized an attending EM physician, an EM resident and two nurses, a structure we consider ideal for the management of a single critically ill patient in our ED. Because the number of eligible attending physicians was greater than the number of residents, each resident participated in two teams.

Two sets of scenarios were designed, consisting of two scenarios each: a medical case and a trauma case.

The first set consisted of:

- Medical patient A - A patient with an acute COPD exacerbation complicated by severe pneumonia, initially requiring noninvasive and subsequently invasive ventilation. Due to patient physiognomy as well as a history of prior tracheostomy, a difficult intubation needed to be anticipated.
- Trauma patient A - A trauma patient with severe brain injury complicated by hemorrhagic shock secondary to intra-abdominal bleeding requiring

activation of the massive transfusion protocol and emergent surgery.

The second set consisted of:

- Medical patient B - A patient with COPD now presenting with a pulmonary embolism necessitating thrombolysis due to clinical instability.
- Trauma patient B - A patient with a difficult airway, a severe, but surgically amenable intra-cranial injury, as well as long bones fractures, overall resulting in hypotension.

Both medical cases had signs of β_2 mimetics overuse (tremor, tachycardia, hypokalemia).

Sets were chosen pseudo-randomly, as participating residents were intended to be exposed to all cases and not to repeat scenarios. Simulations were performed in the ED resuscitation room during work hours. Scenarios utilized a high-fidelity manikin as well as the usual medical equipment and documentation forms from the resuscitation room. The role of different specialties doctors who were called for advice/help was played by one of the instructors, as well as the role of paramedics. The role of the radiologist was played by the radiologist on duty that day. Both a fixed camera, placed on the ceiling of the resuscitation room, as well as a mobile camera recorded all of the exercises. No debriefings took place after the assessment scenarios were completed.

We then held single day (6 to 7 hours) training sessions over a period of ten days. Participants chose the day for their training session at their convenience. Training sessions took place in a recreation of the ED resuscitation room in the simulation center. Each training session started with a lecture focusing on medical errors and CRM principles. No medical teaching was provided. After the lecture, participants had the opportunity to familiarize themselves with the manikin and the simulation setting.

Each training session involved two multi-professional teams, each team consisting of one EM attending physician, one EM resident and two nurses. During the training sessions participants were permitted to change teams as long as the multi-professional composition remained the same. We utilized six critical patient scenarios (two trauma cases and four medical cases), different from those used in the initial assessment. One team completed a scenario while the other team observed the exercise remotely via a high-resolution real time video transmission system. Either as participants or observers, each team was exposed to all cases.

The simulation was run by two instructors (one doctor and one nurse) with CRM background training and one IT technician. Each scenario was followed by an instructor facilitated debriefing. Both technical and non-technical issues relating to team performance and team work were discussed. Identified clinical errors were addressed by either the participants or the instructors. Members of the team who had completed the scenario had priority in providing feedback, but both observers and participants were involved in debriefings.

Two months after receiving training, a final assessment was performed consisting of the same scenarios, teams and setting used in the initial assessment. As in the initial assessment, the scenarios were video-recorded, and no briefing was provided.

Objective measurement of clinical team performance was performed through a checklist that was designed for each scenario (Figure 1). The checklist included essential assessment items for the diagnosis and treatment of a critical patient, with the focus on key actions and decisions. The number of times a critical procedure was performed, as well as time to completion of the critical steps where appropriate, were recorded.

The video recordings were analyzed, and the checklists completed by an assessor blinded to whether the scenarios analyzed were the initial or final assessment.

Statistical analysis

The collected data were organized into several SPSS data files. The variables obtained were binary type and continuous type. For binary data analysis, the likelihood ratio was used to assess the difference in proportions. Continuous variables were tested for normality using Kolmogorov-Smirnov test. Our data followed a non-gaussian distribution, therefore a non-parametric test was used to compare the central tendency for the data series. Wilcoxon signed rank test was used, since we analyzed paired data. The significance level used in all tests was 0.05.

RESULTS

Seventy participants were enrolled in the study and 69 completed the study. One nurse participant resigned her position and she was not available for the final assessment scenario. She was replaced by another nurse who had attended the training. The initial nurse was excluded from the final analysis and the nurse who re-

Checklist medical case A			
Team members:			
	Yes	No	Time
1. Assess consciousness	<input type="checkbox"/>	<input type="checkbox"/>	
2. Assess A	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/> < 2 min
B	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/> < 2 min
C	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/> < 2 min
3. Give oxygen	<input type="checkbox"/>	<input type="checkbox"/>	_____
4. Semi sitting position	<input type="checkbox"/>	<input type="checkbox"/>	_____
5. Monitoring ECG	<input type="checkbox"/>	<input type="checkbox"/>	_____
6. Monitoring BP	<input type="checkbox"/>	<input type="checkbox"/>	_____
7. Monitoring SpO ₂	<input type="checkbox"/>	<input type="checkbox"/>	_____
8. Temperature measurement	<input type="checkbox"/>	<input type="checkbox"/>	_____
9. Focused clinical exam			
• Cardiac auscultation	<input type="checkbox"/>	<input type="checkbox"/>	
• Pulmonary auscultation	<input type="checkbox"/>	<input type="checkbox"/>	
• Jugular veins	<input type="checkbox"/>	<input type="checkbox"/>	
• Sings of PVT/ lower limbs	<input type="checkbox"/>	<input type="checkbox"/>	
10. Focused history			
• Anamnesis	<input type="checkbox"/>	<input type="checkbox"/>	
• Past medical history	<input type="checkbox"/>	<input type="checkbox"/>	
• Treatment	<input type="checkbox"/>	<input type="checkbox"/>	
11. 12 leads ECG	<input type="checkbox"/>	<input type="checkbox"/>	_____
12. Intravenous access	<input type="checkbox"/>	<input type="checkbox"/>	_____
13. Arterial blood gases (ABG)	<input type="checkbox"/>	<input type="checkbox"/>	_____
14. CBC, electrolytes, renal & hepatic tests	<input type="checkbox"/>	<input type="checkbox"/>	
15. CRP	<input type="checkbox"/>	<input type="checkbox"/>	
16. NTproBNP	<input type="checkbox"/>	<input type="checkbox"/>	
17. Troponin	<input type="checkbox"/>	<input type="checkbox"/>	
18. D-dimers	<input type="checkbox"/>	<input type="checkbox"/>	
19. Chest X-ray	<input type="checkbox"/>	<input type="checkbox"/>	
20. Cardiac ultrasound	<input type="checkbox"/>	<input type="checkbox"/>	
21. Give β ₂ mimetics	<input type="checkbox"/>	<input type="checkbox"/>	
22. Noninvasive ventilation	<input type="checkbox"/>	<input type="checkbox"/>	_____
• Choose the mask	<input type="checkbox"/>	<input type="checkbox"/>	
• Set the ventilator correctly	<input type="checkbox"/>	<input type="checkbox"/>	
• Explain to the patient	<input type="checkbox"/>	<input type="checkbox"/>	
• Apply the mask	<input type="checkbox"/>	<input type="checkbox"/>	
• Observe for tolerance & synchronism	<input type="checkbox"/>	<input type="checkbox"/>	
• Observe the patient	<input type="checkbox"/>	<input type="checkbox"/>	
• Observe the ventilator	<input type="checkbox"/>	<input type="checkbox"/>	
23. Repeat ABG after 30 minutes	<input type="checkbox"/>	<input type="checkbox"/>	
24. Give appropriate antibiotic	<input type="checkbox"/>	<input type="checkbox"/>	_____
25. Differential diagnosis			
• Exacerbation of COPD+ pneumonia	<input type="checkbox"/>	<input type="checkbox"/>	
• Acute pulmonary edema	<input type="checkbox"/>	<input type="checkbox"/>	
• Acute coronary syndrome	<input type="checkbox"/>	<input type="checkbox"/>	
• Pulmonary embolism	<input type="checkbox"/>	<input type="checkbox"/>	
• Spontaneous pneumothorax	<input type="checkbox"/>	<input type="checkbox"/>	
26. Final diagnosis			
• Exacerbation of COPD+ pneumonia	<input type="checkbox"/>	<input type="checkbox"/>	
27. Recognize β ₂ -mimetics overdose	<input type="checkbox"/>	<input type="checkbox"/>	

Fig. 1. Example of a checklist used for the assessment of clinical performance

placed her was analyzed only once with the team to whom she was allocated at the beginning of the study.

The male: female ratio was 13:17 for doctors and 11:29 for nurses. Professional experience was variable (average 70 months, minimum 8 and maximum 300 months).

A checklist was completed for each scenario and each team. A total of 40 checklists were collected, 20 for the initial assessment and 20 the for the final one.

For the medical case A, the patient with acute COPD exacerbation complicated by severe pneumonia, critical elements where considered: oxygen administration, sitting position, establishing an intravenous access, appropriate investigations- especially ECG and arterial blood gases, recognition of the need for non-invasive ventilation, recognition of β₂ sympatico-mimetics overdose, appropriate antibiotic administration. Results are reported in table 1 and table 2. A trend towards improvement was seen for these elements. Concerning the time until the procedure was performed, a statistically significant value was reached only for intravenous access. In terms of number of procedures performed, β₂ sympatico-mimetics overdose recognition and assessment of efficiency of NIV through repeated ABG 30 minutes after initiation of it, reached significance level.

For medical case B, the patient with COPD now presenting with a pulmonary embolism, critical elements where: oxygen administration, semi-sitting position, establishing an intravenous access, appropriate investigations- especially ECG, arterial blood gases, angioCT scan/ cardiac ultrasound, recognition of the need for and initiation of thrombolysis, request for cardiologic advice and recognition of β₂ sympatico-mimetics overdose. Results are reported in table 3 and table 4. As for the medical case A, an improvement was seen for most of the elements. Significance level was reached in terms of number of procedures performed for cardiac ultrasound requested and β₂ sympatico-mimetics overdose recognition. Concerning the time until the procedure was performed, significance level was reached for oxygen administration and sitting position. For time until a cardiac ultrasound was requested, the p value couldn't be calculated as the test was requested during the initial assessment scenario only by two teams.

Critical elements for the trauma case A, the patient with severe brain injury complicated by hemorrhagic shock secondary to intra-abdominal bleeding, were considered: 2 intravenous access, FAST ultrasound examination, administration of fluid bolus and O negative PRBCs, recognition of a shock patient requiring urgent surgery (request for surgeon advise early, transfer to the

Table 1. Checklist results for medical case A

Element		Number of teams "Yes"/initial	Number of teams "Yes"/final	Number of teams "No"/initial	Number of teams "No"/final	Likelihood Ratio*
Consciousness		10	10	0	0	
ABC evaluation < 2 minutes		10	10	0	0	
Oxygen administration **		10	10	0	0	
Sitting position **		10	10	0	0	
Monitoring	Electrocardiogram	10	10	0	0	
	Blood pressure	10	10	0	0	
	Oxygen saturation	10	10	0	0	
	Temperature	8	9	2	1	0.528
Focused clinical examination	Cardiac auscultation	1	2	9	1	0.528
	Pulmonary auscultation	10	10	0	0	
	Jugular veins	0	1	10	9	
	Signs of PVT/edema	5	10	5	0	
Focused history	Anamnesis	10	10	0	0	
	Past medical history	10	10	0	0	
	Treatment	8	10	2	0	
12-lead electrocardiogram **		6	10	4	0	
Intravenous line **		10	10	0	0	
Arterial blood gases (ABG) **		10	10	0	0	
Lab tests	CBC, electrolytes, Renal & hepatic tests	9	10	1	0	
	C-reactive protein	1	0	9	10	
	NTproBNP	8	2	8	2	1.000
	Troponin	2	4	8	6	0.326
	D-dimers	6	8	4	2	0.326
Chest radiography		10	10	0	0	
Cardiac ultrasound		1	1	9	9	1.000
Give β_2 mimetics		9	3	1	7	0.004
Noninvasive ventilation **		10	10	0	0	
	Choose the mask	8	7	2	3	0.605
	Set the ventilator correctly	8	9	2	1	0.528
	Explain to the patient	8	10	2	0	0.112
	Apply the mask	10	10	0	0	
	Observe for tolerance & synchronism	6	9	4	1	0.112
	Observe the patient	9	9	1	1	1.000
	Observe the ventilator	7	9	3	1	0.255
Repeat ABG after 30 minutes		4	9	6	1	0.015
Give appropriate antibiotic		8	10	2	0	
Differential diagnosis	Exacerbation of COPD+ pneumonia	10	10	0	0	
	Acute pulmonary edema	4	10	6	0	
	Acute coronary syndrome	0	3	10	7	
	Pulmonary embolism	7	10	3	0	
	Spontaneous pneumothorax	2	6	8	4	0.063
Correct diagnosis		10	10	0	0	
Recognize β_2 mimetics overdose		1	7	9	3	0.004

*Chi-Square Tests; ** Time to achievement recorded; ABC- airway, breathing, circulation, PVT-profound venous thrombosis, CBC- complete blood count, NTproBNP- N-terminal pro b-type natriuretic peptide, COPD-chronic obstructive pulmonary disease

Table 2. Results for medical case A – Time to achievement of critical steps

Time to.... (seconds)	Initial		Final		P value*
	Median	IQR	Median	IQR	
Oxygen administration	65	80	39.5	63	0.203
Sitting position	17	113	6	18	0.074
12 leads electrocardiogram	56	121	136.5	111	0.463
Intravenous line	123.5	114	58.5	54	0.013
Arterial blood gases	142.5	99	70.5	99	0.169
Noninvasive ventilation	308	304	215.5	124	0.386

*Wilcoxon Signed Ranks Test

Table 3. Checklist results for medical case B

Element	Number of teams	Number of teams	Number of teams	Number of teams	Likelihood Ratio*	
	"Yes"/initial	"Yes"/final	"No"/initial	"No"/final		
Consciousness	10	10	0	0		
ABC evaluation < 2 minutes	10	10	0	0		
Oxygen administration **	10	10	0	0		
Sitting position **	9	10	1	0		
Monitoring	Electrocardiogram	10	10	0	0	
	Blood pressure	10	10	0	0	
	Oxygen saturation	10	10	0	0	
	Temperature	6	1	4	9	0.015
Focused clinical examination	Cardiac auscultation	4	8	6	2	0.063
	Pulmonary auscultation	9	8	1	2	0.528
	Jugular veins	1	3	9	7	0.255
	Signs of PVT/edema	9	10	1	0	
Focused history	Anamnesis	10	10	0	0	
	Past medical history	10	10	0	0	
	Treatment	10	10	0	0	
12-lead electrocardiogram**	10	10	0	0		
Intravenous line **	10	10	0	0		
Arterial blood gases (ABG)	10	10	0	0		
Lab tests	CBC, electrolytes, Renal & hepatic tests	10	10	0	0	
	CRP	0	0	10	10	
	NT-proBNP	8	9	2	1	0.528
	Troponin	9	8	1	2	0.528
	D-dimers	9	10	1	0	
Chest radiography	6	4	4	6	0.369	
Fluids bolus	2	9	8	1	0.001	
Give β_2 mimetics	8	2	2	8	0.005	
Request angioCT scan **	10	8	0	2		
Angio-CT scan performed	6	2	4	6	0.132	
Request cardiac ultrasound **	2	7	8	3	0.021	
Request venous Doppler ultrasound	2	0	8	10		
Decide to thrombolysis **	9	10	1	0		
Verify contraindications to thrombolysis	2	7	8	3	0.021	
Start thrombolysis **	9	10	1	0		
Ask for cardiologic advise **	10	10	0	0		
Differential diagnosis	Exacerbation of COPD+ pneumonia	6	9	4	1	0.112
	Acute pulmonary edema	4	5	6	5	0.653
	Acute coronary syndrome	5	10	5	0	
	Pulmonary embolism	10	10	0	0	
	Spontaneous pneumothorax	5	8	5	2	0.155
Correct diagnosis	10	10	0	0		
Recognize β_2 mimetics overdose	2	8	8	2	0.005	

*Chi-Square Test; ** Time to achievement recorded; ABC- airway, breathing, circulation, PVT-profound venous thrombosis, CBC- complete blood count, NT-proBNP- N-terminal pro b-type natriuretic peptide, COPD-chronic obstructive pulmonary disease

Table 4. Results for medical case B – Time to achievement of critical steps

Time to.... (seconds)	Initial		Final		P value*
	Median	IQR	Median	IQR	
Oxygen administration	76	81	51.5	29	0.022
Sitting position	20	245	6.5	15	0.012
12 leads electrocardiogram	160	125	120	30	0.221
intravenous line	115	116	105	48	0.859
Arterial blood gases	159	73	128	144	0.767
AngioCT scan	345	369	501	NA	NA
Cardiac ultrasound	790	NA	520	450	NA
Thrombolysis decision	515	233	510	394	0.515
Start thrombolysis	570	673	606.5	410	0.678
Request cardiologic advice	336	280	300	580	0.333

*Wilcoxon Signed Ranks Test; NA- not available

operating theatre), recognition of the need to intubate and anticipation of a difficult airway, recognition and initiation of treatment for increased ICP. Results are reported in table 5 and table 6. There was a trend towards improvement in all the elements, though not statistically significant, except for recognition and initiation of treatment of high ICP, and the time until airway was secured.

The trauma case B, the patient with a difficult airway, severe TBI and long bones fractures resulting in

hypotension, had the following elements: intravenous access, administration of fluid bolus, FAST ultrasound examination, recognition of severe brain injury requiring urgent measures to decrease the ICP, performance of the CT scan and early involvement of the neurosurgeon, recognition of the need to intubate, anticipation of a difficult airway and management of a “cannot ventilate, cannot intubate” situation (number of direct laryngoscopies, call for help, decide and perform a cri-

Table 5. Checklist results for trauma case A

Element	Number of teams “Yes”/initial	Number of teams “Yes”/final	Number of teams “No”/initial	Number of teams “No”/final	Likelihood Ratio*
Consciousness	10	10	0	0	
A-airway	6	10	4	0	
B-breathing	9	10	1	0	
Primary survey (ABC < 2 minutes)	9	10	1	0	
(DE < 5 minutes)	9	10	1	0	
D GCS	9	10	1	0	
Motor response	7	10	3	0	
Pupillary exam	9	10	1	0	
E-exposure	10	10	0	0	
Oxygen administration	10	10	0	0	
Electrocardiogram	10	10	0	0	
Monitoring	10	10	0	0	
Blood pressure	10	10	0	0	
Oxygen saturation	10	10	0	0	
Temperature	1	5	9	5	0.044
Anamnesis	10	10	0	0	
Past medical history	1	0	9	10	
Allergies	0	0	10	10	
Focused history	0	1	10	9	
Treatment	0	1	10	9	
Mechanism of injury	7	10	3	0	
Time from injury	0	7	10	3	
Last meal	0	1	10	9	
Insert an intravenous line **	10	10	0	0	
ABO+ Rhesus	10	10	0	0	
Glucose level	2	1	8	9	0.528
CBC, electrolytes, renal & hepatic, coagulation tests, creatine kinase, arterial blood gases	10	10	0	0	
Fluids bolus **	7	10	3	0	
FAST ultrasound exam **	10	10	0	0	
12-lead electrocardiogram **	3	4	7	6	0.639
Insert a second intravenous line **	10	10	0	0	
Recognize signs of shock	10	10	0	0	
Transfusion of O negative PRBCs	6	9	4	1	0.121
Recognize signs of increased ICP	6	10	4	0	
Recognize the need for intubation **	10	10	0	0	
Anticipate a difficult airway	3	7	7	3	0.070
Airway secured **	10	10	0	0	
Request the surgeon advise **	10	10	0	0	
Request for a CT scan	3	3	7	7	1.000
Request the neurosurgeon advise	5	8	5	2	0.155
Treat high ICP	2	7	8	3	0.021
Immobilize the fracture	5	10	5	0	
Perform the logroll	1	3	9	7	0.255
Perform a complete secondary survey	1	0	9	10	
Transfer the patient to the operating room **	9	10	1	0	
IOT checklist completed verified	0	2	10	8	

*Chi-Square Tests; ** Time to achievement recorded; D-disability, CBC-complete blood count, a FAST-focused assessment with Sonography in Trauma, PRBCs-packed red blood cells, ICP-intracranial pressure

Table 6. Results for trauma case A – Time to achievement of critical steps

Time to.... (seconds)	Initial		Final		P value*
	Median	IQR	Median	IQR	
First intravenous line	65	49	43.5	45	0.139
Fluids bolus	208	258	116.5	114	0.128
FAST ultrasound exam	174	177	160.5	160	0.721
12 leads electrocardiogram	658	NA	114	52	0.109
Second intravenous line	174	352	158.5	190	0.646
Recognize the need for IOT	149	116	109	199	0.203
Airway secured	401	210	317	78	0.017
Request the surgeon advise	339.5	215	247.5	247	0.241
Transfer into the operating room	580	295	605	70	0.374

*Wilcoxon Signed Ranks Test; NA- not available; FAST-focused assessment with ultrasound in trauma, IOT-otracheal intubation

cothyroidotomy). Results are given in table 7 and table 8. An improvement was seen in most of the elements. Significant improvement was shown in the elements correlated with severe TBI and difficult airway management, in terms of number of procedures performed (recognition and treatment of increased ICP, anticipation of difficult airway) and time until procedure was completed (request for neurosurgeon advice, first attempt to intubate, decision for cricothyroidotomy, airway secured).

■ DISCUSSION

CRM trainings are addressing a set of nontechnical skills, “the cognitive, social and personal resource skills that complement technical skills, and contribute to safe and efficient task performance” [10]. These nontechnical skills can be divided into four domains essential for an effective team: teamwork, task management, decision making and situational awareness, all of them being linked by communication [11]

The CRM key principles that act as a framework for teaching teamwork skills to emergency residents, according to GABA [5] and modified by Carne [12] are:

- Know your environment
- Anticipate, share and review the plan
- Ensure leadership and role clarity
- Communicate effectively
- Call for help early
- Allocate attention wisely- avoid fixation
- Distribute the workload-monitor and support team members.

In our study we assessed whether a single-day training on CRM principles combining didactic and high-fidelity simulated session may improve the clinical performance (objectively measured through a checklist) of an interprofessional emergency team. We found that

most of the measured clinical parameters improved. We hypothesize this is due to an improvement of non-technical skills, as we didn’t teach the clinical skills in the simulation cases explicitly. Also, in each team we had an experienced attending physician, the mean experience for the EM attendings being 149 months (minimum 72 and maximum 300 months), so lack of knowledge was unlikely.

Many of the clinical decisions as well as the evaluation of the patient include the correct usage of CRM principles.

The cases were chosen in such a way that they were typical for an average acute ED patient, but still strongly dependent on correct application of CRM principles. E.g., when dealing with a difficult airway, in our case a “cannot ventilate cannot intubate” situation, the team should anticipate, share the same mental model, distribute the workload and plan ahead while staying alert for sudden changes of situation. All the required skills correspond to CRM principles and indeed improved significantly after the training as shown by the number of teams who anticipated the difficult airway ($p=0.044$), number of attempts to intubate by direct laryngoscopy, the time until decision to perform a cricothyroidotomy was taken and the airway was secured.

Managing a polytrauma patient with hemorrhagic shock, requiring damage control surgery and massive transfusion protocol activation, requires to: anticipate, share the plan, ensure leadership and role clarity, communicate effectively, distribute the workload, call for help early and maintain standards. An improvement was seen in the present study, reflected by number of teams that administrate fluids bolus and O negative PRBCs from the ED storage, monitored the temperature as hypothermia is one of the components of the lethal triad in bleeding patients [13], time until they insert the intravenous lines, gave the fluids bolus, performed the FAST exam, call for surgeon.

Table 7. Checklist results for trauma case B

Element	Number of teams "Yes"/initial	Number of teams "Yes"/final	Number of teams "No"/initial	Number of teams "No"/final	Likelihood Ratio*
Consciousness	10	10	0	0	
A-airway	9	10	1	0	
B-breathing	9	10	1	0	
Primary survey (ABC < 2 minutes)	9	10	1	0	
(DE < 5 minutes) D GCS	8	10	2	0	
Motor response	7	10	3	0	
Pupillary exam	8	10	2	0	
E-exposure	10	10	0	0	
Oxygen administration	8	10	2	0	
Electrocardiogram	10	10	0	0	
Blood pressure	10	10	0	0	
Oxygen saturation	10	10	0	0	
Temperature	0	5	10	5	
Anamnesis	10	10	0	0	
Past medical history	1	5	9	5	0.044
Allergies	0	7	10	3	
Treatment	0	1	10	9	
Mechanism of injury	7	10	3	0	
Time from injury	5	10	5	0	
Last meal	0	0	10	10	
Insert an intravenous line **	10	10	0	0	
ABO+ Rhesus	10	10	0	0	
Glucose level	2	3	8	7	0.605
CBC, electrolytes, renal & hepatic & coagulation tests, creatine kinase, arterial blood gases	10	10	0	0	
Fluids bolus **	3	3	7	7	1.000
FAST ultrasound exam **	5	7	5	3	0.359
12-lead electrocardiogram	1	4	9	6	0.112
Insert a second intravenous line **	8	8	2	2	1.000
Request for CT scan	9	10	1	0	
Request the neurosurgeon **	9	10	1	0	
Recognize signs of increased ICP	3	9	7	1	0.004
Recognize the need for intubation **	10	10	0	0	
Anticipate a difficult airway	5	9	5	1	0.044
Number of direct laryngoscopy attempts (more than two)	6	0	4	10	
Ventilation achieved between intubation attempts	10	10	0	0	
Ask for help **	5	5	5	5	1.000
Use of alternative techniques for intubation	5	0	5	10	
Decide for cricothyroidotomy **	10	10	0	0	
Airway secured **	10	10	0	0	
Treat high ICP	4	9	6	1	0.015
Immobilize the fracture	8	9	2	1	0.528
Give antibiotic	4	10	6	0	
Perform the logroll	4	8	6	2	0.063
Perform a complete secondary survey	3	1	7	9	0.255
IOT checklist completed verified	0	2	10	8	

*Chi-Square Test; ** Time to achievement recorded; D-disability, CBC-complete blood count, a FAST-focused assessment with Sonography in Trauma, ICP-intracranial pressure, IOT-oro-tracheal intubation

Also, the recognition of a condition such as β_2 sympatho-mimetics overdose, in a patient presenting with respiratory distress, where signs as tachycardia and agitation might have different causes, request situational awareness and wise allocation of attention. In both medical cases, a significant improvement was

seen after the training ($p=0.004$ for case A and $p=0.005$ for case B).

Avoiding fixation error was important in order to avoid a wrong diagnosis in medical cases as

they had a similar history and almost the same clinical presentation, except for the blood pressure that was

Table 8. Results for trauma case B – Time to achievement of critical steps

Time to.... (seconds)	Initial		Final		P value*
	Median	IQR	Median	IQR	
First intravenous line	66	85	69.5	80	0.674
Fluids bolus	245	703	114	188	0.317
FAST ultrasound exam	170	875	240	510	0.715
Second intravenous line	150	207	195	552	0.917
Request for neurosurgeon	1200	338	580	465	0.008
Recognize the need for IOT	240	175	145	80	0.017
First IOT attempt	512	179	355	147	0.007
Call for help	682	356	520	186	0.655
Decide for cricothyroidotomy	790	195	446,5	262	0.005
Airway secured	1030	304	602	250	0.005

*Wilcoxon Signed Ranks Test; FAST-focused assessment with ultrasound in trauma, IOT-oro-tracheal intubation

lower but still in normal range initially, for the patient with PE. Their final diagnosis was correct before and after the training, but more differential diagnoses were considered during the final assessment.

The idea of team skills training being associated with better clinical performance and empirically with better patient outcome was already raised by Wright et al [14]. In their study on medical students, run in two settings: a classroom-based and a high-fidelity simulated environment, they found a strong correlation between team skills rating and objective performance measures within the high fidelity simulated environment, but no correlation on the classroom-based environment.

Very few studies measured the impact of CRM training on patient outcome [8]. One of these found a significant decrease in mortality for in hospital pediatric cardiac arrest after simulation CRM training. [15].

A positive association between better teamwork, better nontechnical skills and disposition time and risk for delays to patient care was demonstrated by Pucher et al, on a study carried on fifty real trauma cases [9].

Similar to our study, the efficiency of a relatively brief (4 hours) teamwork training, including a one-hour web didactic session followed by a human patient simulator-based session, on clinical performance of an interdisciplinary trauma team was shown by Steinmann et al [16]. This improvement was seen on simulated cases as well as on real trauma cases.

The effectiveness of teamwork training on clinical performance of an interprofessional team (surgeons, nurses), reflected by the decrease in time from arrival to CT scanner, endotracheal intubation and operating room, was demonstrated also by Capella et al [17]. As

in the previously mentioned study, these results were obtained from real cases.

The present study has several limitations. The study was run in a single center and this limited the number of participants. Due to the very small number of EM residents available during the study period, they were exposed twice to the initial and final assessment in two different teams. Although we used two sets of scenarios and we did not perform a debriefing afterwards, we cannot exclude that their experience with the first team influenced their performance with the second one. It is difficult to appreciate if this had an impact the team clinical performance.

Participants were evaluated two months after the training. We did not assess them immediately after the training, so we do not know if their performance improved or declined over this time period.

Ideally, after a CRM training the actual patient outcome would be measured. This was our initial intention when we implemented this study. Unfortunately, we encountered a technical problem when we tried to analyze real cases. The quality of the image returned by the fixed camera was good, but the quality of the sound was too poor to permit a proper evaluation.

■ CONCLUSION

Our study supports the use of combined CRM training for improving clinical performance of an interprofessional emergency team. Empirically this may improve the patient outcome, but further research is necessary to establish the ideal length of the training, the real impact on patient outcome, the length of the decay phase and the optimal method and time for refresher training.

■ ACKNOWLEDGEMENTS

The authors thank all the participants in the study. Special thanks to Oana Mates, Andreea Soceanu and Bogdan Posteuca for their help in collecting the data.

■ CONFLICT OF INTEREST

None to declare

■ REFERENCES

- Rovamo L, Nurmi E, Mattila MM, et al. Effect of a simulation-based workshop on the multidisciplinary teamwork of newborn emergencies: an interventional study. *BMC Res Notes*. 2015;8:671.
- Hicks MC, Kiss A, Bandiera WG, et al. Crisis Resources for Emergency Workers (CREW II): results of a pilot study and simulation-based crisis resource management course for emergency medicine residents. *CJEM*. 2012;14(6):354-62.
- Petrosoniak A, Hicks CM. Beyond crisis resource management: new frontiers in human factors training for acute care medicine. *Curr Opin Anesthesiol*. 2013;26:699-706.
- Rall M, Gaba D- Human Performance and Patient Safety. In Miller, RD (ed): *Miller's Anesthesia*. Philadelphia: Elsevier Churchill Livingstone. 2005, pp. 3021-72.
- Gaba D, Howard S, Fish K. Simulation-based training in anesthesia crisis resource management (ACRM): a decade of experience. *Simul Gaming*. 2001;32:175-93.
- Morey JC, Simon R, Jay GD, et al. Error Reduction and Performance Improvement in the Emergency Department through Formal Teamwork Training: Evaluation Results of the MedTeams Project. *Health Serv Res*. 2002;37(6):1553-81.
- Chiniara G, Cole G, Brisbin K, et al. Simulation in healthcare: a taxonomy and a conceptual framework for instructional design and media selection. *Med Teach*. 2013;35(8):e1380-95.
- Boet S, Bould MD, Fung L, et al. Transfer of learning and patient outcome in simulated crisis resource management: a systematic review. *Can J Anesth*. 2014;61:571-82.
- Pucher PH, Aggarwal R, Batrick N, et al. Nontechnical skills performance and care processes in the management of the acute trauma patient. *Surgery*. 2014;155(5):902-9.
- Flin R, O'Connor R, Crichton M. *Safety at the Sharp End: A Guide to Nontechnical Skills*. Farnham, UK: Ashgate; 2008.
- Bleetman A, Sanusi S, Dale T, Brace S. Human factors and error prevention in emergency medicine. *Emerg Med J*. 2012;29:389-93.
- Carne B, Kennedy M, Gray T. Crisis resource management in emergency medicine. *Emerg Med Australas*. 2012;24:7-13.
- Simmons JW, Powell MF. Acute traumatic coagulopathy: pathophysiology and resuscitation. *Br J Anaesth*. 2016;117(3):iii31-iii43.
- Wright MC, Phillips-Bute BG, Petrusa ER, Griffin KL, Hobbs GW, Taekman JM. Assessing teamwork in medical education and practice: relating behavioural teamwork ratings and clinical performance. *Med Teach*. 2009;31:30-8.
- Andreatta P, Saxton E, Thompson M, Annich G. Simulation-based mock codes significantly correlate with improved pediatric cardiopulmonary arrest survival rates. *Pediatr Crit Care Med*. 2011;12:33-8.
- Steinemann S, Berg B, Skinner A et al. In situ-multidisciplinary, simulated-based teamwork training improves early trauma care. *J Surg Educ*. 2011;68:472-7.
- Capella J, Smith S, Philp A et al. Teamwork training improves the clinical care of trauma patients. *J Surg Educ*. 2010;67:439-43.