

## Occurrence of antibiotic residues in raw fish *Clarias gariepinus* and *Oreochromis niloticus* from intensive rearing system in Benin

S.E.P Mensah, H. Dakpogan, A. B. Aboh, K. Chabi Sika, M. Abléto, K. Y. K. B Adjahoutonon, O D Koudandé, Pascal Sanders, G.A. Mensah

### ► To cite this version:

S.E.P Mensah, H. Dakpogan, A. B. Aboh, K. Chabi Sika, M. Abléto, et al.. Occurrence of antibiotic residues in raw fish *Clarias gariepinus* and *Oreochromis niloticus* from intensive rearing system in Benin. *Veterinaria*, Veterinary Faculty Sarajevo, 2019, 68 (2), pp.91-94. anses-02336074

**HAL Id: anses-02336074**

**<https://hal-anses.archives-ouvertes.fr/anses-02336074>**

Submitted on 28 Oct 2019

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Occurrence of antibiotic residues in raw fish *Clarias gariepinus* and *Oreochromis niloticus* from intensive rearing system in Benin

Mensah S. E. P<sup>1</sup>, Dakpogan H<sup>3</sup>, Aboh A. B<sup>4</sup>, Chabi Sika K<sup>2</sup>, Abléto M<sup>2</sup>, Adjahoutonon K. Y. K. B<sup>1\*</sup>, Koudandé O. D<sup>1</sup>, Sanders P<sup>5</sup>, Mensah G. A<sup>1</sup>

## Abstract

An observational cross-sectional study was carried out to investigate antibiotic residues in *Clarias gariepinus* and *Oreochromis niloticus* from the intensive rearing system in Southern Benin. The targeted antibiotic families were tetracyclins, amphenicols, beta-lactams and macrolides. One hundred and forty-four (144) samples were used per antibiotic family for residue detection in the fish muscle, making 576 treated samples. The Charm II method based on radioimmunoassay was used. The results showed an overall residue prevalence of 11.1%, which is attributable to tetracyclines alone. There was no residue of the other antibiotic families in the treated fish muscles. The adult fish were the only contaminated age group with 22.2% residue prevalence compared to the fingerlings ( $p < 0.05$ ). Likewise, the antibiotic residues were significantly ( $p < 0.05$ ) more prevalent in *Clarias gariepinus* (16.7%) than in *Oreochromis niloticus* (5.6%). The adult fishes are generally fatter than the young ones, and that can lower the fish body antibiotic elimination capability. This first detection of tetracyclines residues in fish produced in Benin shows the need to rule and control antibiotic use in the developing fish industry to preserve consumers' health.

**Keywords:** Tilapia, African catfish, fish farm, antibiotic residue, Benin

<sup>1</sup> Institut National des Recherches Agricoles du Bénin, Centre de Recherches Agricoles d'Agonkanmey, Laboratoire des Recherches Zootechnique, Vétérinaire et Halieutique (INRAB/CRA-Agonkanmey/LRZVH), 01BP884 Recette Principale Cotonou, Bénin.

<sup>2</sup> Laboratoire Centrale de Sécurité Sanitaire des Aliments/Ministère de l'Agriculture de l'Élevage et de la Pêche (LCSSA/MAEP),

<sup>3</sup> Université Nationale d'Agriculture/Ecole de Gestion et d'Exploitation des Systèmes d'Élevage/Kétou, BP43 Kétou, Bénin.

<sup>4</sup> Université Nationale d'Agriculture/Ecole d'Aquaculture de la Vallée, BP43 Kétou, Bénin.

<sup>5</sup> Agence nationale de sécurité sanitaire (Anses), Laboratoire de Fougères, F-35302 Fougères, France.

\*Corresponding author: bricead@gmail.com

## Introduction

Aquaculture is a worldly fast-growing animal production sector. Worldwide, fish provide 15% of animal production source protein to more than three billion people and play a vital economical role by contributing specifically to unemployment rate reduction (FAO, 2011). In Benin, food safety is a big concern because of the high population increase rate of 3.5% annually (INSAE, 2013). The efficient way of addressing this problem is the intensification of animal production including the fish production.

Fish production in Benin is at its early intensification stage. The use of antibiotics as food supplements for disease prevention and treatment and as growth promoters (Reda *et al.*, 2013; Pham *et al.*, 2015) is a common practice. However, such use of antibiotics without veterinary control lead inevitably to the presence of antibiotic residues in the animal-derived products and by-products (Mensah *et al.*, 2014). The presence of antibiotic residue in fish products was well documented. The utilization of antibiotic products in aquaculture is prejudicial to the aquatic environment and aqualife on one hand, and on the other hand, to the fish products consumers due to the

toxicity risk of antibiotic residues (Cabello, 2006; Olatoye and Basiru, 2013; Dhaouadi *et al.*, 2015). Unfortunately, no scientific work has been done to date on antibiotic residues in intensive fish farming products in Benin.

The current study was carried out in an observational cross-sectional design to assess the occurrence of antibiotic product residue in fish from the intensive rearing system in Southern Benin.

## Material and methods

### Fish sample collection

Fish were collected in an intensive fish production farm in a southern region of Benin. The two fish species produced in this farm were *Clarias gariepinus* (African catfish) and *Oreochromis niloticus* (Tilapia of the Nile River). The farm includes two major farm units located in two different districts. The first unit was the hatchery and catfish fattening site located in Ouèdo, and the second was the tilapia growing farm located in Pahou. The fish were fed with the commercial feed.

One hundred and forty-four (144) fishes were randomly collected: seventy-two (72) per species and thirty-six (36) for each age group, adult and growers.

The collected fish samples were transported in the ice boxes to the Central Laboratory of Sanitary and Safety Food Control of the Ministry of Agriculture, Livestock and Fishery of Benin, and then stored at -20°C until the laboratory analysis.

### Laboratory analysis

Four antibiotic families were targeted in the laboratory analysis (Table 1). These antibiotics were broad spectrum antibiotics likely to be used by the fish farmers. Consequently, four samples were taken from each fish to search for amphenicols, beta-lactams, macrolides and tetracyclines antibiotic residues with a total of 576 fish samples made available for the test.

Charm II screening test was used for antibiotic residues detection in the fish samples. This is a radioimmunoassay with the competition for the specific binding sites between drug residues and radioactive tracers. The amount of tracer that binds to the receptor sites is measured and compared to the previously determined control point (Table 1). Control point is calculated from the average value obtained running 6 tests of negative controls which values must not deviate more than 15% from the average. The control point is obtained by subtracting 40% from the negative control average (Charm Sciences Inc., 2019).

**Table 1:** Detection kit capacity for antibiotic residue (Charm Sciences Inc., 2019)

Antibiotic family	Antibiotics	Cut-off values (ppb)
Amphenicols	Chloramphenicol	10
	Florfenicol	-
Beta-lactams	Amoxillin	50
	Ampicillin	40
	Cefalexin	200
	Cefquinone	200
	Ceftiofur and metabo	400
	Cephapirin	40
	Cloxacillin	500
	Dicloxacillin	300
	Oxacillin	500
	Penicillin G	50
Macrolides	Erythromycin	100
	Lincomycin	400
	Pirlimycin	600
	Spiramycin	400
	Tilmicosin	50
	Tulathromycin	100
Tetracyclines	Tylosin	400
	Chlortetracycline	100
	Oxytetracycline	100
	Tetracycline	25

### Statistical analysis

The Pearson Chi square test (without Yates continuity correction) was used to compare the occurrence of antibiotic residues in the fish muscles between fish age groups (adults and fingerlings) and species (*C. gariepinus* and *O. niloticus*) in R 3.5.1.

## Results

Only tetracycline residues were found in 11.1% of the fish muscle samples regardless of species and age group. All the fish muscle samples were free from amphenicols, macrolides and beta-lactams' residues.

Considering age groups, only the muscle samples of adult fish (22.2%) contained residues of this antibiotic family. No residue was found in fingerlings muscle samples. Otherwise, such tetracycline residues were three times more prevalent ( $p < 0.05$ ) in *Clarias gariepinus* (16.7%) than in *Oreochromis niloticus* (5.6%).

## Discussion and conclusions

In this study, a radioimmunoassay is used for residue detection because it is a fast and reliable method compared with microbiological, immunological or physicochemical methods for antibiotic residue screening and quantification in the animal muscles.

Several studies reported the presence of antimicrobial residues in food of animal origin in Benin (Mensah *et al.*, 2014). However, no investigation or quantitative analysis has been done on the presence of antibiotic residues in fish produced and consumed in this country. In Ibadan, Nigeria, antibiotic residues have been detected using Premi®Test kit in 52.5% of ready-to-eat *C. gariepinus* muscle collected from restaurants and farms (Olatoye and Basiru, 2013). The presence of antibiotic residue in the fish muscle samples of the current study is in line with the results of the above mentioned study, and can be explained by the routine use of antibiotics in fish diets in the intensive production system. This might result from adding antibiotics to the locally-produced foods just before harvesting in order to boost growth (Reda *et al.*, 2013) and prevent disease and stress during live-fish transportation (Romero *et al.*, 2007; Olatoye and Basiru, 2013; Pham *et al.*, 2015).

Tetracyclines residues were the only antibiotic residues found in this study. Similarly, tetracyclines residues had been detected in, respectively, 7.84% and 30% of locally-produced fish in Vietnam (Pham *et al.*, 2015) and Nigeria (Olatoye and Basiru, 2013). Tetracyclines seem to be one of the most commonly used antibiotic families in fish production (Olatoye and Basiru, 2013; Reda *et al.*, 2013).

Moreover, in this study, only adult fish muscles tested positive to antibiotic residues, irrespectively of the species. The body adipose tissue is known to be susceptible to the drug residue in general, so that it can significantly lower the antibiotic pharmacokinetic activities of the body

organs. This can explain the difference between the two age groups, adult and fingerlings, in antibiotic residue prevalence. Two fish species investigated in this study are positive to tetracyclines residues, with a higher prevalence in *C. gariepinus* than *O. niloticus*. This may be attributable to different disease management since drug administration, and/or blood composition of *C. gariepinus* and *O. niloticus* (Hamid *et al.*, 2013), which can influence tetracyclines pharmacokinetic. Likewise, tetracyclines residues have been detected in the samples from those two fish species in the respective concentrations of 1.98 and 3.08 ppm in Ibadan, Nigeria (Olusola *et al.*, 2012). Eventhough the fish samples collected in our study are free from amphenicols, macrolides and beta-lactams residues, chloramphenicol (an amphenicol) residues have been found in the *C. gariepinus* muscle sample in Nigeria (Olusola *et al.*, 2012). The presence of residues of other antibiotic families like quinolones in the *O. niloticus* muscle sample has also been mentioned (Xu *et al.*, 2006; Pham *et al.*, 2015). Withdrawal time of 3 to 42 days depending on the antibiotic used and water temperature, is needed to avoid the presence of antibiotic residues above maximum residue limit (MRL) in fish (Chafer-Pericas *et al.*, 2010).

This first detection of tetracyclines residues in raw *C. gariepinus* and *O. niloticus* produced in Benin proves that antibiotics are misused in fish farming as in other animal farming (Mensah *et al.*, 2011). Fish consumers in this country are then exposed to many risks related to the presence of antibiotic residues in food like antibiotic resistance, carcinogenicity, mutagenicity, nephropathy, hepatotoxicity or allergy (Mensah *et al.*, 2014). Local fish industry should be included in a national system for control of drug residues in food of animal origin to preserve consumers' health.

### Acknowledgements

The authors would like to thank the National Institute of Agricultural Research of Benin for funding this study, and the International Atomic Energy Agency for providing the material.

## References

1. Cabello, F. C. 2006. Heavy use of prophylactic antibiotics in aquaculture: a growing problem for human and animal health and for the environment. *Environ. Microbiol.*, 8(7), 1137-1144.
2. Cháfer Pericás, C., Maquieira, Á., Puchades, R., Company, B., Miralles, J. and Moreno, A., 2010. Multiresidue determination of antibiotics in aquaculture fish samples by HPLC-MS-MS. *Aquacul. Res.*, 41, 217-225.
3. Charm Sciences Inc. 2019. Charm II Test Kits. <http://www.charm.com/products/test-and-kits/antibiotic-tests/charm-ii-test-kits/>. (Accessed 27.4. 2019).
4. Dhaouadi, R., Tarhouni, D.E., Louati, A., 2015. Utilisation des antibiotiques en aquaculture. <http://docplayer.fr/22592327-Utilisation-des-antibiotiques-en-aquaculture.html>. (Accessed 10.3.2018).
5. FAO, 2011. Family poultry communications. International Network for Family Poultry Development, 20 (2), 57 p.
6. Hamid, S. A., Ahmed, F. M., Mohammed, I. A., Ali, S. M., 2013. Physical & chemical characteristics of blood of two fish species (*Oreochromis niloticus* and *Clarias lazera*). *World's Vet. J.*, 3 (1), 17-20.
7. INSAE (Institut National de la Statistique et de l'Analyse Economique), 2013. Résultat provisoires du RGPH 4. 8 p.
8. Mensah, S.E.P., Ahissou, H.H., Koudandé, O.D., Salifou S., Mensah, G.A., Abiola, F.A., 2011. Detection of antibiotics residues in meat of reformed and marketed laying hens in southern Benin. *Int. J. Biol. Chem. Sci.* 5 (6), 2195-2204.
9. Mensah, S.E.P., Koudandé, O.D., Sanders, P., Laurentie, M., Mensah, G.A., Abiola, F.A., 2014. Résidus d'antibiotiques et denrées d'origine animale en Afrique: risques de santé publique. *Rev. Sci. Tech. Off. Int. Epiz.* 33 (3), 1-27.
10. Olatoye, I. O., Basiru, A. 2013. Antibiotic Usage and Oxytetracycline Residue in African Catfish (*Clarias gariepinus*) in Ibadan, Nigeria. *World Journal of Fish and Marine Sciences* 5 (3), 302-309.
11. Olusola, A. V., Folashade, P. A., Ayoade, O. I. 2012. Heavy metal (lead, Cadmium) and antibiotic (*Tetracycline* and *Chloramphenicol*) residues in fresh and frozen fish types (*Clarias gariepinus*, *Oreochromis niloticus*) in Ibadan, Oyo State, Nigeria. *Pak J BiolSci*, 15(18), 895-9.
12. Pham, D.K., Chu, J., Do, N.T., Brose, F., Degand, G., Delahaut, P., De Pauw, E., Douny, C., Van Nguyen, K., Dinh Vu, T., Scippo, M.-L., Wertheim, H.F.L. 2015. Monitoring antibiotic use and residue in freshwater aquaculture for domestic use in Vietnam. *EcoHealth*, 12 (3), 480-489.
13. Reda, R. M., Ibrahim, R. E., Ahmed, E. N. G., El-Bouhy, Z. M. 2013. Effect of oxytetracycline and florfenicol as growth promoters on the health status of cultured *Oreochromis niloticus*. *Egypt. J. Aquat. Res.*, 39(4), 241-248.
14. Romero, R., López, J.C, Gómez, E., Garrido, A., Martínez, J.L., 2007. Simultaneous determination of selected veterinary antibiotics in gilthead seabream (*Sparus aurata*) by liquid chromatography-mass with the spectrometry. *J. Chromatogr. B*, 857, 142-146.
15. Xu, W., Zhu, X., Wang, X., Deng, L., & Zhang, G. 2006. Residues of enrofloxacin, furazolidone and their metabolites in Nile tilapia (*Oreochromis niloticus*). *Aquacul.*, 254(1-4), 1-8.

# Prisustvo antibiotikih rezidua u sirovoj ribi *Clarias gariepinus* i *Oreochromis niloticus* iz intenzivnog uzgoja u Beninu

## SAŽETAK

Izvedeno je opservacijsko presječno istraživanje kako bi se utvrdilo prisustvo antibiotikih rezidua u ribama *Clarias gariepinus* i *Oreochromis niloticus* iz intenzivnog uzgoja u južnom Beninu. Ispitivani su antibiotiči iz skupine tetraciklina, amfenikola, beta-laktama i makrolida. U istraživanju su ispitivana sto četrdeset i četiri (144) uzorka po antibiotičkoj skupini, ukupno 576 uzoraka na prisustvo rezidua u ribljim mišićima. Od metoda je korišten Charm II radioimuno test. Rezultati su pokazali prisustvo rezidua tetraciklina od 11,1%. U ispitivanim uzorcima ribe nisu otkrivene rezidue ostalih antibiotičkih skupina. Jedina kontaminirana skupina riba su bile odrasle jedinke sa prevalencom od 22,2% u odnosu na mlade jedinke ( $p < 0.05$ ). Prevalenca antibiotičkih rezidua je statistički signifikantnija ( $p < 0.05$ ) kod vrste *Clarias gariepinus* (16.7%) u odnosu na *Oreochromis niloticus* (5.6%). Odrasle jedinke riba generalno imaju više masnog tkiva nego mlade jedinke, što može sniziti eliminacijsku sposobnost tijela. Prvo otkrivanje prisustva antibiotičkih rezidua u ribama uzgojenim u Beninu je ukazalo na potrebu kontrole upotrebe antibiotika u ribogojstvu u svrhu zaštite zdravlja potrošača.

**Ključne riječi:** Tilapija, Afrički som, ribnjak, antibiotički reziduum, Benin