

Manuscript version: Author's Accepted Manuscript

The version presented in WRAP is the author's accepted manuscript and may differ from the published version or Version of Record.

Persistent WRAP URL:

<http://wrap.warwick.ac.uk/129478>

How to cite:

Please refer to published version for the most recent bibliographic citation information. If a published version is known of, the repository item page linked to above, will contain details on accessing it.

Copyright and reuse:

The Warwick Research Archive Portal (WRAP) makes this work by researchers of the University of Warwick available open access under the following conditions.

Copyright © and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable the material made available in WRAP has been checked for eligibility before being made available.

Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

Publisher's statement:

Please refer to the repository item page, publisher's statement section, for further information.

For more information, please contact the WRAP Team at: wrap@warwick.ac.uk.

Complete stereoinversion of L-tryptophan by a fungal single-module nonribosomal peptide synthetase

Yang Hai,¹ Matthew Jenner,^{3,4} and Yi Tang^{1,2*}

¹Department of Chemical and Biomolecular Engineering, ²Department of Chemistry and Biochemistry, University of California, Los Angeles, California 90095, USA. ³Department of Chemistry, University of Warwick, Coventry, UK. ⁴Warwick Integrative Synthetic Biology (WISB) Centre, University of Warwick, Coventry, UK.

Supporting Information Placeholder

ABSTRACT: Single-module nonribosomal peptide synthetases (NRPSs) and NRPS-like enzymes activate and transform carboxylic acids in both primary and secondary metabolism; and are of great interests due to their biocatalytic potentials. The single-module NRPS IvoA is essential for fungal pigment biosynthesis. Here, we show that IvoA catalyzes ATP-dependent unidirectional stereoinversion of L-tryptophan to D-tryptophan with complete conversion. While the stereoinversion is catalyzed by the epimerization (E) domain, the terminal condensation (C) domain stereoselectively hydrolyzes D-tryptophanyl-S-phosphopantetheine thioester and thus represents a noncanonical C domain function. Using IvoA, we demonstrate a biocatalytic stereoinversion/deracemization route to access a variety of substituted D-tryptophan analogs in high enantiomeric excess.

Recently, a single-module NRPS, encoded by the gene *ivoA* from *Aspergillus nidulans* with an unusual domain architecture annotated as A-T-C-C* was proposed to acetylate L-tryptophan.⁹ The enzymatic product *N*-acetyl-L-tryptophan was suggested to be further oxidized by a P450 enzyme IvoC and a phenol oxidase IvoB en route to the conidiophore pigment (**Figure 1B**). Although genetic studies provided compelling evidence implicating *ivoA* in the biosynthesis of *N*-acetyl-hydroxytryptophan,^{10,11} the proposed acetyltransferase activity of IvoA is an unlikely fit for a single-module NRPS. Furthermore, the mechanistic proposal for IvoA is at odds with accepted logic of NRPS enzymology for the following reasons: 1) It is metabolically wasteful to activate the carboxy group of a substrate at the expense of one equivalent of ATP in order to accomplish *N*-acetylation by acetyl-CoA; 2) It is against the NRPS directionality rule for a downstream C domain to carry out a condensation reaction (acetylation here) with an upstream T domain as the acceptor.^{1,12}

Nonribosomal peptide synthetases (NRPSs) are modular enzymes employing an assembly-line logic to synthesize a myriad of peptide-based secondary metabolites with diverse structures and biological activities.¹ Single-module NRPS and NRPS-like enzymes adopt similar thiotemplated enzymology with a single set of adenylation (A) and thiolation (T) domain. These enzymes have important functions in transforming carboxylic acid substrates in primary and secondary metabolism;² and have increased interests as biocatalysts due to their functional diversity (**Figure 1**).³ Following selection and thermodynamic activation of the carboxylic acid by the A domain, the substrate is preserved as phosphopantetheine (Ppant) thioester on the T domain. Depending on the type of downstream releasing domains, the thioester intermediates are subjected to a broad range of modifications (**Figure 1a**), including but not limited to: esterification/amidation by a condensation (C) domain in A-T-C;⁴ Dieckman/aldol condensation or cyclization by a thioesterase (TE) domain in A-T-TE;^{5,6} 2- or 4-electron reduction by reductase (R) domain in either A-T-R or A-T-R-R;^{2b,7} 2-electron reduction followed by PLP-dependent aldol condensation in A-T-R-P.⁸ In natural product biosynthetic pathways, these enzymes generate natural product scaffolds with structural diversity that complements the chemical space of canonical nonribosomal peptides, such as dihydroxybenzoquinone, furanone, butyrolactone, and dihydropyrazine (**Figure 1A**).


C ⁰ (H1428A)	0.008 0.002	± 40 ± 10	3.5
-------------------------	----------------	-----------	-----

^aSubstrate inhibition was observed with K_i^{app} of 4 ± 1 (mM).

We next followed the reaction by using chiral-HPLC and complete conversion of L-tryptophan (1 mM) to D-tryptophan was observed in 3 hours (**Figure 2D**). Chiral-resolution allowed us to determine the apparent steady-state kinetics by quantifying D-tryptophan formation (**Table 1**, **Figure S4**). The apparent k_{cat} (38 min^{-1}) is similar to that of other characterized single-module NRPSs and NRPS-like enzymes,^{4,7} while the K_m ($\sim 50 \mu\text{M}$) is close to the intracellular level of L-tryptophan ($12 \mu\text{M}$).¹⁵ Both E and C domains are catalytically important for IvoA, as inactivating either domain by mutating the catalytic histidine residues (H963A and H1428A) substantially compromised the apparent turnover number k_{cat} (420-fold by E⁰ while 4700-fold by C⁰). In contrast, the apparent Michaelis constants were not changed, suggesting that substrate binding (at the A domain) was not affected.

Taken together, these data indicate that IvoA lacks acetyltransferase activity *in vitro*, but instead is a *bona fide* ATP-dependent enzyme catalyzing enantioselective stereo-inversion of L-tryptophan to D-tryptophan. The observed acetylation of D-tryptophan *in vivo* must be carried out by an endogenous acetyltransferase. Because yeast histone acetyltransferase Hpa3 is known to act as a D-amino acid *N*-acetyltransferase for detoxification of D-amino acids,¹⁶ we overexpressed IvoA in the *hpa3*-deleted yeast strain constructed by replacing *hpa3* with *leu2* (SI Methods, Figure S5). Consequently, the culture medium was devoid of *N*-acetyltryptophan, whereas free D-tryptophan (ee = 98%) were still accumulated inside the cells (**Figure S6-S7**). Therefore, we conclude that IvoA does not acetylate tryptophan and the origin of the negligible acetyltransferase activity of IvoA observed *in vitro* may derive from trace amount of contaminated yeast Hpa3.

Scheme 1. Stereo-inversion of amino acid enantiomer into its mirror-image counterpart by common racemases and IvoA.



Distinct from common PLP-dependent or PLP-independent amino acid racemases (**Scheme 1**), which often catalyze bidirectional stereo-inversion and also inevitably lead to racemization (equilibrium constant approaches unity),¹⁷ IvoA catalyzes unidirectional stereo-inversion, completely converting L-tryptophan to its enantiomer D-tryptophan. The complete conversion is driven by coupled ATP hydrolysis, which is thermodynamically favored (**Scheme 1**),¹⁸ and is enabled by the thiotemplate enzymology of IvoA (**Figure 3**). We reason that the activated tryptophan is delivered to the E domain as tryptophanyl-S-Ppant thioester, which undergoes epimerization to give a mixture of D/L-tryptophanyl-S-Ppant diastereoisomers in equilibrium. We propose that dynamic kinetic resolution may be accomplished by the C domain in a releasing step, which stereoselectively hydrolyzes the D-tryptophanyl-S-Ppant thioester to achieve irreversible conversion.




Figure 3. Working model of IvoA.

As mentioned earlier, even though IvoA A domain prefers L-tryptophan, D-tryptophan can still be adenylated and thioesterified (**Figure S2**). In addition, the loaded D-tryptophanyl-S-Ppant underwent epimerization by IvoA E domain as evidenced by similar, yet slower deuterium “wash-in” behavior under multiple-turnover condition (**Figure S8**). The slower turnover measured by deuterium incorporation reflects the lower adenylation efficiency of D-tryptophan. Nonetheless, the occurrence of hydrogen-deuterium exchange at D-tryptophanyl-S-Ppant C α position not only suggests that epimerization is faster than the C-domain catalyzed D-specific tryptophanyl thioester hydrolysis, but also indicates that the D/L-tryptophanyl-S-Ppant equilibrium can be approached from either direction (Figure 3). However, IvoA cannot convert D-tryptophan to L-tryptophan, which suggests that the L-tryptophanyl-S-Ppant is not hydrolyzed by the C domain. A D-specific hydrolytic releasing C domain is therefore the key for unidirectional complete stereo-inversion.

To directly demonstrate the stereoselectivity of IvoA C domain, we purified the standalone IvoA-C and assayed its activity *in vitro*. Addition of IvoA-C in equimolar to either IvoA(C⁰) mutant or IvoA- Δ C truncation mutant successfully rescued the impaired stereo-inversion activity, which proved that the standalone IvoA-C is active (**Figure S9**). We then synthesized both D- and L-tryptophanyl-S-*N*-acetylcysteamine as surrogate substrates mimicking the IvoA T domain bound tryptophanyl-S-Ppant intermediates. However, the enzyme did not catalyze hydrolysis significantly above the background nonenzymatic rate (**Figure S10**). Using D-tryptophanyl-S-pantetheine (D-Trp-pant) also did not improve enzymatic hydrolysis. We reason that the protein:protein interaction between T and C domain is important for substrate recognition, which has been shown in other studies of C domains.¹⁹ Hence, we chose to enzymatically load D/L-tryptophan to IvoA- Δ C(E⁰) by taking advantage of the promiscuous A domain. It is imperative to inactivate the E domain in this truncation mutant in order to minimize the epimerization. The formation of corresponding D/L-tryptophanyl-S-Ppant of IvoA- Δ C was confirmed by intact protein mass spectrometry (**Figure S11**). Free excess D/L-tryptophan substrates were quickly removed from IvoA- Δ C(E⁰) by using desalting spin columns and the loaded D/L-tryptophanyl-S-IvoA- Δ C(E⁰) were immediately subjected to IvoA-C catalyzed hydrolysis. The liberated free tryptophan was then quantified by LC-MS. As shown in **Figure 4**, IvoA-C stereoselectively hydrolyzed D-

tryptophanyl-S-IvoA- $\Delta C(E^0)$ over L-tryptophanyl-S-IvoA- $\Delta C(E^0)$. NRPS C domains that have thioesterase activity are rare, and to date only one example from crocacin PKS-NRPS hybrid assembly-line was known, but did not show stereoselectivity.²⁰ Therefore, the IvoA-C characterized here represents a novel C domain and we classify it as a $^D C_{H2O}$ subtype according to the universally acknowledged nomenclature.¹²





Figure 4. Characterization of IvoA-C activity in vitro by LC-MS.

The verified stereoinversion activity of IvoA prompted us to explore its biocatalytic potential. D-tryptophan and its substituted analogues are important building blocks for many peptide pharmaceuticals, such as FDA approved lanreotide, pasireotide, octreotide, macimorelin, triptorelin, etc. Recently, there is growing interest in developing biocatalytic processes for syntheses of substituted D-tryptophans by stereoinversion and deracemization from the L-enantiomers and *rac*-tryptophans, respectively.²¹ However, to overcome the entropically unfavorable deracemization process ($\Delta G^0 = 0.4$ kcal/mol),²² the current methods are based on multi-step cascade reactions to establish non-equilibrium conditions for enrichment of D-enantiomers.²¹ In contrast, IvoA offers a concise one-step, direct nonredox stereoinversion/deracemization process, and allows us to access a library of D-tryptophan analogues in high enantiomeric excess (ee >99%) at millimolar level. Different substitution groups, either electron-withdrawing or electron-donating, at most positions (4, 5, 6 and 7) on the indole ring can be tolerated (**Table 2**). No conversion of 2-Me-DL-tryptophan is due to inefficient activation by A domain (**Figure S12**), which suggests that substitution at 2-position may interfere with substrate recognition. The poor substrates are generally those with larger substituents (e.g. 5-NO₂, 5-CN, 6-Br, 7-Br), which reflects the size limit by IvoA A domain. In light of recent success in A domain engineering,²³ it is conceivable that the substrate scope can be expanded in the future by enlarging the substrate binding pocket of A domain.

In summary, our biochemical study uncovered the unusual activity of IvoA, and our findings expand the function diversity of single-module NRPSs. The reassigned function of IvoA also provides insight to fungal pigment biosynthesis. By inverting the chirality of tryptophan, IvoA perhaps can modulate amino acid flux to pigment biosynthesis in vivo. Considering the proposed role of IvoB and IvoC, one can speculate that the D-configuration generated by IvoA may be retained in the final uncharacterized conidiophore pigment.

Table 2. Biocatalytic stereoinversion or deracemization of substituted tryptophans.^a



Entry	Substrate		ee (%)
	Stereochemis-try	R	
1	L	H	>99
2	L	5-OMe	>99
3	<i>rac</i>	5-CN	63
4	<i>rac</i>	5-NO ₂	44
5	<i>rac</i>	4-F	>99
6	<i>rac</i>	5-F	>99
7	<i>rac</i>	6-F	>99
8	<i>rac</i>	5-Cl	>99
9	<i>rac</i>	6-Cl	98
10	<i>rac</i>	5-Br	>99
11	<i>rac</i>	6-Br	75
12	<i>rac</i>	7-Br	87
13	<i>rac</i>	2-Me	1.3
14	<i>rac</i>	4-Me	>99
15	<i>rac</i>	5-Me	>99
16	<i>rac</i>	6-Me	97
17	<i>rac</i>	7-Me	>99

^aExpt. Cond.: 1.5 mM substrates, 5 μ M IvoA, 5 mM ATP, 10 mM MgCl₂, 100 mM K₂HPO₄ buffer, pH 7.5.

ASSOCIATED CONTENT

Supporting Information.

This material is available free of charge via the Internet at <http://pubs.acs.org>. Experimental procedures, chromatograms, and spectroscopic data.

AUTHOR INFORMATION

Corresponding Author

Yi Tang yitang@ucla.edu

Notes

No competing financial interests have been declared.

ACKNOWLEDGMENT

This work was supported by the NIH 1R35GM118056 to YT. YH is a Life Sciences Research Foundation fellow sponsored by the Mark Foundation for Cancer Research. MJ is the recipient of a BBSRC Future Leader Fellowship (BB/R01212/1). The Bruker MaxIs II instrument used in this study was funded by the BBSRC (BB/M017982/1). We thank Dr. Dewei Gao for helpful discussion regarding organic synthesis.

REFERENCES

- (1) (a) Fischbach, M. A. and Walsh, C. T. Assembly-line enzymology for polyketide and nonribosomal peptide antibiotics: Logic, machinery, and mechanisms. *Chem. Rev.* **2006**, *106*, 3468-3496. (b) Sieber, S. A. and Mrahiel, M. A. Molecular mechanisms underlying nonribosomal peptide synthesis: approaches to new antibiotics. *Chem. Rev.* **2005**, *105*, 715-738.
- (2) (a) Ehmman, D. E.; Gehring, A. M.; Walsh, C. T. Lysine biosynthesis in *Saccharomyces cerevisiae*: mechanism of α -amino acid reductase (Lys2) involves posttranslational phosphopantetheinylation by Lys2. *Biochemistry* **1999**, *38*, 6176-6177. (b) Hai, Y.; Huang, A. M.; Tang, Y. Structure-guided function discovery of an NRPS-like glycine betaine reductase for choline biosynthesis in fungi. *Proc. Natl. Acad. Sci. USA* **2012**, *109*, 21402-21407. (c) Richardt, A.;

- Kemme, T.; Wagner, S.; Schwarzer, D.; Marahiel, M. A.; Hovemann, B. T. Ebonyl, a novel nonribosomal peptide synthetase for β -alanine conjugation with biogenic amines in *Drosophila*. *J. Biol. Chem.* **2003**, *278*, 41160-41166. (d) Wang, M.; Beissner, M.; Zhao, H. Aryl-aldehyde formation in fungal polyketides: Discovery and characterization of a distinct biosynthetic mechanism. *Chem. Biol.* **2014**, *21*, 257-263.
- (3) (a) Winkler, M. Carboxylic acid reductase enzymes (CARs). *Curr. Opin. Chem. Biol.* **2018**, *43*, 23-29. (b) Ramsden, J. I.; Heath, R. S.; Derrington, S. R.; Montgomery, S. L.; Mangas-Sanchez, J.; Mulholland, K. R.; Turner, N. J. Biocatalytic *N*-alkylation of amines using either primary alcohols or carboxylic acids via reductive amine cascade. *J. Am. Chem. Soc.* **2019**, *141*, 1201-1206.
- (4) (a) Gao, X.; Chooi, Y.-H.; Ames, B. D.; Wang, P.; Walsh, C. T.; Tang, Y. Fungal indole alkaloid biosynthesis: genetic and biochemical investigation of the tryptocalanine pathway in *Penicillium aethiopicum*. *J. Am. Chem. Soc.* **2011**, *133*, 2729-2741. (b) Shinohara, Y.; Takahashi, S.; Osada, H.; Koyama, Y. Identification of a novel sesquiterpene biosynthetic machinery involved in astellolide biosynthesis. *Sci. Rep.* **2016**, *6*, 32865.
- (5) (a) Balibar, C. J.; Howard-Jones, A. R.; Walsh, C. T. Terrequinone A biosynthesis through L-tryptophan oxidation, dimerization and bisprenylation. *Nat. Chem. Biol.* **2007**, *3*, 584-592. (b) Schneider, P.; Weber, M.; Rosenberger, K.; Hoffmeister, D. A one-pot chemoenzymatic synthesis for the universal precursor of anti-diabetics and antiviral bisindolylquinones. *Chem. Biol.* **2007**, *14*, 635-644.
- (6) Hühner, E.; Öqvist, K.; Li, S.-M. Design of α -keto carboxylic acid dimers by domain recombination of nonribosomal peptide synthetase (NRPS)-like enzymes. *Org. Lett.* **2019**, *21*, 498-502.
- (7) Yu, X.; Liu, F.; Zou, Y.; Tang, M.-C.; Hang, L.; Houk, K. N.; Tang, Y. Biosynthesis of strained piperazine alkaloids: uncovering the concise pathway of herquiline A. *J. Am. Chem. Soc.* **2016**, *138*, 13529-13532.
- (8) Baccile J. A.; Spraker, J. E.; Le, H. H.; Brandenburger, E.; Gomez, C.; Bok, J. W.; Macheleidt, J.; Brakhage, A. A.; Hoffmeister, D.; Keller, N. P.; Schroeder, F. C. Plant-like biosynthesis of isoquinoline alkaloids in *Aspergillus fumigatus*. *Nat. Chem. Biol.* **2016**, *12*, 419-424.
- (9) Sung, C. T.; Chang, S.-L.; Entwistle, R.; Ahn, G.; Lin, T.-S.; Petrova, V.; Yeh, H.-H.; Praseuth, M. B.; Chiang, Y. M.; Oakley, B. R.; Wang, C. C. C. Overexpression of a three-gene conidial pigment biosynthetic pathway in *Aspergillus nidulans* reveals the first NRPS known to acetylate tryptophan. *Fungal Genet. Biol.* **2017**, *12*, 419-424.
- (10) Clutterbuck, A. J. A mutational analysis of conidial development in *Aspergillus nidulans*. *Genetics*, **1969**, *63*, 317-327.
- (11) McCorkindale, N. J.; Hayes, D.; Johnston, G. A.; Clutterbuck, A. J. *N*-Acetyl-6-hydroxytryptophan a natural substrate of a monophenol oxidase from *Aspergillus nidulans*. *Phytochemistry*, **1983**, *22*, 1026-1028.
- (12) Bloudoff, K. and Schmeing, T. M. Structural and functional aspects of the nonribosomal peptide synthetase condensation domain superfamily: discovery, dissection and diversity. *Biochem. Biophys. Acta*, **2017**, *1865*, 1587-1604.
- (13) Harvey, C. J. B.; Tang, M.; Schlecht, U.; Horecka, J.; Fischer, C. R.; Lin, H. C.; Naughton, B.; Cherry, J.; Miranda, M.; Li, Y. F.; Chu, A. M.; Hennessy, J. R.; Vandova, G. A.; Inglis, D.; Aiyar, R. S.; Steinmetz, L. M.; Davis, R. W.; Medema, M. H.; Sattely, E.; Khosla, C.; St Onge, R. P., Tang, Y.; Hillenmeyer, M. E. Hex: A heterologous expression platform for the discovery of fungal natural products. *Sci. Adv.* **2018**, *11*, eaar5459.
- (14) Stachelhaus, T.; Walsh, C. T. Mutational analysis of the epimerization domain in the initiation module PheATE of gramicidin S synthetase. *Biochemistry* **2000**, *39*, 5775-5787.
- (15) Bennett, B. D.; Kimball, E. H.; Gao, M.; Osterhout, R.; Van Dien, S. J.; Rabinowitz, J. D. Absolute metabolite concentrations and implied enzyme active site occupancy in *Escherichia coli*. *Nat. Chem. Biol.* **2009**, *5*, 593-599
- (16) (a) Yow, G. Y.; Uo, T.; Yoshimura, T.; Esaki, N. D-amino acid *N*-acetyltransferase of *Saccharomyces cerevisiae*: a close homologue of histone acetyltransferase Hpa2p acting exclusively on free D-amino acids. *Arch. Microbiol.* **2004**, *182*, 396-403. (b) Yow, G. Y.; Uo, T.; Yoshimura, T.; Esaki, N. Physiological role of D-amino acid *N*-acetyltransferase of *Saccharomyces cerevisiae*: detoxification of D-amino acids. *Arch. Microbiol.* **2006**, *185*, 39-46. (c) Sampath, V.; Liu, B.; Tafrov, S.; Srinivasan, M.; Rieger, R.; Chen, E. I.; Sternglanz, R. Biochemical characterization of Hpa2 and Hpa3, two small closely related acetyltransferases from *Saccharomyces cerevisiae*. *J. Biol. Chem.* **2013**, *288*, 21506-21513.
- (17) (a) Tanner, M. E. Understanding Nature's strategies for enzyme-catalyzed racemization and epimerization. *Acc. Chem. Res.* **2002**, *35*, 237-246. (b) Fischer, C.; Ahn, Y.-C.; Vederas, J. C. Catalytic mechanism and properties of pyridoxal 5'-phosphate independent racemases: how enzymes alter mismatched acidity and basicity. *Nat. Prod. Rep.* **2019**, in press, DOI: 10.1039/c9np00017h.
- (18) Calculated under 0.25 M ionic strength at pH 7 according to Alberty, R. A. Calculating apparent equilibrium constants of enzyme-catalyzed reactions at pH 7. *Biochem. Educ.* **2000**, *28*, 12-17.
- (19) (a) Hai, Y. and Tang, Y. Biosynthesis of long-chain *N*-acyl amide by a truncated polyketide synthase-nonribosomal peptide synthetase hybrid megasynthase in fungi. *J. Am. Chem. Soc.* **2018**, *140*, 1271-1274. (b) Gao, X.; Haynes, S. W.; Ames, B. D.; Wang, P.; Vien, L. P.; Walsh, C. T.; Tang, Y. Cyclization of fungal nonribosomal peptides by a terminal condensation-like domain. *Nat. Chem. Biol.* **2012**, *8*, 823-830.
- (20) Müller, S.; Rachid, S.; Hoffmann, T.; Surup, F.; Volz, C.; Zaburanyi, N.; Müller, R. Biosynthesis of crocacin involves an unusual hydrolytic release domain showing similarity to condensation domains. *Chem. Biol. Catal.* **2014**, *21*, 855-865.
- (21) (a) Schnepel, C.; Kemker, I.; Sewald, N. One-pot synthesis of D-halotryptophans by dynamic stereoinversion using a specific L-amino acid oxidase. *ACS Catal.* **2019**, *9*, 1149-1158. (b) Parmegiani, F.; Casamajo, A. R.; Walton, C. J. W.; Galman, J. L.; Turner, N. J.; Chica, R. A. One-pot biocatalytic synthesis of substituted D-tryptophans from indoles enabled by an engineered aminotransferase. *ACS Catal.* **2019**, *9*, 3482-3486. (c) Matsuyama, A.; Mitsuhashi, K.; Tokuyama, S.; Yamamoto, H. D-aminoacylase and gene encoding the same. U. S. Patent 6,887,697 B2, 2005.
- (22) Gruber, C. C.; Lavandera, I.; Faber, K.; Kroutil W. From a racemate to a single enantiomer: deracemization by stereoinversion. *Adv. Synth. Catal.* **2006**, *348*, 1789-1805.
- (23) (a) Ishikawa, F.; Miyanaga, A.; Kitayama, H.; Nakamura, S.; Nakanishi, I.; Kudo, F.; Eguchi, T.; Tanabe, G. An engineered aryl acid adenylation domain with an enlarged substrate binding pocket. *Angew. Chem. Int. Ed. Engl.* **2019**, *58*, 6906-6910. (b) Niquille, D. L.; Hansen, D. A.; Mori, T.; Fercher, D.; Kries, H.; Hilvert, D. Nonribosomal biosynthesis of backbone-modified peptides. *Nat. Chem.* **2018**, *10*, 282-287.




Table of Contents
