

Trends in Endocrinology and Metabolism: invited review

Going back to the biology of FGF21: new insights

Jo E Lewis¹, Francis J P Ebling², Ricardo J Samms³, Kostas Tsintzas²

¹Institute of Metabolic Sciences and MRC-Metabolic Diseases Unit, University of Cambridge, Cambridge, CB0 0QQ, UK

²School of Life Sciences, University of Nottingham Medical School, Queen's Medical Centre, Nottingham, NG7 2UH, UK.

³Lilly Research Laboratories, Indianapolis, USA.

Email correspondence to: kostas.tsintzas@nottingham.ac.uk

MRC/ARUK Centre for Musculoskeletal Ageing
School of Life Sciences
University of Nottingham Medical School

Queen's Medical Centre

Nottingham

NG7 2UH

UK

tel: +44 115 82 30127

Word count: 4786 (excluding Abstract, Highlights, Text boxes and Outstanding questions)

Figures: 2

Abstract

Fibroblast growth factor 21 (FGF21) is a protein highly synthesised in the liver that exerts paracrine and endocrine control of many aspects of energy homeostasis in multiple tissues. In preclinical models of obesity and type 2 diabetes, treatment with FGF21 improves glucose homeostasis and promotes weight loss, and as a result FGF21 has attracted considerable attention as a therapeutic agent for the treatment of metabolic syndrome in humans. An improved understanding of the biological role of FGF21 may help to explain why its therapeutic potential in humans has not been fully realised. This review will cover the complexities in FGF21 biology in rodents and humans, with emphasis on its role in the protection from central and peripheral facets of obesity.

FGF21 – a novel metabolic regulator

Fibroblast growth factor 21 (FGF21) is a hormone shown to play a key role in the regulation of energy homeostasis in both preclinical studies and man. Consequently, leveraging the broad metabolic effects of FGF21 for the treatment of metabolic syndrome has received significant interest from the pharmaceutical industry. The therapeutic potential of FGF21 was first recognised when it was reported that FGF21 reduces plasma glucose and triglycerides in mouse models of obesity, and that overexpression of FGF21 provided resistance to body weight gain in mice on a high fat diet (1,2). However, contemporary data challenge existing concepts, and many facets of FGF21 biology remain divisive and/or poorly understood. Furthermore, only some of the metabolic effects characteristic of FGF21 action in preclinical studies using laboratory rodents are evident in humans, representing a significant challenge for the translational potential of engineered FGF21-based therapeutics. Indeed, several human clinical trials using FGF21 analogues have failed to observe meaningful blood glucose lowering and insulin sensitizing effects, although significant improvements in dyslipidaemia and markers of fatty liver disease have been reported (see text Box 1). Consequently, FGF21 biology is rapidly moving into new arenas – control of macronutrient preference, providing protection from nonalcoholic steatohepatitis and nonalcoholic fatty liver disease (NASH/NALFD respectively), or more broadly as a biomarker for disease. Hence, there is a need to go back to the natural biology of FGF21 with the objective of better understanding its role in these diseases with the aim of maximizing its therapeutic potential.

Box 1 - Clinical trials with FGF21 analogues

Long-acting FGF21 analogues have been utilised in several human clinical trials showing improvement in some, but not all aspects of the metabolic syndrome. In overweight/obese patients with type 2 diabetes (T2D), treatment with LY2405319, a variant of FGF21, produced improvements in fasting insulin levels but only a modest, nonsignificant fasting glucose lowering effect (3). Similarly, treatment with PF-05231023, a long-acting FGF21 analogue, and Pegbelfermin (BMS-986036), a PEGylated human FGF21 analogue, had no effects on glycaemic control and HbA1c in overweight/obese patients with T2D (4-6). The effects of those analogues on body weight was variable, with no change observed after 12 weeks of treatment with BMS-986036 (6), a small reduction of 1.8% in response to 4 weeks of LY2405319 (3) and a modest loss of 5-6% after 4 weeks on PF-05231023 (4). However, all trials reported improvements in dyslipidaemia with significant reductions in plasma triglycerides (up to 50%) and improved plasma lipoprotein profile (reductions of total cholesterol by up to 20% and LDL by ~30%, and increases in HDL by up to 25%). Overall, the effects of FGF21 analogues on aspects of glucose control and body weight loss are modest at best and cannot match the improvements observed with

current pharmacological and lifestyle changes-based treatments targeting weight loss and/or improvement of β -cell function and insulin sensitivity in patients with T2D (7). However, the therapeutic potential of FGF21 analogues may be realised when combined with other current therapies targeting glycaemic control or as a monotherapy to improve plasma lipid profiles and for the treatment of NASH/NAFLD (8) (see text Box No 4).

Since its initial isolation from liver in 2000 (9), FGF21 has been shown to be widely expressed in metabolic organs including the gastrointestinal tract, adipose tissue and pancreas (10). FGF21 has been demonstrated to cross the blood-brain barrier in rats, is detectable in human cerebrospinal fluid [CSF], and there is a linear relationship between serum and CSF levels (11,12). Furthermore, FGF21 biology demonstrates circadian rhythmicity – serum concentrations exhibit a clear diurnal rhythm, and dysregulation correlates with obesity-induced lipid disorders (13). Despite this and its widespread distribution, various strains of FGF21 knockout (KO) mice show variable effects on body weight and composition, and circulating glucose. This in itself is controversial, with some groups reporting fasting hypoglycaemia – a possible consequence of the age of the animals and/or the fasting schedules (14-16). Although these animals showed normal growth, development and fertility, subsequent studies have suggested that FGF21 may have adverse effects on these processes but is not essential (1,16-20). Thus, the physiological role of FGF21 is in the adaption to altered metabolic demand, for example in times of stress; levels of FGF21 are increased under a variety of physiological interventions and pathogenic events, including starvation (acutely in mice, chronically in humans), obesity (in mice and humans) and exercise (in mice and humans) (17,21-23). Furthermore, the effects of treatment with exogenous FGF21 are pleiotropic. Here we will explore the complexities in FGF21 biology and physiology in animal models and adult humans, its pharmacology and metabolic actions from a preclinical and clinical perspective.

Box 2 - The biology of FGF21

FGF21 (gene ID 56636) is located on chromosome 7, contains three coding exons, which produces a single transcript, a pre-protein of 210 amino acids (aa) with a N-terminal signal peptide of 22aa. In humans, FGF21 (gene ID 26291) also contains three exons, however, this produces two transcripts via alternative promoters, both of which encode a pre-protein of 208aa, including a 28aa signal peptide. The sequence identity is circa 80%. Functionally the FGF superfamily is grouped into three subfamilies; intracellular FGFs which lack a signal peptide, FGFs which lack heparin sulphate binding capacity, indicating the potential for their escape into the circulation, exerting systemic action, and FGFs with a high heparin sulphate binding capacity, thus potentially act in an autocrine/paracrine manner (24). At target tissues, FGF21 binds to and activates members of the FGFR

superfamily of receptor tyrosine kinases (25). FGFR activation by FGF21 *in vitro* and *in vivo* is dependent on obligate co-receptor β Klotho (KLB), an FGFR-binding, single-pass transmembrane protein (2,26). Preference for both FGFR1c- and FGFR3c-KLB complexes has been demonstrated, so given the widespread expression of the FGFRs, tissue specificity and sensitivity is mainly governed by the presence and relative expression of KLB (27-30). Activation of these receptor complexes by FGF21 leads to a plethora of rapid signaling events (31,32).

Box 3 - Tissue specific physiology

The highest level of FGF21 expression in the adult mouse is the liver. The liver is regarded as the major source of circulating FGF21; liver-specific deletion of FGF21 created by crossing FGF21-floxed mice (FGF21^{fl/fl}) with an albumin-Cre line results in dramatic reductions in circulating FGF21, and the increase typically observed in response to fasting is lost (33). Peroxisome proliferator-activated receptor α (PPAR α), a nuclear receptor activated by fatty acids, is required for the normal adaptive response to starvation (34). Mice lacking PPAR α accumulate hepatic triglyceride and become hypoketonemic during fasting and starvation (35-37). A downstream target of PPAR α , hepatic FGF21 increases in response to dietary manipulation (fasting, ketogenic diet) and is rapidly suppressed by refeeding in mice (17,18,38). Indeed, FGF21 was offered as the “missing link” in the biology of fasting, inducing adipose tissue lipolysis, liver ketogenesis and metabolic adaptation to the fasting state (39). Furthermore, FGF21^{-/-} mice showed impaired adaptation to ketosis, whilst adenoviral knockdown of FGF21 in the liver caused mice consuming a ketogenic diet to display fatty livers, hypertriglyceridemia and reduced fat oxidation (15). Having established that FGF21 was critical to the adaptive metabolic response to starvation in mice, data emerged demonstrating the induction of FGF21 in response to amino acid deprivation and protein restriction. This is mediated by the eukaryotic translation initiation factor (EIF) 2 α -activating transcription factor (ATF)4-CHOP axis of the ER stress response in mice (40). Mice consuming diets deficient in methionine, leucine and alanine demonstrated increased serum FGF21 concentrations (40-42). Furthermore, consumption of methionine/choline-deficient diets in mice results in NASH and is associated with increased hepatic expression and serum levels of FGF21. FGF21 knock out mice have an exacerbated phenotype on this deficient diet, in part due to attenuated activation of fatty acid oxidation, as well as increased expression of genes involved in lipid uptake and synthesis (43,44). Pharmacological treatment of these mice with FGF21 resolved the phenotype.

Box 4 - Lessons from rodent models provide a liver-centric view.

One of the many paradoxes in the FGF21 field is that hepatic FGF21 expression is also increased in response to nutrient excess, a process described in rodents and humans that is likely mediated via the carbohydrate response element binding protein (ChREBP) that may participate in the pathogenesis of NAFLD (9,21,45-47). FGF21 levels in liver and adipose tissue are also elevated in db/db obese diabetic mice as compared with their lean db/+ littermates, in adult male rhesus macaque monkeys chronically maintained on a high fat diet (HFD), and in serum in overweight/obese humans (48,49). Serum FGF21 is positively correlated with adiposity, fasting circulating insulin and triglycerides – logistic regression analysis demonstrated an independent association between serum FGF21 and metabolic syndrome (48). It was subsequently shown that in response to carbohydrate intake, the

liver produces FGF21 to selectively suppress sugar intake by acting on the paraventricular nucleus of the hypothalamus (50) (see the CNS section below). Therefore, FGF21 is a key regulator of the nutritional state of the organism and co-ordinates the adaptive metabolic response. Whilst the induction of FGF21 in response to changes in macronutrient ingestion is important for metabolic adaptation. Whether the liver is an important site of FGF21 action, however, is debatable. ERK1/2 phosphorylation and mRNA expression of immediate early gene markers downstream of the FGFR-KLB complex that are readily observed in adipose tissue in response to treatment with FGF21 are not so apparent in liver. However, Akt phosphorylation in the liver is reported in lean and ob/ob animals, in which FGF21 suppresses glucose production and increases insulin sensitivity (51). Furthermore, other groups have shown that FGF21 does induce FRS2 and ERK1/2 phosphorylation in the liver of both lean and obese animals – with a time course and dose response similar to adipose tissue (28,52). Furthermore, in HepG2 cells, FGF21 induces ERK1/2 phosphorylation providing evidence that FGF21 may act directly on the hepatocyte (53). Indeed, FGF21 exerts dual effects on hepatic energy metabolism, modulating net flux through both carbohydrate and fat metabolism (54) (see Seasonal Lessons section below). Therefore, by stimulating hepatic fat oxidation, FGF21 may be an attractive target for protection from increased hepatic lipid content and insulin resistance that frequently accompanies obesity and T2D. Indeed, targeting hepatic fat accumulation and insulin resistance is a major therapeutic approach to the management of NAFLD and NASH, two of most prevalent forms of liver disease that are closely associated with obesity and T2D (55). An engineered variant of FGF21 has been shown to attenuate NASH progression in an animal model of NASH (56). Whereas a clinical phase 2 trial (NCT02413372) with a recombinant FGF21 in NASH patients significantly reduced hepatic fat content and biomarkers of fibrosis (57). Therefore, FGF21 may be emerging as a new class of drug for the treatment of fatty liver disease.

Adipose tissue: a key determinant of FGF21 action

Adipose tissue is a highly dynamic organ, which in response to the internal environment, undergoes extensive remodelling. Caloric restriction, for example, promotes mitochondrial biogenesis and fatty acid oxidation, whereas white adipose tissue rapidly expands in response to an obesogenic diet by increasing lipid storage and the number of adipocytes (58,59). These are evolutionary protective mechanisms to ensure adequate glucose supply for the brain, and delivery of free fatty acids (FFA) and amino acids to peripheral organs. Adipose tissue is known to play a crucial role in the buffering of diet-derived FFA in the post-prandial period (60). If this role, however, is impaired due to a disruption of insulin signaling in adipose tissue, excessive lipids accumulate in other tissues such as liver and skeletal muscle, and may result in the development of insulin resistance (61). Therefore insulin-sensitising agents targeting adipose tissue are in high demand.

Initially, FGF21 was shown to stimulate glucose uptake in 3T3-L1 adipocytes via glucose transporter 1 (GLUT1) (2). An effect dependent on ERK1/2 and ELK-1 signaling (27). Indeed, acute administration of FGF21 to ob/ob and db/db mice or to mice with diet-induced obesity rapidly improved insulin sensitivity and decreased plasma glucose levels (51,62). However, the effect on glucose uptake and GLUT1 in adipose tissue was not apparent in chronically treated animals, despite lower blood glucose, which was achieved via the suppression of hepatic glucose production (63). Subsequently, FGF21 was shown to increase glucose uptake in BAT and WAT in lean mice, suggesting a differential response to FGF21 dependent on metabolic status and/or adiposity (64).

Chronic treatment of mice with FGF21 did, however, increase energy expenditure and result in weight loss (65,66), and, as a consequence, improve insulin sensitivity. However, the role of uncoupling respiration via activation of UCP1 to the pharmacological effects of FGF21 is not completely resolved. Studies using UCP1-KO mice have shown that several but not all metabolic endpoints are UCP1 independent (67,68). Whereas UCP1-mediated thermogenesis was not required for FGF21 to lower body weight, improve glycaemic control or reduce circulating insulin and lipids (67,68), the ability of FGF21 to promote postprandial glucose tolerance was impaired in the absence of UCP1 in obese, glucose intolerant mice (69). On the other hand, the ability of FGF21 to augment energy expenditure was significantly attenuated in one (67) but not another study (68). Further research is required to elucidate these seemingly contradictory findings.

FGF21 signaling in adipose tissue was essential for the acute insulin-sensitizing effects of FGF21, specifically in brown adipocytes, but not for its effects on body weight – this effect was via non-adipose tissues (70). Interestingly, BAT is present in humans and correlates negatively with BMI and fasting blood glucose levels (71). However, humans possess relatively less BAT when compared to rodents which may account for the attenuated glucose lowering responses to FGF21 treatment observed in human clinical trials (Please see text Box 1). Given the variable distribution of BAT in different populations (e.g. more commonly associated with women and lean individuals) (71), further targeted clinical work is required to determine the contribution of BAT in the metabolic responses to FGF21 in human subjects.

The increase in circulating FGF21 in response to fasting resulted in the hypothesis that FGF21 induces lipolysis in adipose tissue. Indeed, this was demonstrated *in vivo* (15,18). However, *in vitro*, FGF21 was shown to be a potent repressor of lipolysis in mouse (and human) adipocytes (72,73). Acute doses and chronic treatment with FGF21 reduce FFA (66,72). Further work is required to dissect this duality and determine the precise role of FGF21 in lipolysis. At target tissues, FGF21 binds to and activates members of the FGFR superfamily of receptor tyrosine kinases (25). FGFR activation by FGF21 is dependent on obligate co-receptor β Klotho (KLB), an FGFR-binding, single-pass transmembrane protein (2,26). Mice lacking the FGFR1 receptor in adipose tissue show attenuated responses to the metabolic effects of FGF21 (74,75). Similarly, mice with complete or adipose-specific ablation of KLB are resistant to acute insulin-sensitising and glucose-lowering effects of FGF21 (74,76). In mice, KLB expression is seen in adipose tissue, the liver, gall bladder, colon and pancreas, as well as the suprachiasmatic and paraventricular nuclei of the hypothalamus, and discrete regions of the brain stem (10,19). Indeed, over-expression of KLB in adipose tissue sensitises mice to endogenous FGF21 and offers protection from diet-induced obesity (29). It was subsequently shown that KLB levels are markedly decreased in white adipose tissue during diet-induced obesity. Interestingly, TNF- α represses KLB expression and impairs FGF21 action in adipocytes *in vitro*, a mechanism mediated by TNF- α induced inflammation as the insulin-sensitizing anti-inflammatory drug rosiglitazone restored KLB levels (28). Furthermore, high glucose represses KLB expression and impairs FGF21 action in mouse pancreatic islets (77). Therefore, the concept of endogenous “FGF21 resistance” has been raised – the biology of which requires further investigating as impairment of the pharmacological actions of FGF21 has not been shown. Furthermore, a hugely interesting aspect of FGF21 biology is the recent evidence from our animal studies (see Seasonal Lessons below) that the actions of FGF21 are enhanced in states of increased adiposity. Therefore, increasing KLB expression would be an interesting therapeutic option. Indeed, GLP-1 analogues and glitazones increase KLB expression *in vivo* (78,79) and may therefore increase FGF21 sensitivity, contributing to further metabolic improvements. On the other hand, maintenance of KLB protein expression in white adipose tissue does not alleviate impaired FGF21 signaling in this tissue or increase FGF21 sensitivity *in vivo* (80). Despite these contradictory results on the role of KLB, adipose tissue appears to be important to the biology of FGF21 and its insulin sensitising effects.

The CNS: the final frontier for FGF21

The CNS integrates a vast array of signals from the periphery to regulate feeding behaviour and energy homeostasis - dysregulation results in obesity and T2D. These signals (hormones and nutrients, plus their by-products) act directly on the hypothalamus, the hindbrain or peripheral neurones which terminate in the nucleus of the solitary tract (NTS)(81). As noted above, FGF21 can cross the rodent blood-brain barrier (11) and is detectable in human cerebrospinal fluid with a linear relationship between serum and CSF levels suggesting that circulating FGF21 is the source of central FGF21 (12) – this is key for therapeutics whose primary targets are the CNS. The FGFR subtypes are widely expressed in the CNS, whilst KLB is highly expressed in the suprachiasmatic nucleus in the hypothalamus and in a number of hindbrain regions (19). The film autoradiography/*in situ* hybridization strategy used by Bookout et al 2013 (19) does not exclude the possibility that KLB might be expressed at lower levels or in isolated cells in other regions of the brain, indeed Liang et al., 2014 (14) reported Western blot and immunostaining of KLB in the paraventricular nucleus of the hypothalamus of mice (82). Overexpression of FGF21 lowers circulating levels of insulin and activation of the HPA axis, and suppresses the reproductive axis (14,19,20). Furthermore, ICV injection of FGF21 increases sympathetic nerve activity in BAT – this effect is attenuated in mice lacking central KLB. Interestingly, treatment with a generic beta-blocker attenuated the effects of centrally administered FGF21 but not of FGF21 delivered peripherally. To exert its effects on energy expenditure, weight loss and cholesterol in mouse models of obesity, central action of FGF21 may be required (83,84).

More recently, FGF21 has been shown to suppress the consumption of simple sugars and non-caloric sweeteners via actions on hypothalamic neurons (50). In addition, FGF21 reduces alcohol preference in mice, whilst stimulating water drinking behaviour (85,86). The latter effect is inhibited by β -adrenergic receptor antagonists, suggesting a critical role of the sympathetic nervous system in mediating the effects of FGF21 in those responses. The emergence of FGF21 as a regulator of food preference both in rodents and humans, along with its role in metabolic regulation, raises the interesting prospect of its therapeutic potential in the dual regulation of appetite and hedonic behaviour(s), and energy expenditure in obesity.

Seasonal lessons

Whilst the ready availability of genetic tools have resulted in mice being the dominant animal model in recent years for studying mechanisms underlying energy metabolism, their high metabolic rate, rapid growth and dependence on brown fat for thermogenesis when maintained under standard housing limits their translational value (87,88). Most mammals are seasonal in that they display profound annual cycles in appetite, fat metabolism, reproduction and body weight (89). Investigating the function of FGF21 in this context provides different insights into its biology, because deposition of body fat does not result in metabolic dysfunction. Thus, it becomes possible to study the effects of FGF21 in healthy obesity without the complications of the development of insulin resistance or NAFLD. Siberian hamsters (*Phodopus sungorus*) have been widely used as a seasonal rodent model because they can be readily maintained in animal facilities, and seasonal cycles in energy metabolism generated simply by changing the environmental daylength (90). Under long day conditions, typically with the light phase being 14 hours of light per day or more, these hamsters become hyperphagic and rapidly gain body weight, whereas under short days with the light phase being 8 hours per day or shorter they reduce their food consumption, and lose up to 30% of body weight over the course of a few months. This primarily reflects loss of visceral adipose depots, though significant loss of lean mass also occurs (91). Thus, studies in this kind of seasonal mammal can address the role of FGF21 in natural situations of weight loss and gain, and provide the opportunity to investigate mechanisms of action of FGF21 in naturally occurring states of adiposity and leanness (Figure 1).

When exposed to short photoperiods, male hamsters show a significant increase in circulating FGF21 by 8 weeks that correlates with significant weight loss and decrease in epididymal fat pad mass at this time point (92). Western blot analysis reveals that whilst FGF21 is detectable in liver, white fat, brown fat and in muscle (gastrocnemius), only in the liver and brown fat are there significant increases in FGF21 concentrations as hamsters progress into short photoperiods (92). The close correlation between liver and plasma FGF21 concentrations supports the argument that the liver is the predominant source of circulating FGF21. Similar observations have been made in another seasonal species, the thirteen-lined ground squirrel (*Ictidomys tridecemlineatus*). Serum FGF21 concentrations increase in squirrels exposed to short photoperiods in the fall, the increase being greater in animals in torpor than

during bouts of arousal in winter, and concentrations then fall again as photoperiod is increased in the following spring (93). Again these circulating FGF21 concentrations closely correlate with expression of FGF21 mRNA in the liver, as assessed by quantitative PCR (93). A key question is whether the increases in circulating FGF21 contribute to the behavioural and physiological adaptations induced by short photoperiod. Chronic infusion of hamsters maintained in summer long day photoperiods with FGF21 results in some aspects of the short day response, for example decreased appetite and decreased body weight, reduced weight of epididymal fat depots, and a decrease in respiratory exchange ratio that would be consistent with increased whole body fat oxidation (94,95) (Figure 1). These effects of FGF21 infusion were lost in hamsters tested after 12 weeks exposure to short day winter photoperiods at which point they had reached the nadir of the seasonal body weight cycle, and effects were also attenuated in ageing male Siberian hamsters that had reduced body weight (94). These observations in naturally-altered states of adiposity support the view that a major target of this hormone is white adipose tissue. This conclusion is supported by analysis of perirenal and subcutaneous white adipose tissue in hamsters treated with FGF21. For example, expression of *fgfr1c* and *βklotho* are upregulated in FGF21-treated hamsters, triglyceride lipase is increased and triglyceride content are reduced, and signal transduction events downstream of FGFR1c-βKlotho receptors including ERK1/2 phosphorylation are increased (94).

Although these studies in hamsters indicate that white adipose tissue is the main determinant of whole-body responsiveness to FGF21, they also identify additional targets (Figure 1). The thermogenic genes *ucp1* and *dio2* (deiodinase 2) were significantly upregulated in interscapular brown fat (94), which likely underlies increases in whole body energy expenditure detected in hamsters infused with FGF21 (95). Hepatic energy metabolism is also affected such that fatty acid oxidation is favoured over carbohydrate oxidation. Indeed, FGF21 infusion reduced pyruvate dehydrogenase activity (PDC, the enzyme that catalyses the rate limiting step in glucose oxidation) and acetylcarnitine accumulation (an index of reduced flux through PDC), but increased long-chain acylcarnitine content (suggesting an increased flux through β-oxidation) in the liver (54). These dual effects of FGF21 on hepatic energy metabolism were mediated by increased protein abundance of PDK4 (a negative regulator of PDC), the phosphorylated (inhibited) form of acetyl-CoA carboxylase (a negative regulator of

delivery of fatty acids into the mitochondria), and the transcriptional co-regulators of energy metabolism PGC1 α and SIRT-1 (54).

Intriguingly, it appears likely that systemic FGF21 may also target the brain. There are consistent suppressive effects of FGF21 on appetite in hamsters (94,95). This is in contrast to studies in mice where FGF21 treatment has been associated with increased food intake, presumably a compensatory response to enhanced whole body energy expenditure (96). Whilst these effects do not necessarily indicate a direct action centrally, perhaps reflecting altered signaling from other peripheral signals to the brain, in hamsters the appetite-suppressing actions of FGF21 are associated with decreased levels of peripheral anorectic signals such as leptin and insulin (94). Importantly, peripheral FGF21 induced upregulation of *dio2* in the hamster hypothalamus (97,98). This enzyme that is highly expressed in hypothalamic tanycytes converts the thyroid hormone precursor thyroxine into the biologically active form tri-iodothyronine (T3). Altered local concentrations of T3 in the hypothalamus are considered to be the main driver of season cycles of appetite and energy expenditure (99,100), so this is a likely pathway of action for FGF21 regulating food intake in hamsters. Further evidence from studies in mice for central actions of FGF21 is considered in a previous section. The distribution of β Klotho has not yet been demonstrated in the hamster, but FGFR1c is expressed in the tanycyte cell layer of hamsters (101). Intracerebroventricular treatment of hamsters with an anti-FGFR1c monoclonal antibody that may mimic the actions of FGF21 decreases appetite, increases whole body energy expenditure and fat oxidation, and induces weight loss (101). Given the recent evidence that tanycytes are in effect part of the blood brain barrier in the hypothalamus and are critical in sensing nutrients and peripheral hormones (102), this is a likely target for peripheral FGF21 that requires further validation.

The catabolic effect of FGF21 are consistent with a role in facilitating the hypophagic and lipolytic state that occurs in hamsters in short days, but it should be borne in mind that the circulating concentrations of FGF21 delivered in these studies is in the supraphysiological range (95). Nelson et al tested the role of FGF21 in generating seasonal metabolic responses by overexpressing FGF21 in ground squirrels using an adenoviral approach (103). This increased circulating FGF21 was associated with decreased circulating insulin and FFA concentrations, but was not sufficient to induce winter torpor bouts (93). It is clear from

analysis of hypothalamic gene expression that hamsters do not respond to the decrease in circulating leptin that reflects reduced caloric intake and increased lipolysis in short days with compensatory hyperphagia (104). Perhaps the role of FGF21 in short days then is to promote metabolic adaptation to negative energy balance, rather than be an underlying driver of winter physiology. Given the efficacy of FGF21 in hamsters in an obese but not metabolically compromised state, perhaps the optimal strategy for clinical use of FGF21 in humans would be in obese populations, particularly those with risk factors for cardiovascular and fatty liver diseases, prior to the development of T2D.

Advances in the biology of FGF21 – a human perspective

FGF21 has been implicated in a number of clinical and non-clinical conditions in humans including the metabolic syndrome and T2D, hepatic liver disease and metabolic regulation during exercise (48,105-108). However, there are still unanswered questions remaining around its biological role in humans as a starvation hormone, exercise-induced myokine and postprandial regulator of metabolism, in particular its link to insulin action and the metabolic syndrome.

FGF21 is often described as a key hormone in the biology of fasting in a variety of animal models. However, in humans, neither ketogenic diets (up to 3 months) nor short-term fasting (up to 48h) increase serum FGF21 levels (109). Interestingly, elevations in FGF21 in humans are seen following prolonged fasting (7-10 days); a possible consequence of the divergent metabolic rates and adaptations to fasting between the species (22,110). Since the ketogenic response in humans precedes the induction of FGF21 and ketogenic diets are associated with a reduction in circulating levels of FGF21, it appears that induction of FGF21 is not required to stimulate ketosis in humans but may serve to facilitate metabolic adaptations to prolonged periods of food deprivation (109).

Importantly, recent evidence suggests that FGF21 is an insulin-dependent postprandial hormone in humans (Figure 2). Indeed, short-term dietary carbohydrate, but not fat interventions, markedly increase the fasting circulating levels of FGF21 (47,111). Furthermore, FGF21 secretion has been shown to respond to oral carbohydrate consumption, with fructose reported to increase plasma levels of FGF21 in humans to a greater extent than

glucose (112). Fructose ingestion was associated with elevated levels of circulating insulin, albeit to a lesser extent than glucose ingestion, which makes it difficult to dissociate its insulin-dependent and -independent effects. Using oral glucose tolerance tests and a combination of hyperglycaemic and hyperinsulinaemic clamps, Samms et al., 2017 (111) subsequently demonstrated that insulin rather than glucose *per se* increases both the total and the bioactive form of FGF21 in the postprandial period in humans. This supports observations from earlier cross-sectional studies showing a positive correlation between plasma insulin and total FGF21 levels in human subjects (48,113). The Fibroblast Activation Protein (FAP) is a serine protease that cleaves and inactivates FGF21 (114). Interestingly, patients with T2D have elevated circulating levels of FAP when compared to non-diabetic controls, and this is associated with an attenuated ratio of bioactive to total FGF21 in response to an oral glucose tolerance test, suggesting that chronic states of insulin resistance may lead to inactivation of FGF21 and limit its metabolic effects (111). Indeed, in response to a high fat dietary intervention that is known to alter insulin-stimulated substrate utilization in healthy humans, the normal induction of circulating FGF21 in response to insulin is attenuated (111). Confirming the mechanism by which insulin and/or insulin resistance regulates the hepatic secretion of FGF21 and FAP in humans and the subsequent tissue specific actions of FGF21 require further investigation. Given that patients with T2D have elevated circulating levels of FAP, specific inhibitors for FAP in those patients (in combination with FGF21 mimetics) could be an appealing therapeutic strategy, particularly as FGF21 was shown to improve some but not all aspects of the metabolic syndrome in humans (see text Box 1).

Acute physiological elevations in plasma FFA levels through intravenous infusion of heparin and lipid emulsions was shown to produce modest elevation of circulating FGF21 (115), although in healthy human volunteers undergoing a lipid tolerance test, acute elevation of plasma FFAs was shown to decrease FGF21 (116,117). In contrast, overfeeding with a high fat diet for 5 days resulted in a robust increase in circulating levels of FGF21 (118), possibly as a compensatory/adaptive mechanism to increased fat oxidation under conditions of energy oversupply. This is supported by the observations that short term (6 days) isoenergetic high fat diets do not alter fasting levels of either total or bioactive FGF21 (111). Longer-term studies of the impact of high-fat diets have not been carried out, but obesity is generally associated with high serum FGF21 in humans, suggesting that chronic adiposity may lead to

an FGF21 resistant state (48,113). This notion is particularly intriguing given that a common SNP in KLB gene (rs2608819), the key co-receptor mediating the tissue sensitivity to FGF21, is associated with higher body mass index, and the protein levels of KLB are reduced in the adipose tissue of obese humans which may decrease its responsiveness to FGF21 *in vivo* (113,119). However, the hepatic expression of KLB and FGFR1 and FGFR3 are increased in obese individuals which, in contrast to adipose tissue, may confer an improved hepatic responsiveness to increased FGF21 levels under those conditions. Large-scale RNA-sequencing studies reveal that mouse KLB gene is predominantly expressed in the liver, pancreas and adipose tissue with no KLB expression detected in skeletal muscle (120). In humans, beyond the liver and adipose tissue, KLB is detected in breast, bone marrow and pancreas (120). It is conceivable that tissue differences at the KLB level between mice and humans may contribute to the divergent findings of the metabolic actions of FGF21 described for the two species.

In addition to its role as an insulin-dependent postprandial hormone, studies in mice have shown that FGF21 suppresses consumption of simple sugars and non-caloric sweeteners but not complex carbohydrates, proteins or lipids via signaling through the paraventricular nucleus in the hypothalamus (50). Furthermore, FGF21 administration markedly reduces sweet and alcohol preference in mice and sweet preference in cynomolgus monkeys. In mice, these effects require the FGF21 co-receptor KLB in the CNS and correlate with reductions in dopamine concentrations in the nucleus accumbens (85). These actions of FGF21 may represent a conserved mechanism among species as genetic studies in humans have shown that SNPs in and around the FGF21 gene are associated with macronutrient preference, including carbohydrate, fat and protein intake, and alcohol consumption (44,121,122). The fact that consumption of dietary carbohydrates (such as fructose and sucrose) may increase the fasting levels of FGF21, which in turn may suppress consumption of simple sugars and non-caloric sweeteners, likely indicates the presence of a negative feedback hormonal loop between the liver and the brain (Figure 2). Interestingly, recent studies have also linked variants in the KLB locus with levels of alcohol intake in humans (123). FGF21 has also been shown to stimulate water drinking behaviour in mice in response to alcohol; this response is dependent upon β -adrenergic signaling (86). Furthermore, injection of recombinant FGF21 reduces *ad libitum* intake of alcohol in mice (122), whereas alcohol ingestion acutely and in

the form of binge drinking increases plasma FGF21 in fasting humans (122,124). Therefore, it is likely that alcohol-induced hepatic secretion of FGF21 serves as an adaptive mechanism aiming to inhibit alcohol appetite and stimulate water consumption in humans and therefore protect against long-term alcohol induced hepatic injury (124).

FGF21 has also been suggested to act as an exercise-induced myokine as its serum levels are elevated in response to whole body submaximal exercise (23,125,126). However, recent elegant studies using tissue arteriovenous differences have shown that exercise is a key stimulus in inducing hepatic release of FGF21 in humans (Figure 2). Indeed, during submaximal exercise most of the release of FGF21 into the systemic circulation occurs through the liver rather than skeletal muscle (107). This was confirmed by a recent study showing that maximal eccentric contractions do not stimulate the release of either total or bioactive FGF21 from human skeletal muscle, whereas its FGF21 protein content is unchanged in response to exercise (108). Interestingly, eccentric exercise resulted in significant elevations in arterialised and venous concentrations of its regulatory enzyme FAP, with no apparent contribution to its release from the exercised leg (Figure 2). These findings raise the possibility that increased levels of FAP may play a role in the inactivation of FGF21 during this type of exercise. Collectively, the results from the literature suggest that FGF21 may be an exercise-induced hepatokine but not myokine and that strategies aiming to minimise the release of FAP may be of benefit in maximising the metabolic effects of FGF21, whether administered exogenously or produced in response to exercise. FGF21 was first described as a key starvation hormone in a variety of animal models. However, recent studies in adult humans provide strong evidence that FGF21 is an insulin-dependent postprandial hormone that plays a key role in the consumption of alcohol, water, simple sugars and non-caloric sweeteners. Further understanding of the role of FGF21 in macronutrient preference and regulation of appetite in humans could be instrumental to uncovering its therapeutic potential in obesity, alcoholism, eating disorders, addictive behaviours and other relevant disorders.

Concluding Remarks and Future Perspectives

Nearly two decades of research on FGF21 have identified it as a potent metabolic regulator when manipulated acutely by pharmacological means or by genetic approaches (in mice), and

have characterised in detail its target tissues, receptor complexes and degradation pathways. However, clinical trials with FGF21 analogues in overweight/obese patients with T2D have not progressed to Phase 3 because the improvements in circulating lipid profiles have not been matched by improvements in glycaemic control and body weight loss. Perhaps the therapeutic potential of FGF21 analogues in T2D may be realised when combined with other current therapies targeting glycaemic control. However, the emergence of FGF21 as a regulator of macronutrient preference both in rodents and humans and its role in the protection from fatty liver disease and dyslipidaemia raises the interesting prospect of its therapeutic potential in obesity and some of its comorbidities. This is supported by the efficacy of FGF21 as a metabolic regulator in seasonal rodents that undergo natural cycles of adiposity and weight loss without the presence of metabolic dysfunction. The way forward with FGF21 may be to investigate its efficacy in obesity before T2D develops, targeting populations with specific polymorphisms that may confer an improved (tissue-specific) responsiveness to FGF21 treatment. This strategy may reveal how responsiveness to FGF21 can be improved and so insulin sensitivity maintained in the long term, reshaping our view of FGF21 as prevention rather than cure.

Acknowledgements

Work in the authors' laboratories is funded by the BBSRC (UK) grant BB/M001555/1.

Conflict of interest

RJS is an employee of Eli Lilly and Company.

JEL, FJPE and KT have had prior collaborative studies with Eli Lilly and Company.

References

1. Potthoff MJ, Inagaki T, Satapati S, Ding X, He T, Goetz R, Mohammadi M, Finck BN, Mangelsdorf DJ, Kliewer SA, Burgess SC. FGF21 induces PGC-1 α and regulates carbohydrate and fatty acid metabolism during the adaptive starvation response. *Proc Natl Acad Sci U S A*. 2009;106(26):10853-10858.
2. Kharitonov A, Shiyanova TL, Koester A, Ford AM, Micanovic R, Galbreath EJ, Sandusky GE, Hammond LJ, Moyers JS, Owens RA, Gromada J, Brozinick JT, Hawkins ED, Wroblewski VJ, Li DS, Mehrbod F, Jaskunas SR, Shanafelt AB. FGF-21 as a novel metabolic regulator. *The Journal of clinical investigation*. 2005;115(6):1627-1635.
3. Gaich G, Chien JY, Fu H, Glass LC, Deeg MA, Holland WL, Kharitonov A, Bumol T, Schilske HK, Moller DE. The effects of LY2405319, an FGF21 analog, in obese human subjects with type 2 diabetes. *Cell metabolism*. 2013;18(3):333-340.
4. Talukdar S, Zhou Y, Li D, Rossulek M, Dong J, Somayaji V, Weng Y, Clark R, Lanba A, Owen BM, Brenner MB, Trimmer JK, Gropp KE, Chabot JR, Erion DM, Rolph TP, Goodwin B, Calle RA. A Long-Acting FGF21 Molecule, PF-05231023, Decreases Body Weight and Improves Lipid Profile in Non-human Primates and Type 2 Diabetic Subjects. *Cell metabolism*. 2016;23(3):427-440.
5. Dong JQ, Rossulek M, Somayaji VR, Baltrukonis D, Liang Y, Hudson K, Hernandez-Illas M, Calle RA. Pharmacokinetics and pharmacodynamics of PF-05231023, a novel long-acting FGF21 mimetic, in a first-in-human study. *British journal of clinical pharmacology*. 2015;80(5):1051-1063.
6. Charles ED, Neuschwander-Tetri BA, Pablo Frias J, Kundu S, Luo Y, Tiruchera GS, Christian R. Pegbelfermin (BMS-986036), PEGylated FGF21, in Patients with Obesity and Type 2 Diabetes: Results from a Randomized Phase 2 Study. *Obesity (Silver Spring)*. 2019;27(1):41-49.
7. 8. Obesity Management for the Treatment of Type 2 Diabetes: Standards of Medical Care in Diabetes-2019. *Diabetes care*. 2019;42(Suppl 1):S81-s89.
8. Adams AC, Kharitonov A. FGF21: The center of a transcriptional nexus in metabolic regulation. *Current diabetes reviews*. 2012;8(4):285-293.
9. Nishimura T, Nakatake Y, Konishi M, Itoh N. Identification of a novel FGF, FGF-21, preferentially expressed in the liver. *Biochimica et biophysica acta*. 2000;1492(1):203-206.
10. Fon Tacer K, Bookout AL, Ding X, Kurosu H, John GB, Wang L, Goetz R, Mohammadi M, Kuro-o M, Mangelsdorf DJ, Kliewer SA. Research resource: Comprehensive expression atlas of the fibroblast growth factor system in adult mouse. *Molecular endocrinology (Baltimore, Md)*. 2010;24(10):2050-2064.
11. Hsueh H, Pan W, Kastin AJ. The fasting polypeptide FGF21 can enter brain from blood. *Peptides*. 2007;28(12):2382-2386.
12. Tan BK, Hallschmid M, Adya R, Kern W, Lehnert H, Rande HS. Fibroblast growth factor 21 (FGF21) in human cerebrospinal fluid: relationship with plasma FGF21 and body adiposity. *Diabetes*. 2011;60(11):2758-2762.
13. Yu H, Xia F, Lam KS, Wang Y, Bao Y, Zhang J, Gu Y, Zhou P, Lu J, Jia W, Xu A. Circadian rhythm of circulating fibroblast growth factor 21 is related to diurnal changes in fatty acids in humans. *Clinical chemistry*. 2011;57(5):691-700.
14. Liang Q, Zhong L, Zhang J, Wang Y, Bornstein SR, Triggle CR, Ding H, Lam KS, Xu A. FGF21 maintains glucose homeostasis by mediating the cross talk between liver and brain during prolonged fasting. *Diabetes*. 2014;63(12):4064-4075.

15. Badman MK, Koester A, Flier JS, Kharitonov A, Maratos-Flier E. Fibroblast growth factor 21-deficient mice demonstrate impaired adaptation to ketosis. *Endocrinology*. 2009;150(11):4931-4940.
16. Hotta Y, Nakamura H, Konishi M, Murata Y, Takagi H, Matsumura S, Inoue K, Fushiki T, Itoh N. Fibroblast growth factor 21 regulates lipolysis in white adipose tissue but is not required for ketogenesis and triglyceride clearance in liver. *Endocrinology*. 2009;150(10):4625-4633.
17. Badman MK, Pissios P, Kennedy AR, Koukos G, Flier JS, Maratos-Flier E. Hepatic fibroblast growth factor 21 is regulated by PPARalpha and is a key mediator of hepatic lipid metabolism in ketotic states. *Cell metabolism*. 2007;5(6):426-437.
18. Inagaki T, Dutchak P, Zhao G, Ding X, Gautron L, Parameswara V, Li Y, Goetz R, Mohammadi M, Esser V, Elmquist JK, Gerard RD, Burgess SC, Hammer RE, Mangelsdorf DJ, Kliewer SA. Endocrine regulation of the fasting response by PPARalpha-mediated induction of fibroblast growth factor 21. *Cell metabolism*. 2007;5(6):415-425.
19. Bookout AL, de Groot MH, Owen BM, Lee S, Gautron L, Lawrence HL, Ding X, Elmquist JK, Takahashi JS, Mangelsdorf DJ, Kliewer SA. FGF21 regulates metabolism and circadian behavior by acting on the nervous system. *Nature medicine*. 2013;19(9):1147-1152.
20. Owen BM, Bookout AL, Ding X, Lin VY, Atkin SD, Gautron L, Kliewer SA, Mangelsdorf DJ. FGF21 contributes to neuroendocrine control of female reproduction. *Nature medicine*. 2013;19(9):1153-1156.
21. Uebanso T, Taketani Y, Yamamoto H, Amo K, Ominami H, Arai H, Takei Y, Masuda M, Tanimura A, Harada N, Yamanaka-Okumura H, Takeda E. Paradoxical regulation of human FGF21 by both fasting and feeding signals: is FGF21 a nutritional adaptation factor? *PloS one*. 2011;6(8):e22976.
22. Fazeli PK, Lun M, Kim SM, Bredella MA, Wright S, Zhang Y, Lee H, Catana C, Klibanski A, Patwari P, Steinhilber ML. FGF21 and the late adaptive response to starvation in humans. *The Journal of clinical investigation*. 2015;125(12):4601-4611.
23. Kim KH, Jeong YT, Oh H, Kim SH, Cho JM, Kim YN, Kim SS, Kim DH, Hur KY, Kim HK, Ko T, Han J, Kim HL, Kim J, Back SH, Komatsu M, Chen H, Chan DC, Konishi M, Itoh N, Choi CS, Lee MS. Autophagy deficiency leads to protection from obesity and insulin resistance by inducing Fgf21 as a mitokine. *Nature medicine*. 2013;19(1):83-92.
24. Beenken A, Mohammadi M. The FGF family: biology, pathophysiology and therapy. *Nature reviews Drug discovery*. 2009;8(3):235-253.
25. Lee S, Choi J, Mohanty J, Sousa LP, Tome F, Pardon E, Steyaert J, Lemmon MA, Lax I, Schlessinger J. Structures of beta-klotho reveal a 'zip code'-like mechanism for endocrine FGF signalling. *Glia*. 2018;553(7689):501-505.
26. Kurosu H, Choi M, Ogawa Y, Dickson AS, Goetz R, Eliseenkova AV, Mohammadi M, Rosenblatt KP, Kliewer SA, Kuro-o M. Tissue-specific expression of betaKlotho and fibroblast growth factor (FGF) receptor isoforms determines metabolic activity of FGF19 and FGF21. *The Journal of biological chemistry*. 2007;282(37):26687-26695.
27. Ge X, Chen C, Hui X, Wang Y, Lam KS, Xu A. Fibroblast growth factor 21 induces glucose transporter-1 expression through activation of the serum response factor/Ets-like protein-1 in adipocytes. *The Journal of biological chemistry*. 2011;286(40):34533-34541.

28. Diaz-Delfin J, Hondares E, Iglesias R, Giralt M, Caelles C, Villarroya F. TNF-alpha represses beta-Klotho expression and impairs FGF21 action in adipose cells: involvement of JNK1 in the FGF21 pathway. *Endocrinology*. 2012;153(9):4238-4245.
29. Samms RJ. Overexpression of beta-Klotho in Adipose Tissue Sensitizes Male Mice to Endogenous FGF21 and Provides Protection From Diet-Induced Obesity. 2016;157(4):1467-1480.
30. Jager J, Wang F, Fang B, Lim HW, Peed LC, Steger DJ, Won KJ, Kharitononkov A, Adams AC, Lazar MA. The Nuclear Receptor Rev-erbalpha Regulates Adipose Tissue-specific FGF21 Signaling. *The Journal of biological chemistry*. 2016;291(20):10867-10875.
31. Suzuki M, Uehara Y, Motomura-Matsuzaka K, Oki J, Koyama Y, Kimura M, Asada M, Komi-Kuramochi A, Oka S, Imamura T. betaKlotho is required for fibroblast growth factor (FGF) 21 signaling through FGF receptor (FGFR) 1c and FGFR3c. *Molecular endocrinology (Baltimore, Md)*. 2008;22(4):1006-1014.
32. Kharitononkov A. FGF21 attenuates lipolysis in human adipocytes - a possible link to improved insulin sensitivity. *Endocrinology*. 2008;582(12):1725-1730.
33. Markan KR, Naber MC, Ameka MK, Anderegg MD, Mangelsdorf DJ, Kliewer SA, Mohammadi M, Potthoff MJ. Circulating FGF21 is liver derived and enhances glucose uptake during refeeding and overfeeding. *Diabetes*. 2014;63(12):4057-4063.
34. Eagleson KL, Fairfull LD, Salton SR, Levitt P. Regional differences in neurotrophin availability regulate selective expression of VGF in the developing limbic cortex. *The Journal of neuroscience : the official journal of the Society for Neuroscience*. 2001;21(23):9315-9324.
35. Hashimoto T, Cook WS, Qi C, Yeldandi AV, Reddy JK, Rao MS. Defect in peroxisome proliferator-activated receptor alpha-inducible fatty acid oxidation determines the severity of hepatic steatosis in response to fasting. *The Journal of biological chemistry*. 2000;275(37):28918-28928.
36. Kersten S, Seydoux J, Peters JM, Gonzalez FJ, Desvergne B, Wahli W. Peroxisome proliferator-activated receptor alpha mediates the adaptive response to fasting. *The Journal of clinical investigation*. 1999;103(11):1489-1498.
37. Leone TC, Weinheimer CJ, Kelly DP. A critical role for the peroxisome proliferator-activated receptor alpha (PPARalpha) in the cellular fasting response: the PPARalpha-null mouse as a model of fatty acid oxidation disorders. *Proc Natl Acad Sci U S A*. 1999;96(13):7473-7478.
38. Lundasen T, Hunt MC, Nilsson LM, Sanyal S, Angelin B, Alexson SE, Rudling M. PPARalpha is a key regulator of hepatic FGF21. *Biochemical and biophysical research communications*. 2007;360(2):437-440.
39. Reitman ML. FGF21: a missing link in the biology of fasting. *Cell metabolism*. 2007;5(6):405-407.
40. De Sousa-Coelho AL, Relat J, Hondares E, Perez-Marti A, Ribas F, Villarroya F, Marrero PF, Haro D. FGF21 mediates the lipid metabolism response to amino acid starvation. *Journal of lipid research*. 2013;54(7):1786-1797.
41. Pissios P, Hong S, Kennedy AR, Prasad D, Liu FF, Maratos-Flier E. Methionine and choline regulate the metabolic phenotype of a ketogenic diet. *Molecular metabolism*. 2013;2(3):306-313.
42. Shimizu M, Morimoto H, Maruyama R, Inoue J, Sato R. Selective Regulation of FGF19 and FGF21 Expression by Cellular and Nutritional Stress. *Journal of nutritional science and vitaminology*. 2015;61(2):154-160.

43. Fisher FM, Chui PC, Nasser IA, Popov Y, Cunniff JC, Lundasen T, Kharitononkov A, Schuppan D, Flier JS, Maratos-Flier E. Fibroblast growth factor 21 limits lipotoxicity by promoting hepatic fatty acid activation in mice on methionine and choline-deficient diets. *Gastroenterology*. 2014;147(5):1073-1083.e1076.
44. Tanaka N, Takahashi S, Zhang Y, Krausz KW, Smith PB, Patterson AD, Gonzalez FJ. Role of fibroblast growth factor 21 in the early stage of NASH induced by methionine- and choline-deficient diet. *Biochimica et biophysica acta*. 2015;1852(7):1242-1252.
45. Sanchez J, Palou A, Pico C. Response to carbohydrate and fat refeeding in the expression of genes involved in nutrient partitioning and metabolism: striking effects on fibroblast growth factor-21 induction. *Endocrinology*. 2009;150(12):5341-5350.
46. Fisher FM, Kim M, Doridot L, Cunniff JC, Parker TS, Levine DM, Hellerstein MK, Hudgins LC, Maratos-Flier E, Herman MA. A critical role for ChREBP-mediated FGF21 secretion in hepatic fructose metabolism. *Molecular metabolism*. 2017;6(1):14-21.
47. Lundsgaard AM, Fritzen AM, Sjoberg KA, Myrmel LS, Madsen L, Wojtaszewski JF, Richter EA, Kiens B. Circulating FGF21 in humans is potently induced by short term overfeeding of carbohydrates. *Molecular metabolism*. 2017;6(1):22-29.
48. Zhang X, Yeung DC, Karpisek M, Stejskal D, Zhou ZG, Liu F, Wong RL, Chow WS, Tso AW, Lam KS, Xu A. Serum FGF21 levels are increased in obesity and are independently associated with the metabolic syndrome in humans. *Diabetes*. 2008;57(5):1246-1253.
49. Nygaard EB, Moller CL, Kievit P, Grove KL, Andersen B. Increased fibroblast growth factor 21 expression in high-fat diet-sensitive non-human primates (*Macaca mulatta*). *Int J Obes (Lond)*. 2014;38(2):183-191.
50. von Holstein-Rathlou S, BonDurant LD, Peltekian L, Naber MC, Yin TC, Claflin KE, Urizar AI, Madsen AN, Ratner C, Holst B, Karstoft K, Vandenbeuch A, Anderson CB, Cassell MD, Thompson AP, Solomon TP, Rahmouni K, Kinnamon SC, Pieper AA, Gillum MP, Potthoff MJ. FGF21 Mediates Endocrine Control of Simple Sugar Intake and Sweet Taste Preference by the Liver. *Cell metabolism*. 2016;23(2):335-343.
51. Berglund ED, Li CY, Bina HA, Lynes SE, Michael MD, Shanafelt AB, Kharitononkov A, Wasserman DH. Fibroblast growth factor 21 controls glycemia via regulation of hepatic glucose flux and insulin sensitivity. *Endocrinology*. 2009;150(9):4084-4093.
52. Fisher FM, Estall JL, Adams AC, Antonellis PJ, Bina HA, Flier JS, Kharitononkov A, Spiegelman BM, Maratos-Flier E. Integrated regulation of hepatic metabolism by fibroblast growth factor 21 (FGF21) in vivo. *Endocrinology*. 2011;152(8):2996-3004.
53. Lin Z, Tian H, Lam KS, Lin S, Hoo RC, Konishi M, Itoh N, Wang Y, Bornstein SR, Xu A, Li X. Adiponectin mediates the metabolic effects of FGF21 on glucose homeostasis and insulin sensitivity in mice. *Cell metabolism*. 2013;17(5):779-789.
54. Samms RJ, Fowler MJ, Cooper S, Murphy M, Emmerson P, Coskun T, Adams AC, Kharitononkov A, Ebling FJP, K.O. T. Dual effects of fibroblast growth factor 21 on hepatic energy metabolism. *Journal of Endocrinology*. 2015;227:37-47.
55. Khan R, Bril F, Cusi K, Newsome PN. Modulation of Insulin Resistance in NAFLD. *Hepatology (Baltimore, Md)*. 2018.
56. Muller CP, Levy D, Elliott P, Lee JH, Kang YE, Chang JY, Park KC, Kim HW, Kim JT, Kim HJ, Yi HS, Shong M, Chung HK, Kim KS. An engineered FGF21 variant, LY2405319, can prevent non-alcoholic steatohepatitis by enhancing hepatic mitochondrial function. *Proc Natl Acad Sci U S A*. 2016;8(11):4750-4763.
57. Sanyal A, Charles ED, Neuschwander-Tetri BA, Loomba R, Harrison SA, Abdelmalek MF, Lawitz EJ, Halegoua-DeMarzio D, Kundu S, Noviello S, Luo Y, Christian R. Pegbelfermin

- (BMS-986036), a PEGylated fibroblast growth factor 21 analogue, in patients with non-alcoholic steatohepatitis: a randomised, double-blind, placebo-controlled, phase 2a trial. *Lancet*. 2019;392(10165):2705-2717.
58. Fujii N, Uta S, Kobayashi M, Sato T, Okita N, Higami Y. Impact of aging and caloric restriction on fibroblast growth factor 21 signaling in rat white adipose tissue. *Experimental gerontology*. 2019;118:55-64.
 59. Jeffery E, Wing A, Holtrup B, Sebo Z, Kaplan JL, Saavedra-Pena R, Church CD, Colman L, Berry R, Rodeheffer MS. The Adipose Tissue Microenvironment Regulates Depot-Specific Adipogenesis in Obesity. *Cell metabolism*. 2016;24(1):142-150.
 60. Frayn KN. Adipose tissue as a buffer for daily lipid flux. *Diabetologia*. 2002;45(9):1201-1210.
 61. Shulman GI. Ectopic fat in insulin resistance, dyslipidemia, and cardiometabolic disease. *The New England journal of medicine*. 2014;371(12):1131-1141.
 62. Xu J, Lloyd DJ, Hale C, Stanislaus S, Chen M, Sivits G, Vonderfecht S, Hecht R, Li YS, Lindberg RA, Chen JL, Jung DY, Zhang Z, Ko HJ, Kim JK, Veniant MM. Fibroblast growth factor 21 reverses hepatic steatosis, increases energy expenditure, and improves insulin sensitivity in diet-induced obese mice. *Diabetes*. 2009;58(1):250-259.
 63. Kong LJ, Feng W, Wright M, Chen Y, Dallas-yang Q, Zhou YP, Berger JP. FGF21 suppresses hepatic glucose production through the activation of atypical protein kinase Ciota/lambda. *European journal of pharmacology*. 2013;702(1-3):302-308.
 64. Camporez JP, Jornayvaz FR, Petersen MC, Pesta D, Guigni BA, Serr J, Zhang D, Kahn M, Samuel VT, Jurczak MJ, Shulman GI. Cellular mechanisms by which FGF21 improves insulin sensitivity in male mice. *Endocrinology*. 2013;154(9):3099-3109.
 65. Coskun T, Bina HA, Schneider MA, Dunbar JD, Hu CC, Chen Y, Moller DE, Kharitononkov A. Fibroblast growth factor 21 corrects obesity in mice. *Endocrinology*. 2008;149(12):6018-6027.
 66. Xu J, Stanislaus S, Chinookoswong N, Lau YY, Hager T, Patel J, Ge H, Weiszmann J, Lu SC, Graham M, Busby J, Hecht R, Li YS, Li Y, Lindberg R, Veniant MM. Acute glucose-lowering and insulin-sensitizing action of FGF21 in insulin-resistant mouse models--association with liver and adipose tissue effects. *American journal of physiology Endocrinology and metabolism*. 2009;297(5):E1105-1114.
 67. Samms RJ, Smith DP, Cheng CC, Antonellis PP, Perfield JW, 2nd, Kharitononkov A, Gimeno RE, Adams AC. Discrete Aspects of FGF21 In Vivo Pharmacology Do Not Require UCP1. *Cell reports*. 2015;11(7):991-999.
 68. Veniant MM, Sivits G, Helmering J, Komorowski R, Lee J, Fan W, Moyer C, Lloyd DJ. Pharmacologic Effects of FGF21 Are Independent of the "Browning" of White Adipose Tissue. *Cell metabolism*. 2015;21(5):731-738.
 69. Kwon MM, O'Dwyer SM, Baker RK, Covey SD, Kieffer TJ. FGF21-Mediated Improvements in Glucose Clearance Require Uncoupling Protein 1. *Cell reports*. 2015;13(8):1521-1527.
 70. BonDurant LD, Ameka M, Naber MC, Markan KR, Idiga SO, Acevedo MR, Walsh SA, Ornitz DM, Potthoff MJ. FGF21 Regulates Metabolism Through Adipose-Dependent and -Independent Mechanisms. *Cell metabolism*. 2017;25(4):935-944.e934.
 71. Lee P, Greenfield JR, Ho KK, Fulham MJ. A critical appraisal of the prevalence and metabolic significance of brown adipose tissue in adult humans. *American journal of physiology Endocrinology and metabolism*. 2010;299(4):E601-606.

72. Li X, Ge H, Weizmann J, Hecht R, Li YS, Veniant MM, Xu J, Wu X, Lindberg R, Li Y. Inhibition of lipolysis may contribute to the acute regulation of plasma FFA and glucose by FGF21 in ob/ob mice. *FEBS letters*. 2009;583(19):3230-3234.
73. Arner P, Pettersson A, Mitchell PJ, Dunbar JD, Kharitonov A, Ryden M. FGF21 attenuates lipolysis in human adipocytes - a possible link to improved insulin sensitivity. *FEBS letters*. 2008;582(12):1725-1730.
74. Adams AC, Cheng CC, Coskun T, Kharitonov A. FGF21 requires betaklotho to act in vivo. *PLoS one*. 2012;7(11):e49977.
75. Foltz IN, Hu S, King C, Wu X, Yang C, Wang W, Weizmann J, Stevens J, Chen JS, Nuanmanee N, Gupte J, Komorowski R, Sekirov L, Hager T, Arora T, Ge H, Baribault H, Wang F, Sheng J, Karow M, Wang M, Luo Y, McKeenan W, Wang Z, Veniant MM, Li Y. Treating diabetes and obesity with an FGF21-mimetic antibody activating the betaKlotho/FGFR1c receptor complex. *Science translational medicine*. 2012;4(162):162ra153.
76. Ding X, Boney-Montoya J, Owen BM, Bookout AL, Coate KC, Mangelsdorf DJ, Kliewer SA. betaKlotho is required for fibroblast growth factor 21 effects on growth and metabolism. *Cell metabolism*. 2012;16(3):387-393.
77. So WY, Cheng Q, Chen L, Evans-Molina C, Xu A, Lam KS, Leung PS. High glucose represses beta-klotho expression and impairs fibroblast growth factor 21 action in mouse pancreatic islets: involvement of peroxisome proliferator-activated receptor gamma signaling. *Diabetes*. 2013;62(11):3751-3759.
78. Samson SL, Sathyanarayana P, Jogi M, Gonzalez EV, Gutierrez A, Krishnamurthy R, Muthupillai R, Chan L, Bajaj M. Exenatide decreases hepatic fibroblast growth factor 21 resistance in non-alcoholic fatty liver disease in a mouse model of obesity and in a randomised controlled trial. *Diabetologia*. 2011;54(12):3093-3100.
79. Adams AC, Halstead CA, Hansen BC, Irizarry AR, Martin JA, Myers SR, Reynolds VL, Smith HW, Wroblewski VJ, Kharitonov A. LY2405319, an Engineered FGF21 Variant, Improves the Metabolic Status of Diabetic Monkeys. *PLoS one*. 2013;8(6):e65763.
80. Markan KR, Naber MC, Small SM, Peltekian L, Kessler RL, Potthoff MJ. FGF21 resistance is not mediated by downregulation of beta-klotho expression in white adipose tissue. *Scientific reports*. 2017;6(6):602-610.
81. Heisler LK, Lam DD. An appetite for life: brain regulation of hunger and satiety. *Current opinion in pharmacology*. 2017;37:100-106.
82. Liang Q, Zhong L, Zhang J, Wang Y, Bornstein SR, Triggle CR, Ding H, Lam KS, Xu A. FGF21 maintains glucose homeostasis by mediating the cross talk between liver and brain during prolonged fasting. *Diabetes*. 2014;63:4064-4075.
83. Owen BM, Ding X, Morgan DA, Coate KC, Bookout AL, Rahmouni K, Kliewer SA, Mangelsdorf DJ. FGF21 acts centrally to induce sympathetic nerve activity, energy expenditure, and weight loss. *Cell metabolism*. 2014;20(4):670-677.
84. Douris N, Stevanovic DM, Fisher FM, Cisu TI, Chee MJ, Nguyen NL, Zarebidaki E, Adams AC, Kharitonov A, Flier JS, Bartness TJ, Maratos-Flier E. Central Fibroblast Growth Factor 21 Browns White Fat via Sympathetic Action in Male Mice. *Endocrinology*. 2015;156(7):2470-2481.
85. Talukdar S, Owen BM, Song P, Hernandez G, Zhang Y, Zhou Y, Scott WT, Paratala B, Turner T, Smith A, Bernardo B, Muller CP, Tang H, Mangelsdorf DJ, Goodwin B, Kliewer

- SA. FGF21 Regulates Sweet and Alcohol Preference. *Cell metabolism*. 2016;23(2):344-349.
86. Song P, Zechner C, Hernandez G, Canovas J, Xie Y, Sondhi V, Wagner M, Stadlbauer V, Horvath A, Leber B, Hu MC, Moe OW, Mangelsdorf DJ, Kliewer SA. The Hormone FGF21 Stimulates Water Drinking in Response to Ketogenic Diet and Alcohol. *Cell metabolism*. 2018;27(6):1338-1347.e1334.
 87. Martin B, Ji S, Maudsley S, Mattson MP. "Control" laboratory rodents are metabolically morbid: Why it matters. *Proceedings of the National Academy of Sciences, USA*. 2010;107:6127-6133.
 88. Ebling FJP. On the value of seasonal mammals for identifying mechanisms underlying the control of food intake and body weight. *Hormones and Behavior*. 2014;66:55-61.
 89. Ebling FJ. Hypothalamic control of seasonal changes in food intake and body weight. *Frontiers in Neuroendocrinology*. 2015;37:97-107.
 90. Scherbarth F, Steinlechner S. Endocrine mechanisms of seasonal adaptation in small mammals: from early results to present understanding. *Journal of Comparative Physiology B*. 2010;180:935-952.
 91. Petri I, Diedrich V, Wilson D, Fernández-Calleja J, Herwig A, Steinlechner S, Barrett P. Orchestration of gene expression across the seasons: Hypothalamic gene expression in natural photoperiod throughout the year in the Siberian hamster. *Scientific reports*. 2016;6:29689. doi: 29610.21038/srep29689.
 92. Samms R, Fowler MJ, Cooper S, Emmerson PJ, Coskun T, Adams AC, Kharitononkov A, Tsintzas K, Ebling FJP. Photoperiodic regulation of FGF21 production in the Siberian hamster. *Hormones and Behavior*. 2014;66:180-185.
 93. Nelson BT, Ding X, Boney-Montoya J, Gerard RD, Kliewer SA, Andrews MT. Metabolic hormone FGF21 is induced in ground squirrels during hibernation but its overexpression is not sufficient to cause torpor. *PLoS one*. 2013;8:e53574.
 94. Lewis JE, Samms RJ, Cooper S, Luckett JC, Perkins AC, Adams AC, Tsintzas K, Ebling FJP. Reduced adiposity attenuates FGF21 mediated metabolic improvements in the Siberian hamster. *Scientific reports*. 2017;7:Article number: 4238 doi:4210.1038/s41598-41017-03607-x.
 95. Murphy M, Samms R, Warner A, Bolborea M, Barrett P, Fowler MJ, Brameld JM, Kharitononkov A, Adams AC, Coskun T, Ebling FJP. Increased responses to the actions of fibroblast growth factor 21 on energy balance and body weight in a seasonal model of adiposity. *Journal of Neuroendocrinology*. 2013;25:180-189.
 96. Coskun T, Bina HA, Schneider MA, Dunbar JD, Hu CC, Chen Y, Moller DE, Kharitononkov A. Fibroblast growth factor 21 corrects obesity in mice. *Endocrinology*. 2008;149:6018-6027.
 97. Lewis JE, Brameld JM, Hill P, Cocco C, Noli B, Barrett P, Ebling FJP, Jethwa PH. Hypothalamic over-expression of VGF in the Siberian hamster increases energy expenditure and reduces body weight gain. *PLoS one*. 2017;12(2):e0172724. doi: 0172710.0171371/journal.pone.0172724.
 98. Lewis JE, Samms RJ, Cooper S, Luckett JC, Perkins AC, Adams AC, Tsintzas K, Ebling FJP. Reduced adiposity attenuates FGF21 mediated metabolic improvements in the Siberian hamster. 2017;7(1):4238.
 99. Lewis JE, Ebling FJP. Tanycytes As Regulators of Seasonal Cycles in Neuroendocrine Function. *Frontiers in Neurology*. 2017;8(79).

100. Ebling FJ, Lewis J. Tanycytes and hypothalamic control of energy metabolism. *Glia*. 2018;66:1176-1184.
101. Samms R, Lewis JE, Lory A, Fowler MJ, Cooper S, Warner A, Emmerson P, Adams AC, Luckett JC, Perkins AC, Wilson D, Barrett P, Tsintzas K, Ebling FJ. Antibody-Mediated Inhibition of the FGFR1c Isoform Induces a Catabolic Lean State in Siberian Hamsters. *Current Biology*. 2015;25:2997–3003.
102. Gao Y, Tschöp MH, Luquet S. Hypothalamic Tanycytes: Gatekeepers to Metabolic Control. *Cell metabolism*. 2014;19:173-174.
103. Nelson BT, Ding X, Boney-Montoya J, Gerard RD, Kliewer SA, Andrews MT. Metabolic hormone FGF21 is induced in ground squirrels during hibernation but its overexpression is not sufficient to cause torpor. *PloS one*. 2013;8(1):e53574.
104. Ebling FJP, Barrett P. The regulation of seasonal changes in food intake and body weight. *Journal of Neuroendocrinology*. 2008;20:827-833.
105. Chavez AO, Molina-Carrion M, Abdul-Ghani MA, Folli F, Defronzo RA, Tripathy D. Circulating fibroblast growth factor-21 is elevated in impaired glucose tolerance and type 2 diabetes and correlates with muscle and hepatic insulin resistance. *Diabetes care*. 2009;32(8):1542-1546.
106. Dushay J, Chui PC, Gopalakrishnan GS, Varela-Rey M, Crawley M, Fisher FM, Badman MK, Martinez-Chantar ML, Maratos-Flier E. Increased fibroblast growth factor 21 in obesity and nonalcoholic fatty liver disease. *Gastroenterology*. 2010;139(2):456-463.
107. Hansen JS, Clemmesen JO, Secher NH, Hoene M, Drescher A, Weigert C, Pedersen BK, Plomgaard P. Glucagon-to-insulin ratio is pivotal for splanchnic regulation of FGF-21 in humans. *Molecular metabolism*. 2015;4(8):551-560.
108. Parmar B, Lewis JE, Samms RJ, Ebling FJP, Cheng CC, Adams AC, Mallinson J, Cooper S, Taylor T, Ghasemi R, Stephens FB, Tsintzas K. Eccentric exercise increases circulating fibroblast activation protein alpha but not bioactive fibroblast growth factor 21 in healthy humans. 2018;103(6):876-883.
109. Christodoulides C, Dyson P, Sprecher D, Tsintzas K, Karpe F. Circulating fibroblast growth factor 21 is induced by peroxisome proliferator-activated receptor agonists but not ketosis in man. *The Journal of clinical endocrinology and metabolism*. 2009;94(9):3594-3601.
110. Galman C, Lundasen T, Kharitonov A, Bina HA, Eriksson M, Hafstrom I, Dahlin M, Amark P, Angelin B, Rudling M. The circulating metabolic regulator FGF21 is induced by prolonged fasting and PPARalpha activation in man. *Cell metabolism*. 2008;8(2):169-174.
111. Samms RJ, Lewis JE, Norton L, Stephens FB, Gaffney CJ, Butterfield T, Smith DP, Cheng CC, Perfield JW, 2nd, Adams AC, Ebling FJP, Tsintzas K. FGF21 Is an Insulin-Dependent Postprandial Hormone in Adult Humans. *The Journal of clinical endocrinology and metabolism*. 2017;102(10):3806-3813.
112. Dushay JR, Toschi E, Mitten EK, Fisher FM, Herman MA, Maratos-Flier E. Fructose ingestion acutely stimulates circulating FGF21 levels in humans. *Molecular metabolism*. 2015;4(1):51-57.
113. Gallego-Escuredo JM, Gomez-Ambrosi J, Catalan V, Domingo P, Giralt M, Fruhbeck G, Villarroya F. Opposite alterations in FGF21 and FGF19 levels and disturbed expression of the receptor machinery for endocrine FGFs in obese patients. *Int J Obes (Lond)*. 2015;39(1):121-129.

- 114.** Zhen EY, Jin Z, Ackermann BL, Thomas MK, Gutierrez JA. Circulating FGF21 proteolytic processing mediated by fibroblast activation protein. *The Biochemical journal*. 2016;473(5):605-614.
- 115.** Mai K, Bobbert T, Groth C, Assmann A, Meinus S, Kraatz J, Andres J, Arafat AM, Pfeiffer AF, Mohlig M, Spranger J. Physiological modulation of circulating FGF21: relevance of free fatty acids and insulin. *American journal of physiology Endocrinology and metabolism*. 2010;299(1):E126-130.
- 116.** Schmidlin T, Boender AJ, Frese CK, Heck AJ, Adan RA, Altelaar AF. Diet-induced neuropeptide expression: feasibility of quantifying extended and highly charged endogenous peptide sequences by selected reaction monitoring. *Analytical chemistry*. 2015;87(19):9966-9973.
- 117.** Matikainen N, Taskinen MR, Stenabb S, Lundbom N, Hakkarainen A, Vaaralahti K, Raivio T. Decrease in circulating fibroblast growth factor 21 after an oral fat load is related to postprandial triglyceride-rich lipoproteins and liver fat. *European journal of endocrinology*. 2012;166(3):487-492.
- 118.** Vienberg SG, Brons C, Nilsson E, Astrup A, Vaag A, Andersen B. Impact of short-term high-fat feeding and insulin-stimulated FGF21 levels in subjects with low birth weight and controls. *European journal of endocrinology*. 2012;167(1):49-57.
- 119.** Kaess BM, Barnes TA, Stark K, Charchar FJ, Waterworth D, Song K, Wang WY, Vollenweider P, Waeber G, Mooser V, Zukowska-Szczechowska E, Samani NJ, Hengstenberg C, Tomaszewski M. FGF21 signalling pathway and metabolic traits - genetic association analysis. *European journal of human genetics : EJHG*. 2010;18(12):1344-1348.
- 120.** Petryszak R, Keays M. Expression Atlas update--an integrated database of gene and protein expression in humans, animals and plants. 2016;44(D1):D746-752.
- 121.** Chu AY, Workalemahu T, Paynter NP, Rose LM, Giulianini F, Tanaka T, Ngwa JS, Qi Q, Curhan GC, Rimm EB, Hunter DJ, Pasquale LR, Ridker PM, Hu FB, Chasman DI, Qi L. Novel locus including FGF21 is associated with dietary macronutrient intake. *Human molecular genetics*. 2013;22(9):1895-1902.
- 122.** Soberg S, Sandholt CH, Jespersen NZ, Toft U, Madsen AL, von Holstein-Rathlou S, Grevengoed TJ, Christensen KB, Bredie WLP, Potthoff MJ, Solomon TPJ, Scheele C, Linneberg A, Jorgensen T, Pedersen O, Hansen T, Gillum MP, Grarup N. FGF21 Is a Sugar-Induced Hormone Associated with Sweet Intake and Preference in Humans. *Cell metabolism*. 2017;25(5):1045-1053.e1046.
- 123.** Schumann G, Liu C, O'Reilly P, Gao H, Song P, Xu B, Ruggeri B, Amin N, Jia T, Preis S, Segura Lepe M, Akira S, Barbieri C, Baumeister S, Cauchi S, Clarke TK, Enroth S, Fischer K, Hallfors J, Harris SE, Hieber S, Hofer E, Hottenga JJ, Johansson A, Joshi PK, Kaartinen N, Laitinen J, Lemaitre R, Loukola A, Luan J, Lytikainen LP, Mangino M, Manichaikul A, Mbarek H, Milaneschi Y, Moayyeri A, Mukamal K, Nelson C, Nettleton J, Partinen E, Rawal R, Robino A, Rose L, Sala C, Satoh T, Schmidt R, Schraut K, Scott R, Smith AV, Starr JM, Teumer A, Trompet S, Uitterlinden AG, Venturini C, Verghnaud AC, Verweij N, Vitart V, Vuckovic D, Wedenoja J, Yengo L, Yu B, Zhang W, Zhao JH, Boomsma DI, Chambers J, Chasman DI, Daniela T, de Geus E, Deary I, Eriksson JG, Esko T, Eulenburg V, Franco OH, Froguel P, Gieger C, Grabe HJ, Gudnason V, Gyllensten U, Harris TB, Hartikainen AL, Heath AC, Hocking L, Hofman A, Huth C, Jarvelin MR, Jukema JW, Kaprio J, Kooner JS, Kutalik Z, Lahti J, Langenberg C, Lehtimaki T, Liu Y, Madden PA, Martin N, Morrison A, Penninx B, Pirastu N, Psaty B, Raitakari O, Ridker P, Rose R,

- Rotter JI, Samani NJ, Schmidt H, Spector TD, Stott D, Strachan D, Tzoulaki I, van der Harst P, van Duijn CM, Marques-Vidal P, Vollenweider P, Wareham NJ, Whitfield JB, Wilson J, Wolffenbuttel B, Bakalkin G, Evangelou E, Liu Y, Rice KM, Desrivieres S, Kliewer SA, Mangelsdorf DJ. KLB is associated with alcohol drinking, and its gene product beta-Klotho is necessary for FGF21 regulation of alcohol preference. *2016;113(50):14372-14377.*
- 124.** Desai BN, Singhal G, Watanabe M, Stevanovic D, Lundasen T, Fisher FM, Mather ML, Vardeh HG, Douris N, Adams AC, Nasser IA, FitzGerald GA, Flier JS, Skarke C, Maratos-Flier E. Fibroblast growth factor 21 (FGF21) is robustly induced by ethanol and has a protective role in ethanol associated liver injury. *Glia.* 2017;6(11):1395-1406.
- 125.** Cuevas-Ramos D, Almeda-Valdes P, Meza-Arana CE, Brito-Cordova G, Gomez-Perez FJ, Mehta R, Oseguera-Moguel J, Aguilar-Salinas CA. Exercise increases serum fibroblast growth factor 21 (FGF21) levels. *PloS one.* 2012;7(5):e38022.
- 126.** Tanimura Y, Aoi W, Takanami Y, Kawai Y, Mizushima K, Naito Y, Yoshikawa T. Acute exercise increases fibroblast growth factor 21 in metabolic organs and circulation. *Physiological reports.* 2016;4(12).

Figure legends

Figure 1. Exposure of Siberian hamsters to short days initiates molt from a summer agouti coat to a white winter coat (top), and programs a decrease in appetite and increased catabolism of visceral white fat (WAT) and hence body weight loss as a winter survival strategy. This is associated with increased hepatic FGF21 production. Treatment of hamsters in the fat summer state with FGF21 induces behavioral and physiological changes that resemble the natural short day response, including decreased appetite, promotion of fat oxidation rather than carbohydrate oxidation (associated with decreased pyruvate dehydrogenase activity in liver: PDC), and increased thermogenic capacity of brown adipose tissue (BAT) via upregulation of uncoupling protein 1 (*ucp1*) and deiodinase II (*dio2*) gene expression.

Figure 2. In this figure, blue arrows depict decreasing (downward) or increasing (upward) circulating concentrations of Fibroblast Activated Protein (FAP) and the bioactive form of FGF21 (aFGF21). Red 'stop' signs denote the absence of secretion in response to exercise. Red crosses and black block arrows indicate impairment in relevant biological processes, which when accompanied with questions marks indicate outstanding biological questions that require further experimentation.

In healthy humans (left panel), dietary carbohydrates (CHO) increase both the total and bioactive form of FGF21 in an insulin dependent manner, whereas levels of FAP are unaffected. Exercise is a key stimulus in inducing hepatic, but not skeletal muscle, release of FGF21. Exercise also increases circulating levels of FAP with no apparent contribution from skeletal muscle. It remains to be established whether the liver releases FAP in response to exercise. Adipose tissue is the key target for the metabolic effects of FGF21. At target tissues, FGF21 binds to and activates members of the FGFR superfamily of receptor tyrosine kinases. This activation is dependent on obligate co-receptor β Klotho (KLB). FGF21 also markedly reduces sweet and alcohol preference via signaling through the paraventricular nucleus in the hypothalamus. The fact that consumption of dietary CHO increases the levels of FGF21, which in turn may suppress consumption of simple sugars and non-caloric sweeteners, likely indicates the presence of a negative feedback hormonal loop between the liver and the brain.

Obese individuals with the metabolic syndrome and patients with type 2 diabetes (right panel) have elevated circulating levels of FAP, and this is associated with an attenuated ratio of bioactive to total FGF21 in response to an oral CHO (glucose) tolerance test, suggesting that chronic states of insulin resistance may lead to inactivation of FGF21 and limit its beneficial metabolic effects. KLB, the key co-receptor mediating the tissue sensitivity to FGF21, is associated with higher BMI, and its protein levels are reduced in the adipose tissue of obese humans which may further decrease its responsiveness to FGF21. Strategies aiming to minimise the release of FAP (either metabolic syndrome- or exercise-induced) may be of benefit in maximising the metabolic effects of FGF21. Finally, it remains to be seen whether the negative feedback hormonal loop between the liver and the brain, as related to CHO consumption and sweet and alcohol preference, is in any way disturbed/impaired in obesity and diabetes, and whether it plays a role in appetite regulation under those conditions.

Figure 1




Figure 2

