



Conference on ‘Nutrient–nutrient interaction’ Plenary Lecture 2

Interactions between nutrients in the maternal diet and the implications for the long-term health of the offspring

William D. Rees

The Rowett Institute of Nutrition and Health, The University of Aberdeen, Foresterhill, Aberdeen AB25 2ZD, Scotland, UK

Nutritional science has traditionally used the reductionist approach to understand the roles of individual nutrients in growth and development. The macronutrient dense but micronutrient poor diets consumed by many in the Western world may not result in an overt deficiency; however, there may be situations where multiple mild deficiencies combine with excess energy to alter cellular metabolism. These interactions are especially important in pregnancy as changes in early development modify the risk of developing non-communicable diseases later in life. Nutrient interactions affect all stages of fetal development, influencing endocrine programming, organ development and the epigenetic programming of gene expression. The rapidly developing field of stem cell metabolism reveals new links between cellular metabolism and differentiation. This review will consider the interactions between nutrients in the maternal diet and their influence on fetal development, with particular reference to energy metabolism, amino acids and the vitamins in the B group.

Developmental origins of health and disease: Folic acid: Methionine: Maternal obesity

Classically nutritional science has applied the principles of reductionism to investigate nutritional deficiencies, exploring complex systems by reducing them to the interactions of their component parts. This approach has been highly successful in identifying the roles of individual nutrients and has given us a clear description of the mechanisms underlying the major deficiency syndromes. However, in recent years the consumption of energy-dense but micronutrient-poor foodstuffs has created a situation where both over and undernutrition can exist simultaneously in a single human population⁽¹⁾. Understanding the range of nutrient interactions caused by these imbalanced diets and the long-term consequences for health is an ongoing challenge for nutritional science.

The nutrient interactions associated with obesity are particularly important during pregnancy with nearly half of the women presenting at maternity clinics in developed countries such as Scotland classified as

overweight or obese (Births in Scottish Hospitals (National Statistics Publication for Scotland) November 2016). Studies from both the developed and developing world show an inverse relationship between diet quality and BMI, particularly in pregnant women in poorer socioeconomic groups^(2,3). This is because cheap and readily available foods tend to comprise energy-dense, micronutrient-poor products such as refined carbohydrates and oils⁽⁴⁾. Furthermore, an energy-dense diet increases the demand for micronutrient cofactors such as the vitamins in the B group, which are essential for energy metabolism. Imbalanced, energy-dense, micronutrient-poor diets, therefore, present with an array of nutrient interactions, all of which require adaptations in cellular metabolism and gene expression.

These interactions between metabolism and gene expression are especially important in the 9 months from conception to birth, as the human fetus grows

Abbreviation: PC, phosphatidylcholine.

Corresponding author: William D. Rees, email w.rees@abdn.ac.uk

and develops from a single cell to a recognisable infant. A plethora of changes transforms four totipotent stem cells in the preimplantation embryo into increasingly committed cell lineages in mesoderm, endoderm and ectoderm and finally into the terminally differentiated cell types in the organs. Growth and differentiation depend upon a continuous balanced supply of nutrients, derived in part from the maternal diet but also from nutrients released by mobilisation of the mother's tissues. Indeed this relationship continues into postnatal life, as the production of milk also depends on mobilisation of maternal reserves. Because of this interrelationship between maternal diet and reserves, seemingly, simple nutritional imbalances invoke surprisingly complex interactions between macro and micro-nutrients. Maternal metabolism is thus crucial in delivering a stable and balanced nutrient supply to the developing fetus.

Fetal development and long-term health

Over the past 20 years, it has become accepted that poor early development can have a profound effect on the risk of developing non-communicable diseases later in life. The original studies of the Boyd Orr Cohort⁽⁵⁾ and subsequent studies by Barker and colleagues⁽⁶⁻⁸⁾ firmly established the relationship between poor growth (and by implication nutrition) in early life and an increased risk of non-communicable disease. Subsequently, these observations have been confirmed and extended in populations across the world. Extremes of birthweight, both above and below the mean, are associated with increased risk of the offspring developing a range of diseases including CVD^(7,9), metabolic disease^(8,10), osteoporosis⁽¹¹⁾ and psychiatric disorders^(12,13). The concept of life-course nutrition encapsulates these ideas in a scheme where early fetal and neonatal development is plastic and influenced by environmental factors, principal amongst these being nutrition. Physiological changes introduced during fetal and neonatal life modulate the risk of developing the non-communicable disease later in life. Once an individual is set on a higher risk trajectory, interventions in adult life can only partly rescue or ameliorate the impact of early life insults⁽¹⁴⁾. Indeed, nutritional status may be important prior to conception as there is evidence that nutritional status affects follicle recruitment and oocyte maturation in the female and spermatogenesis in the male, modifying the metabolic competence of the gametes and influencing the subsequent development of the conceptus⁽¹⁵⁾.

Maternal metabolism

Pregnancy can be divided into two periods, an initial anabolic phase followed by a second catabolic phase⁽¹⁶⁾. The anabolic phase covers the first two-thirds of pregnancy. In this period, fetal tissues are a very small part of the mother's body and the available nutrients are directed towards the growth of key maternal organs such as liver and adipose tissue. The nutrient reserves stored in the anabolic phase can then be

mobilised during the catabolic phase of gestation (from about 20 weeks onwards in human subjects), supplementing nutrients from the diet. For example, a large increase in lipolysis in adipose tissue late in gestation provides fatty acids for the fetus⁽¹⁷⁾ and also for the mammary gland in readiness for lactation⁽¹⁸⁾. These maternal reserves of protein and lipids are particularly important in supporting fetal metabolism in cases where the maternal diet is limited or imbalanced.

Maternal micronutrient reserves are also mobilised to meet the increased demands of pregnancy and in many cases are preferentially allocated to the fetus^(19,20). If these stores are exhausted during gestation, the deficit continues into lactation, affecting the metabolism of both the mother and her infant. For example, pregnancy has been estimated to require approximately 740 mg additional iron⁽²¹⁾. However, since an ideal diet only provides approximately 5 mg iron daily, a woman must enter pregnancy with iron stores of at least 300 mg if she is to have sufficient iron to avoid becoming deficient by the time the child is born⁽²²⁾. Many women, and particularly those in developing countries, have insufficient reserves and become iron deficient by the time the baby is born. Fetal and postnatal iron deficiency results in a range of adverse consequences for mother and infant including low birth weight, impaired cognitive development and poor immune function⁽²³⁾. Iron is a key component of enzymes of oxidative metabolism and as the electron-carrying capacity of iron-dependent enzymes is reduced by iron deficiency^(24,25), glucose is increasingly used as a metabolic substrate in place of lipids^(26,27). This is also the case in pregnant animals as an increase in the incorporation of glucose into polar lipids suggest a shift from fat to glucose as the preferred substrate for oxidative metabolism⁽²⁸⁾. As pregnancy progresses there are more pronounced changes in lipid metabolism ultimately resulting in marked changes in both amounts and fatty acid composition of the milk⁽²⁹⁾.

A similar pattern is seen with other essential vitamins or minerals, all of which have important roles as cofactors or prosthetic groups in enzymes of major metabolic pathways. In developing countries, and to a lesser extent in developed countries, multiple micronutrient deficiencies are present, creating a complex array of potential nutrient interactions that are only now beginning to be studied with advanced metabolomics approaches. These complex interactions also make it difficult to evaluate the outcome of interventions. Frequently supplements with one or two components prove to be relatively ineffective when compared with multiple micronutrient supplements⁽³⁰⁾.

Consequences of altered fetal growth and development

Animal models have been extremely important both in exploring nutrient interactions in pregnancy and in understanding the underlying mechanisms. Protein metabolism during pregnancy provides a good example of the complex interactions between nutrients during the anabolic and catabolic phases of pregnancy. One of the most widely studied animal models is the rat low-protein model⁽³¹⁾. In this deceptively simple model, laboratory

rats are fed a reduced protein diet for the duration of gestation and lactation and the offspring are compared with animals fed a control diet containing the recommended amount of protein. In most cases, the low protein diet contains 8 or 9 % protein (w/w) and is compared with a control diet containing 18–20 % w/w protein, which is the amount recommended by the National Research Council and American Institute for Nutrition for reproduction in the rat⁽³²⁾. Because of the bi-phasic nature of pregnancy, protein restriction has a limited effect on growth in the early stages of fetal life⁽³³⁾. The effects of a modest protein restriction on growth only become more noticeable during the lactation phase⁽³⁴⁾ when the mobilisation of maternal reserves is no longer able to compensate for the reduced protein supply^(35,36).

Whilst the requirements for individual amino acids have not been established in human subjects, estimates extrapolated from animal studies suggest the demand for individual amino acids changes with both time and the availability of other nutrients⁽³⁷⁾. Plasma amino acid concentrations in the maternal circulation of the female rat reflect the two phases of protein metabolism in pregnancy; typically, concentrations fall in the first stages of gestation and then increase between days 19 and 21 (Fig. 1). Dietary protein restriction has very modest effects on the free amino acid concentrations because amino acids entering the peripheral circulation are constantly buffered by amino acids released by the degradation of existing tissue proteins⁽³⁸⁾ and only a limited number of amino acids are differentially affected, mostly decreasing. The only notable exception is threonine, one of the indispensable amino acids⁽³⁹⁾. The placenta also responds to a reduced dietary supply of protein and this helps to stabilise amino acid concentrations in the fetal circulation, which are usually higher than in the maternal circulation; however, the profile tends to follow the pattern in the maternal circulation^(33,40).

However, amino acids are more than just the monomers for protein synthesis; they are also important substrates for intermediary metabolism, creating a range of potential nutrient interactions in the low protein model. Animal studies suggest that a low protein diet produces changes in intermediary metabolism, which affect development throughout gestation, even in the pre-, and periconception phases of life before the conceptus has become implanted⁽⁴¹⁾. Changes in intermediary metabolism possibly linked to the oxidation of excess amino acids alters the available concentrations of other important metabolic fuels. For example, concentrations of TAG in the maternal circulation of animals fed a low protein diet are higher than in animals fed sufficient protein and this is likely to increase the supply of fatty acids to fetal tissues and influence their development⁽⁴²⁾.

Interactions between macro-nutrients in the maternal diet and the effect on the postnatal phenotype of the offspring

In adult life, the offspring of female rats fed a low protein diet exhibit changes in insulin action and in time, these

develop into diabetic-like symptoms⁽⁴³⁾. This phenotype, reflected in the area under the insulin release curve when animals are given a glucose challenge⁽⁴⁴⁾, is the result of reduced growth and development of the pancreatic β -cells⁽⁴⁵⁾ and a decrease in the insulin sensitivity of the tissues⁽⁴⁶⁾. However, the extent of this programming is dependent on factors other than just the protein content of the maternal diet⁽⁴⁷⁾. The type of lipid and hence the fatty acid composition of the diet modifies the final outcome such that the offspring of rats fed a low protein diet prepared with maize oil, low in *n*-3 fatty acids, release less insulin in the glucose tolerance test compared with the offspring of animals fed a diet prepared with soya oil^(48,49).

This and other similar results suggest that changes in maternal metabolism, modifying the supply of macronutrients to the developing fetal organs, may be an important part of the programming mechanism. Increases in maternal plasma TAG concentrations are a feature of a number of different models of fetal programming in the rat, including low protein, high fat⁽⁵⁰⁾ and folate deficient diets⁽⁵¹⁾ as well as iron deficiency⁽²⁸⁾, suggesting an important role for lipid metabolism in programming. A proposed mechanism envisages the setting of metabolic set points during development in response to the available nutrient supply⁽⁵²⁾. For example, the regulation of blood glucose by insulin is a closed loop biological control system, maintaining plasma concentrations between the tightly constrained maximum and minimum values. The set points at which insulin or glucagon are released in response to changes in the prevailing concentrations of glucose are learned during development. If the prevailing glucose concentrations are higher than average during development, then a correspondingly higher set point may be established during fetal development affecting both insulin release and tissue insulin sensitivity.

Nutrient interactions caused by maternal obesity during pregnancy

Whilst many prospective studies in human populations or animal models have focused on fetal growth restriction caused by nutrient deficiency, maternal obesity caused by the oversupply of energy is now perceived to be a greater problem. Increasing numbers of women are now presenting at antenatal clinics with a higher than normal BMI. Obesity in pregnancy carries a number of risks for the mother with higher rates of pre-eclampsia and premature birth, while the child is at increased risk of macrosomia and gross malformations such as neural tube defects⁽⁵³⁾. There is also evidence for long-term changes in both metabolism and behaviour in the offspring of obese mothers⁽⁵⁴⁾.

Obesity in human populations is not the simple product of a high-energy diet, the simultaneous presence of a double-burden of obesity and malnutrition has been described in populations worldwide⁽⁵⁵⁾. There is an inverse association between BMI and diet quality^(2,56) and this translates into an inverse association between BMI and micronutrient status^(57,58). Deficiencies in the




Fig. 1. The biphasic nature of amino acid metabolism in the rat during gestation. Concentrations of glycine (top) leucine (middle) and threonine (bottom) in the maternal plasma of rats fed a diet containing high (18 % w/w) protein (open circles dotted line) and low (9 % w/w) protein (closed circles solid line). Adapted from Rees *et al.* (33,39).

morbidly obese include vitamins B₁, B₁₂, folic acid, Vitamin A, Vitamin D and minerals including iron and zinc⁽⁵⁹⁾. The double burden is a product of socio-economic factors that promote the consumption of a diet containing large quantities of refined fat or carbohydrate products. Whilst these low-cost foods are plentiful and have a high-energy content, processing tends to deplete them of key vitamins. For example, milling wheat and other grains remove the vitamin-rich aleurone layer, containing approximately 80 % of total niacin together with considerable quantities of other B vitamins, betaine and choline^(60,61). At present little is known about the range of complex nutrient interactions




Fig. 2. The interactions between amino acids, choline, the methionine cycle and lipid metabolism in the liver. 5-CH₃-THF, 5 methyl tetrahydrofolate; 5,10-CH₂-THF, 5,10 methylene tetrahydrofolate; DHF, dihydrofolate; Pcho, phosphocholine; SAM, S-adenosyl methionine; SAHcy, S-adenosyl homocysteine; THF, tetrahydrofolate.

involving vitamins, minerals and amino acids involved in the metabolism of excessive amounts of energy. Incomplete or diverted biochemical reactions could lead to toxicity such as damage caused by the accumulation of NEFA⁽⁶²⁾. As the prevalence of obesity increases, we cannot consider the consequences of excess fat consumption in isolation. The influence of marginal micro-nutrient deficiency must also be taken into account, even in wealthy Western countries.

Folic acid, the B vitamins and choline in pregnancy

Central in the metabolism of lipids is the B group of vitamins, which include folic acid. There is a direct relationship between maternal folate status and birthweight, illustrating the importance of folic acid for fetal development^(63,64). Folate derivatives are also important for specific processes in development. It is well known that folic acid supplements reduce the risk of neural tube defects in pregnancy, although questions remain over the precise mechanism. Recent evidence suggests that an improvement in the formate supply and a corresponding increase in nucleotide biosynthesis may be an important factor⁽⁶⁵⁾. Formate, produced in the mitochondria is transported into the cytoplasm to provide formyl tetrahydrofolate groups necessary for a range of reactions some of which may be essential for neural tube development⁽⁶⁶⁾.

Fetal cell proliferation requires the production of large quantities of DNA and membrane lipids as well as the synthesis of new protein. All of these processes are dependent on folate-mediated C_1 metabolism and derivatives of folic acid are important intermediates in the methionine cycle, which is in turn linked to lipid and nucleotide metabolism (Fig. 2). The methionine cycle maintains intracellular pools of S-adenosyl methionine, the donor of methyl groups in more than hundred essential metabolic reactions. In addition to methionine, the methionine cycle also depends on a variety of other nutrients, particularly the amino acids glycine-serine and threonine as well as micronutrients such as the B vitamins⁽⁶⁷⁾.

A crucial intermediate in the cycle is the non-protein amino acid homocysteine, concentrations of which are increased when the rate of production exceeds the capacity of the reactions that convert it back to methionine (remethylation) or convert it to cysteine (transulphuration). The importance of C_1 metabolism and the methionine cycle for fetal growth is illustrated by the inverse relationship between plasma homocysteine concentrations and birth weight⁽⁶⁸⁾. Nutrient interactions have a bearing on the rate of homocysteine removal as both remethylation and transulphuration reactions depend on the availability of other amino acids and vitamin cofactors. The converse is also true in that amino acids and cofactors can become depleted depending on whether homocysteine is remethylated or removed by transulphuration. For example, both choline and derivatives of folic acid provide methyl groups that are used to convert homocysteine back to methionine. The flux through the remethylation reactions increases when methionine is limited and is dependent on the availability of choline, betaine and folic acid in the diet⁽⁶⁹⁾. Reducing the supply of folic acid increases the utilisation of choline and reduces the use of serine, glycine and threonine. As a result, the phosphocholine pool is depleted and concentrations of serine, glycine and threonine increase in the maternal plasma⁽⁷⁰⁾. Restricting the amount of methionine in the diet also reduces concentrations of taurine, another non-protein amino acid implicated in fetal programming^(71,72). If additional methionine is provided in the diet then these folate and choline dependent changes in methionine cycling are lost showing that surplus methionine can be used as a source of methyl groups. Homocysteine can be converted by the transulphuration pathway to cysteine, via the intermediate cystathionine in a reaction dependent upon vitamin B_6 , pyridoxal phosphate and the amino acid serine⁽⁷³⁾. This pathway is an important route for the direct use of methionine as a methyl donor and is essential for reproduction as animals lacking the gene coding for cystathionine- β -synthase are infertile as the conceptus fails to implant⁽⁷⁴⁾.

Measurements of methionine turnover using stable isotopes show that in the rat both maternal and fetal livers have active methionine cycles. There is a gradual decline in the remethylation of methionine in the maternal liver as gestation progresses. However, at the same time, methionine is actively produced from homocysteine in the fetal liver, with homocysteine taken up from the maternal circulation and converted to methionine by

the fetal tissues⁽⁷⁵⁾. When animals are fed a diet low in folic acid and choline there is a decrease in the remethylation of homocysteine in the maternal liver; however, to compensate for this there is an increase in the rate of methionine cycling in the fetal liver. In part, this reflects the preferential partitioning of the available folic acid to the fetal tissues but also suggests that there is an increase in the utilisation of methyl groups in fetal tissues, perhaps to produce products normally produced in the maternal compartment.

Impact of limited C_1 metabolism on the metabolism of the mother

One of the most obvious features of pregnant rats fed a methyl-deficient diet is an accumulation of lipid in the maternal liver late in gestation⁽⁵¹⁾. The liver has an important role in this period, processing lipids in the diet alongside those mobilised from the adipose tissue in the catabolic phase of pregnancy. Despite the increased flow of TAG through the hepatocytes, intracellular concentrations are normally low as a lipid is rapidly exported as VLDL, which consist of a lipid core surrounded by an outer layer of phospholipids and lipoproteins. The predominant phospholipid in VLDL is phosphatidylcholine (PC) and this must be produced by the liver to maintain high rates of VLDL secretion. However, diets low in choline and associated methyl donors restrict PC synthesis, and lipid that cannot be exported accumulates as droplets within the hepatocytes. The liver produces PC (Fig. 2) either through the cytidine diphosphocholine pathway (also known as the Kennedy pathway) using choline as a precursor or by the conversion of phosphatidylethanolamine to PC by the enzyme phosphatidylethanolamine methyl transferase^(76,77). Since the phosphatidylethanolamine methyl transferase pathway requires three methyl groups from S-adenosyl methionine for the synthesis of a single PC molecule, it creates a substantial demand for methyl groups⁽⁷⁸⁾. It becomes apparent from the complex interrelationships that changes in the supply of amino acids and choline can interact with maternal micronutrient status to produce a range of different outcomes. Further complexity is introduced by increases in the flow of lipids through the liver, a product of both excess in the mother's diet and mobilisation of excess body fat.

Analysis of fetal tissues suggests that methyl deficiency produces changes in the fetal liver^(79,80), heart⁽⁸¹⁾, gut⁽⁸²⁾ and brain^(83,84). Measurements of methionine flux show that methyl metabolism is protected in the late gestation fetus⁽⁷⁵⁾ suggesting that it is the deficiency or excess of the products of folate-dependent metabolism that influences the regulation of gene expression⁽⁸¹⁾.

Fetal metabolism: the totipotent stem cells of the preimplantation embryo

The process of tissue differentiation is accompanied by widespread changes in gene expression. Actively



transcribed genes are located in euchromatin, which are loosely packaged areas of chromatin accessible for transcription. Genes not required are tightly packaged into heterochromatin, which is less accessible. These changes in chromatin structure require extensive covalent modification of both DNA and the proteins that surround it⁽⁸⁵⁾. Prominent changes include the addition of methyl groups to DNA (DNA methylation) and histone proteins (histone methylation) and the acetylation of histone proteins (histone acetylation). Euchromatin is characterised by the presence of low levels of DNA methylation and hyperacetylated histones whereas heterochromatin has high levels of DNA methylation and hypoacetylation of histones.

It is becoming apparent that the metabolic state of the cell can modify the epigenetic regulation of gene expression during differentiation through changes in the availability of cofactors required for the modification of DNA and histones^(86,87). The most widely studied, thanks to the availability of effective tools to precisely measure changes in the modification of individual bases, are the epigenetic marks created by DNA methylation. Changes in DNA methylation and gene expression in the offspring of animals fed a methyl-deficient diet were first shown in the classical studies of the Agouti mouse⁽⁸⁸⁾. Subsequent studies of human populations have identified a relationship between methyl donors in the maternal diet and DNA methylation in the offspring⁽⁸⁹⁾. The advent of high throughput methods for assessing DNA methylation has made it possible to explore the relationships between maternal nutrition and fetal epigenotype and a range of population-wide studies are currently in progress^(90,91).

The methylation of DNA is only part of the process of cell differentiation; the posttranslational modification of histones also plays a vital part in maintaining the pluripotent state in stem cells. Initially isolated from early embryos, embryonic stem cells are difficult to maintain in tissue culture, as they are prone to spontaneous differentiation. In a breakthrough study, threonine was identified as a critical component of the culture media⁽⁹²⁾ as rodent cells grown in low threonine media rapidly lost the capacity to differentiate. Subsequent studies demonstrated that the catabolism of threonine produces both glycine and acetyl CoA; glycine is used to produce C₁ folate derivatives and acetyl CoA is used for protein acetylation^(93,94). In a similar way, human stem cells use methionine as the principal source of methyl groups for histone methylation⁽⁹⁵⁾. The reversal of epigenetic marking by the removal of methyl groups from histones and DNA is also sensitive to the availability of specific metabolites. Histone demethylases and ten-eleven translocation methylcytosine dioxygenases, which catalyse the first step of DNA demethylation require both the tri-carboxylic acid cycle intermediate α -ketoglutarate and ferric iron^(94,96).

These changes in intermediary metabolism are linked to the function of the mitochondria during cell differentiation. Undifferentiated, or 'naive' embryonic stem cells, have relatively high levels of oxidative phosphorylation. When differentiation is initiated and cells enter the 'primed'

state, oxidative phosphorylation becomes uncoupled and metabolism switches to glycolysis, using glucose for energy and increasing the production of acetate for the acetylation of histones⁽⁹⁷⁾. Energy production from oxidative phosphorylation then increases again as the cells become terminally differentiated. Mitochondria in oocytes and preimplantation embryos have an immature structure with underdeveloped cristae and low rates of oxidative phosphorylation, the classical mitochondrial morphology with well-developed cristae and an active oxidative phosphorylation only appears in terminally differentiated cells⁽⁹⁸⁾. The reprogramming of differentiated somatic cells into induced pluripotent stem cells reverses these changes and restores the glycolytic state. Changes in metabolism occur before changes in gene expression, suggesting that altered metabolism is required for the cell fate change rather than a consequence of it⁽⁹⁶⁾.

Future perspectives

Optimal fetal development is important for long-term health and there is clear evidence that the supply of macro and micro-nutrients in the maternal diet interact to programme fetal development. There are a variety of potential mechanisms including endocrine programming, changes in organ development and epigenetic programming of gene expression. The rapidly developing field of stem cell metabolism is now opening a new window on the differentiation of cells during development. At present, these studies are limited to *in vitro* analysis of cells in culture but clearly show that several metabolic pathways have the potential to influence stem cell growth and differentiation through changes in intracellular pools of intermediary metabolites, such as α -ketoglutarate, S-adenosyl methionine, and acetyl-CoA⁽⁹⁹⁾. It is possible that these changes can produce some profound differences in cellularity of the organs by preventing the premature differentiation of cells. It remains to be seen how these findings will translate into the whole body physiology of the developing fetus, however, it is easy to see potential links between metabolism and the energy-dense but micronutrient-poor diets consumed by many human subjects. Whilst it is possible that maternal metabolism is sufficiently well buffered to protect the developing stem cells there is good evidence from animal models to show that imbalanced diets alter the supply of glucose, amino acids and particularly lipids to the developing fetus. A number of new analytical methods are now needed to explore the relationship between metabolism and chromatin structure. By coupling these with insights into metabolism from the developing field of metabolomics, we may be in a good position to begin to document the complex nutrient interactions that influence chromatin structure, fetal development and long-term health.

Acknowledgements

The author would like to thank colleagues at the Rowett Institute and members of his laboratory, past and

present, for facilitating the development of this research area.

Financial Support

Work in the author's laboratory was supported by The Scottish Government Strategic Research Programme.

Conflict of Interest

None.

Authorship

The author had sole responsibility for all aspects of preparation of the present paper.

References

1. Demaio AR & Branca F (2018) Decade of action on nutrition: our window to act on the double burden of malnutrition. *BMJ Global Health* **3**, Suppl. 1.
2. Laraia BA, Bodnar LM & Siega-Riz AM (2007) Pregravid body mass index is negatively associated with diet quality during pregnancy. *Public Health Nutr* **10**, 920–926.
3. Shin D, Lee KW & Song WO (2016) Pre-pregnancy weight status is associated with diet quality and nutritional biomarkers during pregnancy. *Nutrients* **8**, 162.
4. Moubarac J-C, Batal M, Louzada ML *et al.* (2017) Consumption of ultra-processed foods predicts diet quality in Canada. *Appetite* **108**, 512–520.
5. Martin RM, Gunnell D, Pemberton J *et al.* (2005) Cohort profile: the Boyd Orr cohort—a historical cohort study based on the 65 year follow-up of the Carnegie Survey of Diet and Health (1937–39). *Int J Epidemiol* **34**, 742–749.
6. Barker DJP & Osmond C (1986) Infant mortality, childhood nutrition, and ischaemic heart disease in England and Wales. *Lancet* **327**, 1077–1081.
7. Barker DJP, Osmond C, Winter PD *et al.* (1989) Weight in infancy and death from ischaemic heart disease. *Lancet* **334**, 577–580.
8. Hales CN, Barker DJ, Clark PM *et al.* (1991) Fetal and infant growth and impaired glucose tolerance at age 64. *BMJ: Br. Med. J.* **303**, 1019–1022.
9. Barker DJ, Osmond C, Forsen TJ *et al.* (2005) Trajectories of growth among children who have coronary events as adults. *N Engl J Med* **353**, 1802–1809.
10. Hales CN & Barker DJ (1992) Type 2 (non-insulin-dependent) diabetes mellitus: the thrifty phenotype hypothesis. *Diabetologia* **35**, 595–601.
11. Cooper C, Eriksson JG, Forsen T *et al.* (2001) Maternal height, childhood growth and risk of hip fracture in later life: a longitudinal study. *Osteoporosis International* **12**, 623–629.
12. Thompson C, Syddall H, Rodin I *et al.* (2001) Birth weight and the risk of depressive disorder in late life. *Br J Psychiatry* **179**, 450–455.
13. Veena SR, Gale CR, Krishnaveni GV *et al.* (2016) Association between maternal nutritional status in pregnancy and offspring cognitive function during childhood and adolescence; a systematic review. *BMC Pregnancy Childbirth* **16**, 220.
14. Hanson MA & Gluckman PD (2014) Early developmental conditioning of later health and disease: physiology or pathophysiology? *Physiol Rev* **94**, 1027–1076.
15. Fleming TP, Watkins AJ, Velazquez MA *et al.* (2018) Origins of lifetime health around the time of conception: causes and consequences. *Lancet* **391**, 1842–1852.
16. Herrera E (2000) Metabolic adaptations in pregnancy and their implications for the availability of substrates to the fetus. *Eur J Clin Nutr* **54**, Suppl. 1, S47–S51.
17. Herrera E, Amusquivar E, López-Soldado I *et al.* (2006) Maternal lipid metabolism and placental lipid transfer. *Horm Res Paediatr* **65**, Suppl. 3, 59–64.
18. Vernon RG & Pond CM (1997) Adaptations of maternal adipose tissue to lactation. *J Mammary Gland Biol Neoplasia* **2**, 231–241.
19. Costello AM & Osrin D (2003) Micronutrient status during pregnancy and outcomes for newborn infants in developing countries. *J Nutr* **133**, 5 Suppl. 2, 1757s–1764s.
20. Devakumar D, Fall CHD, Sachdev HS *et al.* (2016) Maternal antenatal multiple micronutrient supplementation for long-term health benefits in children: a systematic review and meta-analysis. *BMC Med* **14**, 90.
21. Fisher AL & Nemeth E (2017) Iron homeostasis during pregnancy. *Am J Clin Nutr* **106**, Suppl. 6, 1567S–1574S.
22. Bothwell TH (2000) Iron requirements in pregnancy and strategies to meet them. *Am J Clin Nutr* **72**, Suppl. 1, 257s–264s.
23. Lozoff B (2007) Iron deficiency and child development. *Food Nutr Bull* **28**, Suppl. 4, S560–S571.
24. Evans TC & Mackler B (1985) Effect of iron deficiency on energy conservation in rat liver and skeletal muscle submitochondrial particles. *Biochem Med* **34**, 93–99.
25. Mackler B, Grace R & Finch CA (1984) Iron deficiency in the rat: effects of oxidative metabolism in distinct types of skeletal muscle. *Pediatr Res* **18**, 499–500.
26. Davis M, Rendina E, Peterson S *et al.* (2012) Enhanced expression of lipogenic genes may contribute to hyperglycemia and alterations in plasma lipids in response to dietary iron deficiency. *Genes Nutr* **7**, 415–425.
27. Borel MJ, Beard JL & Farrell PA (1993) Hepatic glucose production and insulin sensitivity and responsiveness in iron-deficient anemic rats. *Am J Physiol – Endocrinol Metab* **264**, E380–EE90.
28. Hay SM, McArdle HJ, Hayes HE *et al.* (2016) The effect of iron deficiency on the temporal changes in the expression of genes associated with fat metabolism in the pregnant rat. *Physiol Rep* **4**, e12908.
29. O'Connor DL, Picciano MF & Sherman AR (1988) Impact of maternal iron deficiency on quality and quantity of milk ingested by neonatal rats. *Br J Nutr* **60**, 477–485.
30. Ramakrishnan U, Goldenberg T & Allen LH (2011) Do multiple micronutrient interventions improve child health, growth, and development? *J Nutr* **141**, 2066–2075.
31. Langley SC, Browne RF & Jackson AA (1994) Altered glucose tolerance in rats exposed to maternal low protein diets in utero. *Comp Biochem Physiol Physiol* **109**, 223–229.
32. Reeves PG, Nielsen FH & Fahey GC Jr (1993) AIN-93 purified diets for laboratory rodents: final report of the American Institute of Nutrition ad hoc writing committee on the reformulation of the AIN-76A rodent diet. *J Nutr* **123**, 1939–1951.
33. Rees WD, Hay SM, Buchan V *et al.* (1999) The effects of maternal protein restriction on the growth of the rat fetus and its amino acid supply. *Br J Nutr* **81**, 243–250.
34. Guzmán C, Cabrera R, Cárdenas M *et al.* (2006) Protein restriction during fetal and neonatal development in the

- rat alters reproductive function and accelerates reproductive ageing in female progeny. *J Physiol* **572**(Pt 1), 97–108.
35. Naismith DJ & Robinson SM (1987) Adaptations in protein metabolism during lactation in the rat. *Br J Nutr* **58**, 533–538.
 36. Pine AP, Jessop NS, Allan GF *et al.* (1994) Maternal protein reserves and their influence on lactational performance in rats. 4. Tissue protein synthesis and turnover associated with mobilization of maternal protein. *Br J Nutr* **72**, 831–844.
 37. Kwon H, Spencer TE, Bazer FW *et al.* (2003) Developmental changes of amino acids in ovine fetal fluids. *Biol Reprod* **68**, 1813–1820.
 38. Elango R & Ball RO (2016) Protein and amino acid requirements during pregnancy. *Adv Nutr* **7**, 839S–844S.
 39. Rees WD, Hay SM & Antipatis C (2006) The effect of dietary protein on the amino acid supply and threonine metabolism in the pregnant rat. *Reprod Nutr Dev* **46**, 227–239.
 40. Gaccioli F, Lager S, Powell T *et al.* (2013) Placental transport in response to altered maternal nutrition. *J Dev Orig Health Dis* **4**, 101–115.
 41. Kwong WY, Miller DJ, Wilkins AP *et al.* (2007) Maternal low protein diet restricted to the preimplantation period induces a gender-specific change on hepatic gene expression in rat fetuses. *Mol Reprod Dev* **74**, 48–56.
 42. McNeil CJ, Maloney CA, Hay SM *et al.* (2007) Sources of dietary protein and lipid interact to modify maternal and fetal development in the pregnant rat. *Proc Nutr Soc* **66**, Suppl. OCA-B, 21A.
 43. Harding JE, Jaquiere AL, Hernandez CE *et al.* (2011) Animal studies of the effects of early nutrition on long-term health. *Nestle Nutr Workshop Ser Paediatr Programme* **68**, 1–11; discussion –6.
 44. Rees WD, Hay SM, Cruickshank M *et al.* (2006) Maternal protein intake in the pregnant rat programs the insulin axis and body composition in the offspring. *Metabolism* **55**, 642–649.
 45. Reusens B, Theys N, Dumortier O *et al.* (2011) Maternal malnutrition programs the endocrine pancreas in progeny. *Am J Clin Nutr* **94**, Suppl. 6, 1824S–1829S.
 46. Martin-Gronert M & Ozanne S (2012) Mechanisms underlying the developmental origins of disease. *Rev Endocr Metab Disord* **13**, 85–92.
 47. Langley-Evans SC (2000) Critical differences between two low protein diet protocols in the programming of hypertension in the rat. *Int J Food Sci Nutr* **51**, 11–17.
 48. Maloney CA, Lilley C, Czopek A *et al.* (2007) Interactions between protein and vegetable oils in the maternal diet determine the programming of the insulin axis in the rat. *Br J Nutr* **97**, 912–920.
 49. Mennitti LV, Oliveira JL, Morais CA *et al.* (2015) Type of fatty acids in maternal diets during pregnancy and/or lactation and metabolic consequences of the offspring. *J Nutr Biochem* **26**, 99–111.
 50. Williams L, Seki Y, Vuguin PM *et al.* (2014) Animal models of in utero exposure to a high fat diet: a review. *BBA – Mol Basis Disease* **1842**, 507–519.
 51. McNeil CJ, Hay SM, Rucklidge G *et al.* (2008) Disruption of lipid metabolism in the liver of the pregnant rat fed folate deficient and methyl donor deficient diets. *Br J Nutr* **99**, 262–271.
 52. Plagemann A (2011) Maternal diabetes and perinatal programming. *Early Hum Dev* **87**, 743–747.
 53. Poston L, Caleyachetty R, Cnattingius S *et al.* (2016) Preconceptional and maternal obesity: epidemiology and health consequences. *Lancet Diabetes Endocrinol* **4**, 1025–1036.
 54. Mikkelsen SH, Hohwü L, Olsen J *et al.* (2017) Parental body mass index and behavioral problems in their offspring: a Danish national birth cohort study. *Am J Epidemiol* **186**, 593–602.
 55. Wahlqvist ML, Krawetz SA, Rizzo NS *et al.* (2015) Early-life influences on obesity: from preconception to adolescence. *Ann N Y Acad Sci* **1347**, 1–28.
 56. Murakami K & Livingstone MBE (2016) Energy density of meals and snacks in the British diet in relation to overall diet quality, BMI and waist circumference: findings from the National Diet and Nutrition Survey. *Br J Nutr* **116**, 1479–1489.
 57. Kimmons JE, Blanck HM, Tohill BC *et al.* (2006) Associations between body mass index and the prevalence of low micronutrient levels among US adults. *MedGenMed* **8**, 59.
 58. Sen S, Iyer C & Meydani SN (2013) Obesity during pregnancy alters maternal oxidant balance and micronutrient status. *J Perinatol* **34**, 105.
 59. Kaidar-Person O, Person B, Szomstein S *et al.* (2008) Nutritional deficiencies in morbidly obese patients: a new form of malnutrition? *Obes Surg* **18**, 870–876.
 60. Likes R, Madl RL, Zeisel SH *et al.* (2007) The betaine and choline content of a whole wheat flour compared to other mill streams. *J Cereal Sci* **46**, 93–95.
 61. Brouns F, Hemery Y, Price R *et al.* (2012) Wheat aleurone: separation, composition, health aspects, and potential food use. *Crit Rev Food Sci Nutr* **52**, 553–568.
 62. Brookheart RT, Michel CI & Schaffer JE (2009) As a matter of fat. *Cell Metab* **10**, 9–12.
 63. Relton CL, Pearce MS & Parker L (2005) The influence of erythrocyte folate and serum vitamin B12 status on birth weight. *Br J Nutr* **93**, 593–599.
 64. van Uitert EM & Steegers-Theunissen RP (2013) Influence of maternal folate status on human fetal growth parameters. *Mol Nutr Food Res* **57**, 582–595.
 65. Kim J, Lei Y, Guo J *et al.* (2018) Formate rescues neural tube defects caused by mutations in Slc25a32. *Proc Natl Acad Sci USA* **115**, 4690–4695.
 66. Brosnan ME & Brosnan JT (2016) Formate: the neglected member of one-carbon metabolism. *Annu Rev Nutr* **36**, 369–388.
 67. Rees WD (2002) Manipulating the sulfur amino acid content of the early diet and its implications for long-term health. *Proc Nutr Soc* **61**, 71–77.
 68. Bergen NE, Schalekamp-Timmermans S, Jaddoe VW *et al.* (2016) Maternal and neonatal markers of the homocysteine pathway and fetal growth: the generation r study. *Paediatr Perinat Epidemiol* **30**, 386–396.
 69. Wilson FA, van den Borne JJ, Calder AG *et al.* (2009) Tissue methionine cycle activity and homocysteine metabolism in female rats: impact of dietary methionine and folate plus choline. *Am J Physiol Endocrinol Metab* **296**, E702–E713.
 70. Maloney CA, Hay SM & Rees WD (2007) Folate deficiency during pregnancy impacts on methyl metabolism without affecting global DNA methylation in the rat fetus. *Br J Nutr* **97**, 1090–1098.
 71. Boujendar S, Arany E, Hill D *et al.* (2003) Taurine supplementation of a low protein diet fed to rat dams normalizes the vascularization of the fetal endocrine pancreas. *J Nutr* **133**, 2820–2825.
 72. Boujendar S, Reusens B, Merezak S *et al.* (2002) Taurine supplementation to a low protein diet during foetal and early postnatal life restores a normal proliferation and apoptosis of rat pancreatic islets. *Diabetologia* **45**, 856–866.





73. Gregory JF, DeRatt BN, Rios-Avila L *et al.* (2016) Vitamin B6 nutritional status and cellular availability of pyridoxal 5'-phosphate govern the function of the transsulfuration pathway's canonical reactions and hydrogen sulfide production via side reactions. *Biochimie* **126**, 21–26.
74. Nuño-Ayala M, Guillén N, Arnal C *et al.* (2012) Cystathionine β -synthase deficiency causes infertility by impairing decidualization and gene expression networks in uterus implantation sites. *Physiol Genomics* **44**, 702–716.
75. Wilson FA, Holtrop G, Calder AG *et al.* (2012) Effects of methyl-deficient diets on methionine and homocysteine metabolism in the pregnant rat. *Am J Physiol Endocrinol Metab* **302**, E1531–E1E40.
76. Vance DE (2014) Phospholipid methylation in mammals: from biochemistry to physiological function. *Biochim Biophys Acta* **1838**, 1477–1487.
77. Cole LK, Vance JE & Vance DE (2012) Phosphatidylcholine biosynthesis and lipoprotein metabolism. *Biochim Biophys Acta* **1821**, 754–761.
78. Stead LM, Brosnan JT, Brosnan ME *et al.* (2006) Is it time to reevaluate methyl balance in humans? *Am J Clin Nutr* **83**, 5–10.
79. McNeil CJ, Hay SM, Rucklidge GJ *et al.* (2009) Maternal diets deficient in folic acid and related methyl donors modify mechanisms associated with lipid metabolism in the fetal liver of the rat. *Br J Nutr* **102**, 1445–1452.
80. Pooya S, Blaise S, Garcia MM *et al.* (2012) Methyl donor deficiency impairs fatty acid oxidation through PGC-1 α hypomethylation and decreased ER- α , ERR- α and HNF-4 α in rat liver. *J Hepatol* **57**, 344–351.
81. Garcia MM, Gueant-Rodriguez RM, Pooya S *et al.* (2011) Methyl donor deficiency induces cardiomyopathy through altered methylation/acetylation of PGC-1 α by PRMT1 and SIRT1. *J Pathol* **255**, 324–335.
82. Bossenmeyer-Pourié C, Blaise S, Pourié G *et al.* (2010) Methyl donor deficiency affects fetal programming of gastric ghrelin cell organization and function in the rat. *Am J Pathol* **176**, 270–277.
83. El Hajj Chehadeh S, Dreumont N, Willekens J *et al.* (2014) Early methyl donor deficiency alters cAMP signaling pathway and neurosteroidogenesis in the cerebellum of female rat pups. *Am J Physiol Endocrinol Metab* **307**, E1009–E1E19.
84. Geoffroy A, Kerek R, Pourié G *et al.* (2017) Late maternal folate supplementation rescues from methyl donor deficiency-associated brain defects by restoring Let-7 and miR-34 pathways. *Mol Neurobiol* **54**, 5017–5033.
85. Reid MA, Dai Z & Locasale JW (2017) The impact of cellular metabolism on chromatin dynamics and epigenetics. *Nat Cell Biol* **19**, 1298.
86. Shyh-Chang N & Daley George Q (2015) Metabolic switches linked to pluripotency and embryonic stem cell differentiation. *Cell Metab* **21**, 349–350.
87. Dahan P, Lu V, Nguyen RMT *et al.* (2018) Metabolism in pluripotency: both driver and passenger? *J Biol Chem* [Epublication ahead of print version].
88. Waterland RA & Jirtle RL (2003) Transposable elements: targets for early nutritional effects on epigenetic gene regulation. *Mol Cell Biol* **23**, 5293–5300.
89. Dominguez-Salas P, Moore SE, Cole D *et al.* (2013) DNA methylation potential: dietary intake and blood concentrations of one-carbon metabolites and cofactors in rural African women. *Am J Clin Nutr* **97**, 1217–1227.
90. Bianco-Miotto T, Craig JM, Gasser YP *et al.* (2017) Epigenetics and DOHaD: from basics to birth and beyond. *J Dev Orig Health Dis* **8**, 513–519.
91. Kooijman MN, Kruithof CJ, van Duijn CM *et al.* (2016) The generation R study: design and cohort update 2017. *Eur J Epidemiol* **31**, 1243–1264.
92. Wang J, Alexander P, Wu L *et al.* (2009) Dependence of mouse embryonic stem cells on threonine catabolism. *Science* **325**, 435–439.
93. Shyh-Chang N, Locasale JW, Lyssiotis CA *et al.* (2013) Influence of threonine metabolism on S-Adenosylmethionine and histone methylation. *Science* **339**, 222–226.
94. Shyh-Chang N & Ng H-H (2017) The metabolic programming of stem cells. *Genes Dev* **31**, 336–346.
95. Shiraki N, Shiraki Y, Tsuyama T *et al.* (2014) Methionine metabolism regulates maintenance and differentiation of human pluripotent stem cells. *Cell Metab* **19**, 780–794.
96. Tatapudy S, Aloisio F, Barber D *et al.* (2017) Cell fate decisions: emerging roles for metabolic signals and cell morphology. *EMBO Rep* **18**, 2105–2118.
97. Moussaieff A, Kogan NM & Aberdam D (2015) Concise review: energy metabolites: key mediators of the epigenetic state of pluripotency. *Stem cells (Dayton, Ohio)* **33**, 2374–2380.
98. Prigione A, Ruiz-Pérez M, Bukowiecki R *et al.* (2015) Metabolic restructuring and cell fate conversion. *Cell Mol Life Sci* **72**, 1759–1777.
99. Zhang J, Zhao J, Dahan P *et al.* (2018) Metabolism in pluripotent stem cells and early mammalian development. *Cell Metab* **27**, 332–338.