



Adolescent contraception and abortion-related care: a three country comparison and Malawi case study

LSE Research Online URL for this paper: <http://eprints.lse.ac.uk/102441/>

Version: Published Version

Conference or Workshop Item:

Coast, Ernestina, Chiudzu, G, Fetters, T and Tembo, L (2019) Adolescent contraception and abortion-related care: a three country comparison and Malawi case study. In: Malawi College of Medicine Research Dissemination Conference, 2019-11-07 - 2019-11-08, College of Medicine.

Reuse

Items deposited in LSE Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the LSE Research Online record for the item.

Adolescent contraception and abortion-related care: a three country comparison and Malawi case study

PRESENTER: **Dr Grace Chiudzu**

OBJECTIVES:

1. To understand why contraceptive and abortion-related care services are not used more fully by adolescents.
2. To understand the opportunities and barriers to scaling up the most effective implementation strategies for meeting adolescents' needs for contraception and abortion-related services.
3. To maximise the uptake of the research findings by decision-makers, practitioners and service providers.


METHODS

1. Facility-based recruitment of [n=105] 10-19 year-olds seeking either safe abortion care or post-abortion care at a tertiary public sector hospital in Lilongwe.
2. Quantitative and qualitative data collected

OUR SAMPLE


Sample characteristics	
	Malawi [n=105]
	%
Currently has paid work	14
Current student	64
10-14 years	1
15-17 years	58
18-19 years	40
Single	86
Married / cohabiting	8
Divorced / separated	6

Adolescents in Malawi have low levels of contraceptive knowledge and use unsafe abortion for unwanted pregnancy.




Preliminary findings from Malawi:


Many girls made **multiple attempts** to terminate their pregnancies before seeking care at a health facility (n = 96)



The percentage of 10 to 19 year olds who used contraception in the past was far higher than those using it **at the time they became pregnant.**




Among contraceptive users, very few were using **highly effective methods.***




*Percentages sum to more than 100% since multiple methods could be reported.

Knowledge of EC was low and past use of EC was non-existent.



Grace Chiudzu: chiudzug@gmail.com
 Ernestina Coast: e.coast@lse.ac.uk
 Tamara Feters: fetterst@ipas.org
 Luke Tembo: TemboL@ipas.org



Take a picture to find out more