


Contents lists available at [ScienceDirect](http://ScienceDirect)

# Clinical and Translational Radiation Oncology

journal homepage: [www.elsevier.com/locate/ctro](http://www.elsevier.com/locate/ctro)

## Original Research Article

# Magnetic Resonance-based Response Assessment and Dose Adaptation in Human Papilloma Virus Positive Tumors of the Oropharynx treated with Radiotherapy (MR-ADAPTOR): An R-IDEAL stage 2a-2b/Bayesian phase II trial



Houda Bahig<sup>a,b,\*</sup>, Ying Yuan<sup>a,2</sup>, Abdallah S.R. Mohamed<sup>a,1,2,3</sup>, Kristy K. Brock<sup>a,1</sup>, Sweet Ping Ng<sup>a,2,3</sup>, Jihong Wang<sup>a,1,2,3</sup>, Yao Ding<sup>a,2,3</sup>, Kate Hutcheson<sup>a,2,3</sup>, Molly McCulloch<sup>a,2</sup>, Peter A. Balter<sup>a,2</sup>, Stephen Y. Lai<sup>a,2,3</sup>, Abraham Al-Mamgani<sup>c</sup>, Jan-Jakob Sonke<sup>c,1,3</sup>, Uulke A. van der Heide<sup>c,1,3</sup>, Christopher Nutting<sup>d,1,3</sup>, X. Allen Li<sup>e,1</sup>, Jared Robbins<sup>e,1</sup>, Mussadiq Awan<sup>e,1,2,3</sup>, Irene Karam<sup>f,1</sup>, Katherine Newbold<sup>g,1,2</sup>, Kevin Harrington<sup>g,1,2</sup>, Uwe Oelfke<sup>g,1,3</sup>, Shreerang Bhide<sup>d,1</sup>, Marielle E.P. Philippens<sup>h,1</sup>, Chris H.J. Terhaard<sup>h,1</sup>, Andrew J. McPartlin<sup>i,1</sup>, Pierre Blanchard<sup>j,2,3</sup>, Adam S. Garden<sup>a,2,3</sup>, David I. Rosenthal<sup>a,2,3</sup>, Gary B. Gunn<sup>a,2,3</sup>, Jack Phan<sup>a,2,3</sup>, Guillaume Cazoulat<sup>a,2</sup>, Michalis Aristophanous<sup>a,1,2</sup>, Kelli K. McSpadden<sup>a,2</sup>, John A. Garcia<sup>a,2</sup>, Cornelis A.T. van den Berg<sup>h,1</sup>, Cornelis P.J. Raaijmakers<sup>h,1</sup>, Linda Kerkmeijer<sup>h,1</sup>, Patricia Doornaert<sup>k,1</sup>, Sanne Blinde<sup>l,1</sup>, Steven J. Frank<sup>a,1,2</sup>, Clifton D. Fuller<sup>a,1,2,3,\*</sup>

<sup>a</sup> University of Texas MD Anderson Cancer Center, Houston, TX, USA<sup>b</sup> Centre Hospitalier de l'Université de Montréal, Montreal, Quebec, Canada<sup>c</sup> Netherlands Cancer Institute, Amsterdam, Netherlands<sup>d</sup> Royal Marsden NHS Foundation Trust, London, UK<sup>e</sup> Medical College of Wisconsin, Milwaukee, WI, USA<sup>f</sup> Sunnybrook Health Sciences Centre, Toronto, Canada<sup>g</sup> Institute of Cancer Research, London, UK<sup>h</sup> University Medical Center Utrecht, Utrecht, Netherlands<sup>i</sup> The Christie NHS Foundation Trust, Manchester, UK<sup>j</sup> Gustave-Roussy, Villejuif, France<sup>k</sup> VU University Medical Center, de Boelelaan, Netherlands<sup>l</sup> Erasmus MC Cancer Institute, Rotterdam, Netherlands

## ARTICLE INFO

### Article history:

Received 24 May 2018

Revised 9 August 2018

Accepted 22 August 2018

Available online 24 August 2018

### Keywords:

Magnetic resonance imaging guided radiotherapy  
Human papilloma virus  
Oropharyngeal cancer  
Adaptive radiotherapy

## ABSTRACT

**Background:** Current standard radiotherapy for oropharynx cancer (OPC) is associated with high rates of severe toxicities, shown to adversely impact patients' quality of life. Given excellent outcomes of human papilloma virus (HPV)-associated OPC and long-term survival of these typically young patients, treatment de-intensification aimed at improving survivorship while maintaining excellent disease control is now a central concern. The recent implementation of magnetic resonance image – guided radiotherapy (MRgRT) systems allows for individual tumor response assessment during treatment and offers possibility of personalized dose-reduction. In this 2-stage Bayesian phase II study, we propose to examine weekly radiotherapy dose-adaptation based on magnetic resonance imaging (MRI) evaluated tumor response. Individual patient's plan will be designed to optimize dose reduction to organs at risk and minimize locoregional failure probability based on serial MRI during RT. Our primary aim is to assess the non-inferiority of MRgRT dose adaptation for patients with low risk HPV-associated OPC compared to historical control, as measured by Bayesian posterior probability of locoregional control (LRC).

**Methods:** Patients with T1-2 N0-2b (as per AJCC 7th Edition) HPV-positive OPC, with lymph node <3 cm and <10 pack-year smoking history planned for curative radiotherapy alone to a dose of 70 Gy in 33

\* Corresponding authors at: The MD Anderson Cancer Center, 1515 Holcombe Boulevard, Houston, TX 77030, USA.

E-mail addresses: [hbahig@mdanderson.org](mailto:hbahig@mdanderson.org) (H. Bahig), [cdfuller@mdanderson.org](mailto:cdfuller@mdanderson.org) (C.D. Fuller).<sup>1</sup> MR-Linac Consortium Head and Neck Tumor Site Group.<sup>2</sup> M.D. Anderson Cancer Center Head and Neck Quantitative Imaging Working Group.<sup>3</sup> Joint Head and Neck Radiotherapy-MRI Development Cooperative.<https://doi.org/10.1016/j.ctro.2018.08.003>

2405-6308/© 2018 The Authors. Published by Elsevier B.V. on behalf of European Society for Radiotherapy and Oncology.

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

fractions will be eligible. All patients will undergo pre-treatment MRI and at least weekly intra-treatment MRI. Patients undergoing MRgRT will have weekly adaptation of high dose planning target volume based on gross tumor volume response. The stage 1 of this study will enroll 15 patients to MRgRT dose adaptation. If LRC at 6 months with MRgRT dose adaptation is found sufficiently safe as per the Bayesian model, stage 2 of the protocol will expand enrollment to an additional 60 patients, randomized to either MRgRT or standard IMRT.

*Discussion:* Multiple methods for safe treatment de-escalation in patients with HPV-positive OPC are currently being studied. By leveraging the ability of advanced MRI techniques to visualize tumor and soft tissues through the course of treatment, this protocol proposes a workflow for safe personalized radiation dose-reduction in good responders with radiosensitive tumors, while ensuring tumoricidal dose to more radioresistant tumors. MRgRT dose adaptation could translate in reduced long term radiation toxicities and improved survivorship while maintaining excellent LRC outcomes in favorable OPC.

*Trial registration:* [ClinicalTrials.gov](http://ClinicalTrials.gov) ID: NCT03224000; Registration date: 07/21/2017.

© 2018 The Authors. Published by Elsevier B.V. on behalf of European Society for Radiotherapy and Oncology. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## 1. Background

### 1.1. Low risk HPV-Associated OPC

In the last decades, a change in the demographics of head and neck cancers has been observed, with an increasing incidence of human papillomavirus (HPV) – associated oropharyngeal cancers (OPC) reaching 60% of all OPC in the United States [1]. HPV-associated OPC is associated with markedly improved prognosis compared to non-HPV-associated OPC [2,3]. The favorable prognosis, along with the long-term survival and markedly younger age patients with HPV-positive disease, have supported that low-risk HPV-associated OPC may be suitable for treatment de-intensification [4]. Many treatment de-intensification strategies are currently being assessed in the context of clinical trials; these notably include use of minimally invasive surgery such as trans-oral robotic surgery [5], targeted therapies [6], and various radiation dose reduction strategies in the context of upfront radiotherapy [7] or after induction chemotherapy [8,9].

### 1.2. MRgRT dose adaptation

The principle of adaptive radiotherapy planning relies on adjusting the treatment plan based on observed changes over the course of therapy. While tumor responsiveness to radiotherapy in OPC has been shown to be associated with permanent tumor control outcomes [10], intra-treatment tumor shrinkage is observed as early as by fraction 11 [11] and complete response at mid-treatment is observed in as high as 50% of patients with HPV-associated OPC [12]. In a recent study, Lee et al. [7] reported excellent disease control outcomes of 10 Gy dose de-escalation to involved lymph nodes that presented early intra-treatment resolution of hypoxia on 18F-fluoromisonidazole - positron emission tomography (PET).

The increasing use of magnetic resonance imaging (MRI) for head and neck radiotherapy planning has the advantage of improved soft-tissue visualization [13], allowing for more confident assessment of anatomical tumor changes during treatment, at no ionizing radiation cost. The recent introduction of the MR-Linac technology, consisting in the combination of a linear accelerator and a 1.5 Tesla MRI, holds the promise to facilitate such adaptive MR-guided radiotherapy (MRgRT) workflows by mean of daily on-line MRI during radiation treatment [14,15]. In this study, we propose weekly RT dose-adaptation based on MRI-based tumor response. This adaptive radiation protocol will target a highly selected population of favorable risk HPV-associated OPC with a small (<5–7%) probability of locoregional failure with photon monotherapy. Our primary aim will be to assess the non-

inferiority of MRgRT dose adaptation for patients with low risk HPV-associated OPC compared to historical control, as measured by Bayesian posterior probability of locoregional control (LRC). By virtue of its assessment of clinical effectiveness and safety of the use of weekly MRI for dose adaptation, this study is a stage 2a/2b study as per the R-IDEAL framework for systematic technology assessment in radiotherapy [16] (Annex 1 – Supplementary material).

Supplementary data associated with this article can be found, in the online version, at <https://doi.org/10.1016/j.ctro.2018.08.003>.

## 2. Methods and design

### 2.1. Study design

This study uses a novel Phase II Bayesian two-stage adaptive design. The stage 1 of this study will enroll 15 patients to the experimental arm exclusively (MRgRT dose adaptation) to preliminarily assess the efficacy and safety of MRgRT. Bayesian decision rules will be used to make the go/no-go decision to move forward to stage 2, in which 60 patients will be randomized (1:1 ratio) into the MRgRT arm or the standard intensity modulated radiotherapy (IMRT) arm for comparison of LRC. Fig. 1 presents the study scheme. This study is approved by the MD Anderson Institutional Review Board and is registered on [clinicaltrials.gov](http://clinicaltrials.gov) (NCT03224000).


### 2.2. Primary objective

To assess non-inferiority of MRgRT dose adaptation and elective volume de-escalation in low risk HPV-associated OPC treated with radiotherapy alone.

- Primary endpoint:
  - o LRC at 6 months. LRC will be defined from time of treatment completion to disease progression at primary site or regional lymph nodes.

### 2.3. Secondary objectives

1. To compare rates of acute and late toxicities of MRgRT vs. standard IMRT.
2. To assess health – related quality of life outcomes in both treatment groups.
3. To assess disease-related outcomes (2-year progression-free survival, distant metastasis-free survival, and overall survival) in both treatment groups.
4. To validate functional imaging kinetics as a correlate of early treatment response.



**Fig. 1.** Scheme of the 2-stage trial design. ECOG = Eastern Cooperative Oncology Group; MRgRT = Magnetic-Resonance guided radiotherapy; IMRT = Intensity Modulated Radiotherapy.

#### 2.4. Conditions for patient eligibility

- Biopsy proven diagnosis of squamous cell carcinoma of the oropharynx (includes tonsil, soft palate, base of tongue, pharyngeal walls of oropharynx)
- Age  $\geq 18$  years
- Clinical stage T1-2, N0-1 (<3 cm), or small volume N2b (1–3 cm) M0, per AJCC 7th edition
- Positive for HPV by p16 immunohistochemistry or HPV in-situ hybridization
- Lifetime pack-year history of <10 years, and currently non-smoking
- No retropharyngeal nor level IV lymphadenopathy
- No head and neck surgery of the primary tumor or lymph nodes except for incisional or excisional biopsies
- Eastern Cooperative Oncology Group performance status 0–2
- Planned for single modality photon radiotherapy
- For females of child-bearing age, a negative pregnancy test

#### 2.5. Conditions for patient ineligibility

- Previous radiation treatment for head and neck mucosal primary cancers within the past 5 years
- Pregnant or breast-feeding females
- Contraindications to MRI

#### 2.6. Radiation therapy

Detailed description of target volumes and organs at risk (OAR) dose constraints are presented in Annex 2 (Supplementary material).

##### 2.6.1. Dose specification

- MRgRT arm

Patients will receive an individualized prescription of up to 69.96 Gy in 33 fractions with radiotherapy administered once daily, 5 days a week. Initial prescription will cover a high-risk clinical region (CTV<sub>6996</sub>) with 2.12 Gy/day. High-risk clinical region will include the gross tumor volume (GTV) plus an additional 5 mm margin. All uninvolved upper-neck elective nodal volumes outside of the CTV<sub>6996</sub> will be encompassed in an elective irradiation volume deemed CTV<sub>5016</sub>, and prescribed 1.52 Gy/day for a total prescription of 50.16 Gy in 33 fractions.

- Standard IMRT arm

Patients will receive 69.96 Gy in 33 fractions to the high-risk CTV<sub>6996</sub>, with radiotherapy administered once daily, 5 days a week. High-risk clinical region will include the GTV determined on MRI plus an additional 5 mm margin. All uninvolved upper-neck elective nodal volumes outside of the high-risk CTV will be encompassed in a volume deemed CTV<sub>5412</sub>, and prescribed 1.64 Gy/day for a total prescription of 54.12 Gy in 33 fractions.

##### 2.6.2. MRgRT arm: weekly adaptive workflow

Patients in the MRgRT arm will have pre-treatment and weekly CT simulation and MR-simulation at an interval of  $5 \pm 2$  fractions serially on treatment. MRI imaging data will be acquired for all patients in **stage 1** of the trial using a non-hybrid MRI device (MAGNETOM Aera 1.5 T MR scanner; Siemens Healthcare, Erlangen, Germany). In **stage 2**, MRI imaging data will be acquired using the integrated MR-Linac, which combines Philips 1.5 T MRI (Marlin, Finland) with 7 MV photon beam Elekta Linear accelerator. Anatomic T1-weighted and T2-weighted MRI sequences will be acquired in the axial plane with diffusion weighting gradients in three orthogonal directions using multiple b-values in the range of 0–1500 s/mm<sup>2</sup>. The radiation plan adaptation will be done once weekly and could be done online or offline. The diffusion weighted images will be obtained and analyzed retrospectively to correlate apparent diffusion coefficient (ADC) changes during treatment with final treatment response.

With the aim to integrate weekly replanning as the default workflow, weekly adaptation of the GTV (primary or nodal) will be performed when a minimally measurable shrinkage of  $\geq 2$  mm is observed. A new high-risk CTV<sub>6996</sub> will be generated to cover the weekly MRI-based GTV plus a margin of 5 mm. Any region previously involved by the tumor will remain in the CTV<sub>5016</sub> to receive the minimum daily dose of 1.52 Gy/day and ensure a minimum “floor” dose of 50.16 Gy.

On the standard IMRT arm, adaptation may be performed exceptionally in cases where volumetric verification imaging is concerning for inadequate coverage of the target volumes or dose deviation to OAR.

#### 2.7. Radiation planning and IGRT

Patients will have planning CT and MRI positioned in a stable supine position, using immobilization devices as previously described [17]. All patients will be planned with IMRT or

volumetric arc therapy. All treatment plans are to be normalized to provide at least 95% volume coverage of the PTV with the prescribed dose. No voxel within the PTV should receive more than 110% of the prescribed dose. Patients will have daily image guidance using in-room volumetric imaging prior to daily treatment.

### 2.8. Patients' evaluations

Pre-treatment, patients will be evaluated independently by a head and neck surgeon, a medical oncologist and a radiation oncologist. Annex 3 (Supplementary material) summarizes evaluations at different time points of the study.

### 2.9. Statistical considerations

A novel Bayesian two-stage adaptive design will be used for this phase II trial (see Fig. 1). Stage 1 involve only the experimental arm (MRgRT), and 15 patients will be enrolled to preliminarily assess the efficacy and safety of the experimental MRgRT. If MRgRT demonstrates promising, non-inferior LRC with a margin of 5%, compared to the historical LRC rate of 96%, as well as a rate of grade  $\geq 3$  toxicity at 6-months  $< 30\%$ , we will move forward to stage 2, in which 60 patients will be randomized (ratio 1:1) into the MRgRT arm or the standard IMRT arm. In stage 2, an interim analysis will be conducted to monitor the LRC and toxicity after 30 patients are randomized. Bayesian decision rules will be used to make the go/no-go decision for each stage (Annex 4, Supplementary material). The endpoint of LRC at 6 months was selected to allow for rapid Bayesian decision, and is based on evidence that the risk of locoregional recurrence after complete response within 12 weeks of treatment in HPV-associated OPC is associated with  $< 5\%$  locoregional recurrence rate at 5 years [18].

## 3. Discussion

In the context of the rampant incidence of HPV-associated OPC, there is crucial need to improve toxicity profiles in this population of often young, long-term survivors. The overall goal of treatment de-intensification strategies is to maintain the current very good disease control outcomes while reducing treatment-induced morbidity. To this end, it is essential that any dose adaptation strategy be associated with appropriate selection of good radiotherapy responders without compromising the chances of cure of poor responders. By leveraging the ability of advanced MRI techniques to visualize tumor and soft tissue characteristics through the course of treatment, it is possible to identify these subgroups of good responders and offer personalized dose-reduction. The introduction of the MR-Linac technology offers a unique opportunity to facilitate such adaptive MRgRT workflows by mean of daily on-line MRI during treatment [14]. The use of a common hybrid MRI device in the stage 2 of this study will allow the development of a high volume adaptive radiotherapy workflow using standardized imaging across multiple institutions, facilitated by the use of an integrated tool set. However, the results of our study could ultimately be scaled to any MRI device in radiotherapy.

While the use of adaptive MRI-based radiotherapy has the advantage of inherently selecting good responders, there remain concerns regarding the minimal radiobiological dose required at the edge of the shrinking GTV. In a recent study by Hamming-Vrieze et al. [20] the behaviour of tissue surrounding the GTV edge of locally advanced oropharyngeal tumors was assessed by placing fiducial markers at the tumor surface. The authors reported a larger GTV displacement on MRI compared to that assessed using fiducial markers, suggesting that part of the GTV was likely dissolving instead of shrinking and, raising concern of possible under-

dosage of microscopic disease with adaptive field reduction following GTV shrinkage assessed on MRI. In our study, this concern is addressed on one part by maintaining a 5 mm margin around the shrinking GTV for the high dose CTV, and on the other part by preserving an unchanged elective volume throughout the treatment, regardless of tumor response, to ensure a tumoricidal floor dose in all areas previously involved by tumor.

Recent report from an *in silico* study by our group [19] evaluating the dosimetric advantage of the proposed MRgRT dose adaptation approach showed a mean dose reduction to target volume of 11 Gy, and a significant dose reduction to swallowing musculature and thyroid gland which translated into a reduction of normal tissue complication probability of dysphagia  $\geq$  grade 2, feeding tube persistence at 6-month, and hypothyroidism at 1-year post-treatment [19]. As a second step, our proposed clinical study will provide essential information on the clinical feasibility, safety and clinical toxicity reduction of MRgRT dose adaptation in well-selected patients. We propose a conservative approach to this MRgRT dose de-escalation model, whereby only patients with highly favorable, low stage and low volume burden HPV-associated OPC will be eligible to enroll in this study. Should the study demonstrate positive results, transposition of this adaptive model to more advanced HPV-associated OPC would be subject of additional investigation.

## 4. Co-author specific contributions

- HB, ASRM, SP- Manuscript drafting, editing and review of comments, clinical trial activation.
- YY- Statistical analysis.
- AA, CN, JH, JR, MA, IK, KN, KH, SB, AJM, PB, ASG, JP, JPPB, PD, LK, SB, SJF, CHJT: Clinical protocol development.
- KKB, PAB, MM, GC, MA, KKS, JG: Image segmentation, adaptive workflow development, cumulative dose calculation.
- JW, YD, JJS, UAH, XAL, UO, MEPP, CATB, CPJR: MRI sequence development, optimization and standardization.
- SL, KH, GBC, DIR: Toxicity assessment planning and functional outcome evaluation management.
- CDF: corresponding author; primary investigator; conceived, coordinated, and directed all study activities, responsible for data collection, project integrity, manuscript content and editorial oversight and correspondence; direct oversight of trainee personnel (HB, ASRM, SP, MA, YD).

## 5. Funding

This research is supported by the Andrew Sabin Family Foundation; Dr. Fuller is a Sabin Family Foundation Fellow. Dr. Fuller receives funding and salary support from the National Institutes of Health (NIH), including: the National Institute for Dental and Craniofacial Research Award (1R01DE025248-01/R56DE025248-01); a National Science Foundation (NSF), Division of Mathematical Sciences, Joint NIH/NSF Initiative on Quantitative Approaches to Biomedical Big Data (QuBBDD) Grant (NSF 1557679); the NIH Big Data to Knowledge (BD2K) Program of the National Cancer Institute (NCI) Early Stage Development of Technologies in Biomedical Computing, Informatics, and Big Data Science Award (1R01CA214825-01); NCI Early Phase Clinical Trials in Imaging and Image-Guided Interventions Program (1R01CA218148-01); an NIH/NCI Cancer Center Support Grant (CCSG) Pilot Research Program Award from the UT MD Anderson CCSG Radiation Oncology and Cancer Imaging Program (P30CA016672) and an NIH/NCI Head and Neck Specialized Programs of Research Excellence (SPORE) Developmental Research Program Award (P50

CA097007-10). Dr. Fuller has received direct industry grant support and travel funding from Elekta AB.

## References

- [1] Marur S, D'Souza G, Westra WH, Forastiere AA. HPV-associated head and neck cancer: a virus-related cancer epidemic. *Lancet Oncol* 2010;11(8):781–9.
- [2] Ang KK, Sturgis EM. Human papillomavirus as a marker of the natural history and response to therapy of head and neck squamous cell carcinoma. *Semin Radiat Oncol* 2012;22(2):128–42.
- [3] Garden AS, Kies MS, Morrison WH, Weber RS, Frank SJ, Glisson BS, et al. Outcomes and patterns of care of patients with locally advanced oropharyngeal carcinoma treated in the early 21st century. *Radiat Oncol* 2013;8:21.
- [4] Kofler B, Laban S, Busch CJ, Lorincz B, Knecht R. New treatment strategies for HPV-positive head and neck cancer. *European archives of oto-rhino-laryngology: official journal of the European Federation of Oto-Rhino-Laryngological Societies (EUFOS): affiliated with the German Society for Oto-Rhino-Laryngology – Head and Neck Surgery* 2014;271(7):1861–7.
- [5] Masterson L, Moualed D, Liu ZW, Howard JE, Dwivedi RC, Tysome JR, et al. De-escalation treatment protocols for human papillomavirus-associated oropharyngeal squamous cell carcinoma: a systematic review and meta-analysis of current clinical trials. *Eur J Cancer (Oxford, England: 1990)* 2014;50(15):2636–48.
- [6] Hay A, Ganly I. Targeted therapy in oropharyngeal squamous cell carcinoma: the implications of HPV for therapy. *Rare Cancers Ther* 2015;3:89–117.
- [7] Lee N, Schoder H, Beattie B, Lanning R, Riaz N, McBride S, et al. Strategy of using intratreatment hypoxia imaging to selectively and safely guide radiation dose de-escalation concurrent with chemotherapy for locoregionally advanced human papillomavirus-related oropharyngeal carcinoma. *Int J Radiat Oncol Biol Phys* 2016;96(1):9–17.
- [8] Marur S, Li S, Cmelak AJ, Gillison ML, Zhao WJ, Ferris RL, et al. E1308: phase II trial of induction chemotherapy followed by reduced-dose radiation and weekly cetuximab in patients with HPV-associated resectable squamous cell carcinoma of the oropharynx-ECOG-ACRIN cancer research group. *J Clin Oncol: Official J Am Soc Clin Oncol* 2017;35(5):490–7.
- [9] Chen AM, Felix C, Wang PC, Hsu S, Basehart V, Garst J, et al. Reduced-dose radiotherapy for human papillomavirus-associated squamous-cell carcinoma of the oropharynx: a single-arm, phase 2 study. *Lancet Oncol* 2017;18(6):803–11.
- [10] Bataini JP, Jaulerry C, Brunin F, Ponvert D, Ghossein NA. Significance and therapeutic implications of tumor regression following radiotherapy in patients treated for squamous cell carcinoma of the oropharynx and pharyngolarynx. *Head Neck* 1990;12(1):41–9.
- [11] Subesinghe M, Scarsbrook AF, Sourbron S, Wilson DJ, McDermott G, Speight R, et al. Alterations in anatomic and functional imaging parameters with repeated FDG PET-CT and MRI during radiotherapy for head and neck cancer: a pilot study. *BMC Cancer* 2015;15:137.
- [12] Ding Y, Hazle JD, Mohamed AS, Frank SJ, Hobbs BP, Colen RR, et al. Intravoxel incoherent motion imaging kinetics during chemoradiotherapy for human papillomavirus-associated squamous cell carcinoma of the oropharynx: preliminary results from a prospective pilot study. *NMR Biomed* 2015;28(12):1645–54.
- [13] Wippold 2nd FJ. Head and neck imaging: the role of CT and MRI. *J Magn Reson Imaging* 2007;25(3):453–65.
- [14] Kontaxis C, Bol GH, Lagendijk JJ, Raaymakers BW. Towards adaptive IMRT sequencing for the MR-linac. *Phys Med Biol* 2015;60(6):2493–509.
- [15] Raaymakers BW, Lagendijk JJ, Overweg J, Kok JG, Raaijmakers AJ, Kerkhof EM, et al. Integrating a 1.5 T MRI scanner with a 6 MV accelerator: proof of concept. *Phys Med Biol* 2009;54(12):N229–37.
- [16] Verkooijen HM, Kerkmeijer LGW, Fuller CD, Huddart R, Faivre-Finn C, Verheij M, et al. R-IDEAL: a framework for systematic clinical evaluation of technical innovations in radiation oncology. *Front Oncol* 2017;7:59.
- [17] Ding Y, Mohamed ASR, Yang J, Colen RR, Frank SJ, Wang J, et al. Prospective observer and software-based assessment of magnetic resonance imaging quality in head and neck cancer: should standard positioning and immobilization be required for radiation therapy applications? *Pract Radiat Oncol* 2015;5(4):e299–308.
- [18] Ng SP, Johnson JM, Gunn GB, Rosenthal DI, Skinner HD, Phan J, et al. Significance of negative post-treatment 18-FDG PET/CT imaging in patients with p16/HPV-positive oropharyngeal cancer. *Int J Radiat Oncol Biol Phys* 2018.
- [19] Mohamed ASR, Aristophanous M, Blanchard P, Kamal M, Ding T, Cardenas C, et al. Prospective in silico study of the feasibility and dosimetric advantages of MRI-guided dose adaptation for human papillomavirus positive oropharyngeal cancer patients compared with standard IMRT. *Clin Transl Radiat Oncol* 2018 (accepted for publication).
- [20] Hamming-Vrieze O, van Kranen SR, Heemsbergen WD, Lange CAH, van den Brekel MWM, Verheij M, Rasch CRN, Sonke JJ. Analysis of GTV reduction during radiotherapy for oropharyngeal cancer: Implications for adaptive radiotherapy. *Radiother Oncol.* 2017 Feb;122(2):224–8. <https://doi.org/10.1016/j.radonc.2016.10.012>. Epub 2016 Nov 18.