



Huang, J. Y., Larose, T. L., Luu, H. N., Wang, R., Fanidi, A., Alcala, K., Stevens, V. L., Weinstein, S. J., Albanes, D., Caporaso, N. E., Purdue, M. P., Ziegler, R. G., Freedman, N. D., Lan, Q., Prentice, R. L., Pettinger, M., Thomson, C. A., Cai, Q., Wu, J., ... Yuan, J-M. (2019). Circulating markers of cellular immune activation in prediagnostic blood sample and lung cancer risk in the Lung Cancer Cohort Consortium (LC3). *International Journal of Cancer*.  
<https://doi.org/10.1002/ijc.32555>

Peer reviewed version

Link to published version (if available):  
[10.1002/ijc.32555](https://doi.org/10.1002/ijc.32555)

[Link to publication record in Explore Bristol Research](#)  
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via Wiley at <https://onlinelibrary.wiley.com/doi/full/10.1002/ijc.32555> . Please refer to any applicable terms of use of the publisher.

## University of Bristol - Explore Bristol Research

### General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:  
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

**Circulating markers of cellular immune activation in pre-diagnostic blood sample and lung cancer risk in the Lung Cancer Cohort Consortium (LC3)**

Joyce Yongxu Huang<sup>1,2\*</sup>, Tricia L. Larose<sup>3,4\*</sup>, Hung N Luu<sup>1,2</sup>, Renwei Wang<sup>1</sup>, Anouar Fanidi<sup>3,5</sup>, Karine Alcalá<sup>3</sup>, Victoria L. Stevens<sup>6</sup>, Stephanie J. Weinstein<sup>7</sup>, Demetrius Albanes<sup>7</sup>, Neil E. Caporaso<sup>7</sup>, Mark P. Purdue<sup>7</sup>, Regina G. Ziegler<sup>7</sup>, Neal D. Freedman<sup>7</sup>, Qing Lan<sup>7</sup>, Ross L. Prentice<sup>8</sup>, Mary Pettinger<sup>8</sup>, Cynthia A. Thomson<sup>9</sup>, Qiuyin Cai<sup>10</sup>, Jie Wu<sup>10</sup>, William J. Blot<sup>10</sup>, Xiao-Ou Shu<sup>10</sup>, Wei Zheng<sup>10</sup>, Alan A. Arslan<sup>11</sup>, Anne Zeleniuch-Jacquotte<sup>12</sup>, Loïc Le Marchand<sup>13</sup>, Lynn R. Wilkens<sup>13</sup>, Christopher A. Haiman<sup>14</sup>, Xuehong Zhang<sup>15</sup>, Meir J. Stampfer<sup>15,16,17</sup>, Jiali Han<sup>18</sup>, Graham G. Giles<sup>19,20</sup>, Allison M. Hodge<sup>19,20</sup>, Gianluca Severi<sup>19,21,22</sup>, Mikael Johansson<sup>23</sup>, Kjell Grankvist<sup>24</sup>, Arnulf Langhammer<sup>25</sup>, Kristian Hveem<sup>4,25</sup>, Yong-Bing Xiang<sup>26,27</sup>, Hong-Lan Li<sup>27</sup>, Yu-Tang Gao<sup>27</sup>, Kala Visvanathan<sup>28</sup>, Per M Ueland<sup>29,30</sup>, Øivind Midttun<sup>31</sup>, Arve Ulvi<sup>31</sup>, Julie E. Buring<sup>32,33</sup>, I-Min Lee<sup>32,33</sup>, Howard D. Sesso<sup>16,32,33</sup>, J. Michael Gaziano<sup>16,33,34</sup>, Jonas Manjer<sup>35</sup>, Caroline Relton<sup>36</sup>, Woon-Puay Koh<sup>37,38</sup>, Paul Brennan<sup>3\*\*</sup>, Mattias Johansson<sup>3\*\*</sup>, Jian-Min Yuan<sup>1,2\*\*</sup>

\*Co-first authors

\*\*These authors jointly supervised this work

**Corresponding authors:**

Jian-Min Yuan

University of Pittsburgh

UPMC Hillman Cancer Center

UPMC Cancer Pavilion, Suite 4C

5150 Centre Avenue

Pittsburgh, PA 15232

USA

Phone: 412-864-7889

Fax: 412-623-3303

Email: [yuanj@upmc.edu](mailto:yuanj@upmc.edu)

AND

Mattias Johansson

International Agency for Research on Cancer (IARC/WHO)

150 cours Albert Thomas<sup>[1]</sup><sub>SEP</sub>

69372 Lyon cedex 08, France<sup>[1]</sup><sub>SEP</sub>

Phone: +33 (0) 4 72 73 80 23

Fax: +33 (0) 4 72 73 83 42

[Email: JohanssonM@iarc.fr](mailto:JohanssonM@iarc.fr)

**Conflict of Interest:** None

**Affiliations:**

1. Division of Cancer Control and Population Sciences, UPMC Hillman Cancer Center, University of Pittsburgh, Pittsburgh, Pennsylvania, USA.
2. Department of Epidemiology, Graduate School of Public Health, University of Pittsburgh, Pittsburgh, Pennsylvania, USA
3. Genetic Epidemiology Group, International Agency for Research on Cancer, Lyon, France
4. K.G. Jebsen Center for Genetic Epidemiology, Department of Public Health & Nursing, Faculty of Medicine and Health Sciences, Norwegian University of Science and Technology, Trondheim, Norway
5. MRC Epidemiology Unit, University of Cambridge School of Clinical Medicine, Cambridge, United Kingdom
6. Behavior and Epidemiology Research Group, American Cancer Society, Inc. 250 Williams St. Atlanta, GA 30303
7. Division of Cancer Epidemiology and Genetics, National Cancer Institute, NIH, Rockville, MD, USA
8. Division of Public Health Sciences Fred Hutchinson Cancer Research Center 1100 Fairview Ave. N, Seattle, Washington 98109, U.S.A.

9. Department of Health Promotion Science, Mel & Enid Zuckerman College of Public Health, University of Arizona, Tucson, AZ, USA
10. Division of Epidemiology, Department of Medicine, Vanderbilt Epidemiology Center and Vanderbilt-Ingram Cancer Center, Vanderbilt University School of Medicine, Nashville, TN, USA
11. Departments of Obstetrics and Gynecology, Population Health, Environmental Medicine and Perlmutter Cancer Center, New York University School of Medicine, New York, NY,
12. Departments of Population Health and Environmental Medicine and Perlmutter Cancer Centre, New York University School of Medicine, New York, NY, USA
13. Epidemiology Program, University of Hawaii Cancer Center, Honolulu, HI, USA
14. Department of Prevention, Keck School of Medicine, University of Southern California, Los Angeles, CA, USA
15. Channing Division of Network Medicine, Department of Medicine, Brigham and Women's Hospital and Harvard Medical School, Boston, MA.
16. Department of Epidemiology, Harvard T. H. Chan School of Public Health, Boston, MA, USA
17. Department of Nutrition, Harvard T.H. Chan School of Public Health, Boston, MA, USA.
18. Department of Epidemiology, Richard M. Fairbanks School of Public Health, Indiana University, Indianapolis, IN, USA
19. Cancer Epidemiology and Division, Cancer Council Victoria, Melbourne, Australia.
20. Centre for Epidemiology and Biostatistics, Melbourne School of Population and Global Health, The University of Melbourne, Melbourne, Australia
21. Italian Institute for Genomic Medicine (IIGM), Torino, Italy.
22. Centre de Recherche en Epidemiologie et Santé des Populations (CESP) UMR1018 Inserm, Facultés de Médecine Université Paris-Saclay, UPS, UVSQ, Gustave Roussy, 94805, Villejuif, France.

23. Department of Radiation Sciences, Oncology, Umeå University, Umeå, Sweden
24. Department of Medical Biosciences, Clinical Chemistry, Umeå University, Umeå, Sweden
25. HUNT Research Centre, Department of Public Health and Nursing, Norwegian University of Science and Technology, Levanger, Norway
26. State Key Laboratory of Oncogene and Related Genes, Shanghai Cancer Institute, Renji Hospital, Shanghai Jiaotong University School of Medicine, Shanghai, China.
27. Department of Epidemiology, Shanghai Cancer Institute, Renji Hospital, Shanghai Jiaotong University School of Medicine, Shanghai, China.
28. George W Comstock Center for Public Health Research and Prevention Health Monitoring Unit, Department of Epidemiology, Johns Hopkins University Bloomberg School of Public Health, USA
29. Department of Clinical Science, University of Bergen, Bergen, Norway
30. Laboratory of Clinical Biochemistry, Haukeland University Hospital, Bergen, Norway
31. Bevital AS, 5021 Bergen, Norway
32. Division of Preventive Medicine, Department of Medicine, Brigham and Women's Hospital and Harvard Medical School, Boston, MA, USA.
33. Division of Aging, Department of Medicine, Brigham and Women's Hospital and Harvard Medical School, Boston, MA, USA.
34. Boston VA Medical Center, Boston, MA USA
35. Department of Surgery, Skåne University Hospital Malmö Lund University, Malmö Sweden
36. MRC Integrative Epidemiology Unit, Population Health Sciences, Bristol Medical School, University of Bristol, Bristol, United Kingdom
37. Health Services and Systems Research, Duke-NUS Medical School, Singapore
38. Saw Swee Hock School of Public Health, National University of Singapore, Singapore

**Key words:** lung cancer, kynurenine, tryptophan, neopterin, quinolinic acid

**Abbreviations:** ANCOVA, Analysis of Covariance; BMI, body mass index; CI, confidence interval; eGFR, estimated glomerular filtration rate; IDO, Indoleamine 2,3-dioxygenase; IFN-gamma, interferon-gamma; KTR, kynurenine to tryptophan ratio; LC3, Lung Cancer Cohort Consortium; OR, odds ratio; QA, quinolinic acid.

**Article Category:** Original Research Article

## ABSTRACT

Cell-mediated immune suppression may play an important role in lung carcinogenesis. We investigated the associations for circulating levels of tryptophan, kynurenine, kynurenine:tryptophan ratio (KTR), quinolinic acid (QA), and neopterin as markers of immune regulation and inflammation with lung cancer risk in 5,364 smoking-matched case-control pairs from 20 prospective cohorts included in the international Lung Cancer Cohort Consortium. All biomarkers were quantified by mass spectrometry-based methods in serum/plasma samples collected on average 6 years before lung cancer diagnosis. Odds ratios (ORs) and 95% confidence intervals (CIs) for lung cancer associated with individual biomarkers were calculated using conditional logistic regression with adjustment for circulating cotinine. Compared with the lowest quintile, the highest quintiles of kynurenine, KTR, QA and neopterin were associated with a 20-30% higher risk, and tryptophan with a 15% lower risk of lung cancer (all  $P_{\text{trend}} < 0.05$ ). The strongest associations were seen for current smokers, where the adjusted ORs (95% CIs) of lung cancer for the highest quintile of KTR, QA and neopterin were 1.42 (1.15–1.75), 1.42 (1.14–1.76) and 1.45 (1.13–1.86), respectively. A stronger association was also seen for KTR and QA with risk of lung squamous cell carcinoma followed by adenocarcinoma, and for lung cancer diagnosed within the first 2 years after blood draw. This study demonstrated that components of the tryptophan-kynurenine pathway with immunomodulatory effects are associated with risk of lung cancer overall, especially for current smokers. Further research is needed to evaluate the role of these biomarkers in lung carcinogenesis and progression.

## **NOVELTY AND IMPACT**

Immune suppression may play an important role in the development and progression of lung cancer.

Authors analyzed circulating levels of tryptophan metabolism pathway including tryptophan, kynurenine, kynurenine:tryptophan ratio, and quinolinic acid as markers of immune regulation and inflammation and found they were associated with increased risk of developing lung cancer.



## INTRODUCTION

Lung cancer is one of the most common cancers accounting for 2.09 million incident cases and 1.76 million deaths worldwide in 2018 <sup>1</sup>. The 5-year survival for lung cancer cases is only 17.7% in the United States (US) <sup>2</sup>, and is even lower globally <sup>3</sup>. This underscores the importance of improving prevention and treatment to reduce lung cancer morbidity and mortality. Whilst the role of the immune system in the development of lung cancer has been increasingly recognized, the mechanisms by which immune mediators influence risk are only partly understood <sup>4,5</sup>.

Previous epidemiological studies that focused on the associations between circulating cytokines and risk of lung cancer have provided inconsistent results. For example, interleukin-6 and interleukin-8 were associated with increased risk of lung cancer in two prospective studies in the US <sup>6</sup> and Europe <sup>7</sup>, but these same markers were not associated with lung cancer risk in a second US cohort <sup>8</sup> that evaluated the associations between 77 inflammatory markers and lung cancer risk, perhaps due to low statistical power. Also, these previous studies were not well powered to study risk in important sub-groups, such as never smokers. In addition, concentrations of cytokines are generally low in the circulation of healthy individuals who have no active infection or malignancy <sup>9</sup>. Thus, investigations of alternative biomarkers for immune regulation and response with the risk of lung cancer are warranted.

Among the pathways involved in cancer innate and adaptive immune tolerance, the catabolism of tryptophan has increasingly been recognized as playing a fundamental role <sup>10</sup>. Interferon gamma (IFN- $\gamma$ )-inducible indoleamine 2,3-dioxygenase (IDO) catalyzes the first rate-limiting reaction that converts tryptophan to kynurenine, which in turn leads to the depletion of local tryptophan and accumulation of kynurenine and their derivatives (**Figure 1**). This results in a highly tolerogenic microenvironment characterized by reduced T effector lymphocytes and natural killer cells and an increased number of functionally active T regulatory cells and myeloid-derived suppressor cells <sup>11</sup>. The ratio between circulating kynurenine and tryptophan (KTR) can therefore be used as a surrogate of IDO activity and IFN- $\gamma$ -mediated immune regulation in tumor microenvironment <sup>12</sup>. IFN-gamma also stimulates the production of neopterin,

a metabolite of guanosine triphosphate, by macrophages<sup>13</sup>. One previous epidemiological study observed an association between KTR and higher risk of lung cancer<sup>14</sup>. A second prospective study showed associations between KTR or neopterin and risk of cancer overall, but no association was observed for risk of lung cancer specifically, possibly due to lack of statistical power<sup>15</sup>. In addition, the downstream metabolites of the kynurenine pathway such as quinolinic acid (QA) have immuno-regulatory effects<sup>16, 17</sup> and may contribute to the development and progression of lung cancer, but has not been investigated in large epidemiological studies.

The purpose of the current study conducted using 20 prospective cohorts from Asia, Australia, Europe and the US was to comprehensively investigate the associations for circulating concentrations of the tryptophan-kynurenine pathway metabolites and neopterin as markers of IFN- $\gamma$ -induced immune regulation with the risk of developing lung cancer. Our large sample size (5364 case-control pairs) allowed us to further investigate these associations by smoking status, histology, and time from blood draw to diagnosis.

## **MATERIALS AND METHODS**

### **Study population**

The design of the Lung Cancer Cohort Consortium (LC3) including cohort design and follow-up procedures has been reported previously<sup>18</sup>. The current investigation included case-control studies of incident lung cancer cases and individually matched controls nested within 20 prospective cohorts from the US, Europe, Australia, and Asia. At recruitment into each cohort, participants signed informed consent forms, completed questionnaires, had blood sample drawn and anthropometric measurements taken. The LC3 was approved by the Institutional Review Board of each contributing cohort and those of participating registries as required.

### **Selection of cases and controls**

Lung cancer cases were defined according to the International Classification of Diseases for Oncology, Second Edition (ICD-O-2), and included all invasive cancers coded as C34.0 to C34.9. Altogether, 11,399 incident lung cancer cases with pre-diagnostic serum or plasma samples in the members of the US National Cancer Institute Cohort Consortium in 2009 were eligible. From these, the LC3 selected a total of 5,545 lung cancer cases, and to optimize the statistical power in smoking stratified analyses, never and former smoking cases were oversampled. For each case, one control was randomly selected from all eligible participants within the same cohort who were alive and free of cancer (except non-melanoma skin cancer) at the same length of time from enrollment as was the index case at diagnosis. Matching criteria were race (US only), sex, date of blood collection ( $\pm 1$  month, relaxed to  $\pm 3$  months for sets without available controls), and date of birth ( $\pm 1$  year, relaxed to  $\pm 3$  years), as well as smoking status in 5 categories: never smokers, former smokers who had quit smoking for  $<10$  or  $\geq 10$  years, and current smokers who smoked  $<15$  or  $\geq 15$  cigarettes per day. After excluding cases who were not able to be correctly matched on smoking status in 5 categories defined above ( $n=126$  cases), had insufficient serum/plasma samples ( $n=42$ ), or had a revised date of diagnosis prior to blood draw ( $n=13$ ), a total of 5364 lung cancer case-control pairs remained eligible for the current analysis.

### **Biochemical analyses**

Serum or plasma samples from all LC3 study participants were sent on dry ice to the Bevital A/S laboratory (<http://www.bevital.no>) in Bergen, Norway, and were kept at  $-80^{\circ}\text{C}$  until analysis. Concentrations of tryptophan, kynurenine<sup>19</sup>, quinolinic acid (QA), neopterin and cotinine<sup>20</sup> – a biomarker of recent tobacco exposure were determined by mass spectrometry based methods (LC-MS/MS, GC-MS/MS). Biochemical analysis was performed in 96-well plates, each containing 86 study samples, 6 calibration samples, 3 quality control samples, and 1 blank sample. Samples from the index case and the matched control subjects were put next to each other in a random order and always analyzed together in the same batch. The laboratory personnel were blind to the case/control status of the test samples. Between-batch coefficient of variation (CV) of quality-control samples for the five analyzed biomarkers were less than 6%<sup>21</sup>. Our previous studies also showed that tryptophan and kynurenine were stable between different

types of blood tube, between serum and plasma, and over different processing lag time, and had high within-person reproducibility<sup>21, 22</sup>.

### **Data availability**

All data relevant to the present study are available upon request to the corresponding authors.

### **Statistical Analysis**

The KTR ratio was calculated as the kynurenine concentration (nmol/L) divided by the tryptophan concentration ( $\mu\text{mol/L}$ ). We logarithmically transformed (base e) original values of all biomarker concentrations and KTR to normalize their skewed distributions. The pair-wise correlations between biomarkers were assessed using Spearman correlation coefficients. The difference in geometric means of biomarkers among three smoking groups (never, former and current smokers) was assessed using Analysis of Covariance (ANCOVA) in all control subjects with adjustment for cohort, age, sex, and estimated glomerular filtration rate (eGFR; a measurement of kidney function that influences the circulating levels of kynurenine and its metabolites). The eGFR was calculated based on participant's age, gender, and creatinine concentration in plasma or serum according to the previously published method<sup>23</sup>.

Study participants were divided into quintiles based on the distributions of biomarker concentrations among controls within a specific cohort. Odds ratios (ORs) of lung cancer for quintiles of biomarker concentrations were calculated relative to the first quintile using conditional logistic regression<sup>24</sup>. Ordinal values (e.g., 1, 2, 3, 4, and 5) for individual biomarkers were used for testing linear trends across quintiles in the biomarker-lung cancer risk associations.

In addition to matching on cohort, race (US only), sex, date of blood draw, date of birth, and the combination of smoking status with years of quitting (for former smokers) and number of cigarettes per day (for current smokers), the multivariable conditional logistic regression models included the following reported risk factors for lung cancer and determinants of kynurenine metabolites as potential confounders: cotinine concentration (continuous, a biomarker of recent nicotine intake)<sup>25</sup>, educational attainment (six categories), body mass index (BMI) in  $\text{kg/m}^2$  ( $<18.5$ ,  $18.5- <25$ ,  $25- <30$ ,  $\geq 30$ ), and eGFR.

Fully adjusted regression models were used in analyses stratified by smoking status (current, former, never smokers), histological subtypes of lung cancer (adenocarcinoma, large-cell carcinoma, small-cell carcinoma, and squamous cell carcinoma), time between blood draw and lung cancer diagnosis (<2, 2–<5, and ≥5 years), and geographical region (US, Europe/Australia, Asia). Potential effect modification of associations between biomarkers and lung cancer risk by demographic, lifestyle, or other factors were examined by including their product term in the multivariate regression models.

Statistical analyses were carried out using SAS software version 9.3 (SAS Institute, Cary, NC). All *P* values reported are two-sided, and those that were <0.05 were considered to be statistically significant.

## **RESULTS**

### **Baseline characteristics of cases and controls**

The current study sample included 5,364 incident lung cancer cases and 5,364 individually matched controls (**Table 1**). Overall, slightly more participants were male (54.2%). Participants from Europe/Australia (EU/AU) and Asia were also predominantly male (57.9% and 69.2%, respectively) whereas participants from the US were predominantly female (58.7%). Current smokers accounted for nearly half the overall study participants (47%, 2,519 case-control pairs), with former (28.3%, 1,518 case-control pairs) and never smokers (24.7%, 1,327 case-control pairs) contributing approximately one-quarter each. Cases and controls had, on average, similar characteristics including BMI and age at recruitment (60 years).

Median age at lung cancer diagnosis was 69.8 (range 53.6 – 82.0) with little variation across geographic regions. The median time between blood draw and lung cancer diagnosis was 5.2 years for the US, 5.8 years for Asian, and 10 years for cohorts in Australia and Europe. Histologically, the majority of lung cancer cases were adenocarcinoma, followed by squamous cell, small cell and large cell carcinoma. Due to a larger overall sample size, the US cohorts contributed the majority of all adenocarcinoma cases

(50.3%), small cell carcinoma cases (49.8%), and large cell carcinoma cases (64.4%). The proportion of squamous cell cases did not differ substantially by region, with each region contributing approximately one-third of cases.

### **Biomarker distribution in study population**

The geometric means of kynurenine, KTR, QA and neopterin were significantly higher in former smokers than never or current smokers, but no difference between current and never smokers among all control subjects after adjustment for age, sex, eGFR, and cohort (**Supplementary Table 1**). Never smokers had significantly higher concentrations of QA than current smokers (368.7 *versus* 334.7  $\mu\text{mol/L}$ ,  $P < 0.001$ ) and higher neopterin (10.8 *versus* 10.3  $\mu\text{mol/L}$ ,  $P = 0.002$ ). There was no difference in tryptophan concentrations among never, former, and current smokers. Each of five analyzed biomarkers were significantly elevated in overweight and obese control subjects and the associations between these biomarkers and BMI were dose-dependent (**Supplementary Table 2**). Median concentrations of circulating biomarkers did not differ substantially across cohorts within geographic region, with few exceptions. For US cohorts, circulating tryptophan concentrations were 20  $\mu\text{mol/L}$  higher in the American Cancer Society Cancer Prevention Study-II (CPS-II) Nutrition cohort compared to the Women's Health Initiative (WHI) cohort (**Supplementary Table 3**). In addition, circulating neopterin concentrations were different among different cohorts within a region; the highest levels were observed in the Multiethnic Cohort (MEC) among the US cohorts and in the Singapore Chinese Health Study among Asian cohorts. Overall kynurenine, KTR, QA, and neopterin concentrations were positively correlated with each other after adjustment for age and sex (partial Spearman correlation coefficient [ $r$ ] = 0.34–0.66) whereas tryptophan was positively correlated with kynurenine ( $r = 0.45$ ) and QA ( $r = 0.13$ ), but inversely correlated with KTR ( $r = -0.43$ ) and was not correlated with neopterin ( $r = -0.01$ ) (**Supplementary Table 4**).

### **Overall and stratified associations of circulating biomarkers and lung cancer risk**

Odds ratios for quintiles of each biomarker with overall lung cancer after controlling for smoking status, duration and intensity, circulating levels of cotinine, and other potential confounders, are shown in **Table 2**. The OR for the top vs. bottom quartiles was 0.85 (0.75-0.96) for tryptophan, 1.22 (1.06-1.40) for kynurenine, 1.31 (1.14-1.50) for KTR, 1.31 (1.14-1.51) for quinolinic acid as well as for neopterin.

**Table 3** shows the odds ratios for lung cancer associated with higher quintiles of biomarkers in current, former, and never smokers separately (see numbers of cases and controls in **Supplementary Table 5**). For current smokers, ORs (95% CIs) for lung cancer for the highest quintiles of KTR, QA and neopterin were 1.42 (1.15–1.75), 1.42 (1.14–1.75), and 1.45 (1.13–1.86), respectively (all  $P_{\text{trend}} \leq 0.005$ ). The corresponding ORs (95% CIs) for former smokers were 1.32 (1.00–1.74), 1.20 (0.90–1.59), and 1.43 (0.97–1.86) (all  $P_{\text{trend}}$  were borderline significant). There was no association between these biomarkers and lung cancer risk for never smokers (all  $P_{\text{trend}} > 0.16$ ). However, no interaction between any biomarker and smoking status for lung cancer risk was detected (all  $P$ 's for multiplicative interaction  $> 0.05$ ).

When data were analyzed by histological subtype of lung cancer (**Supplementary Table 5**), positive associations were observed for KTR and QA and risk of lung squamous cell carcinoma, and for QA and risk of lung adenocarcinoma (**Table 4**). The associations for other biomarkers with risk of adenocarcinoma or squamous cell carcinoma, and for all biomarkers with large cell and small cell carcinomas were not statistically significant.

In the sensitivity analysis, the associations for blood concentrations of kynurenine, KTR and QA were observed for the risk of lung cancer diagnosed within 2 years after blood draw (**Table 5** and **Supplementary Table 5**). Higher levels of neopterin were associated with higher risk of lung cancer diagnosed within 2 to <5 years after blood draw. The association between QA and lung cancer risk remained, albeit weakened, even 5 or more years after blood collection.

## DISCUSSION

### Principal findings

In the largest prospective epidemiological study, we demonstrated the associations for lower levels of tryptophan and higher levels of kynurenine and its metabolites as well as neopterin with risk of developing lung cancer overall. These associations were strongest among current smokers, to a lesser extent, among former smokers, and null among never smokers. These positive associations were strongest for lung squamous cell cancer, and for lung cancer cases diagnosed within two years of blood draw.

### **Higher circulating KTR concentrations and risk of lung cancer**

Tryptophan is an essential amino acid for immune cell proliferation. Early studies suggested that immune suppressive effect of tryptophan catabolism on T cell is a consequence of decreased concentration of tryptophan<sup>26</sup>. As shown in Figure 1, IDO is the primary enzyme that catalyzes the initial step of the tryptophan metabolism pathway, which converts tryptophan to kynurenine. IDO is upregulated by inflammatory cytokine such as INF- $\gamma$  and tumor necrosis factor alpha (TNF- $\alpha$ )<sup>27, 28</sup>. A variety of cells express IDO, including monocytes, macrophages, dendritic cells, endothelial cells, fibroblasts, and certain cancer cells. IDO in tumor cells serves as an immunosuppressive enzyme that limits T cell responses against tumors<sup>29, 30</sup>. IDO has been found to be over-expressed in several types of cancers including lung cancer<sup>31</sup>. Inhibition of IDO by 1-methyltryptophan significantly delayed the tumor outgrowth in a mouse model of Lewis Lung carcinoma<sup>31</sup>. Clinical studies showed that mRNA expression of IDO was higher in lung cancer tissues than adjacent non-malignant lung tissues of patients<sup>32</sup>.

Emerging data suggest that kynurenine may play a more direct role than the consumption of tryptophan catabolized by IDO in immune regulation and responses in tumor microenvironment<sup>33-35</sup>. Kynurenine can activate transcription factor aryl hydrocarbon receptor (AhR). The activation of AhR induces a number of immunosuppressive phenotypes including the generation of immune-tolerant dendritic cells and regulatory T cells, which collectively foster a tumor immunological microenvironment that is defective in recognizing and eradicating cancer cells<sup>36</sup>. In a recent experimental study, kynureninase administration, which depleted kynurenine but did not impact



the consumption of tryptophan by IDO, significantly reduced tumor growth. Kynureninase treatment also significantly increased CD8<sup>+</sup> tumor infiltrating lymphocytes (TILs) and the production of IFN- $\gamma$ , TNF- $\alpha$  and interleukin-2 by CD8<sup>+</sup> T cells <sup>37</sup>. These data suggest that accumulation of kynurenine renders the immunosuppression in tumor microenvironment. High levels of circulating kynurenine and KTR in humans could be the consequence of enhanced IDO by inflammatory cytokines and/or the reduced metabolism of downstream kynurenine pathway. Previous studies have found that lung cancer patients had higher serum KTR concentration than healthy controls <sup>38</sup>. Our results with higher kynurenine or KTR with higher risk of lung cancer are consistent with findings from the prospective European Prospective Investigation into Cancer and Nutrition (EPIC)<sup>14</sup>, though the present study has much larger sample size and diverse populations. In addition, the stronger associations between kynurenine or KTR concentration and risk of lung cancer for individuals within 2 years of blood draw support the notion that cancer cells at subclinical stage may contribute to the elevation of circulating kynurenine in our patient population.

Our study found that the association for KTR or kynurenine metabolite QA with lung squamous cell carcinoma was stronger than that with lung adenocarcinoma. One possible explanation is the interaction between kynurenine and polycyclic aromatic hydrocarbons (PAH), specifically benzo[a]pyrene (B[a]P), in cigarette smoke, on the activation of AhR. Cigarette smoking is more strongly associated with risk of lung squamous cell carcinoma than adenocarcinoma in humans <sup>39</sup>. Experimental studies have shown that exposure to airborne particulate matters (mainly contains PAH) primarily induced lung squamous cell carcinoma in mice with intact AhR gene but no tumors at all in mice without AhR, suggesting that AhR is critical for the development of PAH-induced lung squamous cell carcinoma<sup>40</sup>. As described above, kynurenine and its downstream metabolites may be able to activate AhR, which would enhance the carcinogenic effect of PAH. These data may explain why our observed associations for KTR and QA were stronger with lung squamous cell carcinoma than that with lung adenocarcinoma.

## **Higher circulating quinolinic acid concentrations and risk of lung cancer**

Ours is the first study to evaluate the association between QA and lung cancer risk. We found that a higher concentration of QA in pre-diagnostic blood samples was associated with higher risk of lung cancer. QA, a downstream metabolite of kynurenine, is a known neurotoxin via stimulation of the presynaptic receptor which induces oxidative stress, and enhances the production of pro-inflammatory cytokines in the brain<sup>41</sup>. In the current study, circulating QA concentrations were highly correlated with KTR ( $r=0.57$ ), which is consistent with the fact that QA concentrations are correlated with IDO expression<sup>17</sup>. Previous studies showed that during inflammation, QA synthesis occurs mainly in immune cells<sup>17</sup>. Given that QA is a precursor of nicotinamide adenine dinucleotide, a coenzyme for redox reactions, accumulation of QA within immune cells could provide substrates for nicotinamide adenine dinucleotide synthesis to meet the enhanced requirements during an immune response<sup>17</sup>. Taken together, the observed association between QA and increased risk of lung cancer could reflect immune response against cancer prior to its clinical presentation. In addition, recent evidence showed that QA can inhibit the proliferation of cancer-killing T and natural killer cells<sup>42</sup>. Therefore, the higher concentrations of QA may promote tumor growth via its role in immune suppression. The association for QA with risk of lung cancer within <2 years of blood draw is stronger than those with longer time intervals, which suggests that this marker may be related to the progression of lung cancer and could be developed as biomarker for early detection of lung cancer.

## **Other notable findings**

In the present analysis, high levels of KTR and QA were associated with higher risk of lung cancer in both current and former smokers, but no risk was observed in never smokers. Former smokers had significantly higher levels of kynurenine, KTR and QA than never and current smokers. Individuals often gained weight after they quit smoking<sup>43</sup>. In our study, all five analyzed biomarkers were significantly associated with BMI in a dose-dependent manner. Our results were consistent with previous study results<sup>44</sup>. It is possible that the alternations of kynurenine metabolism in former smokers may contribute to their continued high risk of lung cancer after smoking cessation. It is interesting to note that the levels of kynurenine metabolites in current smokers were comparable with those in never smokers, but the

associations for these biomarkers with lung cancer risk were seen in current smokers only. These results suggest that cigarette smoking may be a prerequisite for kynurenine pathway to impact on the risk of developing lung cancer, but smoking is less likely to directly confound the kynurenine metabolites-lung cancer risk association.

### **Strengths and limitations**

The strengths of our study include 1) prospective design, 2) usage of pre-diagnostic plasma/serum samples, and 3) large sample size that provided sufficient power for stratified analysis. We also measured concentrations of metabolites of the kynurenine pathway, including QA as a novel inflammatory marker. In addition to matching on smoking status, intensity and duration, we also controlled for circulating cotinine concentration, a biomarker of recent tobacco exposure<sup>44</sup>, and eGFR, a renal function measurement that is strongly related to circulating concentrations of kynurenine and its metabolites<sup>43</sup>. We also measured KTR and neopterin, novel biomarkers for cellular immune activation as shown in prior work to have high intra-class correlation (ICCs, 0.74 and 0.67, respectively) in 4 different sampling visits over 3.5 years<sup>45</sup>. This indicated that a single time point measurement is a relatively reliable biomarker for long-term levels, and these biomarkers may be better than traditional cytokine biomarkers such as IFN-gamma and interleukins whose ICCs were lower<sup>46</sup>. The present study had some limitations. Although our analysis was based on a hypothesis that markers of immune modulation may be important in lung cancer etiology, the specific mechanisms underlying the observed associations are not clear. Given the complexity of immune response and their inter-connectedness, our studied biomarkers had relatively modest associations with lung cancer risk which limits their clinical utility for lung cancer screening and management. As in any observational study, our results could be confounded by other factors, including smoking, which is an established risk factor for lung cancer. Concentrations of all biomarkers except tryptophan varied among 3 groups of smokers – highest in former smokers and lowest in current smokers (**Supplementary Table 1**). Lung cancer risk was only significantly associated with KTR and QA concentrations in former and current smokers. Although smoking status, density and duration were matched for cases and controls in the present study

and circulating cotinine concentration was additionally adjusted for in the statistical analysis, the residual confounding of smoking on the observed biomarker-lung cancer risk associations cannot completely be ruled out.

## **CONCLUSION**

Our study demonstrates that lower circulating concentration of tryptophan and higher concentrations of kynurenine (i.e., higher KTR) and kynurenine downstream metabolite QA, biomarkers for immune regulation are associated with increased risk of lung cancer overall, in particular, among current smokers. Stronger associations for kynurenine, KTR and QA with imminent cancer occurring within the initial years after blood draw suggest that immune suppression in tumor microenvironment may play a more important role in the progression from a subclinical to clinical stage of lung cancer.

## **ACKNOWLEDGEMENTS**

We thank all participants, investigators and staff of the 20 original cohort studies that contributed data and biospecimens to the international Lung Cancer Cohort Consortium (LC3) project. For the Nurses' Health Study and the Health Professionals Follow-up Study, we acknowledge following state cancer registries for their help to ascertain cancer cases: AL, AZ, AR, CA, CO, CT, DE, FL, GA, ID, IL, IN, IA, KY, LA, ME, MD, MA, MI, NE, NH, NJ, NY, NC, ND, OH, OK, OR, PA, RI, SC, TN, TX, VA, WA, WY. The authors assume full responsibility for analyses and interpretation of these data. For the Campaign Against Cancer and Stroke (CLUE I) and the Campaign Against Cancer and Heart Disease (CLUE II) cohorts, we thank the Maryland Cancer Registry, Center for Cancer Surveillance and Control, Department of Health and Mental Hygiene for providing Cancer Incidence Data,

## **FUNDING**

The Lung Cancer Cohort Consortium (LC3) was supported by National Institutes of Health/National Cancer Institute grant No. 1U01CA155340. The work of T. L. Larose presented in this paper was undertaken during a postdoctoral placement at the International Agency for Research on Cancer, within the framework of an agreement between the Research Council of Norway and the Norwegian University of Science and Technology.

We acknowledge the main funding sources to individual participating cohorts (Funding agency, grant number): The Nurses' Health Study and the Health Professionals Follow-up Study (NIH UM1CA167552, UM1CA186107, and P01CA87969); the Shanghai Cohort Study and the Singapore Chinese Health Study (NIH R01CA1144034 and UM1CA182876); the Shanghai Men's Health Study (NIH UM1 CA173640); The Shanghai Women's Health Study (NIH UM1 CA182910); The Southern Community Cohort Study (NIH R01 CA092447 and U01 CA202979); The Women's Health Initiative (NIH contracts HHSN268201100046C, HHSN268201600001C, HHSN268201600002C, HHSN268201600003C, HHSN268201600004C, and HHSN271201600004C); the Women's Health Study (NIH CA047988, CA182913, HL043851, HL080467, and HL099355), the Physicians' Health Study (NIH CA097193, CA34944, CA40360, HL26490, and HL34595), and the New York University Women's Health Study (NIH UM1 CA182934, P30 CA016087, and P30 ES000260). The Alpha-Tocopherol Beta-Carotene Study (HHSN261201500005C and NCI Intramural Research Program); the Prostate, Lung, Colorectal, and Ovarian Cancer Screening Trial (Intramural Research Program, Division of Cancer Epidemiology and Genetics, National Cancer Institute (NCI) and contracts from the Division of Cancer Prevention, NCI); the Melbourne Collaborative Cohort Study (VicHealth and Cancer Council Victoria and Australian National Health and Medical Research Council grants 209057, 396414 and 1074383); The CLUE I and II (the State of Maryland, the Maryland Cigarette Restitution Fund, and the National Program of Cancer Registries); and the HUNT Study is a collaboration between NTNU (Norwegian University of Science

and Technology, Faculty of Medicine and Health Sciences), Trøndelag County Council, Central Norway Health Authority, and the Norwegian Institute of Public Health.

## REFERENCES

1. World Health Organization. Cancer Fact Sheet, vol. 2018, 2018.
2. Howlander N, Noone AM, Krapcho M, et al. SEER Cancer Statistics Review, 1975-2013, National Cancer Institute. Bethesda, MD, [http://seer.cancer.gov/csr/1975\\_2013/](http://seer.cancer.gov/csr/1975_2013/), based on November 2015 SEER data submission, posted to the SEER web site, April., 2016.
3. Dela Cruz CS, Tanoue LT, Matthay RA. Lung cancer: epidemiology, etiology, and prevention. *Clin Chest Med* 2011;32: 605-44.
4. Takahashi H, Ogata H, Nishigaki R, et al. Tobacco smoke promotes lung tumorigenesis by triggering IKKbeta- and JNK1-dependent inflammation. *Cancer Cell* 2010;17: 89-97.
5. Gomes M, Teixeira AL, Coelho A, et al. The role of inflammation in lung cancer. *Adv Exp Med Biol* 2014;816: 1-23.
6. Pine SR, Mechanic LE, Enewold L, et al. Increased levels of circulating interleukin 6, interleukin 8, C-reactive protein, and risk of lung cancer. *J Natl Cancer Inst* 2011;103: 1112-22.
7. Brenner DR, Fanidi A, Grankvist K, et al. Inflammatory Cytokines and Lung Cancer Risk in 3 Prospective Studies. *Am J Epidemiol* 2017;185: 86-95.
8. Shiels MS, Pfeiffer RM, Hildesheim A, et al. Circulating inflammation markers and prospective risk for lung cancer. *J Natl Cancer Inst* 2013;105: 1871-80.
9. Biancotto A, Wank A, Perl S, et al. Baseline levels and temporal stability of 27 multiplexed serum cytokine concentrations in healthy subjects. *PLoS One* 2013;8: e76091.
10. Mbongue JC, Nicholas DA, Torrez TW, et al. The Role of Indoleamine 2, 3-Dioxygenase in Immune Suppression and Autoimmunity. *Vaccines (Basel)* 2015;3: 703-29.
11. Lob S, Konigsrainer A, Rammensee HG, et al. Inhibitors of indoleamine-2,3-dioxygenase for cancer therapy: can we see the wood for the trees? *Nat Rev Cancer* 2009;9: 445-52.
12. Mojic M, Takeda K, Hayakawa Y. The Dark Side of IFN-gamma: Its Role in Promoting Cancer Immuno-evasion. *Int J Mol Sci* 2017;19.
13. Pingle SK, Tumane RG, Jawade AA. Neopterin: Biomarker of cell-mediated immunity and potent usage as biomarker in silicosis and other occupational diseases. *Indian journal of occupational and environmental medicine* 2008;12: 107-11.

14. Chuang SC, Fanidi A, Ueland PM, et al. Circulating biomarkers of tryptophan and the kynurenine pathway and lung cancer risk. *Cancer Epidemiol Biomarkers Prev* 2014;23: 461-8.
15. Zuo H, Tell GS, Vollset SE, et al. Interferon-gamma-induced inflammatory markers and the risk of cancer: the Hordaland Health Study. *Cancer* 2014;120: 3370-7.
16. Fallarino F, Grohmann U, Vacca C, et al. T cell apoptosis by tryptophan catabolism. *Cell death and differentiation* 2002;9: 1069-77.
17. Moffett JR, Namboodiri MA. Tryptophan and the immune response. *Immunology and cell biology* 2003;81: 247-65.
18. Fanidi A, Muller DC, Yuan JM, et al. Circulating Folate, Vitamin B6, and Methionine in Relation to Lung Cancer Risk in the Lung Cancer Cohort Consortium (LC3). *J Natl Cancer Inst* 2018;110.
19. Ueland PM, Midttun O, Windelberg A, et al. Quantitative profiling of folate and one-carbon metabolism in large-scale epidemiological studies by mass spectrometry. *Clin Chem Lab Med* 2007;45: 1737-45.
20. Midttun O, Kvalheim G, Ueland PM. High-throughput, low-volume, multianalyte quantification of plasma metabolites related to one-carbon metabolism using HPLC-MS/MS. *Analytical and bioanalytical chemistry* 2013;405: 2009-17.
21. Midttun O, Theofylaktopoulou D, McCann A, et al. Circulating concentrations of biomarkers and metabolites related to vitamin status, one-carbon and the kynurenine pathways in US, Nordic, Asian, and Australian populations. *Am J Clin Nutr* 2017;105: 1314-26.
22. Hustad S, Eussen S, Midttun O, et al. Kinetic modeling of storage effects on biomarkers related to B vitamin status and one-carbon metabolism. *Clin Chem* 2012;58: 402-10.
23. Levey AS, Stevens LA, Schmid CH, et al. A new equation to estimate glomerular filtration rate. *Annals of internal medicine* 2009;150: 604-12.
24. Breslow N, Day N. *Statistical methods in cancer research, vol.1: The analysis of case-control studies*. Lyon: IARC: IARC Scientific Pub No. 32., 1980.
25. Benowitz NL, Hukkanen J, Jacob P, 3rd. Nicotine chemistry, metabolism, kinetics and biomarkers. *Handb Exp Pharmacol* 2009: 29-60.
26. Mellor AL, Munn DH. IDO expression by dendritic cells: tolerance and tryptophan catabolism. *Nat Rev Immunol* 2004;4: 762-74.



27. Hissong BD, Byrne GI, Padilla ML, et al. Upregulation of interferon-induced indoleamine 2,3-dioxygenase in human macrophage cultures by lipopolysaccharide, muramyl tripeptide, and interleukin-1. *Cell Immunol* 1995;160: 264-9.
28. Currier AR, Ziegler MH, Riley MM, et al. Tumor necrosis factor-alpha and lipopolysaccharide enhance interferon-induced antichlamydial indoleamine dioxygenase activity independently. *J Interferon Cytokine Res* 2000;20: 369-76.
29. Munn DH, Sharma MD, Lee JR, et al. Potential regulatory function of human dendritic cells expressing indoleamine 2,3-dioxygenase. *Science* 2002;297: 1867-70.
30. Uyttenhove C, Pilotte L, Theate I, et al. Evidence for a tumoral immune resistance mechanism based on tryptophan degradation by indoleamine 2,3-dioxygenase. *Nat Med* 2003;9: 1269-74.
31. Munn DH, Mellor AL. IDO and tolerance to tumors. *Trends Mol Med* 2004;10: 15-8.
32. Karanikas V, Zamanakou M, Kerenidi T, et al. Indoleamine 2,3-dioxygenase (IDO) expression in lung cancer. *Cancer biology & therapy* 2007;6: 1258-62.
33. Routy JP, Routy B, Graziani GM, et al. The Kynurenine Pathway Is a Double-Edged Sword in Immune-Privileged Sites and in Cancer: Implications for Immunotherapy. *Int J Tryptophan Res* 2016;9: 67-77.
34. Dagenais-Lussier X, Aounallah M, Mehraj V, et al. Kynurenine Reduces Memory CD4 T-Cell Survival by Interfering with Interleukin-2 Signaling Early during HIV-1 Infection. *J Virol* 2016;90: 7967-79.
35. Cheong JE, Sun L. Targeting the IDO1/TDO2-KYN-AhR Pathway for Cancer Immunotherapy - Challenges and Opportunities. *Trends Pharmacol Sci* 2018;39: 307-25.
36. Mezrich JD, Fechner JH, Zhang X, et al. An interaction between kynurenine and the aryl hydrocarbon receptor can generate regulatory T cells. *J Immunol* 2010;185: 3190-8.
37. Triplett TA, Garrison KC, Marshall N, et al. Reversal of indoleamine 2,3-dioxygenase-mediated cancer immune suppression by systemic kynurenine depletion with a therapeutic enzyme. *Nat Biotechnol* 2018;36: 758-64.
38. Suzuki Y, Suda T, Furuhashi K, et al. Increased serum kynurenine/tryptophan ratio correlates with disease progression in lung cancer. *Lung Cancer* 2010;67: 361-5.
39. Khuder SA. Effect of cigarette smoking on major histological types of lung cancer: a meta-analysis. *Lung Cancer* 2001;31: 139-48.

40. Matsumoto Y, Ide F, Kishi R, et al. Aryl hydrocarbon receptor plays a significant role in mediating airborne particulate-induced carcinogenesis in mice. *Environ Sci Technol* 2007;41: 3775-80.
41. Lugo-Huitron R, Ugalde Muniz P, Pineda B, et al. Quinolinic acid: an endogenous neurotoxin with multiple targets. *Oxid Med Cell Longev* 2013;2013: 104024.
42. Frumento G, Rotondo R, Tonetti M, et al. Tryptophan-derived catabolites are responsible for inhibition of T and natural killer cell proliferation induced by indoleamine 2,3-dioxygenase. *The Journal of experimental medicine* 2002;196: 459-68.
43. Theofylaktopoulou D, Midttun O, Ulvik A, et al. A community-based study on determinants of circulating markers of cellular immune activation and kynurenines: the Hordaland Health Study. *Clin Exp Immunol* 2013;173: 121-30.
44. Boffetta P, Clark S, Shen M, et al. Serum cotinine level as predictor of lung cancer risk. *Cancer Epidemiol Biomarkers Prev* 2006;15: 1184-8.
45. Midttun O, Townsend MK, Nygard O, et al. Most blood biomarkers related to vitamin status, one-carbon metabolism, and the kynurenine pathway show adequate preanalytical stability and within-person reproducibility to allow assessment of exposure or nutritional status in healthy women and cardiovascular patients. *J Nutr* 2014;144: 784-90.
46. Epstein MM, Breen EC, Magpantay L, et al. Temporal stability of serum concentrations of cytokines and soluble receptors measured across two years in low-risk HIV-seronegative men. *Cancer Epidemiol Biomarkers Prev* 2013;22: 2009-15.

**Table 1.** Baseline and clinical characteristics of study participants overall and by continent, the Lung Cancer Cohort Consortium (LC3) Study

Baseline and Clinical Characteristics	US cohorts		EU/AU cohorts		Asian cohorts		Overall	
	No.(%) of participants in group		No.(%) of participants in group		No.(%) of participants in group		No.(%) of participants in group	
	Cases (n=2,400)	Matched controls (n=2,400)	Cases (n=1,189)	Matched controls (n=1,189)	Cases (n=1,775)	Matched controls (n=1,775)	Cases (n=5,364)	Matched controls (n=5,364)
Sex								
Men	991 (41.3%)	991 (41.3%)	688 (57.9%)	688 (57.9%)	1,229 (69.2%)	1,229 (69.2%)	2,908 (54.2%)	2,908 (54.2%)
Women	1,409 (58.7%)	1,409 (58.7%)	501 (42.1%)	501 (42.1%)	546 (30.8%)	546 (30.8%)	2,456 (45.8%)	2,456 (45.8%)
Smoking status								
Never	569 (23.7%)	569 (23.7%)	156 (13.1%)	156 (13.1%)	602 (33.9%)	602 (33.9%)	1,327 (24.7%)	1,327 (24.7%)
Former	1,007 (42.0%)	1,007 (42.0%)	335 (28.2%)	335 (28.2%)	176 (9.9%)	176 (9.9%)	1,518 (28.3%)	1,518 (28.3%)
Current	824 (34.3%)	824 (34.3%)	698 (58.7%)	698 (58.7%)	997 (56.2%)	997 (56.2%)	2,519 (47.0%)	2,519 (47.0%)
Education								
Less than high school	237 (9.9%)	215 (9%)	662 (55.6%)	598 (50.2%)	898 (50.6%)	883 (49.7%)	1,797 (33.5%)	1,696 (31.6%)
Completed high school	357 (14.9%)	374 (15.6%)	159 (13.4%)	180 (15.2%)	243 (13.7%)	230 (13.0%)	759 (14.1%)	784 (14.6%)
Vocational school	422 (17.6%)	435 (18.1%)	180 (15.2%)	200 (16.8%)	289 (16.3%)	279 (15.7%)	891 (16.6%)	914 (17.0%)
Some college	402 (16.8%)	393 (16.4%)	107 (9%)	129 (10.9%)	171 (9.6%)	196 (11%)	680 (12.7%)	718 (13.4%)
College graduate	357 (14.9%)	319 (13.3%)	63 (5.3%)	64 (5.4%)	104 (5.9%)	113 (6.4%)	524 (9.8%)	496 (9.2%)
Graduate studies	574 (23.9%)	637 (26.5%)	10 (0.8%)	8 (0.7%)	62 (3.5%)	65 (3.7%)	646 (12%)	710 (13.2%)
Unknown	51 (2.1%)	27 (1.1%)	8 (0.7%)	10 (0.8%)	8 (0.5%)	9 (0.5%)	67 (1.2%)	46 (0.9%)
Body Mass Index <sup>a</sup>								
< 18.5	30 (1.3%)	31 (1.3%)	14 (1.2%)	10 (0.8%)	157 (8.8%)	113 (6.4%)	201 (3.7%)	154 (2.9%)
18.5-24.9	1,088 (45.3%)	1,020 (42.5%)	521 (43.8%)	435 (36.6%)	1,203 (67.8%)	1,192 (67.2%)	2,812 (52.4%)	2,647 (49.3%)
25.0-29.9	841 (35%)	858 (35.8%)	468 (39.4%)	536 (45.1%)	369 (20.8%)	424 (23.9%)	1,678 (31.3%)	1,818 (33.9%)
≥ 30.0	378 (15.8%)	430 (17.9%)	185 (15.6%)	206 (17.4%)	46 (2.6%)	46 (2.6%)	609 (11.4%)	682 (12.7%)
Unknown	63 (2.6%)	61 (2.5%)	1 (0.1%)	2 (0.2%)	0 (0.0%)	0 (0.0%)	64 (1.2%)	63 (1.2%)
	<b>Continuous variables, median (5th-95th percentile)</b>							
Age at recruitment (years)	60 (42-74)	60 (42-74)	60 (45-70)	60 (45-70)	60 (46-72)	60 (46-71)	60 (44-72)	60 (44-72)
Circulating concentrations for biomarkers								
Tryptophan, μmol/L	63.9 (41.3-89.1)	64.4 (43.7-90.5)	67.8 (48.9-92.7)	68.1 (50.1-91.1)	67.3 (48.6-91.2)	67.5 (49.1-90.1)	66.0 (44.9-90.8)	66.5 (46.2-90.7)
Kynurenine, μmol/L	1.51 (1.00-2.37)	1.53 (1.02-2.34)	1.52 (1.06-2.19)	1.52 (1.07-2.18)	1.49 (1.08-2.18)	1.48 (1.09-2.14)	1.50 (1.04-2.25)	1.51 (1.05-2.22)
Kynurinine:tryptophan ratio (nmol/μmol)	22.6 (16.6-38.6)	23.6 (16.6-37.0)	22.3 (16.4-33.8)	22.0 (16.5-32.1)	21.9 (15.9-34.1)	21.9 (16.0-32.1)	22.6 (16.2-36.2)	22.6 (16.4-34.5)
Quinolinic acid, nmol/L	364 (200-789)	363 (207-741)	341 (201-633)	334 (202-605)	350 (207-651)	350 (216-605)	354 (203-708)	353 (208-685)
Neopterin, nmol/L	12.0 (5.74-25.0)	11.8 (5.66-25.5)	10.2 (4.78-20.9)	10.3 (4.38-19.5)	10.6 (5.29-24.0)	10.7 (5.28-24.6)	11.1 (5.31-24.0)	11.0 (5.14-24.6)
	<b>Clinical characteristics, case participants only</b>							
Age at diagnosis, median (range), years	70 (55-83)		69 (54-80)		69 (52-80)		69.8 (53.6-82.0)	
Time from blood draw to diagnosis (years)	5.2 (1-15.5)		10.0 (1.5-16.0)		5.8 (0.7-16.5)		6.3 (1.0-16.0)	
Histology, No. (%)								
Large cell carcinoma	112 (4.6%)		46 (4.0%)		16 (1.0%)		174 (3.3%)	
Small cell carcinoma	245 (10.4%)		150 (12.5%)		99 (5.5%)		492 (9.2%)	
Squamous cell carcinoma	291 (11.9%)		231 (19.5%)		319 (17.9%)		836 (15.5%)	
Adenocarcinoma	1,034 (42.7%)		419 (34.5%)		615 (34.6%)		2,056 (38.4%)	
Missing / Unknown	735 (31.4%)		357 (29.5%)		726 (41%)		1,806 (33.6%)	

<sup>a</sup> Body mass index is calculated as weight in kilograms divided by height in meters squared

**Table 2:** Odds ratios of lung cancer incidence comparing higher quintiles with the lowest quintile of circulating biomarkers  
The Lung Cancer Cohort Consortium (LC3) Study

Biomarkers and risk estimates	Quintiles of biomarker <sup>a</sup>					<i>P</i> <sub>trend</sub>
	Q1	Q2	Q3	Q4	Q5	
Tryptophan						
Cases/controls	1199/1073	1,020/1,073	1,057/1,073	1,095/1,073	993/1,072	
Adjusted odds ratio (95% CI) <sup>a</sup>	1.00	<b>0.86 (0.76-0.97)</b>	<b>0.87 (0.77-0.98)</b>	<b>0.86 (0.76-0.97)</b>	<b>0.85 (0.75-0.96)</b>	<b>0.019</b>
Kynurenine						
Cases/controls	1083/1073	1,044/1,073	1,148/1,086	1,020/1,061	1,069/1,071	
Adjusted odds ratio (95% CI) <sup>a</sup>	1.00	1.05 (0.92-1.18)	1.02 (0.89-1.16)	1.01 (0.89-1.15)	<b>1.22 (1.06-1.40)</b>	<b>0.033</b>
KTR <sup>b</sup>						
Cases/controls	1082/1073	1,028/1,073	1,020/1,073	1,034/1,073	1,197/1,072	
Adjusted odds ratio (95% CI) <sup>a</sup>	1.00	0.97 (0.86-1.10)	0.96 (0.84-1.09)	1.01 (0.89-1.15)	<b>1.31 (1.14-1.50)</b>	<b>&lt;0.001</b>
Quinolinic acid						
Cases/controls	1158/1073	1,033/1,074	902/1,072	1,123/1,073	1,148/1,072	
Adjusted odds ratio (95% CI) <sup>a</sup>	1.00	0.95 (0.84-1.08)	0.94 (0.83-1.06)	1.09 (0.96-1.25)	<b>1.31 (1.14-1.51)</b>	<b>&lt;0.001</b>
Neopterin						
Cases/controls <sup>c</sup>	1022/1072	1,065/1,071	1,087/1,070	1,051/1,069	1,128/1,071	
Adjusted odds ratio (95% CI) <sup>a</sup>	1.00	<b>1.12 (0.99-1.27)</b>	1.09 (0.96-1.24)	1.12 (0.98-1.28)	<b>1.31 (1.14-1.51)</b>	<b>0.001</b>

<sup>a</sup> All models were adjusted for educational attainment (categorical), body mass index (kg/m<sup>2</sup>) (categorical), serum cotinine concentrations (continuous), estimated glomerular filtration rate (continuous), and cohort; Bold figures indicate the 95% confidence intervals (CIs) of odds ratio did not include one or P values for trend were less than 0.05.

<sup>b</sup> KTR, kynurenine to tryptophan ratio.

<sup>c</sup> Eleven case-control pairs were excluded due missing value.

**Table 3:** Odds ratios of lung cancer incidence comparing higher quintiles with the lowest quintile of circulating biomarkers stratified by smoking status

The Lung Cancer Cohort Consortium (LC3) Study

Smoking status and biomarker	Adjusted Odds Ratio (95% Confidence Interval) by quintiles of biomarker <sup>a</sup>					P trend
	Q1	Q2	Q3	Q4	Q5	
Current smokers						
Tryptophan	1.00	<b>0.80 (0.66-0.98)</b>	<b>0.79 (0.65-0.96)</b>	<b>0.81 (0.66-0.98)</b>	<b>0.78 (0.63-0.95)</b>	<b>0.046</b>
Kynurenine	1.00	0.86 (0.72-1.04)	1.04 (0.86-1.26)	1.05 (0.86-1.28)	1.16 (0.93-1.46)	0.057
KTR <sup>b</sup>	1.00	0.94 (0.79-1.11)	0.95 (0.79-1.13)	1.04 (0.87-1.26)	<b>1.42 (1.15-1.75)</b>	<b>0.005</b>
Quinolinic acid	1.00	0.97 (0.83-1.14)	1.04 (0.87-1.24)	<b>1.26 (1.04-1.53)</b>	<b>1.42 (1.14-1.76)</b>	<b>&lt;0.001</b>
Neopterin	1.00	1.10 (0.91-1.34)	1.21 (0.98-1.48)	<b>1.28 (1.02-1.61)</b>	<b>1.45 (1.13-1.86)</b>	<b>0.003</b>
Former smokers						
Tryptophan	1.00	<b>0.73 (0.58-0.92)</b>	<b>0.81 (0.64-1.03)</b>	0.82 (0.65-1.04)	<b>0.74 (0.58-0.94)</b>	0.077
Kynurenine	1.00	1.12 (0.84-1.49)	1.23 (0.95-1.60)	1.02 (0.78-1.34)	1.08 (0.82-1.41)	0.955
KTR <sup>b</sup>	1.00	1.06 (0.80-1.41)	1.18 (0.89-1.54)	1.13 (0.86-1.49)	<b>1.32 (1.00-1.74)</b>	<b>0.035</b>
Quinolinic acid	1.00	1.00 (0.74-1.33)	0.82 (0.62-1.08)	1.17 (0.89-1.55)	1.20 (0.90-1.59)	<b>0.037</b>
Neopterin	1.00	1.14 (0.86-1.50)	1.14 (0.85-1.54)	1.01 (0.74-1.37)	1.34 (0.97-1.86)	0.196
Never smokers						
Tryptophan	1.00	1.00 (0.80-1.26)	1.04 (0.81-1.32)	1.16 (0.88-1.53)	0.87 (0.64-1.18)	0.911
Kynurenine	1.00	1.06 (0.85-1.33)	1.12 (0.88-1.43)	1.05 (0.80-1.38)	1.17 (0.85-1.59)	0.406
KTR <sup>b</sup>	1.00	1.00 (0.79-1.27)	0.92 (0.72-1.19)	0.92 (0.71-1.19)	1.17 (0.88-1.54)	0.562
Quinolinic acid	1.00	0.89 (0.69-1.13)	<b>0.70 (0.54-0.90)</b>	0.92 (0.71-1.20)	1.07 (0.79-1.44)	0.707
Neopterin	1.00	1.09 (0.85-1.40)	1.18 (0.89-1.55)	1.30 (0.97-1.74)	1.19 (0.86-1.63)	0.168

<sup>a</sup> All models were adjusted for educational attainment (categorical), body mass index (kg/m<sup>2</sup>) (categorical), serum cotinine concentrations (continuous), estimated glomerular filtration rate (continuous), and cohort; Bold figures indicate the 95% confidence intervals of odds ratio did not include one or P values for trend were less than 0.05.

<sup>b</sup> KTR, kynurenine to tryptophan ratio.

**Table 4:** Odds ratios of lung cancer incidence by histological subtype comparing higher quintiles with the lowest quintile of circulating biomarkers

The Lung Cancer Cohort Consortium (LC3) Study

Histological subtype and biomarker	Adjusted Odds Ratio (95% Confidence Interval) by quintiles of biomarker <sup>a</sup>					P trend
	Q1	Q2	Q3	Q4	Q5	
Large cell carcinoma						
Tryptophan	1.00	1.88 (0.94-3.74)	1.94 (0.95-3.98)	1.97 (0.90-4.31)	1.96 (0.80-4.81)	0.142
Kynurenine	1.00	1.62 (0.73-3.57)	1.25 (0.54-2.88)	2.05 (0.96-4.36)	2.28 (0.92-5.68)	<b>0.048</b>
KTR <sup>b</sup>	1.00	<b>0.40 (0.17-0.90)</b>	0.74 (0.34-1.61)	1.02 (0.47-2.18)	0.90 (0.40-2.04)	0.339
Quinolinic acid	1.00	1.04 (0.49-2.19)	0.65 (0.31-1.37)	1.12 (0.50-2.48)	1.68 (0.68-4.12)	0.344
Neopterin	1.00	1.33 (0.51-3.49)	1.70 (0.69-4.15)	1.11 (0.41-3.02)	1.97 (0.66-5.87)	0.390
Small cell carcinoma						
Tryptophan	1.00	0.77 (0.51-1.17)	0.74 (0.48-1.13)	<b>0.57 (0.37-0.88)</b>	0.82 (0.53-1.25)	0.189
Kynurenine	1.00	0.87 (0.58-1.32)	1.02 (0.67-1.55)	1.01 (0.64-1.59)	0.99 (0.62-1.57)	0.838
KTR <sup>b</sup>	1.00	0.65 (0.43-1.01)	1.08 (0.70-1.65)	0.74 (0.47-1.16)	1.13 (0.71-1.80)	0.447
Quinolinic acid	1.00	0.83 (0.54-1.28)	0.79 (0.51-1.25)	1.37 (0.88-2.13)	1.32 (0.81-2.14)	0.071
Neopterin	1.00	1.20 (0.75-1.92)	1.07 (0.64-1.80)	0.89 (0.53-1.50)	1.29 (0.71-2.36)	0.823
Squamous cell carcinoma						
Tryptophan	1.00	<b>0.67 (0.49-0.93)</b>	<b>0.68 (0.48-0.96)</b>	0.72 (0.51-1.01)	0.76 (0.54-1.06)	0.304
Kynurenine	1.00	<b>0.71 (0.50-1.00)</b>	1.21 (0.86-1.69)	1.04 (0.73-1.47)	1.22 (0.84-1.77)	0.066
KTR <sup>c</sup>	1.00	1.06 (0.77-1.46)	0.99 (0.71-1.38)	1.01 (0.72-1.41)	<b>1.68 (1.17-2.43)</b>	<b>0.023</b>
Quinolinic acid	1.00	<b>1.58 (1.15-2.16)</b>	<b>1.38 (0.99-1.93)</b>	<b>1.56 (1.11-2.20)</b>	<b>1.99 (1.35-2.91)</b>	<b>0.003</b>
Neopterin	1.00	1.61 (1.14-2.26)	1.21 (0.83-1.75)	1.36 (0.92-2.01)	1.34 (0.88-2.04)	0.468
Adenocarcinoma						
Tryptophan	1.00	0.96 (0.79-1.17)	0.98 (0.80-1.20)	<b>1.26 (1.01-1.56)</b>	0.89 (0.70-1.12)	0.764
Kynurenine	1.00	1.04 (0.85-1.27)	1.14 (0.93-1.39)	1.00 (0.80-1.24)	1.09 (0.86-1.39)	0.615
KTR <sup>b</sup>	1.00	1.02 (0.83-1.24)	1.00 (0.82-1.23)	1.00 (0.81-1.24)	1.12 (0.89-1.40)	0.426
Quinolinic acid	1.00	0.85 (0.68-1.05)	0.85 (0.68-1.06)	1.01 (0.80-1.27)	<b>1.36 (1.05-1.74)</b>	<b>0.009</b>
Neopterin	1.00	1.04 (0.84-1.29)	1.13 (0.90-1.43)	1.19 (0.92-1.52)	1.27 (0.97-1.66)	0.059

<sup>a</sup> All models were adjusted for educational attainment (categorical), body mass index (kg/m<sup>2</sup>) (categorical), serum cotinine concentrations (continuous), estimated glomerular filtration rate (continuous), and cohort; Bold figures indicate the 95% confidence intervals of odds ratio did not include one or P values for trend were less than 0.05.

<sup>b</sup> KTR, kynurenine to tryptophan ratio.

**Table 5:** Odds ratios of lung cancer incidence comparing higher quintiles with the lowest quintile of circulating biomarkers stratified by time from blood draw to cancer diagnosis, The Lung Cancer Cohort Consortium (LC3) Study

Time from blood draw to cancer diagnosis and biomarkers	Adjusted Odds Ratio (95% Confidence Interval) by quintiles of biomarker <sup>a</sup>					P trend
	Q1	Q2	Q3	Q4	Q5	
<b>&lt; 2 years</b>						
Tryptophan	1.00	<b>0.56 (0.37-0.85)</b>	<b>0.65 (0.43-0.99)</b>	0.73 (0.47-1.14)	0.72 (0.45-1.13)	0.542
Kynurenine	1.00	1.21 (0.78-1.89)	1.22 (0.76-1.96)	1.15 (0.71-1.87)	<b>1.86 (1.13-3.08)</b>	<b>0.032</b>
KTR <sup>b</sup>	1.00	1.14 (0.72-1.80)	0.95 (0.60-1.52)	0.98 (0.61-1.57)	<b>1.92 (1.17-3.14)</b>	<b>0.024</b>
Quinolinic acid	1.00	1.20 (0.76-1.90)	0.86 (0.53-1.38)	<b>1.62 (1.00-2.61)</b>	<b>2.28 (1.38-3.77)</b>	<b>&lt;0.001</b>
Neopterin	1.00	1.18 (0.73-1.89)	1.22 (0.74-2.01)	1.44 (0.82-2.51)	1.52 (0.85-2.72)	0.154
<b>2–4.9 years</b>						
Tryptophan	1.00	1.02 (0.75-1.39)	1.03 (0.75-1.41)	0.97 (0.70-1.35)	0.83 (0.59-1.17)	0.279
Kynurenine	1.00	1.02 (0.72-1.44)	0.97 (0.69-1.36)	0.95 (0.66-1.37)	1.13 (0.76-1.66)	0.689
KTR <sup>b</sup>	1.00	0.98 (0.71-1.35)	0.93 (0.67-1.29)	1.08 (0.78-1.51)	1.29 (0.90-1.84)	0.142
Quinolinic acid	1.00	0.87 (0.62-1.21)	0.77 (0.55-1.07)	1.10 (0.79-1.54)	1.24 (0.85-1.79)	0.136
Neopterin	1.00	0.98 (0.70-1.38)	1.05 (0.72-1.54)	<b>1.76 (1.16-2.68)</b>	<b>1.72 (1.10-2.67)</b>	<b>0.003</b>
<b>≥ 5 years</b>						
Tryptophan	1.00	0.85 (0.70-1.03)	0.84 (0.70-1.02)	0.93 (0.77-1.13)	0.85 (0.69-1.04)	0.413
Kynurenine	1.00	0.95 (0.79-1.13)	1.13 (0.94-1.35)	1.05 (0.87-1.28)	1.17 (0.94-1.45)	0.092
KTR <sup>b</sup>	1.00	0.99 (0.84-1.17)	1.08 (0.91-1.29)	1.09 (0.91-1.31)	1.21 (0.98-1.48)	0.058
Quinolinic acid	1.00	0.96 (0.81-1.13)	0.96 (0.8-1.15)	<b>1.21 (1.00-1.45)</b>	<b>1.28 (1.03-1.59)</b>	<b>0.005</b>
Neopterin	1.00	1.07 (0.89-1.28)	1.09 (0.89-1.33)	1.09 (0.87-1.35)	1.22 (0.96-1.56)	0.158

<sup>a</sup> All models were adjusted for educational attainment (categorical), body mass index (kg/m<sup>2</sup>) (categorical), serum cotinine concentrations (continuous), estimated glomerular filtration rate (continuous), and cohort; Bold figures indicate the 95% confidence intervals of odds ratio did not include one or P values for trend were less than 0.05.

<sup>b</sup> KTR, kynurenine to tryptophan ratio.

**Supplementary Table 1:** Geometric means<sup>a</sup> of biomarkers by smoking status in total control subjects

## The Lung Cancer Cohort Consortium (LC3) Study

Biomarkers	Never smokers (95% CI)	Former smokers (95% CI)	Current smokers (95% CI)	<i>P</i> for Former vs Never	<i>P</i> for Current vs Never	<i>P</i> for Current vs Former	<i>P</i> for 3 smoking groups
No. of subjects	1,327	1,518	2,519				
Tryptophan (μmol/L)	65.6 (64.7-66.4)	65.8 (65.1-66.2)	65.1 (64.6-65.7)	0.632	0.429	0.123	0.293
Kynurenine (μmol/L)	1.48 (1.46-1.50)	<b>1.56</b> <b>(1.54-1.58)</b>	1.48 (1.48-1.50)	<b>&lt;0.001</b>	0.999	<b>&lt;0.001</b>	<b>&lt;0.001</b>
KTR (nmol/μmol) <sup>b</sup>	22.7 (22.4-23.0)	<b>23.7</b> <b>(23.5-24.0)</b>	22.8 (22.6-23.1)	<b>&lt;0.001</b>	0.478	<b>&lt;0.001</b>	<b>&lt;0.001</b>
Quinolinic Acid (nmol/L)	368.7 (361.2-376.5)	<b>387.0</b> <b>(380.5-393.6)</b>	334.7 (330.0-339.5)	<b>&lt;0.001</b>	<0.001	<b>&lt;0.001</b>	<b>&lt;0.001</b>
Neopterin <sup>c</sup> (nmol/L)	10.8 (10.5-11.0)	11.1 (10.9-11.3)	10.3 (10.1-10.5)	0.095	<b>0.002</b>	<b>&lt;0.001</b>	<b>&lt;0.001</b>

<sup>a</sup>Geometric means were derived from analysis of covariance (ANCOVA) with adjustment for age, gender, estimated glomerular filtration rate (eGFR), and cohort indicators that also provided *P* values for pairwise and overall difference in biomarkers concentrations among 3 smoking categories. Bold figures indicate the 95% confidence intervals of geometric means in former smokers did not overlap with those in never or current smokers or *P* values for comparing geometric means between two smoking groups were less than 0.05.

<sup>b</sup>KTR: Kynurenine: tryptophan ratio.

<sup>c</sup>Seven subjects (1 never smoker, 3 former smokers, and 3 current smokers) were excluded due to missing data.



**Supplementary Table 2:** Geometric means of biomarkers by body mass index in total control subjects,  
the Lung Cancer Cohort Consortium (LC3) Study

Body mass index (kg/m <sup>2</sup> )	No. of subjects	Geometric Mean <sup>a</sup> (95% CI)				
		Tryptophan (μmol/L)	Kynurenine (μmol/L)	KTR <sup>b</sup> (nmol/μmol)	Neopterin (nmol/L)	Quinolinic Acid (nmol/L)
<18.5	153	61.1 (59.3-63.0)	1.42 (1.36-1.46)	23.1 (22.3-23.9)	10.5 (10.0-11.1)	317.7 (302.5-333.7)
18.5-24.9	2,654	64.6 (64.1-65.1)	1.46 (1.46-1.48)	22.6 (22.4-22.8)	10.4 (10.3-10.6)	336.3 (332.1-340.6)
25.0-29.9	1,874	66.4 (65.8-67.0)	1.54 (1.52-1.54)	23.1 (22.9-23.4)	10.7 (10.5-10.9)	369.6 (364.5-374.9)
≥30.0	683	66.7 (65.8-67.7)	1.64 (1.60-1.66)	24.5 (24.1-24.8)	11.3 (11.0-11.6)	415.4 (405.8-425.2)
<i>P</i> <sub>trend</sub>		<b>&lt;0.001</b>	<b>&lt;0.001</b>	<b>&lt;0.001</b>	<b>&lt;0.001</b>	<b>&lt;0.001</b>

<sup>a</sup> Geometric means were derived from analysis of covariance (ANCOVA) with adjustment for age, gender, smoking status, estimated glomerular filtration rate (eGFR) and cohort indicators; Bold figures indicate that P values for trend in geometric means across body mass index levels were less than 0.05.

<sup>b</sup>KTR: Kynurenine: tryptophan ratio.

**Supplementary Table 3: Median (20<sup>th</sup>-80<sup>th</sup> percentiles) of circulating biomarker concentrations among control subjects from different cohorts, The Lung Cancer Cohort Consortium (LC3) Study**

<b>Cohorts by Region<sup>a</sup></b>	<b>N</b>	<b>Tryptophan, (μmol/L)</b>	<b>Kynurenine, (μmol/L)</b>	<b>KTR<sup>b</sup> (nmol/μmol)</b>	<b>Quinolinic acid (nmol/L)</b>	<b>Neopterin (nmol/L)</b>
<b>US cohorts</b>						
CLUE I & II	191	68.3 (57.2-80.7)	1.58 (1.29-1.88)	22.9 (19.3-27.6)	353 (278-468)	13.6 (10.3-19.5)
CPS-II	182	71.6 (60.4-86.8)	1.81 (1.47-2.16)	24.5 (20.7-30.6)	454 (340-614)	9.71 (7.60-12.9)
HPFS	155	67.4 (60.0-77.3)	1.64 (1.37-1.92)	23.9 (20.3-29.9)	362 (273-497)	9.34 (6.87-12.2)
MEC	174	67.6 (57.9-79.2)	1.63 (1.40-1.92)	24.2 (20.6-28.7)	383 (283-534)	17.3 (13.3-22.3)
NHS	345	69.3 (58.7-81.0)	1.52 (1.29-1.80)	21.7 (18.4-27.5)	349 (261-475)	13.6 (10.5-18.1)
NYUWHS	171	66.0 (57.8-77.5)	1.44 (1.23-1.84)	22.3 (19.2-26.3)	331 (250-430)	14.0 (11.0-18.4)
PHS	81	64.2 (57.2-71.3)	1.60 (1.37-1.83)	25.0 (21.8-29.1)	397 (324-521)	7.16 (5.85-9.86)
PLCO	450	64.7 (53.2-78.8)	1.68 (1.35-2.05)	25.1 (21.1-31.3)	433 (317-598)	12.4 (8.84-17.2)
SCCS	226	54.6 (45.6-65.5)	1.24 (1.05-1.46)	22.5 (18.2-27.3)	325 (232-473)	8.83 (7.03-13.2)
WHI	241	51.3 (45.7-57.7)	1.28 (1.07-1.55)	24.8 (20.9-29.9)	308 (244-412)	13.4 (9.33-17.7)
WHS	184	68.7 (59.4-79.3)	1.50 (1.24-1.83)	22.0 (18.7-26.0)	363 (280-494)	6.66 (4.72-8.82)
<b>EU/AU cohorts</b>						
ATBC	200	76.1 (65.7-86.8)	1.62 (1.37-1.93)	21.5 (19.0-24.2)	367 (291-453)	12.9 (10.3-16.6)
HUNT	193	73.7 (63.1-84.0)	1.59 (1.38-1.92)	22.0 (18.8-27.2)	319 (244-449)	9.51 (6.29-13.7)
MCCS	354	68.6 (59.7-78.1)	1.54 (1.28-1.81)	22.2 (18.8-26.4)	345 (274-451)	11.2 (8.81-14.1)
MDCS	198	63.2 (55.9-74.1)	1.40 (1.16-1.71)	21.8 (18.5-27.0)	324 (245-422)	10.7 (7.51-14.3)
NSHDS	244	61.9 (58.7-81.0)	1.41 (1.19-1.69)	22.7 (19.4-27.1)	306 (241-407)	5.89 (4.42-8.05)
<b>Asian cohorts</b>						
SCHS	422	66.9 (56.9-77.7)	1.58 (1.34-1.89)	24.0 (20.1-28.0)	365 (285-472)	16.9 (13.2-21.9)
SCS	513	75.4 (64.8-86.2)	1.53 (1.30-1.81)	20.3 (17.7-23.7)	331 (263-422)	10.8 (8.36-14.3)
SMHS	421	64.4 (56.6-73.6)	1.43 (1.24-1.68)	22.1 (19.1-26.8)	343 (267-455)	8.53 (6.13-11.9)
SWHS	419	64.2 (56.3-73.1)	1.38 (1.18-1.64)	21.7 (18.5-25.5)	363 (296-460)	8.28 (6.37-11.8)

<sup>a</sup>US: United States; WHI: The Women's Health Initiative; SCCS: The Southern Community Cohort Study; NYUWHS: The New York University Women's Health Study; CPS-II: The American Cancer Society Cancer Prevention Study-II Nutrition Cohort; CLUE I: The Campaign Against Cancer and Stroke; CLUE II: The Campaign Against Cancer Heart Disease; MEC: The Multiethnic Cohort; WHS: Women's Health Study; PHS: Physician's Health Study; NHS: The Nurses' Health Study; HPFS: Health Professional's Follow-up Study; EU: European; AU: Australian; MCCS: The Melbourne Collaborative Cohort Study; MDCS: The Malmö Diet and Cancer Study; NSHDS: The Northern Sweden Health and Disease Study; ATBC: The Alpha-Tocopherol, Beta-Carotene Cancer Prevention Study; HUNT: The Nørd-Trøndelag Health Study; SMHS: The Shanghai Men's Health Study; SWHS: The Shanghai Women's Health Study; SCHS: The Singapore Chinese Health Study; SCS: The Shanghai Cohort Study.

<sup>b</sup>KTR: Kynurenine: tryptophan ratio.

**Supplemental Table 4:** Partial Spearman correlation coefficients between two circulating biomarkers among all control subjects after adjustment for age and sex  
The Lung Cancer Cohort Consortium (LC3) Study

Biomarkers	Kynurenine	KTR <sup>a</sup>	Neopterin	Quinolinic acid
Tryptophan	<b>0.45</b>	<b>-0.43</b>	-0.01	<b>0.13</b>
Kynurenine		<b>0.55</b>	<b>0.34</b>	<b>0.66</b>
KTR <sup>a</sup>			<b>0.35</b>	<b>0.53</b>
Neopterin				<b>0.34</b>

<sup>a</sup> KTR, Kynurenine to tryptophan ratio (nmol/μmol).

Bold figures indicate that P values for the correlation coefficients were less than 0.05.

**Supplemental Table 5: Number of subjects (cases/controls) in stratified analyses**  
The Lung Cancer Cohort Consortium (LC3) Study

<b>Subset</b>	<b>Q1</b>	<b>Q2</b>	<b>Q3</b>	<b>Q4</b>	<b>Q5</b>
<b>Smoking status</b>					
<b>Current smokers</b>					
Tryptophan	494/419	441/456	475/496	556/564	553/584
Kynurenine	549/515	517/554	545/537	493/498	415/415
KTR <sup>a</sup>	648/626	535/570	465/507	457/457	414/359
Quinolinic acid	748/698	544/582	429/469	452/435	346/335
Neopterin	544/561	532/541	516/508	466/466	455/437
<b>Former smokers</b>					
Tryptophan	329/272	278/313	305/305	304/303	302/325
Kynurenine	178/191	215/208	329/289	325/351	471/479
KTR <sup>a</sup>	165/182	232/248	301/294	318/333	502/461
Quinolinic acid	155/151	206/212	240/303	383/354	534/498
Neopterin	215/228	270/260	301/299	307/349	422/379
<b>Never smokers</b>					
Tryptophan	376/382	301/304	277/272	235/206	138/163
Kynurenine	356/367	312/311	274/260	202/212	183/177
KTR <sup>a</sup>	272/265	261/255	254/272	259/283	281/252
Quinolinic acid	255/224	283/280	233/300	288/284	268/239
Neopterin	263/283	263/270	270/263	278/254	251/255
<b>Histological subtype<sup>b</sup></b>					
<b>Large cell carcinoma</b>					
Tryptophan	26/39	42/40	44/37	36/33	26/25
Kynurenine	30/37	35/35	30/34	42/36	37/32
KTR <sup>a</sup>	32/27	27/44	34/33	38/32	43/38
Quinolinic acid	29/26	41/36	26/42	41/40	37/30
Neopterin	21/24	29/34	53/40	33/43	37/32
<b>Small cell carcinoma</b>					
Tryptophan	105/80	90/94	95/102	89/112	113/104
Kynurenine	91/85	87/96	100/96	95/92	119/123
KTR <sup>a</sup>	104/92	80/104	99/85	87/105	122/106
Quinolinic acid	116/112	75/90	80/99	110/89	111/102
Neopterin	90/92	94/85	90/93	103/121	114/100
<b>Squamous cell carcinoma</b>					
Tryptophan	181/144	135/154	158/166	171/184	191/188
Kynurenine	145/135	133/174	200/173	170/175	188/179
KTR <sup>a</sup>	164/167	162/164	156/174	157/180	197/151
Quinolinic acid	150/179	186/160	145/157	174/177	181/163
Neopterin	136/153	190/157	170/183	171/170	169/172
<b>Adenocarcinoma</b>					
Tryptophan	454/453	422/447	426/439	423/343	331/374
Kynurenine	470/471	422/416	425/395	366/397	373/377
KTR <sup>a</sup>	418/411	413/410	391/396	395/405	439/434
Quinolinic acid	462/408	407/420	330/394	437/397	420/437
Neopterin	418/432	384/398	404/393	404/399	443/431
<b>Time from blood draw to cancer diagnosis<sup>c</sup></b>					
<b>&lt;2 years</b>					
Tryptophan	100/73	91/116	104/107	89/87	88/89
Kynurenine	74/80	90/92	89/93	88/101	131/106
KTR <sup>a</sup>	75/79	87/88	87/104	84/99	139/102
Quinolinic acid	76/79	82/87	64/102	108/97	142/107
Neopterin	84/94	78/78	87/87	88/81	133/130
<b>2-4.9 years</b>					
Tryptophan	155/149	177/169	183/173	164/164	153/177
Kynurenine	140/132	161/158	187/193	170/184	174/165
KTR <sup>a</sup>	161/164	156/159	148/159	168/170	199/180
Quinolinic acid	168/150	142/151	134/170	202/191	186/170
Neopterin	177/182	153/173	129/152	158/130	213/193

<b>≥ 5 years</b>					
Tryptophan	418/369	508/535	543/562	626/603	495/521
Kynurenine	502/507	548/560	590/555	506/530	444/438
KTR <sup>a</sup>	620/621	539/566	518/515	488/486	425/402
Quinolinic acid	610/570	548/581	465/526	540/510	427/403
Neopterin	609/621	549/537	514/512	464/488	450/428

<sup>a</sup>KTR: Kynurenine: tryptophan ratio.

<sup>b</sup>1,806 cases were other or unknown histological types of cancer, thus these cases and their individually matched controls were excluded from the analysis.

<sup>c</sup>1,470 cases had missing information on the time interval from blood draw to cancer diagnosis, thus these cases and their individually matched controls were excluded from the analysis.

**Figure 1: The kynurenine pathway of tryptophan metabolism**

