



Manlik, O., Krützen, M., Kopps, A. M., Mann, J., Bejder, L., Allen, S. J., ... Sherwin, W. B. (2019). Is MHC diversity a better marker for conservation than neutral genetic diversity? A case study of two contrasting dolphin populations. *Ecology and Evolution*, 9(12), 6986-6998.
<https://doi.org/10.1002/ece3.5265>

Publisher's PDF, also known as Version of record

License (if available):
CC BY

Link to published version (if available):
[10.1002/ece3.5265](https://doi.org/10.1002/ece3.5265)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the final published version of the article (version of record). It first appeared online via Wiley at <https://doi.org/10.1002/ece3.5265> . Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/pure/about/ebr-terms>

Is MHC diversity a better marker for conservation than neutral genetic diversity? A case study of two contrasting dolphin populations

Oliver Manlik^{1,2}  | Michael Krützen³ | Anna M. Kopps² | Janet Mann⁴ | Lars Bejder^{5,6} | Simon J. Allen⁷ | Celine Frère⁸ | Richard C. Connor⁹ | William B. Sherwin^{2,6}

¹Biology Department, College of Science, United Arab Emirates University, Al Ain, United Arab Emirates

²Evolution and Ecology Research Centre, School of Biological, Earth and Environmental Sciences, University of New South Wales, Sydney, New South Wales, Australia

³Department of Anthropology, University of Zurich, Zurich, Switzerland

⁴Department of Biology and Department of Psychology, Georgetown University, Washington, District of Columbia

⁵Marine Mammal Research Program, Hawai'i Institute of Marine Biology, University of Hawai'i at Manoa, Kaneohe, Honolulu

⁶Aquatic Megafauna Research Unit, School of Veterinary and Life Sciences, Murdoch University, Murdoch, Western Australia, Australia

⁷School of Biological Sciences, University of Bristol, Bristol, United Kingdom

⁸Faculty of Science, Health, Education and Engineering, University of the Sunshine Coast, Sippy Downs, Queensland, Australia

⁹Biology Department, UMASS-Dartmouth, Dartmouth, Massachusetts

Correspondence

Oliver Manlik, Biology Department, College of Science, United Arab Emirates University, P.O. Box 15551, Al Ain, United Arab Emirates.

Email: oliver.manlik@uaeu.ac.ae

Funding information

WV Scott Foundation; National Science Foundation, Grant/Award Number: 0918308, 0941487 and 1559380; Sea World Research and Rescue Foundation, Grant/Award Number: RM09107; NSERC; National Geographic Society; PADI Foundation; Australian Research Council; Shark Bay Shire; United Arab Emirates University

Abstract

Genetic diversity is essential for populations to adapt to changing environments. Measures of genetic diversity are often based on selectively neutral markers, such as microsatellites. Genetic diversity to guide conservation management, however, is better reflected by adaptive markers, including genes of the major histocompatibility complex (MHC). Our aim was to assess MHC and neutral genetic diversity in two contrasting bottlenose dolphin (*Tursiops aduncus*) populations in Western Australia—one apparently viable population with high reproductive output (Shark Bay) and one with lower reproductive output that was forecast to decline (Bunbury). We assessed genetic variation in the two populations by sequencing the MHC class II DQB, which encompasses the functionally important peptide binding regions (PBR). Neutral genetic diversity was assessed by genotyping twenty-three microsatellite loci.

We confirmed that MHC is an adaptive marker in both populations. Overall, the Shark Bay population exhibited greater MHC diversity than the Bunbury population—for example, it displayed greater MHC nucleotide diversity. In contrast, the difference in microsatellite diversity between the two populations was comparatively low.

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2019 The Authors. *Ecology and Evolution* published by John Wiley & Sons Ltd.

Our findings are consistent with the hypothesis that viable populations typically display greater genetic diversity than less viable populations. The results also suggest that MHC variation is more closely associated with population viability than neutral genetic variation. Although the inferences from our findings are limited, because we only compared two populations, our results add to a growing number of studies that highlight the usefulness of MHC as a potentially suitable genetic marker for animal conservation. The Shark Bay population, which carries greater adaptive genetic diversity than the Bunbury population, is thus likely more robust to natural or human-induced changes to the coastal ecosystem it inhabits.

KEYWORDS

adaptive genetic variation, bottlenose dolphin, cetacean, conservation genetics, major histocompatibility complex, microsatellites

1 | INTRODUCTION

A loss of genetic diversity is often associated with reduced fitness and can negatively impact population viability (Chapman, Nakagawa, Coltman, Slaters, & Sheldon, 2009; Frankham, Ballou, & Briscoe, 2010; Reed & Frankham, 2003). Until recently, studies that assessed genetic diversity in wild animal populations typically used adaptively neutral genetic markers, such as microsatellites. However, neutral genetic markers offer little insight into the adaptive potential to cope with natural and artificial change (Allendorf & Luikart, 2008; Hedrick, 2005; Holderegger, Kamm, & Gugerli, 2006). Therefore, to assess genetic diversity that captures information relevant to the conservation of populations, it is prudent to use genetic markers linked to ecologically important traits (Manlik, Schmid-Hempel, & Schmid-Hempel, 2017; Piertney & Webster, 2010; van Tienderen, Haan, Linden, & Vosman, 2002).

One such adaptive marker is the major histocompatibility complex (MHC) (reviewed by Sommer, 2005). The MHC plays an important role in responding to antigens and initiating an immune response in vertebrates. Major histocompatibility complex variation has been associated with various fitness traits, including factors important for population viability, such as resistance to parasites, survival, and reproductive success (Hedrick, 2003; Kalbe et al., 2009; Kurtz et al., 2004; Sepil, Lachish, Hink, & Shelton, 2013; Sepil, Lachish, & Sheldon, 2013; Thoss, Ilmonen, Musolf, & Penn, 2011; Wegner, Kalbe, Milinski, & Reusch, 2008). High levels of MHC diversity observed across a variety of vertebrate species are commonly explained by balancing selection (Garrigan & Hedrick, 2003). Balancing selection maintains high levels of MHC diversity by two possible, not mutually exclusive, mechanisms: frequency-dependent selection (Borghans, Beltman, & Boer, 2004) and heterozygote advantage (Doherty & Zinkernagel, 1975). The frequency-dependent selection model suggests that MHC diversity is pathogen-mediated, because rare MHC variants are selected for by host-pathogen co-evolution. In contrast, heterozygote advantage explains balancing selection due to heterozygotes having greater fitness than homozygotes.

Compared to terrestrial vertebrates, relatively little is known about MHC diversity in cetaceans, and the extent to which cetacean MHC diversity is associated with population viability remains uncertain. The vaquita (*Phocoena sinus*) population, endemic to the Gulf of California, showed low levels of MHC II variation (Munguia-Vega et al., 2007) and is now considered functionally extinct (Taylor et al., 2017). In contrast, the extinct baiji (*Lipotes vexillifer*) of the Yangtze River exhibited very high MHC diversity (Xu et al., 2012; Yang, Yan, Zhou, & Wei, 2005). Reduced MHC diversity may not necessarily adversely affect population viability (Radwan, Biedrzycka, & Babik, 2010). Caveats for many of these studies are that they had no baseline measure of genetic diversity in a conspecific viable population or no comparison of MHC and other types of genetic variation. No study to date has compared MHC and neutral genetic diversity of conspecific cetacean populations that differ with respect to population parameters and viability forecasts.

In this study, we used two genetic markers, MHC and neutral microsatellites, to assess genetic diversity of two contrasting bottlenose dolphin (*Tursiops aduncus*) populations—one in Shark Bay (SB) and another off Bunbury (BB), Western Australia (Figure 1). These two populations, more than 1,000 km apart (Figure 2), are not connected by dispersal. Each population exhibits limited genetic exchange with its neighboring populations (Allen et al., 2016; Manlik et al., 2018). The two populations differ greatly with respect to population viability. A comparative population viability analysis showed that the SB population appeared stable, but the BB population was forecast to decline with a high probability of extinction, unless supported by immigration (Manlik et al., 2016). The large difference in viability between the two populations was best explained by considerable differences in reproductive rates (Manlik et al., 2016). Besides this difference in reproductive rates, the two populations also differ with respect to anthropogenic pressure (Manlik et al., 2016). The SB population occurs in a remote UNESCO World Heritage area with markedly lower anthropogenic activity, whereas BB inhabits waters adjacent to an expanding regional city and port with comparatively high vessel traffic (Manlik, 2019; Manlik et al.,



FIGURE 1 Mother and calf bottlenose dolphin (*Tursiops aduncus*) in Shark Bay, a UNESCO World Heritage Site in Western Australia. Photograph: Ewa Krzyszczyk

2016; Nicholson, Bejder, Allen, Krützen, & Pollock, 2012; Smith, Frère, Kobryn, & Bejder, 2016; Sprogis et al., 2018).

SB and BB also differ with respect to reported population sizes. SB population size was estimated by aerial surveys to be about 2,000–3,000 individuals (minimum estimates; Preen, Marsh, Lawler, Prince, & Shepherd, 1997) in a 14,900 km² area, but other studies investigating various sections of SB suggest that the population may be much larger (e.g. Nicholson et al., 2012). BB population size was assessed to be approximately 260 individuals for the 120 km² area (Manlik et al., 2016). However, smaller seasonal abundance estimates have been reported for BB (Smith, Pollock, Waples, Bradley, & Bejder, 2013; Sprogis et al., 2016). Different methodologies to estimate population sizes, and the issue of connectivity, make comparison difficult, but all studies suggest that SB is substantially larger than BB.

The aim of this study was to compare MHC II DQB genetic diversity and microsatellite diversity between these two contrasting dolphin populations. Given that only few MHC studies have been conducted on populations with differing reproductive success or population forecasts, this provided a rare opportunity to compare MHC and neutral genetic diversity between two natural populations with considerable differences in viability. If MHC variation reflects differences in fitness, and given the large difference in reproductive output between the two populations (Manlik et al., 2016), we would expect to observe a larger inter-population difference in MHC diversity than in microsatellite diversity. Additionally, to assess evolutionary and ecologically relevant genetic variation, we evaluated signals of selective pressure on MHC II DQB. We did this by assessing nonsynonymous versus synonymous nucleotide substitutions (Nei & Gojobori, 1986), whether substitutions occurred at codons expressing antigen-binding residues, and by performing a Tajima's *D* test (Tajima, 1989).

2 | MATERIALS AND METHODS

2.1 | Sample collection and DNA extraction

Between 1997 and 2013, we opportunistically collected skin samples from free-ranging bottlenose dolphins in Shark Bay (SB) and off Bunbury (BB), Western Australia (Figure 2), using a biopsy system designed for small cetaceans (Krützen et al., 2002). Tissue samples were stored in a saturated NaCl/20% (v/v) dimethyl sulfoxide solution for DNA stabilization (Amos & Hoelzel, 1991). We isolated genomic DNA following standard phenol–chloroform protocol (Davis, Dibner, & Battey, 1986), or alternatively using the Genra Puregene Tissue Kit (Qiagen).




FIGURE 2 (a) Shark Bay, a UNESCO World Heritage area, is about 13,000 km² in size and is divided by the Peron Peninsula, which bisects the bay into a western and an eastern gulf. Sampling sites included a 300 km² area (circled) north of Monkey Mia and an area of ca. 260 km² (circled) in the western gulf. (b) The inset shows the relative location of the study sites (Shark Bay & Bunbury). The coastal study area of Bunbury covers about 120 km² and extends approximately 1.5 km offshore with a linear distance of 50 km. The study site includes the coastal areas, embayment, Leschenault Inlet and outer harbors (5 km²), estuary and river mouth (15 km²). Transects of the outer-water Bunbury study site are shown. These figures are modified from Figure S1 of Manlik et al. (2016)

Sampling in SB included two sites in western and eastern SB (Figure 2), that are connected by extensive gene flow (Krützen, Sherwin, Berggren, & Gales, 2004) and appear to form one large continuous population. A total of 686 and 125 dolphins were biopsied in SB and off BB, respectively. Sex of individuals was determined by various methods, as described by Sprogis et al. (2016), including genetic sexing (Baker et al., 1998). We performed chi-square tests to assess whether the numbers of males and females in the samples were significantly different from those in the surveyed populations or different from an expected 50:50 male to female ratio. Sex ratios for surveyed individuals versus sampled individuals were not significantly different (SB: $\chi^2 = 0.42$, $p = 0.515$; BB: $\chi^2 = 0.16$, $p = 0.693$) nor were the ratios of sampled individuals significantly different from 50:50 (SB: $\chi^2 = 0.10$, $p = 0.757$; BB: $\chi^2 = 1.7$, $p = 0.190$). To assess whether it was justified to pool samples collected from eastern and western SB, we estimated subpopulation fixation index (F_{ST}) based on microsatellite data, using GENALEX 6.501 (Peakall & Smouse, 2006, 2012), and compared MHC and microsatellite diversity between the two sampling locations.

Data were collected under research permits (SF005997; SF006538; SF007046; SF007596; SF008480; SF009119) licensed by the Western Australian Department of Environment and Conservation (now the Western Australian Department of Parks and Wildlife). This study was carried out in accordance with the Murdoch University Animal Ethics Committee approval (W2076/07; W2307/10; W2342/10).

2.2 | Amplification, Sanger sequencing, and sequence variant determination

To characterize MHC genetic variants of the two populations, we amplified and sequenced the MHC II DQB exon 2 (hereafter MHC DQB), which encompasses the functionally important PBR (Baker et al., 2006; Hayashi et al., 2003; Hoelzel, Stephens, & O'Brien, 1999; Murray, Malik, & White, 1995; Seddon & Ellegren, 2002), and which is the region under strongest selection (Hughes & Nei, 1989). Amplification was performed using the universal primer pair DQB1 and DQB2, 5'CATGTGCTACTTCACGTTCCGG 3' (forward), 5'CTGGTAGTTGTGTCTCCACAC 3' (reverse), which were originally designed by Tsuji, Aizawa, and Sasaki (1992), and previously used to amplify cetacean MHC (Caballero et al., 2010; Du, Zheng, Wu, Zhao, & Wang, 2010; Hayashi et al., 2003, 2006; Heimeier et al., 2009; Moreno-Santillán, Lacey, Gendron, & Ortega, 2016; Murray et al., 1995; Vassilakos, Natoli, Dahlheim, & Hoelzel, 2009).

PCR for MHC DQB was performed using 1.25 μ M primers, 0.2 mM dNTPs, 1.0 mM $MgCl_2$, 1.25 U GoTaq DNA polymerase (Promega), and 20–100 ng (5 μ l) of template DNA in a total volume of 25 μ l. Thermal cycling was conducted on an Eppendorf Mastercycler (ep gradient S) with an initial denaturing temperature of 95°C for 15 min, 30–35 cycles of denaturation at 95°C (1 min), and annealing at 55°C (30 s), followed by an elongation step at 72°C for 1 min.

PCR products were visualized by electrophoresis on a 1.5% agarose gel (1 \times TBE buffer) stained with GelRed™ (Biotium). All MHC amplicons were sequenced in the forward and reverse direction using Big Dye 3.1 on a 3730xl DNA Analyzer (Applied Bioscience) at the Ramaciotti Centre of the University of New South Wales.

MHC DQB Sanger sequences (172 bp; forward and reverse) were aligned with CLUSTALW (Thompson, Higgins, & Gibson, 1994) in GENEIOUS 6.1 (Drummond et al., 2010). Double-peaks were called using the "Heterozygotes" plugin (GENEIOUS) based on the default threshold of 50% peak height and double-checked by visual inspection. Subsequently, MHC DQB sequence variants were inferred by reconstructing haplotype phases from the unphased sequence alignment data using the coalescent-based Bayesian method PHASE (Stephens & Donnelly, 2003; Stephens, Smith, & Donnelly, 2001) in DNASP version 5.10.01 (Librado & Rozas, 2009) with 100 iterations, 1 thinning interval and 100 burn-in iterations. PHASE was shown to be reliable for reconstructing haplotypes (Stephens & Donnelly, 2003), including MHC haplotypes (Bos, Copurenko, Williams, & DeWoody, 2008; Bos, Turner, & DeWoody, 2007; Silva & Edwards, 2009). After haplotype reconstruction, the MHC DQB alignments in DNASP contained sequences for 276 SB and 65 BB individuals. We performed a BLASTN search to compare inferred MHC DQB sequence variants to sequences in the NCBI database.

2.3 | Assessing signals of selection acting on MHC DQB

To assess signals of selection, we compared rates of nonsynonymous (d_N) and synonymous (d_S) substitutions within the 172-bp MHC DQB region. We used the Nei–Gojobori method (Nei & Gojobori, 1986) for a codon-based test of positive selection (two-sided z-test) implemented in MEGA version 7.08 (Kumar, Stecher, & Tamura, 2016) to test whether $d_N > d_S$ for (a) all codons of the entire sequence; (b) codons of the putative peptide binding region (PBR), that is, variable codons that code for amino acids that have been reported to bind to antigens; and (c) putative nonpeptide binding regions (non-PBR). Variance estimation for the z-test was based on 1,000 bootstrap replicates. Additionally, we used DNASP to perform Tajima's *D* test (Tajima, 1989), which detects departure from selective neutrality or historical changes in population size.

2.4 | Assessment of MHC sequence diversity

After alignment in GENEIOUS, we compared sequence variation using DNASP 5.10.01 (Librado & Rozas, 2009). We recorded the following measures of sequence variation: (a) nucleotide diversity (π), as described by Nei (1987) (equation 10.5), (b) haplotype diversity (H_d) (Nei, 1987; equation 8.4), (c) Watterson mutation estimator (Θ_W), according to Watterson (1975) (equation 1.4), and (d) the mutation parameter, theta (Θ_{Eta}) per nucleotide site from the total number of mutations (Nei, 1987; equation 10.3).

Sampling variances and standard deviations were calculated for nucleotide diversity and haplotype diversity according to Nei (1987)

and for Watterson mutation estimator according to Tajima (1993). We also calculated standard errors of the mean between the three SB conservative samples and across all subsamples (SB: 19 subsamples; BB: 5 subsamples). We used *t* tests to compare the mean π , H_d , Θ_{W^*} and Θ_{Eta} values between the two populations across all subsamples.

2.5 | Assessment of microsatellite diversity

All sampled BB individuals were previously genotyped for 25 polymorphic microsatellite loci (Manlik et al., 2018). We followed the same procedure and checks for genotyping individuals of the SB population as described in Manlik et al. (2018): We used previously tested primers for polymorphic microsatellite loci (Hoelzel, Potter, & Best, 1998; Kopps et al., 2014; Krützen, Valsecchi, Connor, & Sherwin, 2001; Nater, Kopps, & Krützen, 2009; Shinohara, Domingo-Roura, & Takenaka, 1997). All primer sequences used in this study are listed in Dryad/Table S1. Microsatellite amplification was performed using the Qiagen Multiplex KitTM in three multiplex PCR reactions as described in Manlik et al. (2018). Fragment analysis of PCR amplicons was performed on a 3730XL DNA Analyzer (Applied Biosystems), employing a Genescan-500 LIZTM size standard. Alleles were scored using GENEMAPPER 4.0 (Applied Biosystems) and the microsatellite plugin for GENEIOUS 6.0 (Drummond et al., 2010). We used MicroChecker version 2.2.3 (van Oosterhout, Hutchinson, Wills, & Shipley, 2004) to test for scoring errors due to stuttering and the presence of large-allele dropouts across all loci and populations. The software INEst version 2.0 (Chybicki & Burczyk, 2009) was used to estimate the frequency of null alleles at microsatellite loci in each population. Linkage disequilibrium for all microsatellite locus pairs was tested with GENEPOP version 4.5.1 (Rousset, 2008). We used GenAIEx 6.501 (Peakall & Smouse, 2006, 2012) to test all loci for departures from Hardy-Weinberg equilibrium (HWE).

Microsatellite diversity was summarized by measuring observed heterozygosity (H_o), expected heterozygosity relative to HWE (H_e), the number of effective alleles (A_e), and Shannon's Index (1H) (Brown & Weir, 1983; Sherwin, Chao, Jost, & Smouse, 2017), using GenAIEx 6.501. We used paired *t* tests to compare the mean values of these measures between the two populations across the microsatellite loci.

2.6 | Sampling for comparison of inter-population genetic diversity

Due to the sample-size difference between SB and BB, we used three sampling approaches to compare genetic diversity between the two populations:

1. Maximum sampling: We sampled the maximum number of individuals for which we obtained MHC DQB sequences or microsatellite genotypes. For SB, this approach included 276 individuals for which we obtained MHC DQB sequences and 667 individuals for which we genotyped for the microsatellite

loci. For BB, the maximum sampling included MHC sequences of 65 individuals and microsatellite genotypes of 84 individuals.

2. Conservative sampling: We first reduced the maximum sample set to only include individuals for which we had both MHC DQB and microsatellite data. This resulted in 55 samples for BB and 239 for SB. In order to compare equal sample sizes that reflect the demography of the two populations, we further subsampled the SB data to include the same number of each of the age classes (calves, juveniles, and adults) and sexes that were found in the conservative BB sample (Dryad/Table S2). From the SB sample set that included 239 individuals, we obtained three subsamples (SB samples 1–3), each containing the same numbers for each of the age classes and sexes found in the BB conservative sample. We did this by randomly choosing from the SB samples 2 calves, 15 juveniles, and 38 adults, of which 32 were males and 23 were females. Individuals of unknown age classes or sexes were excluded. Each of the individuals was only sampled once, for example individuals included in SB sample 1 were not included in SB sample 2 or 3.
3. Subsampling: In order to allow for statistical comparisons of MHC diversity measures between SB and BB, we subsampled both populations by randomly choosing 11 samples from each conservative sampling set of each population. Each sample was only included once in each subsample. This generated 19×11 (209) subsamples for SB and 5×11 (55) subsamples for BB.

Other methods, such as rarefaction, are often used to investigate the effect of sample size, but we believe our three sampling approaches address this more thoroughly.

3 | RESULTS

Pooling the eastern and western SB datasets was justified because the subpopulation fixation index (F_{ST}) comparing the two sampling sites in SB showed very little differentiation ($F_{ST} = 0.006$; Dryad/Table S3), indicating that the two sites represent one population. Also, MHC and microsatellite diversity of the two SB sampling sites were similar (Dryad/Table S3).

3.1 | Sequence variants of MHC DQB

Forward and reverse MHC DQB sequences of a total 341 individuals (SB: 276; BB: 65) were analyzed. Totals of 186 and 43 MHC DQB sequence variants were inferred by haplotype reconstruction for SB and BB, respectively. We did not detect any patterns in the sequences that indicated multiple allelism (i.e. having more than two alleles or sequence variants per amplicon/individual), gene duplications, stop codons, or frameshifts. Comparing MHC DQB sequences of seven mother-father-offspring trios (Kopps, 2007) did not reveal any patterns that were inconsistent with single-locus Mendelian inheritance. Nonetheless, we refrain from classifying these inferred sequence variants as novel MHC alleles,

TABLE 1 The estimated rates of nonsynonymous (d_N) and synonymous (d_S) substitutions (\pm standard errors of the mean) for putative peptide binding regions (PBR) and nonpeptide binding regions (non-PBR) and their ratios for DQB exon 2 in the Shark Bay (SB) and Bunbury (BB) dolphin population

Pop.	Sites	N	d_N	d_S	d_N/d_S	z	p
SB	PBR	17	6.334 \pm 1.196	3.974 \pm 1.439	1.59	4.563	0.0000061
	Non-PBR	39	1.539 \pm 0.205	1.659 \pm 0.446	0.93	0.276	ns
	All	56	3.500 \pm 0.558	3.095 \pm 0.778	1.13	2.976	0.0018
BB	PBR	17	2.297 \pm 0.536	1.362 \pm 0.723	1.69	3.033	0.0015
	Non-PBR	39	0.244 \pm 0.0849	0.197 \pm 0.134	1.24	1.518	ns
	All	56	0.993 \pm 0.224	0.694 \pm 0.309	1.43	3.023	0.0015

Note: N is the number of codons in each category. The p-value is the significance value for the difference between d_N and d_S , using a two-sided z-test.

which are commonly confirmed by sequencing clones (Marsh et al., 2010) or by re-genotyping all individuals with rare haplotypes (Ahmad et al., 2002). Thirty-nine individuals were homozygous for all 172 bps for one of four unique sequences (Dryad/Figure S1). All sequences showed high similarity (98%–100%) to published MHC DQB alleles in dolphins (Dryad/Table S4).

3.2 | Signals of selection acting on MHC

We detected signals of selection acting on MHC DQB. In both populations, nonsynonymous (d_N) substitution rates were significantly greater than the synonymous (d_S) substitutions rates in the entire 172-bp region and in the putative PBR, but not in the non-PBR (Table 1). About 82% of the variable nucleotide sites (18 out of 22) are within codons that have been associated with the PBR (Dryad/Figure S1). Notably, the large majority of variable nucleotide sites were detected within the PBR (Dryad/Figure S1). Tajima's D was near zero for MHC DQB in both populations (Table 2). However, in the SB population it significantly departed from zero ($D = -1.82$, $p < 0.05$) when considering only nonsynonymous substitutions (Table 2).

3.3 | MHC sequence diversity

Regardless of the sampling approach (maximum, conservative, or subsampling), dolphins of SB showed greater MHC DQB diversity than those of BB, except with respect to Hd , which showed no significant difference (Figure 3; Table 3). On the basis of the conservative sampling approach, mean π of the SB population was 0.066 ($SE = 0.0022$), substantially greater than that of BB ($\pi = 0.053$, $SE = 0.0018$; Figure 3a). In comparison with BB, the SB population showed larger Θ_W (SB: $\Theta_W = 0.0801$, $SE = 0.0047$; BB: $\Theta_W = 0.0496$, $SE = 0.0029$; Figure 3c). Theta (Θ_{Eta}) was substantially larger for SB

($\Theta_{Eta} = 0.100$, $SE = 0.0077$) than for BB ($\Theta_{Eta} = 0.063$, $SE = 0.0049$; Figure 3d). Based on the subsampling approach, π , Θ_W and Θ_{Eta} were significantly greater for SB than for BB (Figure 3; Table 5).

3.4 | Microsatellite diversity

Neither population showed evidence for scoring errors due to stuttering or large-allele dropouts for any of the microsatellite loci. We also did not detect evidence for null alleles for any of the loci. Linkage disequilibrium tests with Genepop showed that, among all comparisons of pairs of microsatellite loci, one pair appeared linked (Tur4_105 & MK8), but this linkage was only observed for BB, so it was unlikely to be a result of physical linkage. Departures from HWE expectations were observed for two microsatellite loci, Tur4_98 (SB) and KWM12 (SB & BB), after Bonferroni correction. Those two loci were removed from subsequent analysis. Consequently, all subsequent results are based on 23 loci.

In contrast to the MHC results, microsatellite diversity showed no significant differences between SB and BB. However, there was a nonsignificant trend of SB being genetically more diverse than BB with respect to H_o , H_e , A_e , and 1H (Figure 4a–d; Tables 4 and 5). The conservative sampling approach showed average numbers of alleles per microsatellite locus of 5.68 in SB and 4.30 in BB; H_o of 0.546 in SB and 0.588 in BB ($t = 0.6482$, $df = 22$; $p = 0.5236$; Figure 4a; Tables 4 and 5); H_e of 0.578 in SB and 0.559 in BB ($t = 0.5508$, $df = 22$, $p = 0.5873$; Figure 4b; Tables 3 and 5); A_e per locus of 2.98 ($SE = 0.35$) for SB and 2.70 ($SE = 0.23$) for BB ($t = 1.011$, $df = 22$, $p = 0.3231$; Figure 4c; Tables 4 and 5); 1H of 1.17 in SB and 1.04 in BB ($t = 1.752$, $df = 22$, $p = 0.0938$; Figure 4d; Tables 4 and 5). There was also no significant difference between the SB subsamples with respect to the microsatellite measures of genetic diversity (Dryad/Table S5). Results of paired t tests for all sampling approaches are shown in Table 5.

TABLE 2 Results of Tajima's D tests performed on all nucleotide sites and nonsynonymous sites within the MHC DQB of the Shark Bay (SB) and Bunbury (BB) dolphin population

MHC DQB region	SB			BB			
	D	p	Sig.	D	p	Sig.	
All sites	-1.54	0.10	$p > 0.05$	ns	-0.55	$p > 0.10$	ns
Nonsynonymous sites	-1.82	<0.05	sig.	-0.81	$p > 0.10$	ns	ns




FIGURE 3 MHC DQB genetic diversity of dolphins in Shark Bay (SB; blue bars) and Bunbury (BB; red bars) (a) nucleotide diversity (π), (b) haplotype diversity (Hd), (c) Watterson mutation estimator from variable sites (Θ_W), and (d) theta per site from Eta (Θ_{Eta}). Each pairing of SB-BB bars represents the results on the basis of each of the three sampling approaches: maximum sampling (SB^a, BB^a), conservative sampling (SB^b, BB^b), and subsampling (SB^c, BB^c). Whiskers depict the respective standard errors of the mean, which are only shown for means across subsamples. Significant values based on *t* tests: sig. * $p < 0.05$; sig. *** $p < 0.0005$; n.s. = nonsignificant ($p > 0.05$)

4 | DISCUSSION

Compared to selectively neutral genetic variation, variation of adaptive genes, such as those of the MHC, is a better proxy for genetic diversity relevant to population viability (Oliver & Piertney, 2012; Sommer, 2005; Ujvari & Belov, 2011). A loss of adaptive genetic diversity reduces reproductive success and survival in the short-term and ultimately diminishes the evolutionary potential of populations


to adapt to environmental changes (Frankham, 2005; Frankham et al., 2010). Our results show that the more stable SB population, which displayed greater reproductive success (Manlik et al., 2016), harbors greater MHC diversity compared with the BB population that was forecast to decline. It is important to note that this was the case, regardless of the sampling approach. Our finding that microsatellites do not show any significant differences between SB and BB suggests that the higher MHC diversity in SB is unlikely due to differences in

TABLE 3 MHC II DQB diversity measures based on conservative and maximum sampling approaches

Pop	Sampling approach	<i>n</i>	π	<i>Hd</i>	Θ_W	Θ_{Eta}	<i>Eta</i>
Bunbury	BB max.	65	0.0538 (0.0025)	0.939 (0.013)	0.0481 (0.013)	0.0652	61
	BB cons.	55	0.0535 (0.0025)	0.941 (0.014)	0.0496 (0.014)	0.0629	57
	BB subsampling mean (5 × 11)	11	0.0533 (0.0010)	0.914 (0.015)	0.0514 (0.039)	0.0581	35.2
Shark Bay	SB max.	276	0.0660 (0.0014)	0.932 (0.0084)	0.0903 (0.019)	0.1376	163
	SB cons. 1	55	0.0665 (0.0026)	0.943 (0.016)	0.0772 (0.020)	0.0948	86
	SB cons. 2	55	0.0701 (0.0032)	0.955 (0.014)	0.0893 (0.023)	0.1147	104
	SB cons. 3	55	0.0625 (0.0195)	0.886 (0.025)	0.0739 (0.020)	0.0893	81
	SB cons. mean (sample 1–3)	55	0.0664	0.928	0.0801	0.0996	90.3
	SB cons. SE (sample 1–3)	55	0.00219	0.0213	0.00469	0.00770	70.0
	SB subsampling mean (5 × 11)	11	0.0663 (0.011)	0.910 (0.011)	0.0625 (0.022)	0.0706	43.8

Note: Measure of MHC II DQB diversity for all samples and subsamples for Shark Bay (SB) and Bunbury (BB). BB max. and SB max are based on the maximum number of samples for which we obtained MHC sequences. BB cons. is based on the samples for which we also have microsatellite data. SB samples 1–3 are based on subsamples that include the same number of calves, juveniles, adults, and females as in the BB cons. set of samples and for which we also have microsatellite data. SB cons. The final two rows show the mean values and standard errors of the mean, respectively, across all three subsamples (SB samples 1–3). Tabulated are *n* = the sample size; π = nucleotide diversity; *Hd* = haplotype diversity; Θ_W = Watterson mutation estimator; Θ_{Eta} = the mutation parameter theta based on number of mutations, *Eta*. Standard deviations for π , *Hd*, and Θ_W are shown in parentheses.

FIGURE 4 Microsatellite diversity (23 polymorphic loci), of dolphins in Shark Bay (SB; blue bars) and Bunbury (BB; red bars): (a) observed heterozygosity (H_o), (b) expected heterozygosity (H_e), (c) effective number of alleles (A_e), and (d) Shannon index (1H). Each pairing of SB-BB bars represents the results on the basis of each of the three sampling approaches: maximum sampling (SB^a, BB^a), conservative sampling (SB^b, BB^b), and subsampling (SB^c, BB^c). Whiskers depict the standard errors of the mean across the 23 loci. Significant values based on paired *t* tests: n.s. = nonsignificant ($p > 0.05$)



population size, because the resultant genetic drift is expected to affect MHC and microsatellites equally (although see Eimes et al., 2011). Therefore, it seems likely that other interactions, such as differential fitness or parasite pressure, are driving the observed MHC pattern.

The number of sequence variants we detected in both populations is unusually high, but a high number of single-locus MHC class II variants have been detected in other cetacean populations as well (e.g. Xu et al., 2012). As mentioned in the Methods, we refrain from classifying the sequence variants as novel MHC alleles, but having

TABLE 4 Microsatellite diversity measures

Pop	Sampling approach	<i>n</i>	H_o	H_e	A_e	1H
Bunbury	BB max.	84	0.59 ^a (0.041) ^a	0.57 ^a (0.039) ^a	2.76 ^a (0.24) ^a	1.058 ^a (0.09) ^a
	BB cons.	55	0.588 (0.0465)	0.559 (0.0424)	2.696 (0.228)	1.037 (0.0938)
	BB subsampling mean (5 × 11)	11	0.590 (0.0465)	0.540 (0.0410)	2.577 (0.204)	0.978 (0.0884)
Shark Bay	SB max.	676	0.579 (0.0378)	0.598 (0.0383)	3.149 (0.379)	1.220 (0.104)
	SB cons. 1	55	0.551 (0.0422)	0.561 (0.0414)	2.886 (0.347)	1.135 (0.105)
	SB cons. 2	55	0.566 (0.0425)	0.587 (0.0382)	3.027 (0.352)	1.184 (0.101)
	SB cons. 3	55	0.576 (0.0404)	0.585 (0.0389)	3.026 (0.354)	1.185 (0.104)
	SB cons. Mean (SB cons. 1–3)	55	0.564 (0.0402)	0.578 (0.0388)	2.980 (0.348)	1.168 (0.102)
	SB cons. SE (SB cons. 1–3)	55	0.0073	0.0084	0.047	0.017
	SB subsampling mean (19 × 11)	11	0.572 (0.0393)	0.561 (0.0361)	2.800 (0.275)	1.083 (0.0895)

Note: Measure of microsatellite diversity based on the three sampling approaches (maximum, conservative and subsampling) for Shark Bay (SB) and Bunbury (BB). BB max.* and SB max. are based on the maximum number of samples for which we obtained microsatellite genotypes. BB cons. are based on the samples for which we also have MHC II DQB data. SB cons. samples 1–3 are based on the conservative sampling that include the same number of calves, juveniles, adults, and females as in the BB cons. set of samples and for which we also have microsatellite data. SB cons. The rows “SB cons. Mean” and “SB cons. SE” show the mean values and standard errors of the mean, respectively, across the three conservative SB subsamples (SB samples 1–3). The rows “subsampling mean” show the mean values based on the subsampling approach. Tabulated are *n* = the sample size; H_o = observed heterozygosity; H_e = expected heterozygosity based on Hardy-Weinberg expectations; A_e = the number of effective alleles; 1H = Shannon index value. Standard errors across the microsatellites are shown in parentheses.

^aMicrosatellite diversity measures based on the BB maximum sampling approach are from Manlik et al. (2018).

TABLE 5 Results of *t* tests comparing genetic diversity measures between Shark Bay and Bunbury for MHC and microsatellites based on the various sampling approaches

MHC subsampling	<i>t</i>	<i>df</i>	<i>p</i>
π	4.303	22	0.0003
<i>H_d</i>	0.191	22	0.8503
Θ_W	2.299	22	0.0314
Θ_{Eta}	2.234	22	0.0359
msat maximum sampling	<i>t</i>	<i>df</i>	<i>p</i>
<i>H_o</i>	0.1364	22	0.8928
<i>H_e</i>	1.015	22	0.3213
<i>A_e</i>	1.182	22	0.2498
¹ <i>H</i>	2.073	22	0.0501
msat conservative sampling	<i>t</i>	<i>df</i>	<i>p</i>
<i>H_o</i>	0.6482	22	0.5236
<i>H_e</i>	0.5485	22	0.5882
<i>A_e</i>	1.011	22	0.3231
¹ <i>H</i>	1.752	22	0.0938
msat subsampling	<i>t</i>	<i>df</i>	<i>p</i>
<i>H_o</i>	0.4743	22	0.6400
<i>H_e</i>	0.6572	22	0.5179
<i>A_e</i>	1.003	22	0.3268
¹ <i>H</i>	1.529	22	0.1406

Note: Results for microsatellites (msat) are based on paired *t* tests comparing diversity values across 23 loci. Significant *p*-values (*p* < 0.05) are shown in bold.

followed the same methodology of inferring sequence variants for both populations allowed us to compare MHC sequence variation between the two populations. Further confirmation of alleles could be achieved by sequencing clones (Marsh et al., 2010) or by re-genotyping all individuals with rare haplotypes (Ahmad et al., 2002).

4.1 | Potential factors contributing to the inter-population differences in MHC diversity

Differences in MHC diversity between the two populations might be related to fitness. Adult females in SB displayed higher reproductive success than BB females (Manlik et al., 2016), and preliminary data suggest that SB females with greater reproductive success also exhibit greater MHC DQB diversity than females with low reproductive success (Manlik, 2016). Another selective effect associated with the inter-population difference in MHC diversity is differences in pathogen communities. High levels of MHC diversity can be maintained by balancing selection due to MHC's function in binding to pathogen-derived antigens (Eizaguirre, Lenz, Kalbe, & Milinski, 2012; Takahata & Nei, 1990; Wegner, Reusch, & Kalbe, 2003). The signal that we detected by the d_N/d_S analyses relates to long periods, with time for mutations to accumulate very slowly, at a rate of about 10^{-9}

per generation per nucleotide site. These patterns were originally proposed for differentiation between species, but the same patterns are expected for variation within a single population, though weaker (Kryazhimskiy & Plotkin, 2008). The higher ratios of nonsynonymous to synonymous substitutions that we observed in the MHC DQB region of both populations are consistent with balancing selection (Kimura, 1977; Yang & Bielawski, 2000). There are studies on numerous vertebrate taxa that show an association between pathogen load, infectivity, and MHC diversity (e.g., Paterson, Wilson, & Pemberton, 1998; Sepil, Lachish, Hink, et al., 2013; Wegner et al., 2008). Vassilakos et al. (2009) proposed that differential pathogen pressure across the range of cetacean populations could explain geographic variation in MHC diversity.

Although the d_N/d_S analyses can detect balancing selection over long periods, on shorter time scales, there might be other influences, such as bottlenecks, or directional selection due to a recent change in pathogen load; these can be detected by Tajima's *D*, with the proviso that because it is sensitive to demographic and selective effects, they could cancel each other out. Bottleneck dolphin mortalities due to pathogens, such as the cetacean morbillivirus, have been reported in Western Australia (Stephens et al., 2014), and outbreaks are associated with high mortality (van Bressemer et al., 2014; Di Guardo & Mazzariol, 2014). If this mortality is selective, then it could give a signal with Tajima's *D*, unless counteracted by some demographic effect. However, little is known about pathogen communities across geographic locations, including the two sites of this study. Other factors are unlikely to explain the differences in MHC diversity between SB and BB: Age and sex are unlikely because our sample sizes had equal numbers of each age class and sex; effects of mate choice (Kamiya, O'Dwyer, Westerdahl, Senior, & Nakagawa, 2014; Yamazaki & Beauchamp, 2007) are possible but unlikely because both SB and BB exhibit a promiscuous mating system (Connor, Richards, Smolker, & Mann, 1996; Smith et al., 2016). Regardless, the difference in MHC diversity between the two populations likely also confers a differential potential to respond to pathogen pressure.

The diverse function and variability of MHC genes reflect evolutionary adaptive processes and thus make them suitable candidates to evaluate genetic diversity relevant to conservation. In this study, we compared MHC genetic diversity and microsatellite diversity of two contrasting bottlenose dolphin populations. We revealed signals of selective processes acting on the MHC DQB in both populations. In comparison with the BB population, the more stable SB population exhibited larger MHC diversity. This is congruent with our hypothesis that the difference in reproductive output and viability between the two populations (Manlik et al., 2016) may be better reflected by adaptive genes of the MHC than putatively neutral microsatellite loci. However, it is important to point out that it is not possible to make conclusive population-level inferences because we compared only two populations. Our results do not allow us to differentiate between cause and effect: Low MHC diversity could be driving population decline, and population decline could be diminishing MHC diversity. Those two explanations are not mutually

exclusive. Both explanations would warrant monitoring MHC diversity of wild animal populations, either as an indicator (reflecting population declines) or as potential driver of population viability (causing population declines). In either case, the greater inter-population difference in MHC diversity relative to microsatellite diversity adds to the growing body of evidence pointing to MHC diversity as a suitable marker for the conservation of vertebrates. Our results also suggest that the SB population, harboring larger MHC diversity, might have a greater potential to respond to a larger variety of pathogens, which would make it more resilient to environmental change.

ACKNOWLEDGMENTS

The Sea World Research and Rescue Foundation Inc. (#RG104136) primarily funded this work. Additional funding came from the WV Scott Foundation. Data were accumulated over many studies, funding sources including NSF (#0918308, 0941487, 1559380, USA), NSERC (Canada), National Geographic Society, PADI Foundation, ARC (Australia), Shark Bay Shire, and in-kind support from Monkey Mia Resort, Shark Bay Resources, tour operators, the Shark Bay and Bunbury communities, volunteers and the Australian Coast Radio Monitors WA Inc. Data collection in Bunbury were facilitated through the South West Marine Research Program and its partners: Bemax Cable Sands, Bunbury Dolphin Discovery Centre, Bunbury Port Authority, City of Bunbury, Cristal Global, WA Department of Parks and Wildlife, Iluka, Millard Marine, Naturaliste Charters, South West Development Commission, WA Plantation Resources and Worsley Alumina. The Shark Bay Dolphin Research Project, the Dolphin Innovation Project and the Dolphin Alliance Project (Shark Bay Dolphin Research Alliance) provided Shark Bay data. This paper represents HIMB and SOEST contribution numbers 1756 and 10700, respectively. We thank D McElligott for sample collection, as well as V Foroughirad, E Krzyszczyk, H Raudino, K Sprogis, S Wild and S Wittwer for checking demographic parameters of dolphins in the databases. We thank CS Baker and anonymous reviewers for valuable comments. Also, thanks to Ewa Krzyszczyk, who allowed us to use her photograph of bottlenose dolphins in Shark Bay (Figure 1).

CONFLICT OF INTERESTS

None declared.

AUTHORS' CONTRIBUTIONS

O.M., W.B.S., M.K., J.M. C.F., L.B., and R.C.C. devised this study. O.M. performed all MHC sequencing and analyses, genotyped microsatellite loci, assessed microsatellite diversity, performed molecular sexing analyses, and wrote the original draft. M.K., A.M.K., and S.J.A. collected most of the biopsy samples. M.K. and A.M.K. extracted DNA, genotyped microsatellite loci, and performed molecular sexing analyses. L.B. and S.J.A. contributed data (identity, sex, age, etc.)

on the Bunbury dolphins. J.M., R.C.C., C.F., M.K., A.M.K., and S.J.A. contributed data (identity, sex, age, etc.) on the Shark Bay dolphins. W.B.S. supervised O.M. on this project and contributed to the genetic analyses. All authors reviewed and edited the manuscript.

DATA ACCESSIBILITY

- MHC DNA sequence alignment available in Dryad/Figure S1 (Manlik et al., 2019; <https://doi.org/10.5061/dryad.73k278d>).
- Microsatellite data: previously uploaded in supplement of published article (Manlik et al., 2018): <https://doi.org/10.1111/mms.12555>. Additional microsatellite data now in Dryad/Table S1, available from the Dryad Digital Repository: Manlik et al. (2019) <https://doi.org/10.5061/dryad.73k278d>.
- Various other data available in Dryad/Appendix (<https://doi.org/10.5061/dryad.73k278d>; Manlik et al., 2019), including: conservative sampling design and numbers (Table S2), MHC and microsatellite diversity measures, including F_{ST} (Table S3), BLASTN results (Table S4), diversity measures and statistics for microsatellite subsamples (Table S5).

ORCID

Oliver Manlik  <https://orcid.org/0000-0002-0924-3768>

REFERENCES

- Ahmad, T., Marshall, S. E., Mulcahy-Hawes, K., Orchard, T., Crawshaw, J., Armuzzi, A., ... Bruce, M. (2002). High resolution MIC genotyping: Design and application to the investigation of inflammatory bowel disease susceptibility. *Tissue Antigens*, 60, 164–179. <https://doi.org/10.1034/j.1399-0039.2002.600207.x>
- Allen, S. J., Bryant, K. A., Kraus, R. H., Loneragan, N. R., Kopps, A. M., Brown, A. M., ... Krützen, M. (2016). Genetic isolation between coastal and fishery-impacted, offshore bottlenose dolphins (*Tursiops* spp.) populations. *Molecular Ecology*, 25, 2735–2753.
- Allendorf, F. W., & Luikart, G. (2008). *Conservation and the genetics of populations*. Malden, MA: Blackwell Publishing.
- Amos, W., & Hoelzel, A. R. (1991). Long term preservation of whale skin for DNA analysis. In A. R. Hoelzel (Ed.), *Genetic ecology of whales and dolphins. Report of the International Whaling Commission special issue 13* (pp. 99–104). Cambridge, UK: International Whaling Commission.
- Baker, C. S., Flórez-González, L., Abernethy, B., Rosenbaum, H. C., Slade, R. W., Capella, J., & Bannister, J. L. (1998). Mitochondrial DNA variation and maternal gene flow among humpback whales of the southern hemisphere. *Marine Mammal Science*, 14, 721–737. <https://doi.org/10.1111/j.1748-7692.1998.tb00758.x>
- Baker, C. S., Vant, M. D., Dalebout, M. L., Lento, G. M., O'Brien, S. J., & Yuhki, N. (2006). Diversity and duplication of DQB and DRB-like genes of the MHC in baleen whales (suborder: Mysticeti). *Immunogenetics*, 58, 283–296. <https://doi.org/10.1007/s00251-006-0080-y>
- Borghans, J. A. M., Beltman, J. B., & de Boer, R. J. (2004). MHC polymorphism under host-pathogen coevolution. *Immunogenetics*, 55, 732–739. <https://doi.org/10.1007/s00251-003-0630-5>
- Bos, D. H., Copurenko, D., Williams, R. N., & DeWoody, A. J. (2008). Inferring population history and demography using

- microsatellites, mitochondrial DNA, and major histocompatibility complex (MHC) genes. *Evolution*, 62, 1458–1468. <https://doi.org/10.1111/j.1558-5646.2008.00364.x>
- Bos, D. H., Turner, S. M., & DeWoody, A. J. (2007). Haplotype inference from diploid sequence data: Evaluating the performance of Bayesian methods using non-neutral MHC sequences. *Heredity*, 144, 228–234.
- Brown, A. H. D., & Weir, B. S. (1983). Measuring genetic variability in plant populations, in isozymes. In S. D. Tanksley, & T. J. Orton (Eds.), *Plant genetics and breeding, Part A* (pp. 219–239). Amsterdam, The Netherlands: Elsevier Science Publishing.
- Caballero, S., Heimeier, D., Trujillo, F., Vianna, J. A., Barrios-Garrido, H., Montiel, M. G., ... Baker, C. S. (2010). Initial description of major histocompatibility complex variation at two class II loci (DQA-DQB) in *Sotalia fluviatilis* and *Sotalia guianensis*. *Latin American Journal of Aquatic Mammals*, 8, 81–95. <https://doi.org/10.5597/lajam00156>
- Chapman, J. R., Nakagawa, S., Coltman, D. W., Slaters, J., & Sheldon, B. C. (2009). A quantitative review of heterozygosity–fitness correlations in animal populations. *Molecular Ecology*, 18, 2746–2765.
- Chybicki, I. J., & Burczyk, J. (2009). Simultaneous estimation of null alleles and inbreeding coefficients. *Journal of Heredity*, 100, 106–113. <https://doi.org/10.1093/jhered/esn088>
- Connor, R. C., Richards, A. F., Smolker, R. A., & Mann, J. (1996). Patterns of female attractiveness in Indian Ocean bottlenose dolphins. *Behaviour*, 133, 37–69. <https://doi.org/10.1163/156853996X00026>
- Davis, L. G., Dibner, M. D., & Battey, J. F. (1986). *Basic methods in molecular biology* (1st ed.). New York, NY: Elsevier Science Publishing.
- Di Guardo, G., & Mazzariol, S. (2014). Cetacean morbillivirus in northern and southern hemispheres. *Frontiers in Microbiology*, 5, 211. <https://doi.org/10.3389/fmicb.2014.00211>
- Doherty, P. C., & Zinkernagel, R. M. (1975). A biological role for the major histocompatibility antigens. *Lancet*, 1, 1406–1409. [https://doi.org/10.1016/S0140-6736\(75\)92610-0](https://doi.org/10.1016/S0140-6736(75)92610-0)
- Drummond, A. J., Ashton, B., Buxton, S., Cheung, M., Cooper, A., Duran, C., ... Wilson, A. (2010). Geneious version 5.5 & 6. Retrieved from <http://www.geneious.com>
- Du, H., Zheng, J., Wu, M., Zhao, Q., & Wang, D. (2010). High MHC DQB variation and asymmetric allelic distribution in the endangered Yangtze finless porpoise, *Neophocaena phocaenoides asiaorientalis*. *Biochemical Genetics*, 48, 433–449. <https://doi.org/10.1007/s10528-009-9327-2>
- Eimes, J. A., Bollmer, J. L., Whittingham, L. A., Johnson, J. A., van Oosterhout, C., & Dunn, P. O. (2011). Rapid loss of MHC class II variation in a bottlenecked population is explained by drift and loss of copy number variation. *Journal of Evolutionary Biology*, 24, 1847–1856. <https://doi.org/10.1111/j.1420-9101.2011.02311.x>
- Eizaguirre, C., Lenz, T. L., Kalbe, M., & Milinski, M. (2012). Rapid and adaptive evolution of MHC genes under parasite selection in experimental vertebrate populations. *Nature Communications*, 3, 621. <https://doi.org/10.1038/ncomms1632>
- Frankham, R. (2005). Genetics and extinctions. *Biological Conservation*, 126, 131–140. <https://doi.org/10.1016/j.biocon.2005.05.002>
- Frankham, R., Ballou, J. D., & Briscoe, D. A. (2010). *Introduction to conservation genetics* (2nd ed.). Cambridge, UK: Cambridge University Press.
- Garrigan, D., & Hedrick, P. (2003). Detecting adaptive molecular polymorphism: Lessons from the MHC. *Evolution*, 57, 1707–1722.
- Hayashi, K., Nishida, S., Yoshida, H., Goto, M., Pastene, L. A., & Koike, H. (2003). Sequence variation of the DQB allele in the cetacean MHC. *Mammal Study*, 28, 89–96. <https://doi.org/10.3106/mamma1study.28.89>
- Hayashi, K., Yoshida, H., Nishida, S., Goto, M., Kanda, N., Baba, Y., & Koike, H. (2006). Genetic variation of the MHC DQB locus in the finless porpoise (*Neophocaena phocaenoides*). *Zoological Science*, 23, 147–153.
- Hedrick, P. (2003). The major histocompatibility complex (MHC) in declining populations: An example of adaptive variation. In W. V. Holt, A. R. Pickard, J. C. Rodger, & D. E. Wildt (Eds.), *Reproduction science and integrated conservation* (pp. 97–113). Cambridge, UK: Cambridge University Press.
- Hedrick, P. W. (2005). *Genetics of populations* (3rd ed.). Boston, MA: Science Books International.
- Heimeier, D., Baker, C. S., Russell, K., Duignan, P. J., Hutt, A., & Stone, G. S. (2009). Confirmed expression of MHC class I and class II genes in the New Zealand endemic Hector's dolphin (*Cephalorhynchus hectori*). *Marine Mammal Science*, 25, 68–90.
- Hoelzel, A. R., Potter, C. W., & Best, P. B. (1998). Genetic differentiation between parapatric 'nearshore' and 'offshore' populations of bottlenose dolphin. *Proceedings of the Royal Society of London, Series B: Biological Sciences*, 265, 1177–1183.
- Hoelzel, A. R., Stephens, J. C., & O'Brien, S. J. (1999). Molecular genetic diversity and evolution at the MHC DQB locus in four species of pinipeds. *Molecular Biology and Evolution*, 16, 611–618.
- Holderegger, R., Kamm, U., & Gugerli, F. (2006). Adaptive vs neutral genetic diversity: Implications for landscape genetics. *Landscape Ecology*, 21, 797–807. <https://doi.org/10.1007/s10980-005-5245-9>
- Hughes, A. L., & Nei, M. (1989). Nucleotide substitution at the major histocompatibility complex class II loci: Evidence for overdominant selection. *Proceedings of the National Academy of Sciences of the United States of America*, 86, 958–962.
- Kalbe, M., Eizaguirre, C., Dankert, I., Reusch, T. B. H., Sommerfeld, R. D., Wegner, K. M., & Milinski, M. (2009). Lifetime reproductive success is maximized with optimal major histocompatibility complex diversity. *Proceedings of the Royal Society of London, Series B: Biological Sciences*, 276, 925–934. <https://doi.org/10.1098/rspb.2008.1466>
- Kamiya, K., O'Dwyer, O., Westerdaal, H., Senior, A., & Nakagawa, S. (2014). Quantitative review of MHC-based mating preference: The role of diversity and dissimilarity. *Molecular Ecology*, 23, 5151–5163.
- Kimura, M. (1977). Preponderance of synonymous changes as evidence for the neutral theory of molecular evolution. *Nature*, 267, 275–276. <https://doi.org/10.1038/267275a0>
- Kopps, A. M. (2007). Who is the one? Paternity assessment in bottlenose dolphins (*Tursiops* sp.) – A species with multilevel male-alliances. MSc thesis, University of Zurich.
- Kopps, A. M., Ackermann, C. Y., Sherwin, W. B., Allen, S. J., Bejder, L., & Krützen, M. (2014). Cultural transmission of tool use combined with habitat specializations leads to fine-scale genetic structure in bottlenose dolphins. *Proceedings of the Royal Society of London, Series B: Biological Sciences*, 281, 20133245. <https://doi.org/10.1098/rspb.2013.3245>
- Krützen, M., Barré, L. M., Möller, L. M., Heithaus, M. R., Simms, C., & Sherwin, W. B. (2002). A biopsy system for small cetaceans: Darting success and wound healing in *Tursiops* spp. *Marine Mammal Science*, 18, 863–878. <https://doi.org/10.1111/j.1748-7692.2002.tb01078.x>
- Krützen, M., Sherwin, W. B., Berggren, P., & Gales, N. (2004). Population structure in an inshore cetacean revealed by microsatellite and mtDNA analysis: Bottlenose dolphins (*Tursiops* sp.) in Shark Bay, Western Australia. *Marine Mammal Science*, 20, 28–47. <https://doi.org/10.1111/j.1748-7692.2004.tb01139.x>
- Krützen, M., Valsecchi, R., Connor, R., & Sherwin, W. B. (2001). Characterization of microsatellite loci in *Tursiops aduncus*. *Molecular Ecology Notes*, 1, 170–172. <https://doi.org/10.1046/j.1471-8278.2001.00065.x>
- Kryazhimskiy, S., & Plotkin, J. B. (2008). The population genetics of dN/dS. *Plos Genetics*, 4, e1000304. <https://doi.org/10.1371/journal.pgen.1000304>
- Kumar, S., Stecher, G., & Tamura, K. (2016). MEGA7: Molecular evolutionary genetics analysis version 7.0 for bigger datasets. *Molecular Biology and Evolution*, 33, 1870–1874. <https://doi.org/10.1093/molbev/msw054>

- Kurtz, J., Kalbe, M., Aeschlimann, P. B., Häberli, M. A., Wegner, K. M., Reusch, T. B. H., & Milinski, M. (2004). Major histocompatibility complex diversity influences parasite resistance and innate immunity in sticklebacks. *Proceedings of the Royal Society of London, Series B: Biological Sciences*, 271, 197–204. <https://doi.org/10.1098/rspb.2003.2567>
- Librado, P., & Rozas, J. (2009). DnaSP v5: A software for comprehensive analysis of DNA polymorphism data. *Bioinformatics*, 25, 1451–1452. <https://doi.org/10.1093/bioinformatics/btp187>
- Manlik, O. (2016). Fitness & major histocompatibility complex diversity of two bottlenose dolphin populations. PhD thesis, University of New South Wales.
- Manlik, O. (2019). The importance of reproduction for the conservation of slow-growing animal populations. In P. Comizzoli, J. Brown, & W. V. Holt (Eds.), *Reproductive science in animal conservation—Progress and prospects* (2nd ed.). New York, NY: Springer.
- Manlik, O., Chabanne, D., Daniel, C., Bejder, L., Allen, S. J., & Sherwin, W. B. (2018). Demography and genetics suggest reversal of dolphin source-sink dynamics, with implications for conservation. *Marine Mammal Science*, <https://doi.org/10.1111/mms.12555>
- Manlik, O., Krützen, M., Koppes, A. M., Mann, J., Bejder, L., Allen, S. J., ... Sherwin, W. B. (2019). Data from: Is MHC diversity a better marker for conservation than neutral genetic diversity? A case study of two contrasting dolphin populations. *Dryad Digital Repository*. <https://doi.org/10.5061/dryad.73k278d>
- Manlik, O., McDonald, J. A., Mann, J., Smith, H. C., Bejder, L., Krützen, M., ... Sherwin, W. B. (2016). The relative importance of reproduction and survival for the conservation of two dolphin populations. *Ecology and Evolution*, 6, 3496–3512. <https://doi.org/10.1002/ece3.2130>
- Manlik, O., Schmid-Hempel, R., & Schmid-Hempel, P. (2017). Parasite infection of specific host genotypes relates to changes in prevalence in two natural populations of bumblebees. *Infection, Genetics and Evolution*, 56, 125–132. <https://doi.org/10.1016/j.meegid.2017.11.019>
- Marsh, S. G. E., Albert, E. D., Bodmer, W. F., Bontrop, R. E., Dupont, B., Erlich, H. A., ... Trowsdale, J. (2010). Nomenclature for factors of the HLA system, 2010. *Tissue Antigens*, 77, 291–455. <https://doi.org/10.1111/j.1399-0039.2010.01466.x>
- Moreno-Santillán, D. D., Lacey, E. A., Gendron, D., & Ortega, O. (2016). Genetic variation at exon 2 of the MHC class II DQB locus in blue whale (*Balaenoptera musculus*) from the Gulf of California. *PLoS ONE*, 11, e0141296. <https://doi.org/10.1371/journal.pone0141296>
- Munguia-Vega, A., Esquer-Garrigos, Y., Rojas-Bracho, L., Vazquez-Juarez, R., Castro-Prieto, A., & Flores-Ramirez, S. (2007). Genetic drift vs. natural selection in a long-term small isolated population: Major histocompatibility complex class II variation in the Gulf of California endemic porpoise (*Phocoena sinus*). *Molecular Ecology*, 16, 4051–4065.
- Murray, B. W., Malik, S., & White, B. N. (1995). Sequence variation at the major histocompatibility complex locus DQ β in beluga whales (*Delphinapterus leucas*). *Molecular Biology and Evolution*, 12, 582–593.
- Nater, A., Koppes, A. M., & Krützen, M. (2009). New polymorphic tetranucleotide microsatellites improve scoring accuracy in the bottlenose dolphin *Tursiops aduncus*. *Molecular Ecology Notes*, 9, 531–534. <https://doi.org/10.1111/j.1755-0998.2008.02246.x>
- Nei, M. (1987). *Molecular evolutionary genetics*. New York, NY: Columbia University Press.
- Nei, M., & Gojobori, T. (1986). Simple methods for estimating the number of synonymous and nonsynonymous nucleotide substitutions. *Molecular Biology and Evolution*, 3, 418–426.
- Nicholson, K., Bejder, L., Allen, S. J., Krützen, M., & Pollock, K. H. (2012). Abundance, survival and temporary emigration of bottlenose dolphins (*Tursiops* sp.) off Useless Loop in the western gulf of Shark Bay, Western Australia. *Marine and Freshwater Research*, 63, 1059–1068. <https://doi.org/10.1071/MF12210>
- Oliver, M. K., & Piertney, S. B. (2012). Selection maintains MHC diversity through a natural population bottleneck. *Molecular Biology and Evolution*, 29, 1713–1720. <https://doi.org/10.1093/molbev/mss06>
- Paterson, S., Wilson, K., & Pemberton, J. M. (1998). Major histocompatibility complex variation associated with juvenile survival and parasite resistance in a large unmanaged ungulate population (*Ovis aries* L.). *Proceedings of the National Academy of Sciences of the United States of America*, 95, 3714–3719. <https://doi.org/10.1073/pnas.95.7.3714>
- Peakall, R., & Smouse, P. E. (2006). GenAIE6: Genetic analysis in Excel. Population genetic software for teaching and research. *Molecular Ecology Notes*, 6, 288–295. <https://doi.org/10.1111/j.1471-8286.2005.01155.x>
- Peakall, R., & Smouse, P. E. (2012). GenAIE6.5: Genetic analysis in Excel. Population genetic software for teaching and research—an update. *Bioinformatics*, 28, 2537–2539. <https://doi.org/10.1093/bioinformatics/bts460>
- Piertney, S. B., & Webster, L. M. (2010). Characterizing functionally important and ecologically meaningful genetic diversity using a candidate gene approach. *Genetica*, 138, 419–432. <https://doi.org/10.1007/s10709-008-9322-2>
- Preen, A. R., Marsh, H., Lawler, I. R., Prince, R. I. T., & Shepherd, R. (1997). Distribution and abundance of dugongs, turtles, dolphins and other megafauna in Shark Bay, Ningaloo Reef and Exmouth Gulf, western Australia. *Wildlife Research*, 24, 185–208. <https://doi.org/10.1071/WR95078>
- Radwan, J., Biedrzycka, A., & Babik, W. (2010). Does reduced MHC diversity decrease viability of vertebrate populations? *Biological Conservation*, 143, 537–544. <https://doi.org/10.1016/j.biocon.2009.07.026>
- Reed, D. H., & Frankham, R. (2003). Correlation between fitness and genetic diversity. *Conservation Biology*, 17, 230–237. <https://doi.org/10.1046/j.1523-1739.2003.01236.x>
- Rousset, F. (2008). genepop'007: A complete re-implementation of the genepop software for Windows and Linux. *Molecular Ecology Resources*, 8, 103–106. <https://doi.org/10.1111/j.1471-8286.2007.01931.x>
- Seddon, J. M., & Ellegren, H. (2002). MHC class II genes in European wolves: A comparison with dogs. *Immunogenetics*, 54, 490–500. <https://doi.org/10.1007/s00251-002-0489-x>
- Sepil, I., Lachish, S., Hink, A. E., & Shelton, B. (2013). MHC supertypes confer both qualitative and quantitative resistance to avian malaria infections in a wild bird population. *Proceedings of the Royal Society of London, Series B: Biological Sciences*, 280, 20130134. <https://doi.org/10.1098/rspb.2013.0134>
- Sepil, I., Lachish, S., & Sheldon, B. (2013). MHC-linked survival and lifetime reproductive success in a wild population of great tits. *Molecular Ecology*, 22, 384–396.
- Sherwin, W. B., Chao, A., Jost, L., & Smouse, P. E. (2017). Information theory broadens the spectrum of molecular ecology and evolution. *Trends in Ecology and Evolution*, 32, 948–963. <https://doi.org/10.1016/j.tree.2017.09.012>
- Shinohara, M., Domingo-Roura, X., & Takenaka, O. (1997). Microsatellites in the bottlenose dolphin *Tursiops truncatus*. *Molecular Ecology*, 6, 695–696.
- Silva, M. C., & Edwards, S. V. (2009). Structure and evolution of a new avian MHC class II B gene in a sub-antarctic seabird, the thin-billed prion (Procellariiformes: *Pachyptila belcheri*). *Journal of Molecular Evolution*, 68, 279–291. <https://doi.org/10.1007/s00239-009-9200-2>
- Smith, H. C., Frère, C., Kobryn, H., & Bejder, L. (2016). Dolphin sociality, distribution and calving as important behavioural patterns informing management. *Animal Conservation*, 19, 462–471. <https://doi.org/10.1111/acv.12263>
- Smith, H. C., Pollock, K. H., Waples, K., Bradley, S., & Bejder, L. (2013). Use of the robust design to estimate seasonal abundance and demographic parameters of a coastal bottlenose dolphin (*Tursiops aduncus*)

- population. *PLoS ONE*, 8, e76574. <https://doi.org/10.1371/annotation/369119db-d9ca-4473-9390-89ee0c2a532f>
- Sommer, S. (2005). The importance of immune gene variability (MHC) in evolutionary ecology and conservation. *Frontiers in Zoology*, 2, 16.
- Sprogis, K. R., Christiansen, F., Raudino, H. C., Kobryn, H. T., Wells, R. S., & Bejder, L. (2018). El Niño Southern Oscillation influences the abundance and movements of a marine top predator in coastal waters. *Global Change Biology*, 24, 1085–1096. <https://doi.org/10.1111/gcb.13892>
- Sprogis, K. R., Pollock, K. H., Raudino, H. C., Allen, S. J., Kopps, A. M., Manlik, O., ... Bejder, L. (2016). Sex-specific patterns in abundance, temporary emigration and survival of Indo-Pacific bottlenose dolphins (*Tursiops aduncus*) in coastal and estuarine waters. *Frontiers in Marine Science*, 3, 12. <https://doi.org/10.3389/fmars.2016.00012>
- Stephens, M., & Donnelly, P. (2003). A comparison of Bayesian methods for haplotype reconstruction from population genotype data. *American Journal of Human Genetics*, 73, 1162–1169. <https://doi.org/10.1086/379378>
- Stephens, M., Smith, N. J., & Donnelly, P. (2001). A new statistical method for haplotype reconstruction from population data. *American Journal of Human Genetics*, 68, 978–989. <https://doi.org/10.1086/319501>
- Stephens, N., Duignan, P., Wang, J., Bingham, J., Finn, H., Bejder, L., ... Holyoake, C. (2014). Cetacean morbillivirus in coastal Indo-Pacific bottlenose dolphins, Western Australia. *Emerging Infectious Diseases*, 20, 666–670. <https://doi.org/10.3201/eid2004.131714>
- Tajima, F. (1989). Statistical method for testing the neutral mutation hypothesis by DNA polymorphism. *Genetics*, 123, 586–595.
- Tajima, F. (1993). Measurement of DNA polymorphism. In M. Takahata, & A. G. Clark (Eds.), *Mechanisms of molecular evolution* (pp. 37–59). Sunderland, MA: Sinauer Associates Inc.
- Takahata, N., & Nei, M. (1990). Allelic genealogy under overdominant and frequency-dependent selection and polymorphism of major histocompatibility complex loci. *Genetics*, 124, 967–978.
- Taylor, B. L., Rojas-Bracho, L., Moore, J., Jaramillo-Legorreta, A., Ver Hoef, J. M., Cardenas-Hinojosa, G., ... Hammond, P. S. (2017). Extinction is imminent for Mexico's endemic porpoise unless fishery bycatch is eliminated. *Conservation Letters*, 10, 588–595. <https://doi.org/10.1111/conl.12331>
- Thompson, J. D., Higgins, D. G., & Gibson, T. J. (1994). CLUSTAL W: Improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position specific gap penalties and weight matrix choice. *Nucleic Acids Research*, 22, 4673–4680. <https://doi.org/10.1093/nar/22.22.4673>
- Thoss, M., Ilmonen, K., Musolf, K., & Penn, D. J. (2011). Major histocompatibility complex heterozygosity enhances reproductive success. *Molecular Ecology*, 20, 1546–1557. <https://doi.org/10.1111/j.1365-294X.2011.05009.x>
- Tsuji, K., Aizawa, M., & Sasaki, T. (1992). *HLA 1991: Proceedings of the eleventh international major histocompatibility workshop and conference*. Oxford, UK: Oxford University Press.
- Ujvari, B., & Belov, K. (2011). Major Histocompatibility Complex (MHC) markers in conservation biology. *International Journal of Molecular Sciences*, 12, 5168–5186. <https://doi.org/10.3390/ijms12085168>
- van Bresse, M. F., Duignan, P. J., Banyard, A., Barbieri, M., Colegrove, K. M., de Guise, S., ... Wellehan, J. F. (2014). Cetacean morbillivirus: Current knowledge and future directions. *Viruses*, 6, 5145–5181. <https://doi.org/10.3390/v6125145>
- van Oosterhout, C., Hutchinson, W. F., Wills, D. P. M., & Shipley, P. (2004). Micro-checker: Software for identifying and correcting genotyping errors in microsatellite data. *Molecular Ecology Notes*, 4, 535–538. <https://doi.org/10.1111/j.1471-8286.2004.00684.x>
- van Tienderen, P. H., de Haan, A. A., van der Linden, G., & Vosman, B. (2002). Biodiversity assessment using markers for ecologically important traits. *Trends in Ecology and Evolution*, 17, 577–582. [https://doi.org/10.1016/S0169-5347\(02\)02624-1](https://doi.org/10.1016/S0169-5347(02)02624-1)
- Vassilakos, D., Natoli, A., Dahlheim, M., & Hoelzel, A. R. (2009). Balancing and directional selection at Exon-2 of the MHC DQB1 locus among populations of odontocete cetaceans. *Molecular Biology and Evolution*, 26, 681–689.
- Watterson, G. A. (1975). On the number of segregating sites in genetic models without recombination. *Theoretical Population Biology*, 7, 256–276. [https://doi.org/10.1016/0040-5809\(75\)90020-9](https://doi.org/10.1016/0040-5809(75)90020-9)
- Wegner, K. M., Kalbe, M., Milinski, M., & Reusch, T. B. H. (2008). Mortality selection during the 2003 European heat wave in three-spined sticklebacks: Effects of parasites and MHC genotype. *BMC Evolutionary Biology*, 8, 124. <https://doi.org/10.1186/1471-2148-8-124>
- Wegner, K. M., Reusch, T. B., & Kalbe, M. (2003). Multiple parasites are driving major histocompatibility complex polymorphism in the wild. *Journal of Evolutionary Biology*, 16, 224–232. <https://doi.org/10.1046/j.1420-9101.2003.00519.x>
- Xu, S., Ju, J., Zhou, X., Wang, L., Zhou, K., & Yang, G. (2012). Considerable MHC diversity suggests that the functional extinction of baiji is not related to population genetic collapse. *PLoS ONE*, 7, e30423. <https://doi.org/10.1371/journal.pone.0030423>
- Yamazaki, Y., & Beauchamp, G. K. (2007). Genetic basis for MHC-dependent mate choice. *Advances in Genetics*, 59, 129–145.
- Yang, G., Yan, J., Zhou, K., & Wei, F. (2005). Sequence variation and gene duplication at MHC DQB loci of baiji (*Lipotes vexillifer*), a Chinese river dolphin. *Journal of Heredity*, 96, 310–317. <https://doi.org/10.1093/jhered/esi055>
- Yang, Z., & Bielawski, J. P. (2000). Statistical methods for detecting molecular adaptation. *Trends in Ecology and Evolution*, 15, 496–503. [https://doi.org/10.1016/S0169-5347\(00\)01994-7](https://doi.org/10.1016/S0169-5347(00)01994-7)

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of the article.

How to cite this article: Manlik O, Krützen M, Kopps AM, et al. Is MHC diversity a better marker for conservation than neutral genetic diversity? A case study of two contrasting dolphin populations. *Ecol Evol*. 2019;9:6986–6998. <https://doi.org/10.1002/ece3.5265>