



May, C. J., Welsh, G. I., Chesor, M., Lait, P. J., Schewitz-Bowers, L. P., Lee, R. W. J., & Saleem, M. A. (2019). Human Th17 cells produce a soluble mediator that increases podocyte motility via signaling pathways that mimic PAR-1 activation. *American Journal of Physiology - Renal Physiology*, 317(4), F913-F921.
<https://doi.org/10.1152/ajprenal.00093.2019>

Peer reviewed version




Link to published version (if available):
[10.1152/ajprenal.00093.2019](https://doi.org/10.1152/ajprenal.00093.2019)

[Link to publication record in Explore Bristol Research](#)
PDF-document

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

A**pJNK Densitometry (n=3)****pp38 MAPK Densitometry (n=3)****B**

Th0 Control

Th17 Control

Th0 5 Min

Th17 5 Min

Th0 15 Min

Th17 15 Min

Th0 30 Min

Th17 30 Min

Th0 45 Min

Th17 45 Min

Th0 60 Min

Th17 60 Min

68 kDa






← p Paxillin s178

← p Paxillin s85


38 kDa







← Beta Actin






A**B****C**





0 hour



12 hours

A**B****C****D**

A15 μ M PAR-1 Agonist**B****C****D****E**

A**B****C****D****E**