

Cognition with magnetic resonance imaging findings and social activities in patients with multiple sclerosis in a Japanese cohort

Journal:	<i>Clinical and Experimental Neuroimmunology</i>
Manuscript ID	CENI-2018-0031.R1
Wiley - Manuscript type:	Original Article
Date Submitted by the Author:	23-Sep-2018
Complete List of Authors:	Niino, Masaaki; Hokkaido Medical Center, Department of Clinical Research Fukazawa, Toshiyuki; Sapporo Neurology Clinic, Kira, Jun-ichi; Kyushu University, Department of Neurology; Kyushu University, Department of Neurology Okuno, Tatsusada; Osaka University, Department of Neurology Mori, Masahiro; Graduate School of Medicine, Chiba University, Neurology Sanjo, Nobuo; Tokyo Medical and Dental University, Department of Neurology and Neurological Science OHASHI, TAKASHI; Tokyo Joshi Ika Daigaku Yachiyo Iryo Center, Neurology Fukaura, Hikoaki; Iwate Medical University, Department of Neurology Fujimori, Juichi; Tohoku Medical and Pharmaceutical University, Department of Neurology Shimizu, Yuko; Tokyo Women's Medical University School of Medicine, Department of Neurology Mifune, Nobuhiro; Kochi University of Technology, Miyazaki, Yusei; Hokkaido Medical Center, Department of Neurology Takahashi, Eri; Hokkaido Medical Center, Department of Clinical Research Kikuchi, Seiji; Hokkaido Medical Center Langdon, Dawn; University of London Benedict, Ralph; University at Buffalo School of Medicine and Biomedical Sciences, Department of Neurology Matsui, Makoto; Kanazawa Medical University, Neurology
Keywords:	cognition, employment, multiple sclerosis
Abstract:	Objective: We investigated the association of magnetic resonance imaging (MRI) findings and social activities with cognition in Japanese patients with multiple sclerosis (MS). Methods: Cognition was evaluated by the Brief International Cognitive Assessment for MS (BICAMS), using previously published data for 156 Japanese patients with MS. The BICAMS results were analyzed with available MRI data, focusing on hyperintense lesions on T2/fluid-attenuated inversion recovery images. Logistic regression analysis was used to assess associations between the BICAMS scores and social activities (i.e., "student," "employed full time," "employed part time," "homemaker," and "unemployed because of MS"). The independent variables were the

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

	<p>BICAMS scores and the Expanded Disability Status Scale, gender, age at examination, education, and disease duration. The dependent variable was the "social activity."</p> <p>Results: Analysis of variance indicated that patients with MS and more cerebral lesions on MRI had lower scores in all three domains of the BICAMS (the Symbol Digit Modalities Test, the second edition of the California Verbal Learning Test, and the revised Brief Visuospatial Memory Test). Scores of all three domains were also significantly lower in patients with cerebellar lesions. Concerning social activities, patients who were unemployed because of MS had lower BICAMS scores compared with employed patients. However, the BICAMS domain scores did not independently affect the other social activities.</p> <p>Conclusions: Higher numbers of cerebral lesions and the presence of cerebellar lesions evaluated by MRI affect cognitive function based on the BICAMS. Cognitive function may affect social activities in patients with MS.</p>

SCHOLARONE™
Manuscripts

For Review Only

Cognition with magnetic resonance imaging findings and social activities in patients with multiple sclerosis in a Japanese cohort

Masaaki Niino,¹ Toshiyuki Fukazawa,² Jun-ichi Kira,³ Tatsusada Okuno,⁴ Masahiro Mori,⁵ Nobuo Sanjo,⁶ Takashi Ohashi,⁷ Hikoaki Fukaura,⁸ Juichi Fujimori,⁹ Yuko Shimizu,¹⁰ Nobuhiro Mifune,¹¹ Yusei Miyazaki,¹ Eri Takahashi,¹ Seiji Kikuchi,¹² Dawn Langdon,¹³ Ralph HB Benedict¹⁴ and Makoto Matsui¹⁵

¹Department of Clinical Research, Hokkaido Medical Center, Sapporo, Japan, ²Sapporo Neurology Hospital, Sapporo, Japan, ³Department of Neurology, Neurological Institute, Graduate School of Medical Sciences, Kyushu University, Fukuoka, Japan, ⁴Department of Neurology, Osaka University Graduate School of Medicine, Suita, Japan, ⁵Department of Neurology, Graduate School of Medicine, Chiba University, Chiba, Japan, ⁶Department of Neurology and Neurological Science, Tokyo Medical and Dental University, Tokyo, Japan, ⁷Department of Neurology, Tokyo Women's Medical University Yachiyo Medical Center, Chiba, Japan, ⁸Department of Neurology, Saitama Medical Center, Saitama Medical University, Saitama, Japan, ⁹Department of Neurology, Tohoku Medical and Pharmaceutical University, Sendai, Japan, ¹⁰Department of Neurology, Tokyo Women's Medical University School of Medicine, Tokyo, Japan, ¹¹School of Economics and Management, Kochi University of Technology, Kochi, Japan, ¹²Department of Neurology, Hokkaido Medical Center, Sapporo, Japan, ¹³Royal Holloway, University of London, Egham, UK, ¹⁴Department of Neurology, School of Medicine and Biomedical Sciences, University at Buffalo, Buffalo, NY, USA, ¹⁵Department of Neurology, Kanazawa Medical University, Ishikawa, Japan

Correspondence

Masaaki Niino, MD, PhD, Department of Clinical Research, Hokkaido Medical Center, Yamanote 5-jo 7-chome, Nishi-ku, Sapporo 063-0005, Japan

Tel: +81-11-611-8111

Fax: +81-11-611-5820

Email: niino@hok-mc.hosp.go.jp

Abstract

Objective: We investigated the association of magnetic resonance imaging (MRI) findings and social activities with cognition in Japanese patients with multiple sclerosis (MS).

Methods: Cognition was evaluated by the Brief International Cognitive Assessment for MS (BICAMS), using previously published data for 156 Japanese patients with MS. The BICAMS results were analyzed with available MRI data, focusing on hyperintense lesions on T2/fluid-attenuated inversion recovery images. Logistic regression analysis was used to assess associations between the BICAMS scores and social activities (i.e., “student,” “employed full time,” “employed part time,” “homemaker,” and “unemployed because of MS”). The independent variables were the BICAMS scores and the Expanded Disability Status Scale, gender, age at examination, education, and disease duration. The dependent variable was the “social activity.”

Results: Analysis of variance indicated that patients with MS and more cerebral lesions on MRI had lower scores in all three domains of the BICAMS (the Symbol Digit Modalities Test, the second edition of the California Verbal Learning Test, and the revised Brief Visuospatial Memory Test). Scores of all three domains were also significantly lower in patients with cerebellar lesions. Concerning social activities, patients who were unemployed because of MS had lower BICAMS scores compared with employed patients. However, the BICAMS domain scores did not independently affect the other social activities.

Conclusions: Higher numbers of cerebral lesions and the presence of cerebellar lesions evaluated by MRI affect cognitive function based on the BICAMS. Cognitive function may affect social activities in patients with MS.

Keywords: cognition, employment, multiple sclerosis

Running Title: BICAMS in MS

Introduction

Patients with multiple sclerosis (MS) often report cognitive impairment, yet this symptom garnered little interest in the clinical and neuroscience literature until the early 1980s. At that time, neuroimaging technology developed to the point where it could provide reliable metrics of neuropathology, and psychometric procedures improved to enable reliable quantification of cognitive abilities.¹ Neuropsychological investigations have shown that 40%–65% of patients with MS experience cognitive impairment, typically with prominent involvement of memory, sustained attention, and information processing speed, rather than global cognitive decline.² In Japan, we have previously reported on cognitive dysfunction in MS using the Brief Repeatable Battery of Neuropsychological tests (BRB-N), which revealed that impaired cognitive function, particularly in information processing speed, was present among Japanese patients.³

Given the essential role of magnetic resonance imaging (MRI) in MS diagnosis and disease surveillance, MS research is often at the forefront of novel and innovative MRI tools for investigating cognitive deficits.⁴ Many of the cognitive deficits in patients with MS can be related to a disconnection syndrome caused by involvement of white matter tracts,⁵ with research showing the importance of white matter lesion sites, microstructural injury, gray matter lesions, and cortical and subcortical gray matter brain atrophy.⁴ Our previous study using BRB-N demonstrated that patients with MS who had more cerebral lesions on MRI had lower scores in most of the BRB-N tests; moreover, the BRB-N scores were significantly lower in patients with brainstem and cerebellar lesions.⁶ Conflating the clinical, radiological, and pathological data, both gray and white matter pathology appear to play unique and likely additive roles in the pathogenesis of MS-related cognitive impairment. However, MRI studies to date have been limited by poor resolutions for detecting disease in brain areas critical for cognition.⁷

Most individuals with MS (>90%) have a history of working before diagnosis, and approximately 60% are employed at time of diagnosis; however, only a small percentage (20%–40%) remain employed after diagnosis.⁸ Improvements in mortality rates and in the preservation of physical capabilities have led to increasing interest in cognitive outcomes and quality of life in these patients.¹ Cognitive impairment limits activities of daily living⁹ and may negatively affect employment status. In fact, declining performance over time in tests of attention and verbal memory are predictive of impaired employment status.¹⁰ Our previous study indicated that patients categorized as being unemployed because of MS had lower BRB-N scores than other for social activity groups.⁶

Recently, we validated the Japanese version of the Brief International Cognitive Assessment for MS (BICAMS). This test comprises the Symbol Digit Modalities Test (SDMT),¹¹ and elements of the second edition of

1
2
3 the California Verbal Learning Test (CVLT2)¹² and the revised Brief Visuospatial Memory Test (BVMTR).¹³ In the
4 present study, we aimed to evaluate the associations of MRI findings and working status with cognitive function
5 among Japanese patients with MS based on the BICAMS data reported in that validation study.
6
7
8
9
10
11

12 **Materials and Methods**

13 *Patients with MS*

14
15
16 The clinical characteristics of the patients included in this study have been described in an earlier report.¹⁴
17 Briefly, 156 Japanese patients with MS (107 females and 49 males) were included, but those with neuromyelitis
18 optica spectrum disorders were excluded. The clinical profiles of the study participants are shown in Table 1.
19
20
21
22

23 *The BICAMS*

24
25 The BICAMS¹⁵ was performed for neuropsychological evaluations. This test is actually a battery three
26 individual tests: the SDMT for cognitive processing speed, the first five recall trials of the CVLT2 for
27 auditory/verbal learning and memory, and the first three recall trials of the BVMTR for visual or spatial memory.¹⁵
28
29 The test battery was administered in the following order, which was fixed: the SDMT, the five CVLT2 trials, and
30 the three BVMTR trials.¹⁶
31
32
33
34
35
36

37 *Assessment of MRI findings*

38
39 MRI scans obtained within a year of performing the BICAMS tests were evaluated for each patient (one
40 patient had an MRI performed 1.5 years before undergoing the BICAMS test). In terms of cerebral lesions, patients
41 were classified into three groups: ≥ 9 lesions, 4–8 lesions, and ≤ 3 lesions. We also checked whether there were
42 brainstem and cerebellar lesions. The MRI scans were obtained by axial slices (5 or 6 mm intervals) at 1.5 or 3
43 Tesla for each subject, and the scan sets per subject contained T2-weighted images and/or T2-weighted
44 fluid-attenuated inversion recovery images.
45
46
47
48
49
50
51

52 *Social activities*

53
54 When receiving the BICAMS tests, participants were requested to provide information on their status
55 with regards the following work status: “student,” “employed full time,” “employed part time,” “homemaker,” and
56
57
58
59
60

1
2
3 “unemployed because of MS”.

6 ***Statistical analysis***

8 All data are expressed as means \pm standard deviation. Statistical analyses were performed using either
9 SAS 9.4 (SAS Institute Inc., Cary, NC) or GraphPad Prism (GraphPad Software, Inc., La Jolla, CA, USA). Raw
10 data for the three component tests (i.e., the SDMT, CVLT2, and BVMTR) were analyzed to identify associations
11 with the MRI findings. Cerebral, brainstem, and cerebellar lesions on MRI were then evaluated by analysis of
12 variance (ANOVA) followed by Tukey’s multiple comparison test. Next, logistic regression analysis was used to
13 assess the presence of causal relationships between social activities and the BICAMS scores. The independent
14 variables were the BICAMS scores, the Expanded Disability Status Scale (EDSS) scores, gender (female = 0; male
15 = 1), age at examination, duration of education, and disease duration, whereas the dependent variables were the
16 social activities, as specified above. *P* values of <0.05 were considered statistically significant.
17
18
19
20
21
22
23
24
25
26
27
28

29 **Results**

30 ***MRI findings and BICAMS scores***

31 *Cognitive function and cerebral, brainstem, or cerebellar lesions*

32
33 In our cohort, 117 patients had ≥ 9 cerebral lesions, 30 had 4–8 lesions, and 9 had ≤ 3 or lesions. There
34 was a trend for lower BICAMS scores in association with higher numbers of cerebral lesions. Only differences
35 between patients with ≥ 9 lesions and with 4–8 lesions were statistically significant in each battery of the BICAMS
36 (Fig. 1a). We also identified 102 and 54 patients without brainstem lesions, respectively. Of note, patients with
37 brainstem lesions had significantly lower scores in all three test batteries of the BICAMS (Fig. 1b). Finally, there
38 were 55 and 101 patients with and without cerebellar lesions, respectively. Patients with cerebellar lesions had
39 significantly lower scores in all three test batteries of the BICAMS (Fig. 1c).
40
41
42
43
44
45
46
47

48 We used the general linear model to assess the presence of causal relationships between MRI lesions and
49 BICAMS scores to control for the effects of age at examination, sex, and EDSS. The results revealed significant
50 differences in BICAMS scores among the three groups of cerebral lesions (SDMT: $F [2, 150] = 5.66, P < 0.05$;
51 CVLT2: $F [2, 150] = 7.01, P < 0.05$; BVMTR: $F [2, 150] = 5.2, P < 0.05$) and between the groups with and without
52 cerebellar lesions (SDMT: $F [1, 151] = 7.56, P < 0.05$; CVLT2: $F [1, 151] = 5.42, P < 0.05$; BVMTR: $F [1, 151] =$
53
54
55
56
57
58
59
60

1
2
3 12.25, $P < 0.05$). In contrast, there was no significant difference in BICAMS scores between the groups with and
4 without brainstem lesions (SDMT: $F [1, 151] = 1.9, P = 0.171$; CVLT2: $F [1, 151] = 1.95, P = 0.165$; BVMTR: $F [1,$
5 $151] = 0.77, P = 0.382$).
6
7
8
9

10 *ANOVA model of the main effects of cerebral, brainstem, or cerebellar lesions*

11
12 We further explored whether each lesion site was associated with decreased BICAMS scores after
13 controlling for the effects of the other two lesions. Brainstem lesions were not associated with the BICAMS.
14 However, cerebral lesions were associated with decreased CVLT2 scores and cerebellar lesions were associated
15 with decreased BVMTR scores (Table 2).
16
17
18
19

20 *Social activities and BICAMS scores*

21
22 Among the patients, 60 were employed full time, 35 were employed part time, 36 were homemakers, and
23 25 were unemployed because of MS, but there were no students. The raw scores for the three BICAMS test
24 batteries by social activity class are shown in Fig. 2. One-way ANOVA was performed with social activity as the
25 independent variables and each BICAMS test as dependent variables. Multiple comparison of patients with MS
26 using Tukey's honestly significant difference test revealed significant differences between the "unemployed
27 because of MS" group and the other social activity groups for each component of the BICAMS ($P < 0.05$) (Fig. 2).
28
29
30
31
32
33
34

35 Several potential factors affect social activities and needed to be considered in the analysis. Initially,
36 logistic regression analysis was performed with the three BICAMS tests, the EDSS scores, gender, age at
37 examination, education duration, and disease duration as independent variables and with each social activity as the
38 dependent variables. Subsequently, a model was generated to explain the relationship of a dependent variable with
39 several significant independent variables selected using a stepwise method. A higher estimated value signified that
40 the social activity group, such as "employed full time" and "unemployed because of MS," had a higher significance
41 with a parameter, especially with EDSS (Table 3). Other factors, such as gender and education, also had
42 independent effects on social activities (Table 3), but the none of the components of the BICAMS affected the
43 social activities status.
44
45
46
47
48
49
50
51
52
53
54
55

56 **Discussion**

57
58
59
60

1
2
3 Given that brain MRI is a cornerstone of MS diagnosis, differential diagnosis, and disease monitoring, it
4 is reasonable to ask whether neuroimaging could also be used to predict cognitive function and impairment in this
5 group.¹⁷ Indeed, patients with more severe MS disease burdens (i.e., greater lesion load and cerebral atrophy on
6 MRI) are known to be at increased risk of cognitive impairment.¹⁸ Several studies in Western countries have also
7 demonstrated that patients with greater lesion frequencies and volumes have significantly greater cognitive
8 impairments compared with patients who have lower lesion burdens.¹⁹⁻²¹

9
10 The current study using the BICAMS obtained similar results to those in our previous study using the
11 BRB-N, indicating that cognitive impairment worsened as the number of cerebral lesions increased in patients with
12 MS.⁶ Although patients who had ≥ 9 cerebral lesions in the current study had significantly lower scores in each
13 BICAMS test than patients with 4–8 lesions, no significant differences in scores were found between patients who
14 had ≤ 3 cerebral lesions and patients who had > 3 lesions. This could be explained by the small sample size of the
15 group with ≤ 3 lesions. Significant correlations have been shown between cognitive function and brain volumes,
16 especially gray matter volume.^{22,23} However, a longitudinal study over 17 years demonstrated that atrophy of the
17 corpus callosum was strongly associated with decreased performance in cognitively demanding information
18 processing tasks.²⁴ Studies of normal-appearing white matter involvement have lent further support to the notion of
19 a disconnection syndrome in patients with MS, which could be the cause of cognitive and physical deficits.⁵ Further
20 research is needed to clarify the associations between cognitive dysfunctions and MRI findings and to understand
21 cognitive impairment in patients with MS.

22
23 Our previous study using the BRB-N demonstrated that patients with brainstem or cerebellar lesions had
24 lower scores than patients without those lesions.⁶ Similar results were obtained in the current study when using the
25 BICAMS, although there was no significant difference in BICAMS scores in patients with and without brainstem
26 lesions after adjusting for age at examination, sex, and EDSS. It may be that patients with cerebellar lesions have
27 more cerebral lesions, and that those patients had lower scores compared with patients without cerebellar lesions.
28 However, in the previous study using the BRB-N, cerebellar lesions were associated with decreased SDMT and
29 Paced Auditory Serial Addition Test scores after controlling for the effects of cerebral and brainstem lesions.⁶ By
30 contrast, in the current study, cerebellar lesions were not statistically associated with decreased SDMT scores after
31 controlling for the effects of those lesions, though the same tendency was shown ($P = 0.097$).

32
33 As discussed in our previous study, the cerebellum may be important in cognitive function and
34 information processing.²⁵⁻²⁷ In the current study, after controlling for the effects of cerebral and brainstem lesions,

1
2
3 cerebellar lesions were associated with decreased BVMTR scores that evaluate visual–spatial episodic memory. It
4 is known that short-term memory impairments associated with cerebellar deficits include difficulty learning and
5 spontaneously recalling new information; these, in turn, reflect deficient strategies for organizing verbal or visual
6 spatial material for encoding, as well as difficulty in locating information in memory stores.²⁷ A significant
7
8 association has also been shown between the total gray and white matter cerebellar volumes and the visual–spatial
9
10 subdomain in a non-faller group of patients with MS.²⁸ In the current study, we did not investigate the cerebellar
11
12 volume in the included patients. However, it is suggested that the cerebellum plays an important role in the control
13
14 of cognitive functions, being strongly interconnected with the prefrontal, superior temporal, posterior parietal, and
15
16 limbic cortices. Further research is needed to clarify the associations between cognitive impairment and cerebellar
17
18 lesions in MS.
19

20
21 Lower SDMT scores were demonstrated in the group of patients who were “unemployed because of MS”
22
23 in our previous study.⁶ The estimated value of SDMT in those data was -0.0627 , which was statistically significant,
24
25 but the effect was small. In the current study, using the same logistic regression analysis as used in our previous
26
27 study and using SDMT, EDSS, and male sex as independent variables and each social activity as the dependent
28
29 variable, we found that the estimated value of SDMT was -0.0409 in the group of patients who were “unemployed
30
31 because of MS”. Those data are not so different from the previous data, which implies that some sample sizes or
32
33 different groups may generate statistically significant or not significant results.
34

35
36 In our previous study, >60% of Japanese patients with MS experienced a change in employment status,
37
38 which is known to affect quality of life significantly.²⁹ Cognitive impairment also has a substantial influence on
39
40 activities of daily living and work capacity,¹ and the worsening of cognitive function could be associated with a
41
42 loss of employment.¹⁰ Using the BRB-N, we previously showed that patients in the “unemployed because of MS”
43
44 group had the lowest scores relative to those in the other social activity groups, and this was confirmed in the
45
46 current study using the BICAMS. Both studies revealed that EDSS scores were highly and negatively associated
47
48 with the “unemployed because of MS” group,⁶ and in addition, the current study demonstrated a positive
49
50 association for the EDSS score and the “employed full time” group. Constructing a model to explain the
51
52 relationship of a dependent variable with several significant independent variables via the stepwise method failed to
53
54 show that cognitive impairments identified by the BICAMS affected social activities in patients with MS. However,
55
56 it has been considered that cognitive dysfunction in MS contributes significantly to unemployment, accidents,
57
58 impaired daily function, and loss of social contacts.³⁰ Prospective studies are needed to evaluate the associations of
59
60

1
2
3 cognitive impairments with social activities. Further, social environments and systems differ considerably among
4 countries and regions and may affect the activities of patients with MS, necessitating that further studies be
5 undertaken in different countries or regions.
6
7

8 In conclusion, our current study demonstrated that cognitive impairment evaluated by the BICAMS is
9 associated with greater numbers of brain lesions on MRI and lower scores on the BICAMS domains. This seems to
10 apply to patients with MS who are unemployed because of their condition, rather than those in other social activity
11 groups. Cognitive impairment is an important aspect of MS that can produce functional handicaps and adversely
12 affect a patient's quality of life.³¹ Currently, there is no simple approach for improving cognitive function in
13 patients with MS. Social engagement and quality of life are important considerations for patients with MS, and
14 there is a growing evidence base to suggest that clinicians need to pay greater attention to how these are affected by
15 cognitive dysfunctions.
16
17
18
19
20
21
22
23
24
25
26

27 **Acknowledgments**

28 We thank Dr. Tomiki Sumiyoshi, Department of Clinical Epidemiology, Translational Medical Center, National
29 Center of Neurology and Psychiatry for his great help and suggestions with the Japanese version of CVLT2. We
30 also thank Ms. Eri Sato at Sapporo Neurology Hospital, Ms. Yumi Kon at Hokkaido Medical Center, Dr. Koji
31 Shinoda at Department of Neurology, Neurological Institute, Graduate School of Medical Sciences, Kyushu
32 University, Dr. Yuji Nakatsuji at Department of Neurology, Toyama University Hospital, and Dr. Takanori Yokota
33 at Department of Neurology and Neurological Science, Tokyo Medical and Dental University for their great help
34 with this study.
35
36
37
38
39
40
41
42
43
44
45

46 **Disclosure of ethical statement**

47 The study protocol was approved by the Ethics Committee of each participating site, and all participants provided
48 written informed consent.
49
50
51
52
53
54

55 **Conflicts of interest**

56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

None declared.

Funding

This study was supported, in part, by a Health and Labour Sciences Research Grant on Rare and Intractable Diseases (Validation of Evidence-based Diagnosis and Guidelines, and Impact on QOL in Patients with Neuroimmunological Diseases) from the Ministry of Health, Labour and Welfare of Japan.

For Review Only

References

1. Benedict RH, Zivadinov R. Risk factors for and management of cognitive dysfunction in multiple sclerosis. *Nat Rev Neurol*. 2011; 7: 332-42.
2. Bobholz JA, Rao SM. Cognitive dysfunction in multiple sclerosis: a review of recent developments. *Curr Opin Neurol*. 2003; 16: 283-88.
3. Niino M, Mifune N, Kohriyama T, et al. Apathy/depression, but not subjective fatigue, is related with cognitive dysfunction in patients with multiple sclerosis. *BMC Neurol*. 2014; 14: 3.
4. Sumowski JF, Benedict R, Enzinger C, et al. Cognition in multiple sclerosis: State of the field and priorities for the future. *Neurology*. 2018; 90: 278-88.
5. Rocca MA, Amato MP, De Stefano N, et al. Clinical and imaging assessment of cognitive dysfunction in multiple sclerosis. *Lancet Neurol*. 2015; 14: 302-17.
6. Niino M, Mifune N, Kohriyama T, et al. Association of cognitive impairment with magnetic resonance imaging findings and social activities in patients with multiple sclerosis. *Clin Exp Neuroimmunol*, 2014; 5: 328-35.
7. DeLuca GC, Yates RL, Beale H, Morrow SA. Cognitive impairment in multiple sclerosis: clinical, radiologic and pathologic insights. *Brain Pathol*. 2015; 25: 79-98.
8. Pompeii LA, Moon SD, McCrory DC. Measures of physical and cognitive function and work status among individuals with multiple sclerosis: a review of the literature. *J Occup Rehabil*. 2005; 15: 69-84.
9. Goverover Y, Genova HM, Hillary FG, DeLuca J. The relationship between neuropsychological measures and the Timed Instrumental Activities of Daily Living task in multiple sclerosis. *Mult Scler*. 2007; 13: 636-44.
10. Morrow SA, Drake A, Zivadinov R, Munschauer F, Weinstock-Guttman B, Benedict RH. Predicting loss of employment over three years in multiple sclerosis: clinically meaningful cognitive decline. *Clin Neuropsychol*. 2010; 24: 1131-45.
11. Smith A. Symbol digit modalities test: Manual, Los Angeles: Western Psychological Services, 1982.
12. Delis DC, Kramer JH, Kaplan E, Ober BA. California Verbal Learning Test Manual: Second Edition, Adult Version, San Antonio, Psychological Corporation, 2000.
13. Benedict RH. Brief Visuospatial Memory Test - Revised: Professional Manual, Odessa, Florida: Psychological Assessment Resources, Inc, 1997.
14. Niino M, Fukazawa T, Kira J, et al. Validation of the Brief International Cognitive Assessment for Multiple

- 1
2
3 Sclerosis in Japan. *Mult Scler J Exp Transl Clin.* 2017; 3: 2055217317748972.
- 4
5 15. Langdon DW, Amato MP, Boringa J, et al. Recommendations for a Brief International Cognitive Assessment
6
7 for Multiple Sclerosis (BICAMS). *Mult Scler.* 2012; 18: 891-8.
- 8
9 16. Benedict RH, Amato MP, Boringa J, et al. Brief International Cognitive Assessment for MS (BICAMS):
10
11 international standards for validation. *BMC Neurol.* 2012; 12: 55.
- 12
13 17. Paul F. Pathology and MRI: exploring cognitive impairment in MS. *Acta Neurol Scand.* 2016; 134: 24-33.
- 14
15 18. Sumowski JF, Leavitt VM. Cognitive reserve in multiple sclerosis. *Mult Scler.* 2013; 19: 1122-7.
- 16
17 19. Patti F, Amato MP, Trojano M, et al. Cognitive impairment and its relation with disease measures in mildly
18
19 disabled patients with relapsing-remitting multiple sclerosis: baseline results from the Cognitive Impairment in
20
21 Multiple Sclerosis (COGIMUS) study. *Mult Scler.* 2009; 15: 779-88.
- 22
23 20. Filippi M, Rocca MA, Benedict RH, et al. The contribution of MRI in assessing cognitive impairment in
24
25 multiple sclerosis. *Neurology.* 2010; 75: 2121-8.
- 26
27 21. Giorgio A, De Stefano N. Cognition in multiple sclerosis: relevance of lesions, brain atrophy and proton MR
28
29 spectroscopy. *Neurol Sci.* 2010; 31: S245-8.
- 30
31 22. Fenu G, Loreface L, Arru M, et al. Cognition in multiple sclerosis: Between cognitive reserve and brain
32
33 volume. *J Neurol Sci.* 2018; 386: 19-22.
- 34
35 23. De Stefano N, Airas L, Grigoriadis N, et al. Clinical relevance of brain volume measures in multiple sclerosis.
36
37 *CNS Drugs.* 2014; 28: 147-56.
- 38
39 24. Granberg T, Martola J, Bergendal G, et al. Corpus callosum atrophy is strongly associated with cognitive
40
41 impairment in multiple sclerosis: Results of a 17-year longitudinal study. *Mult Scler.* 2015; 21: 1151-8.
- 42
43 25. Valentino P, Cerasa A, Chiriaco C, et al. Cognitive deficits in multiple sclerosis patients with cerebellar
44
45 symptoms. *Mult Scler.* 2009; 15: 854-9.
- 46
47 26. Lesage E, Apps MA, Hayter AL, et al. Cerebellar information processing in relapsing-remitting multiple
48
49 sclerosis (RRMS). *Behav Neurol.* 2010; 23: 39-49.
- 50
51 27. Bodranghien F, Bastian A, Casali C, et al. Consensus Paper: Revisiting the Symptoms and Signs of Cerebellar
52
53 Syndrome. *Cerebellum.* 2016; 15: 369-91.
- 54
55 28. Kalron A, Allali G, Achiron A. Cerebellum and cognition in multiple sclerosis: the fall status matters. *J Neurol.*
56
57 2018; 265: 809-16.
- 58
59 29. Kikuchi H, Mifune N, Niino M, et al. Impact and characteristics of quality of life in Japanese patients with
60

1
2
3 multiple sclerosis. *Qual Life Res.* 2011; 20: 119-31.

4
5 30. Rao SM. Neuropsychology of multiple sclerosis. *Curr Opin Neurol.* 1995; 8: 216-20.

6
7 31. McNicholas N, O'Connell K, Yap SM, Killeen RP, Hutchinson M, McGuigan C. Cognitive dysfunction in
8 early multiple sclerosis: a review. *QJM.* 2018; 111: 359-64.
9

10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Review Only

Figure Legends

Fig. 1. MRI lesions and BICAMS scores

Patients were divided into three cerebral lesion groups (≥ 9 , 4–8, and ≤ 3) based on MRI findings. The group with ≥ 9 lesions had significantly lower scores than the group with 4–8 lesions (Fig. 1a). Patients with brainstem lesions had lower scores in all BICAMS tests compared with those who had no brainstem lesions (Fig. 1b). Patients with cerebellar lesions had lower scores in all BICAMS tests (Fig. 1c). Main bars represent mean values and error bars represent standard deviations. ($*P < 0.05$)

Abbreviations: BICAMS, Brief International Cognitive Assessment for MS; BVMTR, the revised Brief Visuospatial Memory Test; CVLT2, the second edition of the California Verbal Learning Test; MRI, magnetic resonance imaging; MS, multiple sclerosis; SDMT, Symbol Digit Modalities Test.

Fig. 2. Social activities and BICAMS scores

The raw BICAMS test scores are shown for each social activity group (employed full time, employed part time, homemaker, or unemployed because of MS). There were significant differences between the “unemployed because of MS” group and the other social activities groups for all BICAMS tests. Main bars represent mean values and error bars represent standard deviations. ($*P < 0.05$)

Abbreviations: BICAMS, Brief International Cognitive Assessment for MS; BVMTR, the revised Brief Visuospatial Memory Test; CVLT2, the second edition of the California Verbal Learning Test; MRI, magnetic resonance imaging; MS, multiple sclerosis; SDMT, Symbol Digit Modalities Test.

Table 1. Clinical profiles of the included patients with MS

	Patients with MS
Number (Female/Male)	156 (107/49)
Age at examination (years) † (range)	41.4 ± 9.3 (23–70)
Duration of education (years) †, ‡ (range)	14.1 ± 1.9 (12–22)
Age at onset (years) † (range)	31.0 ± 9.2 (10–64)
EDSS † (range)	2.4 ± 2.0 (0.0–8.0)
Clinical course	
Primary-progressive	1
Relapsing-remitting	137
Secondary-progressive	18
MRI findings	
Cerebral lesions	
≥9 lesions	117
4–8 lesions	30
≤3 lesions	9
Brainstem lesions	
Positive	102
Negative	54
Cerebellar lesions	
Positive	55
Negative	101
Social activities	
Student	0
Employed full time	60
Employed part time	35
Homemaker	36
Unemployed because of MS	25

† mean ± standard deviation (SD),

Table 2. ANOVA model in which only the main effects of cerebral, brainstem or cerebellar lesions are hypothesized

	SDMT	CVLT2	BVMTR
Cerebral lesions			
<i>F</i> value (2, 151)	1.80	3.51	1.85
<i>P</i> value	0.169	<0.05	0.16
Effect size (partial η^2)	0.023	0.044	0.024
Brainstem lesions			
<i>F</i> value (1, 151)	1.14	0.39	0
<i>P</i> value	0.287	0.535	0.98
Effect size (partial η^2)	0.008	0.003	0.000
Cerebellar lesions			
<i>F</i> value (1, 151)	2.79	1.38	7.11
<i>P</i> value	0.097	0.243	<0.05
Effect size (partial η^2)	0.018	0.009	0.045

Table 3. Logistic regression analysis on social activities

Social activity	Parameter	Estimated value	Standard error	<i>P</i> value	Odds ratio	95% Wald confidence limits for odds ratio
Employed full time	EDSS	-0.2804	0.1130	0.0131	0.756	0.605–0.943
	Gender	1.344	0.4163	0.0012	3.834	1.696–8.671
	Age	-0.0564	0.0231	0.0146	0.945	0.903–0.989
	Education	0.2520	0.1037	0.0150	1.287	1.050–1.577
Employed part time	Gender	-1.232	0.5187	0.0175	0.393	0.106–0.806
Homemaker	Gender	-2.409	0.7564	0.0014	0.090	0.020–0.396
	Education	-0.2675	0.1275	0.0360	0.765	0.596–0.983
Unemployed because of MS	EDSS	0.6504	0.1337	<0.0001	1.916	1.475–2.490
	Gender	1.510	0.5827	0.0096	4.527	1.445–14.183
	Duration	0.0731	0.0351	0.0373	1.076	1.004–1.153

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Fig1a

75x61mm (300 x 300 DPI)

only




Fig1b

78x64mm (300 x 300 DPI)




Fig1c

78x65mm (300 x 300 DPI)




Fig2

137x114mm (300 x 300 DPI)