

Clinical and genomic features of adult and paediatric acute leukaemias with ophthalmic manifestations

Lisa Stenman Skarsgård ,^{1,2} Mattias K. Andersson,³ Marta Persson,³ Ann-Cathrine Larsen,⁴ Sarah E. Coupland,^{5,6} Göran Stenman,³ Steffen Heegaard ^{2,4}

To cite: Skarsgård LS, Andersson MK, Persson M, et al. Clinical and genomic features of adult and paediatric acute leukaemias with ophthalmic manifestations. *BMJ Open Ophthalmology* 2019;**4**:e000362. doi:10.1136/bmjophth-2019-000362

Received 25 June 2019
 Revised 27 August 2019
 Accepted 7 September 2019



© Author(s) (or their employer(s)) 2019. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹Department of Surgery, Ostfold Hospital Trust, Fredrikstad, Norway

²Department of Pathology, Copenhagen University Hospital, Rigshospitalet, Copenhagen, Denmark

³Sahlgrenska Cancer Center, Department of Pathology, University of Gothenburg, Gothenburg, Sweden

⁴Department of Ophthalmology, Copenhagen University Hospital, Rigshospitalet, Glostrup, Denmark

⁵Department of Molecular and Clinical Cancer Medicine, University of Liverpool, Liverpool, UK

⁶Liverpool Clinical Laboratories, Royal Liverpool University Hospital, Liverpool, UK

Correspondence to

Dr Steffen Heegaard; sthe@sund.ku.dk

ABSTRACT

Objective To describe the clinicopathological and genomic features of nine patients with primary and secondary orbital/ocular manifestations of leukaemia.

Methods All orbital/ocular leukaemic specimens from 1980 to 2009 were collected from the Danish Register of Pathology. In six cases, medical records and formalin-fixed, paraffin-embedded blocks were available. Three cases from the Department of Pathology, Royal Liverpool University Hospital, were also included. Immunophenotypes and MYB oncoprotein expression were ascertained by immunohistochemistry. Genomic imbalances were analysed with comparative genomic hybridisation arrays and oncogene rearrangements with fluorescence in situ hybridisation.

Results Four patients had B-cell precursor acute lymphoblastic leukaemia (BCP-ALL) and five had acute myeloid leukaemia (AML). Two patients with BCP-ALL and one with AML had primary orbital manifestations of leukaemia. Common symptoms were proptosis, displacement of the eye, and reduced eye mobility in patients with orbital leukaemias and pain, and reduced visual acuity in patients with ocular leukaemias. All patients with primary orbital lesions were alive up to 18 years after diagnosis. All but one patient with secondary ophthalmic manifestations died of relapse/disseminated disease. *ETV6* and *RUNX1* were rearranged in BCP-ALL, and *RUNX1* and *KMT2A* in AML. Genomic profiling revealed quiet genomes (0–7 aberrations/case). The MYB oncoprotein was overexpressed in the majority of cases.

Conclusions Leukaemias with and without ophthalmic manifestations have similar immunophenotypes, translocations/gene fusions and copy number alterations. Awareness of the clinical spectrum of leukaemic lesions of the eye or ocular region is important to quickly establish the correct diagnosis and commence prompt treatment.

INTRODUCTION

Leukaemic infiltrates can occur in the eye or ocular region as a primary manifestation of leukaemia or as an infiltration of systemic disease.^{1–2} Ophthalmic manifestations are more common in acute leukaemias than in chronic leukaemias.^{1–3} Typical symptoms of patients with leukaemic infiltrates in the ophthalmic region include eyelid oedema and

Key message

What is already known about this subject?

- ▶ Leukaemic infiltrates may occur in the eye or ocular region, and in rare cases it is the first presenting symptom of leukaemia.
- ▶ Leukaemias with ophthalmic manifestations are rarely biopsied or examined extensively, and therefore our knowledge about these lesions is limited.

What are the new findings?

- ▶ This is the first comprehensive, integrated clinicopathological, cytogenetic and genomic analyses of acute leukaemias with ophthalmic manifestations.
- ▶ We show for the first time that leukaemias with eye and/or ocular manifestations have similar genetic and molecular profiles compared with other non-site-specific leukaemias.

How might these results change the focus of research or clinical practice?

- ▶ Our findings further emphasise the broad clinical spectrum of leukaemic lesions with ophthalmic manifestations and the need to consider leukaemia in the differential diagnosis of patients with proptosis, reduced mobility and/or displacement of the eye.
- ▶ Awareness of the clinical spectrum of leukaemic lesions of the eye or ocular region is important to quickly establish the correct diagnosis and commence prompt treatment.

swellings, chemosis and exophthalmos. Globe displacement by a leukaemic mass may restrict ocular mobility and in some cases reduce visual acuity.⁴ Orbital tumours may present as a diffuse infiltrate or as a large single mass. Solid infiltrates can involve all structures in the extraocular region, including the ocular adnexa and the optic nerve,^{4–6} and constitute approximately 2% of all malignancies in this region.⁵

Intraocular manifestations of leukaemias are most frequently seen clinically in the retina, but on histopathological examination, leukaemic lesions of the choroid are equally as common.^{4–6} Patients with intraocular

leukaemic manifestations often have reduced vision, and occasionally also pain or decreased mobility of the eye. Retinal detachment, chemosis, retinitis, glaucoma, uveitis or hypopyon are observed on clinical examination.^{4,6}

Ophthalmic signs and symptoms may also result from side effects associated with leukaemia, such as anaemia, thrombocytopenia, hyperviscosity, immunosuppression and infections.^{1,4,6} When a primary leukaemia arises in the eye or ocular adnexa, subsequent involvement of the peripheral blood or bone marrow usually occurs within 1 year of the ocular disease.⁷

Leukaemias with ophthalmic manifestations are rarely biopsied or examined extensively,^{1,5} and therefore our knowledge about these lesions is still limited. In this study, we describe in detail the clinical, cytogenetic, and genomic features of nine cases of primary and secondary ophthalmic leukaemias.

MATERIALS AND METHODS

Patient material

All cases of ocular and ocular adnexal leukaemic lesions from 1980 to 2009 were collected from the Danish Register of Pathology. In six cases, medical records and formalin-fixed, paraffin-embedded (FFPE) tissue blocks were available. Also included were three leukaemias with ophthalmic manifestations from the Department of Pathology, Royal Liverpool University Hospital, UK. The following information was collected from the patients' medical records: age, sex, diagnosis, histopathological findings, cytogenetic data, molecular genetic data (when available), location of the lesion, symptoms, treatment and clinical follow-up.

Histopathology and immunohistochemistry

The orbital and ocular biopsies were evaluated on sections stained with H&E and periodic acid-Schiff. For immunohistochemistry, FFPE tissue blocks were cut into sections 4 µm thick and mounted on slides. Stains were performed using the streptavidin-biotin method. Antibodies against CD3, CD10, CD13, CD15, CD20, CD33, CD34, CD43, CD45, CD79α, CD117, MPO, TLC1, BCL-2, TdT and lysozyme were used in most cases. The expression of the MYB oncoprotein (SPM-175; Santa-Cruz, Dallas, Texas, USA) was studied as described.⁸ Samples were scored as MYB positive when >25% of the neoplastic cells stained positive for MYB. Negative and positive controls were included in all stains, and internal positive controls were evaluated where appropriate. For analyses of immunostainings, positive tumour cells were counted in five high-power fields (×400).

Array comparative genomic hybridisation (arrayCGH) analysis

Genomic DNA from FFPE blocks was isolated with the DNeasy Blood and Tissue Kit (Qiagen, Hilden, Germany). In seven cases, there was enough DNA available for arrayCGH analysis with Human Genome CGH Microarray 244K oligonucleotide arrays (Agilent Technologies, Santa Clara, California, USA).⁸ Data were

analysed with NEXUS Copy Number V.7.0 Discovery Edition (BioDiscovery, El Segundo, California, USA).⁸ The settings for aberration calls were 1.5 for amplification, 0.3 for gain, -0.3 for loss and -1.5 for homozygous deletion. The FASST2 segmentation algorithm was used to define non-random regions of copy number alterations (CNAs) across the genome at a significance threshold of $p=1.0E-8$. In samples from cases 2 and 5, the settings for aberration calls were 1.5 for amplification, 0.5 for gain, -0.5 for loss and -1.5 for homozygous deletion at a significance threshold of $p=1.0E-18$. The accuracy of each aberration call was confirmed manually.

Fluorescence in situ hybridization (FISH)

Rearrangements of *ETV6* and *KMT2A* were analysed on 5 µm FFPE sections with FISH dual-colour break probes (Leica Biosystems, Wetzlar, Germany). The protocols for pre-treatment, hybridisation and post-hybridisation washes were as recommended by the manufacturer. Fluorescence signals were digitised, processed and analysed with the Isis FISH imaging system V.5.5 (MetaSystems, Altlußheim, Germany). At least 50 nuclei were scored for each probe and case.

Patient and public involvement

Patients and the public were not involved in the design, conduct and reporting of the research. However, permission was obtained to include photographs of two of the patients in the publication.

RESULTS

Clinical characteristics of primary ophthalmic leukaemias

We identified three cases of acute leukaemias with primary ophthalmic manifestations in the Danish Register of Pathology from 1980 to 2009. The clinical, cytogenetic and molecular genetic findings are summarised in table 1.

Case 1 was an otherwise healthy 5-year-old boy with left-sided exophthalmos, a swollen lacrimal gland and a bluish-red discolouration of the eyelid. The eye examination, including visual acuity, was normal. A CT scan showed a homogeneous mass in the left orbit and displacement of the optic nerve. The lateral rectus muscle was surrounded by neoplastic tissue. Physical examination revealed enlarged lymph nodes on the neck and testis. Analysis of peripheral blood showed anaemia, and bone marrow and testis biopsies revealed lymphoblastic leukaemia cells positive for CD3, CD10 and CD79α. The cellular morphology and immunoprofile were consistent with BCP-ALL. The patient responded well to chemotherapy (NOPHO ALL-92 protocol) and was still in complete remission at the last follow-up, 13 years after diagnosis.

Case 2 was a 9-year-old girl with swelling and redness of the left eyelid. Her visual acuity was normal, but she had ptosis and decreased mobility of the eyelid (figure 1A). The patient was initially treated with antibiotics, but the lesion continued to enlarge. A preauricular lymph node

Table 1 Clinical and cytogenetic findings and gene rearrangements/mutations in nine cases of acute leukaemia with ophthalmic manifestations

Case	Age (years)/sex	Diagnosis	Location	Karyotype/chromosome translocation	Gene rearrangement/mutation	MYB expression	Clinical follow-up
1	5/M	BCP-ALL	Superior orbital region (left)*	46, XY, t(2;3)(p11;q29) [11]/46 XY [16]	No <i>ETV6</i> rearrangement†	+	NED after 13 years
2	9/F	BCP-ALL	Superior orbital region (left)*	47, XX, t(12;21)(p13;q22), +21	<i>ETV6</i> rearrangement†	+	NED after 5 years
3	17/M	BCP-ALL	Bilateral uveal and retinal leukaemic infiltrates, optic nerve invasion (left)	NDA	<i>ETV6</i> rearrangement†	-	Orbital lesion after 1 year, DOD after 1.3 years
4	32/M	BCP-ALL	Leukaemic infiltrate of the iris (right)	46, XY [25]	NDA	NDA	Relapses after 6 and 27 years, ocular lesion after 28 years, DOD after 29 years
5	1/M	AML FAB M5	Inferior orbital region (left)*	46, XY, t(9;1)(p22;q23) [7]	<i>KMT2A</i> rearrangement†	+	NED after 18 years
6	40/F	AML FAB M4	Orbital region (left)	46XX, -7, +11, inv(16)(p13q22)	No <i>KMT2A</i> rearrangement†	+	Orbital lesion after 2 years, DOD after 5 years
7	68/M	AML FAB M1	Inferior orbital region (left)	46, XY [25]	No <i>KMT2A</i> rearrangement†	+	Relapse after 2 years, orbital lesion after 3 years, DOC after 3.5 years
8	70/F	AML FAB M2	Retinal and subretinal infiltrate (left)	NDA	<i>FLT3</i> ITD mutation <i>NPM1</i> mutation	NDA	Ocular lesion after 9 months, relapse 1.5 years, DOD after 2 years
9	68/F	CLL, high-grade transformation to AML FAB M2	Choroid, conjunctiva, and anterior orbital region (right)	t(8;21)(q22;q22)	<i>RUNX1-RUNX1T1</i> gene fusion	-	DOD

*Primary ophthalmic lesion.

†FISH analysis.

AML, acute myeloid leukaemia; BCP-ALL, B-cell precursor acute lymphoblastic leukaemia; CLL, chronic lymphocytic leukaemia; DOC, dead of other causes; DOD, dead of disease; F, female; ITD, internal tandem duplication in juxtamembrane domain; M, male; NDA, no data available; NED, no evidence of disease.




Figure 1 (A) 9-year-old girl (case 2) with left-sided proptosis, discoloration of the upper eye lid and ptosis. (B) Patient in (A) after 2 months of treatment. (C) 1-year-old boy (case 5) with left-sided proptosis and oedema of both eyelids. (D) MRI scan of the orbits (coronal view) of the patient in (C) shows a homogeneous mass involving the inferior half of the left orbit (asterisk).

and several submandibular nodes were swollen. The eyeball was displaced downward and medially. A CT scan revealed a mass in the superior orbital region involving the eyelid. A bone marrow biopsy showed infiltration of malignant lymphoblastic cells positive for CD10, CD20, CD79 α , CD43, TdT and BCL-2. The cellular morphology and immunoprofile were consistent with BCP-ALL. She responded well to chemotherapy (NOPHO ALL 2000 protocol) (figure 1B) and was still in complete remission at the latest follow-up, 5 years after diagnosis.

Case 5 was a 1-year-old boy with fever and left-sided proptosis of a few weeks' duration. He had oedema of both eyelids, bluish discoloration of the inferior lid and proptosis (4 mm) (figure 1C). CT and MRI showed a large homogeneous mass involving the inferior half of the left orbit, extending from the inferior lid to the orbital apex (figure 1D). Orbital and bone marrow biopsies revealed leukaemic infiltrates with neoplastic cells positive for Sudan black B, lysozyme, CD43 and CD45. The microscopic findings and the immunoprofile were consistent with AML, FAB type M5. Blood analysis showed pancytopenia. The patient had several blood transfusions and received chemotherapy (NOPHO AML 1993 protocol). He is still in remission, 18 years after diagnosis.

None of the three paediatric patients with primary orbital presentation of disease had any evidence of leukaemic infiltrates in the retina or choroid.

Clinical characteristics of secondary ophthalmic leukaemias

Six patients were diagnosed with systemic leukaemia before ophthalmic symptoms were present. Orbital/ocular manifestations were evident at a mean age of 49.2 years (range, 17–70 years). The leukaemic lesions were bilateral in one case, affected the left eye in three cases and the right eye in two. The average duration of symptoms before medical consultation was 5 weeks (range, 2–9 weeks). All patients with orbital tumours had proptosis, displacement of the eye and restricted eye mobility. Two patients with ocular manifestations had leukaemic infiltrates in the retina and subretinally; one of these patients also had involvement of the uvea and infiltration of the optic nerve. Two additional patients had infiltrates in the choroid, iris and anterior segment. Two patients with ocular lesions had pain from the eye; one also had blurred vision, and an examination revealed a visual acuity of 0.25, vitreous haze and papilloedema. One patient had a palpable mass. The lesions were diagnosed as BCP-ALL in two cases (figure 2A, B) and AML




Figure 2 (A) Lymphoblastic cells with irregular nuclei and sparse cytoplasm infiltrating the iris (H&E staining) in a patient with B-cell precursor acute lymphoblastic leukaemia (case 4); (B) lymphoblastic cells from case 4 are strongly immunoreactive for terminal deoxynucleotidyl transferase (TdT); (C) myeloblastic cells with an eosinophilic cytoplasm and indistinct cell boundaries and lymphocytes infiltrating the orbital fat tissue (H&E staining) in a patient with acute myeloid leukaemia Fab M1 (case 7); (D) myeloblastic cells from case 7 are strongly immunoreactive for myeloperoxidase; (E–H) expression of the MYB oncoprotein in ocular leukaemic lesions from case 1, B-cell precursor acute lymphoblastic leukaemia (E), case 2, B-cell precursor acute lymphoblastic leukaemia (F), case 5, acute myeloid leukaemia Fab M5 (G), and case 7, acute myeloid leukaemia Fab M1 (H).

in three (figure 2C, D); one patient had a history of chronic lymphocytic leukaemia (CLL) with high-grade transformation to AML (table 1). All six patients received chemotherapy; two patients also received allogeneic bone marrow transplants, and two had radiotherapy. Four of the six patients died of relapse and/or disseminated disease 4–36 months after the onset of eye symptoms, one died of disseminated disease an unknown number of months after diagnosis and one died of heart failure 1.5 years after relapse.

Expression of the MYB oncoprotein

Two of the three analysed BCP-ALLs showed strong nuclear immunostaining for the MYB oncoprotein in the majority of leukaemic cells (table 1 and figure 2E, F). Three of four analysed AML samples also had strong

nuclear staining in the majority of neoplastic cells (table 1 and figure 2G, H). The MYB oncoprotein was not expressed in one BCP-ALL (case 3) and one AML (case 9).

Cytogenetic and molecular genetic characteristics

Cytogenetic information was available from three patients with BCP-ALL and four with AML. The karyotypic alterations are shown in table 1. Two of the three ALLs had abnormal karyotypes, and one had no cytogenetic changes (case 4). Case 2 had the classical t(12;21) (p13;q22) translocation associated with paediatric BCP-ALL, whereas case 1 had an uncommon t(2;3) (p11;q29) seen in a small subset of BCP-ALL. The case with the t(12;21) had a rearrangement of *ETV6* consistent with an *ETV6*–*RUNX1* gene fusion. FISH analysis also revealed an *ETV6* rearrangement in case 3 (figure 3A); case 1 had no evidence of *ETV6* rearrangement. Similarly, three of the four AMLs had abnormal karyotypes: case 5 had a t(9;11) (p22;q23) typical of the M5 subtype; case 6 had an inv(16) (p13q22), monosomy 7, and trisomy 11; and case 9 had a t(8;21) (q22;q22) resulting in a *RUNX1*–*RUNX1T1* fusion. The fourth AML had an apparently normal karyotype (case 7). FISH analysis revealed that neither case 6 nor case 7 had any rearrangements of *KMT2A*, whereas case 5 had a rearranged *KMT2A* allele (figure 3B). Nucleotide sequence analysis revealed that case 8 (AML) had an *FLT3* internal tandem duplication mutation and an exon 12 *NPM1* mutation (data not shown).

Genomic profiling

Genome-wide arrayCGH yielded analysable results from six of seven leukaemic patients with ocular involvement (table 2), three of which had primary ocular lesions (cases 1, 2 and 5). One BCP-ALL (case 1) and one AML (case 6) had no CNAs; the four other cases had an average of 3.3 CNAs per case (range 1–7) (table 2). One homozygous deletion, including the tumour suppressor *CDKN2A*, was detected in a BCP-ALL (case 3) (figure 3C). Case 2 (BCP-ALL) had gain of 21q21.1–q22.3, including the *RUNX1*, *ERG* and *ETS2* oncogenes (figure 3D). Interestingly, this case had also gain of a 0.5 Mb segment in 12p13.2 and a breakpoint in *ETV6*, consistent with an *ETV6*–*RUNX1* gene fusion. There were no high-level gene amplifications and no recurrent CNAs.

DISCUSSION

Here, we present a comprehensive clinical and genomic profiling study of nine leukaemias with orbital and/or ocular manifestations. Three were primary orbital manifestations of the leukaemia, representing all such cases histologically analysed in Denmark from 1980 to 2009. The remaining six cases had secondary orbital/ocular lesions. Four of our patients had BCP-ALL, and five had AML, one of which was originally a CLL. Transformation of CLL to high-grade AML is an uncommon event




Figure 3 FISH and arrayCGH analyses of acute leukaemias with ophthalmic manifestations. (A) FISH analysis showing a rearranged *ETV6* allele (split red and green signals indicated by arrowheads) in a B-cell precursor acute lymphoblastic leukaemia (case 3). (B) FISH analysis showing a rearranged *KMT2A* allele (split red and green signals indicated by arrowheads) in a patient with acute myeloid leukaemia FAB M5 and a t(9;11) translocation (case 5). (C) ArrayCGH analysis showing homozygous loss of the tumour suppressor gene *CDKN2A* (arrow) in a B-cell precursor acute lymphoblastic leukaemia (case 3). (D) ArrayCGH analysis showing gain of 21q21.1–q22.3, including the *RUNX1*, *ERG* and *ETS2* oncogenes, and loss of the terminal end of 21q in a B-cell precursor acute lymphoblastic leukaemia (case 2).

and is associated with a poor prognosis.⁹ Thus, all nine ophthalmic lesions in this study were acute leukaemias.

The average age at diagnosis of our patients with primary ophthalmic lesions was 5 years (24.2 years for all our patients), and there were similar numbers of males and females. The patients with orbital tumours presented with proptosis, displacement of the eye and reduced eye mobility. Two patients with ocular infiltrations had pain, and one also had reduced visual acuity. Notably, none of the paediatric patients with primary orbital manifestations had leukaemic infiltrates in the retina or choroid, whereas four of six patients with secondary ophthalmic leukaemias had such infiltrates. Taken together, our findings further emphasise the broad clinical spectrum of leukaemic lesions that may manifest in the eye or ocular

region,^{1 3 4 6 7 10 11} and the need to consider leukaemia in the differential diagnosis of patients with proptosis, reduced mobility and/or displacement of the eye.^{1 4 6}

The molecular mechanisms by which leukaemic cells give rise to extramedullary dissemination remain elusive. Thus, it is unclear why certain leukaemias present with orbital and/or ocular manifestations. There is, however, evidence suggesting that chemokines may be involved in organ-specific homing of neoplastic cells.¹² Interestingly, ALL cells frequently express the chemokine receptors CXCR4 and CXCR3 and CLL cells the CXCR4 and CCR7 receptors.¹² There is also a recent report demonstrating that $\alpha 6$ integrin, which frequently is overexpressed in ALL, interacts with laminin and mediates the migration of ALL cells to the central nervous system.¹³ Further

Table 2 ArrayCGH analysis of seven cases of acute leukaemias with ophthalmic manifestations

Case	Diagnosis	CNA†	Cytoband	Chromosome region	Length (bp)	Number of genes	Candidate genes
1	BCP-ALL*			No CNAs			
2	BCP-ALL*	Gain	5q33.3	157 399 690–158 503 889	1 104 199	2	
		Gain	12p13.2	11 547 615–12 038 149	490 525	2	<i>ETV6</i>
		Gain	12q14.3	66 617 004–67 627 932	1 010 928	5	
		Loss	16p13.3	5 063 719–6 565 236	1 501 517	9	
		Loss	17q11.2	28 211 019–29 173 479	962 460	16	
		Gain	21q21.1–q22.3	21 672 155–43 977 574	22 305 419	209	<i>RUNX1, ERG, ETS2</i>
3	BCP-ALL	Loss	1pter–p35.23	0–7 329 451	7 329 451	156	
		Gain	1q21.1–qter	142 764 722–249 250 621	106 485 899	1267	
		Loss	9p21.3	20 612 727–22 192 890	1 580 163	30	<i>CDKN2A</i>
5	AML FAB M5*	Loss	11q13.5–qter	75 898 626–135 006 516	59 107 890	493	
		Loss	20q13.33	59 008 978–60 021 837	1 012 859	5	
6	AML FAB M4			No CNAs			
7	AML FAB M1	Loss	7q21.2–q36.3	92 294 039–156 451 959	64 157 920	618	
9	AML FAM M2			NA			

*Primary ophthalmic lesion.

AML, acute myeloid leukaemia; BCP-ALL, B-cell precursor acute lymphoblastic leukaemia; CNA, copy number alteration; NA, not analysable because of poor array quality.

studies are, however, needed to elucidate the exact mechanisms behind dissemination of acute leukaemias to the eye or ocular region.

The three patients with primary leukaemic orbital lesions were in complete remission 6, 9 and 18 years, respectively, after diagnosis. In previous studies, patients with orbital or ocular involvement of leukaemia had a poor prognosis and short overall survival since eye involvement often indicates recurrent disease.^{2 3} In our patients diagnosed with leukaemia before orbital/ocular lesions occurred, the median survival was 1.06 years (range, 2 months to 2 years). In a comprehensive study from the Children's Oncology Group including 1459 paediatric patients with AML, those with involvement of orbital and central nervous system sites had a significantly better survival than patients with AML outside the central nervous system, those with leukaemia in the cerebrospinal fluid and those with no extramedullary leukaemia; the overall survival of the patients with AML with orbital involvement in this study was 92%.¹⁴

Cytogenetic data were available from three of four BCP-ALLs and four of five AMLs (table 1). Five cases had abnormal karyotypes and two had apparently normal karyotypes. The cytogenetic aberrations in these cases are similar to those in leukaemias without ophthalmic manifestations.^{15 16} FISH analysis revealed *ETV6* rearrangements in two of three BCP-ALLs (table 1),

consistent with *ETV6*-associated translocations. Indeed, case 2 had a t(12;21) translocation commonly associated with the *ETV6*–*RUNX1* gene fusion seen in approximately 25% of paediatric ALLs.¹⁷ Patients with this fusion usually have a favourable prognosis.¹⁷ Similarly, FISH analysis revealed a rearrangement of *KMT2A* in one of three AML samples analysed. This case had a t(9;11)(p22;q23) known to result in a *KMT2A*–*MLLT3* fusion.¹⁸ The prognostic significance of the t(9;11) is controversial. Recent studies suggest that this translocation carries an intermediate risk.^{19 20} Our patient with the t(9;11) was alive with no evidence of disease 18 years after diagnosis.

The genomic profiles of the orbital/ocular leukaemic lesions were further characterised by high-resolution arrayCGH (table 2). One BCP-ALL and one AML had no CNAs; the remaining four cases had rather quiet genomes. These findings are consistent with studies of leukaemic cells in bone marrow and/or peripheral blood,^{21–24} and with the observation that fusion gene-driven neoplasms often have few other genomic alterations.²⁵ Notably, case 2 had gain of 21q21.1–q22.3, including the *RUNX1*, *ERG* and *ETS2* oncogenes, and case 3 had a 1.5 Mb homozygous deletion in 9p21.3 including the *CDKN2A* tumour suppressor gene. This gene is frequently deleted in ALL and is associated with a poor prognosis and a poor response to treatment.^{26 27}

MYB encodes a transcription factor that is important in the control of cell division, apoptosis and differentiation of haematopoietic stem/progenitor cells.^{28,29} *MYB* is also an oncogene that is activated and overexpressed in subsets of acute leukaemias and in certain solid tumours.^{28–31} Here, we show for the first time that *MYB* is overexpressed in ophthalmic lesions in patients with acute leukaemias. In these cases, *MYB* is likely to be an important driver of leukaemogenesis and, therefore, also a potential therapeutic target. Further studies of *MYB* in acute leukaemias may therefore lead to better treatments for these malignancies.

In summary, we present the first comprehensive, integrated clinical, cytogenetic and genomic analyses of nine acute leukaemias with ophthalmic manifestations. These leukaemias did not differ significantly from those without clinically visible ophthalmic manifestations with regard to immunophenotype, cytogenetic aberrations, gene fusions and CNAs. Awareness of the clinical spectrum of leukaemic lesions of the eye or ocular region is important to quickly establish the correct diagnosis and commence prompt treatment.

Acknowledgements We thank Therese Carlsson for excellent technical assistance.

Contributors LSS and SH planned and designed the study; LSS, A-CL and SC collected patient data; SH and SC performed pathology reviews; MP and MA performed experiments; LSS, MP, MA, SH, A-CL and GS collected and analysed data; LSS and GS drafted the manuscript. All authors contributed to the revision of the manuscript and approval of the final version.

Funding This study was supported by a grant from Synoptik-Fonden, Denmark and the Swedish Children's Cancer Foundation.

Competing interests None declared.

Patient consent for publication Obtained.

Ethics approval This study was approved by the Local Scientific Ethics Committee (journal no. H-4-2013-003), the Danish Data Protection Agency (journal no. 2012-41-0747) and the Health Research Authority of England (REC Ref. 11/NW0759).

Provenance and peer review Not commissioned; externally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Lisa Stenman Skarsgård <http://orcid.org/0000-0002-3662-6676>
Steffen Heegaard <http://orcid.org/0000-0001-5906-7670>

REFERENCES

- Kincaid MC, Green WR. Ocular and orbital involvement in leukemia. *Surv Ophthalmol* 1983;27:211–32.
- Russo V, Scott IU, Querques G, et al. Orbital and ocular manifestations of acute childhood leukemia: clinical and statistical analysis of 180 patients. *Eur J Ophthalmol* 2008;18:619–23.
- Bidar M, Wilson MW, Laquis SJ, et al. Clinical and imaging characteristics of orbital leukemic tumors. *Ophthalmic Plast Reconstr Surg* 2007;23:87–93.
- Sharma T, Grewal J, Gupta S, et al. Ophthalmic manifestations of acute leukaemias: the ophthalmologist's role. *Eye* 2004;18:663–72.
- Johansen S, Heegaard S, Bogeskov L, et al. Orbital space-occupying lesions in Denmark 1974–1997. *Acta Ophthalmol Scand* 2000;78:547–52.
- Reddy SC, Jackson N, Menon BS. Ocular involvement in leukemia—a study of 288 cases. *Ophthalmologica* 2003;217:441–5.
- Murthy R, Vemuganti GK, Honavar SG, et al. Extramedullary leukemia in children presenting with proptosis. *J Hematol Oncol* 2009;2.
- von Holstein SL, Fehr A, Persson M, et al. Adenoid cystic carcinoma of the lacrimal gland: Myb gene activation, genomic imbalances, and clinical characteristics. *Ophthalmology* 2013;120:2130–8.
- Tambaro FP, Garcia-Manero G, O'Brien SM, et al. Outcomes for patients with chronic lymphocytic leukemia and acute leukemia or myelodysplastic syndrome. *Leukemia* 2016;30:325–30.
- Hmidi K, Zaouali S, Messaoud R, et al. Bilateral orbital myeloid sarcoma as initial manifestation of acute myeloid leukemia. *Int Ophthalmol* 2007;27:373–7.
- Maka E, Lukáts O, Tóth J, et al. Orbital tumour as initial manifestation of acute myeloid leukemia: granulocytic sarcoma: case report. *Pathol Oncol Res* 2008;14:209–11.
- Zlotnik A, Burkhardt AM, Homey B. Homeostatic chemokine receptors and organ-specific metastasis. *Nat Rev Immunol* 2011;11:597–606.
- Yao H, Price TT, Cantelli G, et al. Leukaemia hijacks a neural mechanism to invade the central nervous system. *Nature* 2018;560:55–60.
- Johnston DL, Alonzo TA, Gerbing RB, et al. Superior outcome of pediatric acute myeloid leukemia patients with orbital and CNS myeloid sarcoma: a report from the Children's Oncology Group. *Pediatr Blood Cancer* 2012;58:519–24.
- Harrison CJ, Johansson B. Acute lymphoblastic leukemia. In: Heim S, Mitelman F, eds. *Cancer cytogenetics*. New Jersey: Wiley-Blackwell, 2010: 237–40.
- Johansson B, Harrison CJ. Acute myeloid leukemia. In: Heim S, Mitelman F, eds. *Cancer cytogenetics*. New Jersey: Wiley-Blackwell, 2010: 48–9.
- Hunger SP, Mullighan CG. Acute lymphoblastic leukemia in children. *N Engl J Med* 2015;373:1541–52.
- Iida S, Seto M, Yamamoto K, et al. MLLT3 gene on 9p22 involved in t(9;11) leukemia encodes a serine/proline rich protein homologous to MLLT1 on 19p13. *Oncogene* 1993;8:3085–92.
- von Neuhoff C, Reinhardt D, Sander A, et al. Prognostic impact of specific chromosomal aberrations in a large group of pediatric patients with acute myeloid leukemia treated uniformly according to trial AML-BFM 98. *J Clin Oncol* 2010;28:2682–9.
- Döhner H, Weisdorf DJ, Bloomfield CD. Acute myeloid leukemia. *N Engl J Med* 2015;373:1136–52.
- Yasar D, Karadogan I, Alanoglu G, et al. Array comparative genomic hybridization analysis of adult acute leukemia patients. *Cancer Genet Cytogenet* 2010;197:122–9.
- van der Veken LT, Buijs A. Array CGH in human leukemia: from somatics to genetics. *Cytogenet Genome Res* 2011;135:260–70.
- Zakaria Z, Ahid MFM, Ismail A, et al. Chromosomal aberrations in ETV6/RUNX1-positive childhood acute lymphoblastic leukemia using 244K oligonucleotide array comparative genomic hybridization. *Mol Cytogenet* 2012;5:41.
- Sarhadi VK, Lahti L, Scheinin I, et al. Targeted resequencing of 9p in acute lymphoblastic leukemia yields concordant results with array CGH and reveals novel genomic alterations. *Genomics* 2013;102:182–8.
- Andersson MK, Stenman G. The landscape of gene fusions and somatic mutations in salivary gland neoplasms—implications for diagnosis and therapy. *Oral Oncol* 2016;57:63–9.
- Braun M, Pastorczak A, Fendler W, et al. Biallelic loss of *CDKN2A* is associated with poor response to treatment in pediatric acute lymphoblastic leukemia. *Leuk Lymphoma* 2017;58:1162–71.
- Fang Q, Yuan T, Li Y, et al. Prognostic significance of copy number alterations detected by multi-link probe amplification of multiple genes in adult acute lymphoblastic leukemia. *Oncol Lett* 2018;15:5359–67.
- Ramsay RG, Gonda TJ. Myb function in normal and cancer cells. *Nat Rev Cancer* 2008;8:523–34.
- Stenman G, Andersson MK, Andrén Y. New tricks from an old oncogene: gene fusion and copy number alterations of Myb in human cancer. *Cell Cycle* 2010;9:2986–95.
- Pattabiraman DR, Gonda TJ. Role and potential for therapeutic targeting of Myb in leukemia. *Leukemia* 2013;27:269–77.
- Andersson MK, Afshari MK, Andrén Y, et al. Targeting the oncogenic transcriptional regulator Myb in adenoid cystic carcinoma by inhibition of IGF1R/AKT signaling. *J Natl Cancer Inst* 2017;109.