



ELSEVIER

Contents lists available at [ScienceDirect](http://ScienceDirect)

## Data in Brief

journal homepage: [www.elsevier.com/locate/dib](http://www.elsevier.com/locate/dib)

## Data Article

## Data supporting the cardiac mitochondria calcium handling in female normotensive and spontaneously hypertensive rats



Alejandro Ciocci Pardo, Gustavo J. Rinaldi, Susana M. Mosca\*

*Centro de Investigaciones Cardiovasculares "Dr. Horacio E. Cingolani", CCT-CONICET, Facultad de Ciencias Médicas, Universidad Nacional de La Plata, Argentina*

## ARTICLE INFO

*Article history:*

Received 22 December 2015

Received in revised form

4 February 2016

Accepted 20 February 2016

Available online 27 February 2016

*Keywords:*

Cardiac mitochondria

Calcium

Female WKY

Female SHR

## ABSTRACT

In association with the published article "Mitochondrial calcium handling in normotensive and spontaneously hypertensive rats: correlation with systolic blood pressure levels" [1], this data article contains information about calcium handling of cardiac mitochondria isolated from female of both rats strains (WKY and SHR). Dataset of mitochondrial permeability transition pore (mPTP) resistance to opening  $\text{Ca}^{2+}$ -mediated,  $\text{Ca}^{2+}$  retention capacity (CRC), time constants and mitochondrial membrane potential ( $\Delta\Psi_m$ ) are showed.

© 2016 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license

(<http://creativecommons.org/licenses/by/4.0/>).

## 1. Specifications Table

|                            |                                         |
|----------------------------|-----------------------------------------|
| Subject area               | <i>Cardiovascular</i>                   |
| More specific subject area | <i>Mitochondria</i>                     |
| Type of data               | <i>Table and figures</i>                |
| How data was acquired      | <i>Spectrofluorometer Hitachi F4500</i> |
| Data format                | <i>Mean ± SE</i>                        |

DOI of original article: <http://dx.doi.org/10.1016/j.mito.2014.12.003>

\* Corresponding author.

E-mail address: [smosca@med.unlp.edu.ar](mailto:smosca@med.unlp.edu.ar) (S.M. Mosca).

<http://dx.doi.org/10.1016/j.dib.2016.02.056>

2352-3409/© 2016 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

|                       |                                                                                                                                                                                              |
|-----------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Experimental factors  | <i>Isolated mitochondria from hearts of female normotensive (WKY) and spontaneously hypertensive rats (SHR).</i>                                                                             |
| Experimental features | mPTP resistance to opening $\text{Ca}^{2+}$ -mediated, $\text{Ca}^{2+}$ -retention capacity, time constants and mitochondrial membrane potential were tested to assess the calcium handling. |
| Data source location  | <i>La Plata (Buenos Aires), Argentina</i>                                                                                                                                                    |
| Data accessibility    | <i>Data are within this article</i>                                                                                                                                                          |

## 2. Value of the data

- The data provide information about calcium handling of cardiac mitochondria obtained from female normotensive and hypertensive rats.
- The data can be compared to data obtained in male rats of both strains [1].
- The data can be used in the development of further experiments to analyze the influence of sex.

## 3. Data

Systolic blood pressure (SBP), body weight (BW), heart weight (HW) and hypertrophy index (HI, calculated as HW and BW ratio) of female WKY and SHR are displayed in Table 1. The mPTP resistance to opening, measured as light scattering decrease, after different  $\text{Ca}^{2+}$  concentrations in samples derived from female of both rats strains was depicted in Fig. 1.  $\text{Ca}^{2+}$  retention capacity (CRC), the time of each pulse (PT) and exponential decay constant (EDC), obtained after addition of successive pulses of  $10 \mu\text{M}$   $\text{Ca}^{2+}$  are shown in Figs. 2 and 3. Rhodamine fluorescence changes used for the membrane potential ( $\Delta\Psi\text{m}$ ) determination in cardiac mitochondria of WKY and SHR hearts appear in Fig. 4.

## 4. Experimental design, materials and methods

### 4.1. Animal model

We used female normotensive (Wistar Kyoto, WKY) and spontaneously hypertensive rats (SHR) of 5 months of age. All procedures followed during this investigation conform to the Guide for the Care and Use of Laboratory Animals published by the US National Institutes of Health (NIH Publication no. 85-23, revised 1996) and to the guidelines laid down by the Animal Welfare Committee of La Plata School of Medicine.


### 4.2. Characteristics of WKY and SHR

Values of SBP, BW, HW and HI in WKY and SHR are shown in Table 1.


**Table 1**  
Values of SBP, BW, HW and HI for female WKY and SHR.

|            | WKY         | SHR          |
|------------|-------------|--------------|
| SBP (mmHg) | 118 ± 7     | 160 ± 5*     |
| BW (g)     | 217 ± 2     | 218 ± 7      |
| HW (mg)    | 595 ± 30    | 732 ± 40*    |
| HI         | 2.77 ± 0.16 | 3.24 ± 0.14* |

\*  $p < 0.05$  SHR vs WKY.




**Fig. 1.** Light scattering decrease (LSD) in arbitrary units (a.u.) after the addition of different  $\text{Ca}^{2+}$  concentrations (20, 50, 100, 200, 500 and 1000  $\mu\text{M}$ ) to samples of mitochondria isolated from female WKY and SHR hearts. Inset: typical trace of light scattering for SHR and WKY at  $\text{Ca}^{2+}$  200  $\mu\text{M}$ . Note that only at  $\text{Ca}^{2+}$  50  $\mu\text{M}$  LSD was significantly lesser in SHR than WKY. \* $p < 0.05$ .




**Fig. 2.** Changes of calcium green fluorescence after  $\text{Ca}^{2+}$  addition in samples of mitochondria from female WKY and SHR hearts. Inset:  $\text{Ca}^{2+}$  retention capacity (CRC) in both rat strains. Note that no statistically significant difference between WKY and SHR was detected.

#### 4.3. Isolation of rat heart mitochondria

Mitochondria were obtained by differential centrifugation following the method described by Mela and Seitz [2].



**Fig. 3.** Pulse time (PT) and exponential decay constant (EDC) of WKY and SHR female. Both parameters were greater in SHR than WKY. \* $p < 0.05$ .



**Fig. 4.** Typical traces of fluorescence changes and mean values of mitochondrial membrane potential ( $\Delta\Psi_m$ ) of samples obtained from female WKY and SHR hearts.  $\Delta\Psi_m$  was similar in both rats strains.

#### 4.4. Mitochondrial permeability transition pore (mPTP) resistance to opening $Ca^{2+}$ -mediated

The mPTP resistance to opening was assessed by addition of different  $Ca^{2+}$  concentrations to samples of 0.3 mg/mL of isolated mitochondria from hearts of WKY and SHR female. The changes were observed as decreases of light scattering and followed using a temperature-controlled Hitachi F4500 spectrofluorometer operating with continuous stirring at excitation and emission wavelengths of 520 nm [3]. Light scattering decrease (LSD) was calculated for each sample by taking the difference of scattered light between before and after the addition of  $CaCl_2$  (Fig. 1).

#### 4.5. Calcium retention capacity (CRC)

CRC (Fig. 2) was defined as the amount of  $Ca^{2+}$  required triggering a massive  $Ca^{2+}$  release by isolated cardiac mitochondria [4]. Briefly, successive pulses of 10  $\mu M$   $Ca^{2+}$  were added to samples of 0.3 mg/mL of isolated mitochondria of hearts from WKY and SHR. After sufficient  $Ca^{2+}$  loading, extramitochondrial  $Ca^{2+}$  concentration abruptly increased, which was recorded by fluorescence changes of Calcium green-5N with excitation and emission wavelengths set at 506 and 532 nm, respectively. Pulse time (PT, s) and exponential decay constant (EDC) of  $Ca^{2+}$  influx to mitochondria were also measured (Fig. 3).

#### 4.6. Mitochondrial membrane potential ( $\Delta\Psi_m$ )

Mitochondrial potential changes were evaluated by measuring rhodamine-123 (RH-123) fluorescence quenching under the buffer described above containing RH-123 0.1  $\mu\text{M}$  [5] and using 0.1 mg/mL of isolated mitochondria.  $\Delta\Psi_m$  was calculated following the instructions previously detailed by Scaduto and Grotyohann [6], using the Nernst–Guggenheim equation (Fig. 4).

#### Acknowledgments

This work was supported in part by the Grant PIP 0229 from Consejo Nacional de Investigaciones Científicas y Tecnológicas (CONICET) of Argentina to Dr. Susana M. Mosca.

#### Appendix A. Supplementary material

Supplementary data associated with this article can be found in the online version at <http://dx.doi.org/10.1016/j.dib.2016.02.056>.

#### References

- [1] A.C. Pardo, G.J. Rinaldi, S.M. Mosca, Mitochondrial calcium handling in normotensive and spontaneously hypertensive rats: correlation with systolic blood pressure levels, *Mitochondrion* 20 (2015) 75–81.
- [2] L. Mela, S. Seitz, Isolation of mitochondria with emphasis on heart mitochondria from small amounts of tissue, *Methods Enzymol.* 55 (1979) 39–46.
- [3] C.P. Baines, C.X. Song, Y.T. Zheng, G.W. Wang, J. Zhang, O.L. Wang, Y. Guo, R. Bolli, E.M. Cardwell, P. Ping, Protein kinase Cepsilon interacts with and inhibits the permeability transition pore in cardiac mitochondria, *Circ. Res.* 92 (8) (2003) 873–880.
- [4] F.N. Obame, C. Plan-Mercier, R. Assaly, R. Zini, J.L. Dubois-Randle, A. Berdeux, D. Morin, Cardioprotective effect of morphine and a blocker of glycogen synthase kinase 3 beta, SB216763[3-(2,4-dichlorophenyl)-4-(1-methyl-1H-indol-3-yl)-1H-pyrrole-2,5-dione], via inhibition of the mitochondrial permeability transition pore, *J. Pharmacol. Exp. Ther.* 326 (2008) 252–258.
- [5] R.K. Emaus, R. Grunwald, J.J. Lemasters, Rhodamine 123 as a probe of transmembrane potential in isolated rat liver mitochondria: spectral and metabolic properties, *Biochim. Biophys. Acta* 850 (1986) 436–448.
- [6] R.C. Scaduto Jr., L.W. Grotyohann, Measurement of mitochondrial membrane potential using fluorescent rhodamine derivatives, *Biophys. J.* 76 (1999) 469–477.