

Gut Microbiota and the Alteration of Immune Balance in Skin Diseases: From Nutraceuticals to Fecal Transplantation

Aleksandra Vojvodic¹, Zorica Peric-Hajzler², Dusica Matovic², Petar Vojvodic³, Tatjana Vlaskovic-Jovicevic³, Goran Sijan⁴, Sanja Dimitrijevic⁵, Nenad Stepic⁶, Uwe Wollina⁷, Badr Abdulwahab E. Badr⁸, Ashraf Badawi⁹, Mohamad Goldust¹⁰, Michael Tirant¹⁰, Van Thuong Nguyen¹¹, Massimo Fioranelli¹², Torello Lotti¹³

¹Department of Dermatology and Venereology, Military Medical Academy, Belgrade, Serbia; ²Military Medical Academy, Belgrade, Serbia; ³Clinic for Psychiatric Disorders "Dr. Laza Lazarevic", Belgrade, Serbia; ⁴Clinic for Plastic Surgery and Burns, Military Medical Academy, Belgrade, Serbia; ⁵Department of Gynecology, Military Medical Academy, Belgrade, Serbia; ⁶Chief of Clinic for Plastic Surgery and Burns, Military Medical Academy, Belgrade, Serbia; ⁷Department of Dermatology and Allergology, Städtisches Klinikum Dresden, Dresden, Germany; ⁸Faculty of Medicine – King Abdulaziz University, Jeddah, Saudi Arabia; ⁹Laser Institute, Cairo University, Cairo, Egypt; ¹⁰G. Marconi University, Rome, Italy; ¹¹National Hospital of Dermatology and Venereology, Hanoi, Vietnam; ¹²Department of Nuclear Physics, Sub-nuclear and Radiation, G. Marconi University, Rome, Italy; ¹³Department of Dermatology, G. Marconi University, Rome, Italy

Abstract

Citation: Vojvodic A, Peric-Hajzler Z, Matovic D, Vojvodic P, Vlaskovic-Jovicevic T, Sijan G, Dimitrijevic S, Stepic N, Wollina U, Badr BAE, Badawi A, Goldust M, Tirant M, Nguyen VY, Fioranelli M, Lotti T. Gut Microbiota and the Alteration of Immune Balance in Skin Diseases: From Nutraceuticals to Fecal Transplantation. Open Access Maced J Med Sci. 2019 Sep 30; 7(18):3034-3038. <https://doi.org/10.3889/oamjms.2019.827>

Keywords: Microbiota; Cytokines; Treatment; Inflammatory; Skin diseases; Immunology

***Correspondence:** Massimo Fioranelli. Department of Nuclear Physics, Sub-nuclear and Radiation, G. Marconi University, Rome, Italy. E-mail: massimo.fioranelli@gmail.com

Received: 10-Jul-2019; **Revised:** 08-Aug-2019; **Accepted:** 09-Aug-2019; **Online first:** 12-Sep-2019

Copyright: © 2019 Aleksandra Vojvodic, Zorica Peric-Hajzler, Dusica Matovic, Petar Vojvodic, Tatjana Vlaskovic-Jovicevic, Goran Sijan, Sanja Dimitrijevic, Nenad Stepic, Uwe Wollina, Badr Abdulwahab E. Badr, Ashraf Badawi, Mohamad Goldust, Michael Tirant, Van Thuong Nguyen, Massimo Fioranelli, Torello Lotti. This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0)

Funding: This research did not receive any financial support

Competing Interests: The authors have declared that no competing interests exist

The P.N.E.I. (Psycho-Neuro-Endocrine-Immunology) approach is represented by the interdisciplinary concept of bidirectional cross-talk between the psycho-neuro-endocrine and immune systems, which can influence the immune response. The well-known Gut-Brain Axis and the Gut-Skin Axis can be merged in a bigger network- the Gut-Brain-Skin Axis, with complex regulation by cytokines, neuro-peptides, neuro-hormones and another messenger (signalling) molecules and maybe the most important modulator of the Gut-Brain-Skin Axis/ the gut microbiota. The role of gut bacterial homeostasis is very important, and the homeostatic imbalance of the immune response may be a relevant etiologic/pathophysiologic factor for extra-intestinal and intestinal inflammatory, allergic and autoimmune diseases. The Low Dose Cytokines Medicine (LDM) is an innovative therapeutic approach. It is based on the most advanced knowledge in molecular biology and low dose pharmacology with the primary outcome. The SKA (Sequential Kinetic Activation) technology, codified and standardised by GUNA S.p.a. -Italy- makes the low doses of signalling molecules able to be active even below the minimum dose classically considered as effective and the significative efficacy of orally administered low-dose signalling molecules is the most representative aspect of LDM. The Physiologic Nutraceuticals and the Low Dose Medicine are two of the most promising approaches for the treatment of skin diseases based on the rebalance of the immune response and the recovery of gut dysbiosis.

Introduction

The P.N.E.I. (Psycho-Neuro-Endocrine-Immunology) approach is represented by the interdisciplinary concept of bidirectional cross-talk between the psycho-neuro-endocrine and immune systems, which can influence the immune response. Also, the immune system can influence the neuroendocrine functions, and this very complex interplay is mediated by a network of a lot of different

hormones, neuropeptides, growth factors, cytokines and other signalling molecules [1], [2], [3].

Skin and gut roles and relations with other organs and tissues are examples of the P.N.E.I. functioning. They are complex immune and neuroendocrine organs integrated into the whole immune-endocrine systems and the most important contact organs of all animals with the environment, with a lot of common characteristics: colonized by specific microbial strains, with a large number of vascular and neural structures, constantly exposed to

a heavy antigenic charge, represented by bacterial flora, with the tolerance of the commensal microbiota. The well-known Gut-Brain Axis and the Gut-Skin Axis can be merged in a larger network-the Gut-Brain-Skin Axis, with complex regulation by cytokines, neuropeptides, neuro-hormones and other messenger (signaling) molecules and maybe the most important modulator of the Gut-Brain-Skin Axis/the gut microbiota [2], [3], [4], [5].

Gut microbiota

It is well known that the gastrointestinal tract is colonised by about 100 trillion bacterial population (microbiota) composed of non-commensal (pathogenic) and commensal (nonpathogenic) species. Microbiota differs in composition and also in the function based on numerous factors such as the specific GI segment but also the diet of a subject, the gender, the age, etc. The nucleus of 57 species is common to all individuals, but each human being has a specific and unique microbiota. The dominant bacterial *fila*, constituting over 90% of the species present in the human intestine, are *Bacteroides* and *Firmicutes*, but also *Lactobacilli*, *Bacteroides*, *Proteobacteria*, *Bifidobacteria* and *Streptococci* are common [3], [4], [5], [6], [7].

The role of gut bacterial homeostasis is very important, and the homeostatic imbalance of the immune response may be a relevant etiologic factor for extra-intestinal and intestinal inflammatory, allergic and autoimmune diseases. This role is not limited to its systems but the whole organism [1], [4], [8].

Gram-negative bacteria, which contain lipopolysaccharide are considered as the main proinflammatory strains, responsible for the expression of pro-inflammatory cytokines such as Interleukin-1 (IL-1) and Interleukin-6 (IL-6) [2], [9].

Formation of biofilm is identified as one of the main forms of chronic intestinal, an adherent conglomerate to the intestinal epithelium having a composition like the extracellular matrix, in which bacteria and fungi proliferate and resid. The antibiotic treatment can affect mostly the bacterial infection caused by bacteria external to the biofilm, and an immune response is not able to be fully effective on biofilm. So, biofilm could be one of the most important etiologic agents of numerous chronic inflammatory diseases, promote the persistence of a Low-Grade Chronic Inflammation (LGCI) that can lead to the Leaky Gut Syndrome, connected with the onset of autoimmune and allergic diseases [1], [10], [11], [12], [13].

The physiological inflammation is supported by Th1-related cytokines-IL-1, TNF- α and IL-6 and these cytokines induce the production of other pro-

inflammatory agents-adhesive molecules, chemokines, growth factors and other mediators such as prostaglandins and nitric oxide (NO) with the stimulation of the leukocyte at the site of inflammation. IL-6 acts as the secondary mediator, responsible for maintaining the inflammatory response itself. Inflammation leads to the increasing level of Interleukin-10 (IL-10), which is the most important Th2 anti-inflammatory cytokine involved in the inflammation resolution phenomenon [3], [4], [12], [13], [14].

LGCI also has systemic effects, and it is often responsible for the disruption of skin immune homeostasis and systemic diseases such as obesity, Type II Diabetes, Vitiligo, Atopic Dermatitis, Psoriasis, BPCO (Broncho-Pulmonary-Chronic-Obstructive Pulmonary Disease) or the RRI (Recurrent Respiratory Infections), Rheumatoid Arthritis, etc.

The key risk factor for the development of allergic sensitisation is the perturbation of microbiota development during the first years of life (due to maternal consumption of antimicrobials during pregnancy, formula feeding or antimicrobial exposure or caesarean birth) [15], [16], [17].

The presence of a shift ("Th1/Th2 shift") of the immunological balance as a consequence of an imbalance between the cytokines expressed by Th1/Th17 and T_{reg}/Th2 lymphocyte subpopulations is one of the most important etiologic factors for a large number of dermatologic diseases. Th1 cytokines hyper-production is strictly linked with inflammatory and autoimmune skin diseases such as Psoriasis and Vitiligo, with the central role of a class of memory T-cells characterised by the presence of the Cutaneous Lymphocyte Antigen (CLA) on their surface and responsible for skin-homing T cell.

The presence of systemic LGCI is a strong trigger for CLA⁺ T-cells [17], [18], [19], [20].

As two of the most important etiologic factors for many of chronic dermatologic inflammatory, autoimmune and allergic diseases, gut dysbiosis (biofilm) and LGCI are currently considered therapeutic targets, but there are no standard therapeutic opportunities to treat LGCI. Also, and very important is that the chronic use of active anti-inflammatory molecules designed for the treatment of acute inflammatory phenomena shows an unfavourable efficacy/adverse effects ratio. Anti-cytokine therapy was proposed for the treatment of inflammatory and autoimmune diseases. The need of high doses of active molecules to reach the therapeutic goal and the low compliance of systemic administration performed by injective routes are the most important and limiting pitfalls connected with the use of high dosage cytokines and other signal molecules [21], [22], [23].

The Low Dose Cytokines Medicine (LDM) is an innovative therapeutic approach. It is based on the

most advanced knowledge in molecular biology and low dose pharmacology with the primary outcome.

The SKA (Sequential Kinetic Activation) technology, codified and standardised by GUNA S.p.a. -Italy- makes the low doses of signalling molecules able to be active even below the minimum dose classically considered as effective and the significative efficacy of orally administered low-dose signalling molecules is the most representative aspect of LDM [17], [22], [23], [24].

The SKA of low dose signalling molecules activates some units of cellular (or plasmatic) receptors by their low concentration.

The researchers performed a double-blind multicenter versus placebo clinical study was conducted on a dermatologic disease in 2014 for treating patients with Psoriasis Vulgaris in order to test the LDM approach with the oral administration of low dose SKA activated cytokines with the great effectiveness of the treatment assessing improvement of the psoriatic lesions and of quality of life (PASI (Psoriasis Area Severity Index) and ii) DLQI (Dermatology Life Quality Index). This study showed the effectiveness and the safety of the SKA low dose cytokines treatment and also that the SKA low dose cytokines treatment has a long action term, which extends over the months following the end of treatment [25], [26], [27], [28].

Lotti T. and colleagues, in 2015, were organised a retrospective clinical study and the effectiveness of adopted treatments for Vitiligo were compared and analysed the clinical data collected from patients who received different treatments for Vitiligo. Patients were treated with low-dose SKA IL-10, IL-4; b-FGF and anti-IL-1 antibodies, topical treatments with cortisone cream alone or in association with both groups of low dose molecules and nbUVB radiation alone or in association with both groups of molecules low dose. Other patients were treated with oral intake of extracts from Ginkgo biloba and with exposure to natural sunlight

The treatment with low dose SKA b-FGF registered an improvement in 74% of the patients. The combination of low doses SKA IL-4, IL-10 and anti-IL-1 antibodies registered an improvement in 77% of patients. The greatest improvement was registered with the association of treatments with low dose SKA molecules and UV-B treatment which has led to improvements in 92/93. No adverse effects have been reported, and the safety and efficacy of low dose SKA were verified [29], [30], [31]. The randomised, double-blinded, controlled clinical trial included children with mild to medium IgE-mediated and non-IgE mediated Atopic Dermatitis in an acute phase of the disease was conducted using low dose SKA (IL-12 and IFN- γ). The main outcome was the reduction of the AD quantified as a reduction of SCORAD index of at least 30%; in the same group, the decrease of SCORAD value continues during the follow-up period reaching

64%. The treated group showed a significant reduction of the intake of drugs for symptoms control (antihistamines and topical corticosteroids) with progressive improvement of the quality of life. No adverse effects have been reported [32], [33], [34].

Conclusions

The disruption of gut and skin homeostatic equilibrium has not only local but also systemic negative outcomes. Gut dysbiosis and inflammatory phenomena are one of the most important etiological components of many dermatologic diseases.

A rebalance action of the gut microbiota and a rebalancing of the inflammatory immune response, for the treatment of many diseases, not adequately managed with currently available therapies, are needed.

The study of biofilm formation and consolidation dynamics and the search for substances, of both natural and synthetic origin, able to intervene effectively on them, could be extremely useful.

It is now possible to design an effective treatment protocol for dysbiosis using medical devices and food supplements, also useful to prevent both dysbiosis and biofilm formation. In selected cases, faecal transplantation is needed.

The availability of Low Dose SKA signalling molecule is the most important point of LDM. Oral administration of low dose SKA signalling molecules represents the innovative core of the entire strategy for the treatment of dermatological diseases characterised by an immune imbalance and LGCI such as in Psoriasis Vulgaris, Vitiligo and Atopic Dermatitis.

The Physiologic Nutraceuticals and the Low Dose Cytokines Medicine and faecal transplantation two of the most promising approaches for the treatment of skin diseases based on the rebalance of the immune response and the recovery of gut dysbiosis.

References

1. Haroon E, Raison CL, Miller AH. Psychoneuroimmunology meets neuropsychopharmacology: translational implications of the impact of inflammation on behavior. *Neuropsychopharmacology*. 2012; 37(1):137-62. <https://doi.org/10.1038/npp.2011.205> PMID:21918508 PMCID:PMC3238082
2. Bacchus W, Aubel D, Fussenegger M. Biomedically relevant circuit-design strategies in mammalian synthetic biology. *Mol Syst*

- Biol. 2013; 9:691. <https://doi.org/10.1038/msb.2013.48> PMID:24061539 PMCID:PMC3792348
3. Tojo R, Suárez A, Clemente MG, de los Reyes-Gavilán CG, Margolles A, Gueimonde M, Ruas-Madiedo P. Intestinal microbiota in health and disease: role of bifidobacteria in gut homeostasis. *World J Gastroenterol.* 2014; 20(41):15163-76. <https://doi.org/10.3748/wjg.v20.i41.15163> PMID:25386066 PMCID:PMC4223251
4. Myles IA, Williams KW, Reckhow JD, Jammeh ML, Pincus NB, Sastalla I, Saleem D, et al. Transplantation of human skin microbiota in models of atopic dermatitis. *JCI Insight.* 2016; 1(10):e86955. <https://doi.org/10.1172/jci.insight.86955> PMID:27478874 PMCID:PMC4963067
5. Pickard JM, Zeng MY, Caruso R, Núñez G. Gut microbiota: Role in pathogen colonization, immune responses, and inflammatory disease. *Immunol Rev.* 2017; 279(1):70-89. <https://doi.org/10.1111/imr.12567> PMID:28856738 PMCID:PMC5657496
6. O'Neill CA, Monteleone G, McLaughlin JT, Paus R. The gut-skin axis in health and disease: A paradigm with therapeutic implications. *Bioessays.* 2016. <https://doi.org/10.1002/bies.201600008> PMID:27554239
7. Underhill DM, Iliev ID. The mycobiota: interactions between commensal fungi and the host immune system. *Nat Rev Immunol.* 2014; 14(6):405-16. <https://doi.org/10.1038/nri3684> PMID:24854590 PMCID:PMC4332855
8. Lynch SV. Gut Microbiota and Allergic Disease. *New Insights. Ann Am Thorac Soc.* 2016; 13(1):S51-4.
9. Chen L, Wen YM. The role of bacterial biofilm in persistent infections and control strategies. *Int J Oral Sci.* 2011; 3(2):66-73. <https://doi.org/10.4248/IJOS11022> PMID:21485310 PMCID:PMC3469879
10. Brauner A, Söderhäll M, Jacobson SH, Lundahl J, Andersson U, Andersson J. Escherichia coli-induced expression of IL-1 alpha, IL-1 beta, IL-6 and IL-8 in normal human renal tubular epithelial cells. *Clin Exp Immunol.* 2001; 124(3):423-8. <https://doi.org/10.1046/j.1365-2249.2001.01533.x> PMID:11472403 PMCID:PMC1906084
11. Huang YJ, Marsland BJ, Bunyavanich S, O'Mahony L, Leung DY, Muraro A, Fleisher TA. The microbiome in allergic disease: Current understanding and future opportunities-2017 PRACTALL document of the American Academy of Allergy, Asthma & Immunology and the European Academy of Allergy and Clinical Immunology. *J Allergy Clin Immunol.* 2017; 139(4):1099-1110. <https://doi.org/10.1016/j.jaci.2017.02.007> PMID:28257972 PMCID:PMC5899886
12. Lee SY, Lee E, Park YM, Hong SJ. Microbiome in the Gut-Skin Axis in Atopic Dermatitis. *Allergy Asthma Immunol Res.* 2018; 10(4):354-362. <https://doi.org/10.4168/aa.2018.10.4.354> PMID:29949831 PMCID:PMC6021588
13. Hueston CM, Deak T. The inflamed axis: the interaction between stress, hormones, and the expression of inflammatory-related genes within key structures comprising the hypothalamic-pituitary-adrenal axis. *Physiol Behav.* 2014; 124:77-91. <https://doi.org/10.1016/j.physbeh.2013.10.035> PMID:24184413
14. Opazo MC, Ortega-Rocha EM, Coronado-Arrázola I, Bonifaz LC, Boudin H, Neunlist M, Bueno SM, et al. Intestinal Microbiota Influences Non-intestinal Related Autoimmune Diseases. *Front Microbiol.* 2018; 9:432. <https://doi.org/10.3389/fmicb.2018.00432> PMID:29593681 PMCID:PMC5857604
15. Mancuso P. The role of adipokines in chronic inflammation. *Immunotargets Ther.* 2016; 5:47-56. <https://doi.org/10.2147/ITT.S73223> PMID:27529061 PMCID:PMC4970637
16. Kraljević Pavelić S, Simović Medica J, Gumbarević D, Filošević A, Pržulj N, Pavelić K. Critical Review on Zeolite Clinoptilolite Safety and Medical Applications in vivo. *Front Pharmacol.* 2018; 9:1350. <https://doi.org/10.3389/fphar.2018.01350> PMID:30538633 PMCID:PMC6277462
17. Van den Wijngaard R, Wankowicz-Kalinska A, Le Poole C, Tigges B, Westerhof W, Das P. Local immune response in skin of generalized vitiligo patients. Destruction of melanocytes is associated with the prominent presence of CLA+ T cells at the perilesional site. *Lab Invest.* 2000; 80(8):1299-309. <https://doi.org/10.1038/labinvest.3780138> PMID:10950121
18. Pallešchi G, Carbone A, Zanello PP, Mele R, Leto A, Fuschi A, Al Salhi Y, et al. Prospective study to compare antibiotics versus the association of N-acetylcysteine, D-mannose and Morinda citrifolia fruit extract in preventing urinary tract infections in patients submitted to urodynamic investigation. *Arch Ital Urol Androl.* 2017; 89(1):45-50. <https://doi.org/10.4081/aiua.2017.1.45> PMID:28403598
19. Prasai TP, Walsh KB, Bhattarai SP, Midmore DJ, Van TT, Moore RJ, Stanley D. Biochar, Bentonite and Zeolite Supplemented Feeding of Layer Chickens Alters Intestinal Microbiota and Reduces Campylobacter Load. *PLoS One.* 2016; 11(4):e0154061. <https://doi.org/10.1371/journal.pone.0154061> PMID:27116607 PMCID:PMC4845986
20. Yeo CK, Vikhe YS, Li P, Guo Z, Greenberg P, Duan H, Tan NS, et al. Hydrogel Effects Rapid Biofilm Debridement with ex situ Contact-Kill to Eliminate Multidrug Resistant Bacteria in vivo. *ACS Appl Mater Interfaces.* 2018; 10(24):20356-20367. <https://doi.org/10.1021/acsami.8b06262> PMID:29806938
21. Marcum ZA, Hanlon JT. Recognizing the Risks of Chronic Nonsteroidal Anti-Inflammatory Drug Use in Older Adults. *Ann Longterm Care.* 2010; 18(9):24-27.
22. Renukuntla J, Vadlapudi AD, Patel A, Boddu SH, Mitra AK. Approaches for enhancing oral bioavailability of peptides and proteins. *Int J Pharm.* 2013; 447(1-2):75-93. <https://doi.org/10.1016/j.ijpharm.2013.02.030> PMID:23428883 PMCID:PMC3680128
23. Biancotto A, Wank A, Perl S, Cook W, Olnes MJ, Dagur PH, et al. Baseline levels and temporal stability of 27 multiplexed serum cytokine concentrations in healthy subjects. *PLoS One.* 2013; 8(12):e76091. <https://doi.org/10.1371/journal.pone.0076091> PMID:24348989 PMCID:PMC3861126
24. Roberti ML, Ricottini L, Capponi A, Sciauzero E, Vicenti P, et al. Immunomodulating treatment with low dose Interleukin-4, Interleukin-10 and Interleukin-11 in psoriasis vulgaris. *J Biol Regul Homeost Agents.* 2014; 28(1):133-9.
25. Luchetti P. Increasing of visual function in patients with retinal atrophy treated with drugs of Low Dose Medicine. Monocentric retrospective observational study. *Minerva Medica Oftalmologica.* 2014; 56(3-4):53-61.
26. Cardani D, Dusio GF, Luchini P, Sciarabba M, Solimene U, Rumio C. Oral Administration of Interleukin-10 and Anti-IL-1 Antibody Ameliorates Experimental Intestinal Inflammation. *Gastroenterology Research.* 2013; 6:124-133. <https://doi.org/10.4021/gr556w> PMID:27785242 PMCID:PMC5074810
27. Barygina V, Becatti M, Lotti T, Taddei N, Fiorillo C. Low dose cytokines reduce oxidative stress in primary lesional fibroblasts obtained from psoriatic patients. *J Dermatol Sci.* 2016; 83(3):242-4. <https://doi.org/10.1016/j.jdermsci.2016.06.002> PMID:27317477
28. Lotti T, Hercogova J, Wollina U, et al. Vitiligo: successful combination treatment based on oral low dose cytokines and different topical treatments. *J Biol Regul Homeost Agents.* 2015; 29(1):53-8.
29. Radice E, Bellone G, Miranda V. Enhancement of the immunomodulatory functions of ex vivo-generated dendritic cells from early-stage colon cancer patients by consecutive exposure to low doses of sequential kinetic activated IL-4 and IL-12. A preliminary study. *Translational Oncology.* 2015; 8:327-38. <https://doi.org/10.1016/j.tranon.2015.06.005> PMID:26310379 PMCID:PMC4562983
30. Lotti T, et al. Successful combination treatment for Psoriasis with phototherapy and Low Dose Cytokines. A spontaneous observational retrospective clinical study. *Der Hautarzt* 2015; 66(11):849-54. <https://doi.org/10.1007/s00105-015-3698-6> PMID:26449546

31. Genazzani AD, Despini G, Chierchia E, Benedetti C, Prati A. Pharmacological and integrative treatment of stress-induced hypothalamic amenorrhea. In *Frontiers in Gynecological Endocrinology*. Springer, Cham, 2016:69-84. https://doi.org/10.1007/978-3-319-23865-4_9
32. Kim JE, Kim HS. Microbiome of the Skin and Gut in Atopic Dermatitis (AD): Understanding the Pathophysiology and Finding Novel Management Strategies. *J Clin Med*. 2019; 8(4):E444. <https://doi.org/10.3390/jcm8040444> PMID:30987008 PMCid:PMC6518061
33. Barygina V, Becatti M, Lotti T, Moretti S, Taddei N, et al. Treatment with low-dose cytokines reduces oxidative-mediated injury in perilesional keratinocytes from vitiligo skin. *J Dermatol Sci*. 2015; 79(2): 163-70. <https://doi.org/10.1016/j.jdermsci.2015.05.003> PMID:26051876
34. Fiorito F, Cantiello A, Granato GE, Navas L, Diffidenti C, De Martino L, Maharajan V, et al. Clinical improvement in feline herpesvirus 1 infected cats by oral low dose of interleukin-12 plus interferon-gamma. *Comp Immunol Microbiol Infect Dis*. 2016; 48:41-7. <https://doi.org/10.1016/j.cimid.2016.07.006> PMID:27638118