



J. bio-sci. 19: 43-52, 2011
<http://www.banglajol.info/index.php/JBS/index>

ISSN 1023-8654

YIELD RESPONSE OF DIALLELIC CROSSED MAIZE (*ZEA MAYS* L.) GENOTYPES TO VARYING NITROGEN REGIMES IN NIGERIA

OB Bello^{*}, MS Afolabi[†], SA Ige², SY Abdulmalik³, MA Azeez⁴, J Mahmud⁵
Department of Biological Sciences, Fountain University, Osogbo, Osun State, Nigeria

¹*Department of Crop Science, Landmark University, Omuaran, Kwara State, Nigeria*

²*Department of Agronomy, University of Ilorin, Ilorin, Nigeria*

³*Department of Agronomy, Ibrahim Badamasi Babangida University, Lapai, Niger State, Nigeria*

⁴*Department of Plant Biology, Ladoke Akintola University of Technology, Ogbomosho, Nigeria*

⁵*Lower Niger River Basin, Ilorin, Kwara State, Nigeria*

Abstract

Context: Development of early maturing maize cultivars that remain productive under low N fertilizer farming system, consistent with the farmers' technologies is a prerequisite to improving adoption of new varieties without increasing production cost.

Objective: To assess the performance of ten early open pollination maize varieties (OPVs) and their F₁ hybrids for grain yield and nitrogen use efficiency (NUE), and also identify productive cultivars under low N fertilizer regimes.

Materials and Methods: The trials were set up in a split plot arrangement with three N fertilizer levels (0, 45 and 90 kg N ha⁻¹) as main plot and the genotypes as sub-plot. Each plot within N level was four-row, laid out in a randomized complete block design of four replications. Ten OPVs were crossed in a half diallel to generate 45 F₁ hybrids during 2004 and 2005 growing seasons. Planting were carried out on 20th July, 2005 and 2nd July, 2006. Agronomic characters studied were grain yield, maize establishment count, days to 50% tasselling and silking as well as plant and ear heights.

Results: The year 2005 growing season was better for all observed characters amongst all the genotypes than the year 2006. Although, expressions of these traits in the hybrids were relatively higher than the OPVs including the grain yield. The total increase in grain yield observed was 1.72 t ha⁻¹ and 1.95 t ha⁻¹ for OPVs and hybrids respectively on application of 90 kg ha⁻¹ over no N-application. However, NUE was optimum at 45 kg N ha⁻¹ in both groups. Grain yield and NUE correlated positively with growth characters measured except for days to 50% silking. Higher genetic gains were recorded for plant and ear heights.

Conclusion: Two drought tolerant varieties (Acr 90 Pool 16-Dt and Tze Comp3 Dt) that combined well with specific cultivars for grain yield and NUE probably have gene pools for low N-tolerance.

Keywords: Nitrogen use efficiency, diallel crosses, maize cultivars, open pollinated varieties

Introduction

Maize (*Zea mays* L.) is an important staple food crops and provides both bulks of raw materials for the livestock and many agro-allied industries in the world (Bello *et al.* 2010, Undie *et al.* 2012). Maize has a wide range of uses than any other cereals because of its world-wide distribution, high yielding nature, ease of processing, readily digestible and relatively lower price of the grain. In recent times, importance of maize production has been widely publicized among farmers of Nigeria. The soils however, are fragile, primarily of kaolinitic Alfisols with low organic matter, effective cation exchange capacity and plant nutrients (Carsky and Iwuafor 1999) creates low yield problems in this agro-climate conditions. The attendant problems of the soils frequently deficient in nitrogen (N) due to high rate of leaching, denitrification, decomposition of organic matter and erosive action of rainfall had not only made nitrogen the most limiting nutrient to maize production in Nigeria (Carsky and Iwuafor 1999, Yusuf *et al.* 2003) but also enhance low soil moisture retention. Despite the soil being low in plant available nutrients, N deficiency is further exacerbated from its continual depletion

^{*}Corresponding author Email: obbellow2002@yahoo.com

from the soil pool by removal of N-containing crop residues from the farm. Nitrogen depletion in maize-based systems in some farmer's fields in West African savannah is estimated to be 36-80 kg N ha⁻¹ per year (Sanginga *et al.* 2001). In view of the fact that farmers hardly adopt option that could provide supplemental N to maize crops (perhaps manuring, mulching, crop rotation and fallowing schemes), the knock-out effect is that maize production by small holder farmers is normally at a lower capacity (with decrease in the total maize yield) and at a subsistence level solely to meet family consumption. Despite fertilizers recommendations (60 to 120 kg N ha⁻¹) for resource poor farmers in the West Africa moist savannah (Carsky and Iwuafor 1999), less than 20 kg N ha⁻¹ used in Nigeria was considered grossly inadequate and 50 kg nutrient ha⁻¹ fertilizer use was recommended across sub-Saharan Africa by African Heads of States at the Fertilizer Summit held in 2006 at Abuja, Nigeria (Yusuf *et al.* 2009).

The southern Guinea savannah (SGS) of Nigeria has great potential for the expansion of maize production beyond the present level due to its bimodal rainfall pattern, high solar radiation and favourable temperature during the growing season. However, the zone is mitigated with problems such as erratic nature of rainfall pattern, low soil nutrient status and water holding capacity due to fragility of the soil that consistently prone to drought (Fakorede *et al.* 2001, Bello 2008). Hybrid maize is well known in the region for its high demand for plant nutrients and other production inputs. The farmers have been indoctrinated with the belief that required amount of inorganic fertilizers need to be applied for maximum performance of hybrid maize and good economic returns. This extra production cost discourages most farmers engaging in hybrid maize production in the country (Kogbe and Adediran 2003). However, as a result of increasing cost of N fertilizer due to reduction in government subsidy, poor infrastructure for production and distribution, resulted to inadequate or untimely supply of fertilizers to farmers. Lack of cash or access to credit and perception of cost-benefit ratio also made N fertilizers sometimes unavailable or are too expensive and often beyond the reach of resource poor farmers who constitute the bulk of maize growers in the moist savannah of West Africa (Juo and Mano 1996). Maize cultivars that are productive under conditions of low N availability are therefore highly desirable.

In the development of improved maize productivity in a sustainable fashion in areas with low N fertility, two basic approaches can be taken. The first, approach is to develop innovative agronomic practices that efficiently utilized N from organic matter and N inputs from biological fixation and atmospheric deposition. The second approach is to work with population with the reservoir of genes for low N tolerance (Smith *et al.* 1994). One strategy for improving the productivity of maize under sub-optimal N fertility is to use the second approach. The low N tolerance cultivars are superior in the utilization of available N, either due to enhanced N-uptake capacity or because of more efficient use of absorbed N for grain production (Lafitte and Edmeades 1994). On the other hand, combination with the first approach will lead to high maize yields. Evaluation of open pollination maize varieties (OPVs) and their F₁ hybrids in a diallel cross however could provide a worthwhile knowledge on those genotypes that could be used either directly for cultivation or for extraction of inbred lines for low N tolerant hybrid development. A corollary is that these genotypes can readily fit into environment with erratic rainfall and low water holding capacity, since they are bred for marginal environments of West and Central Africa. Thus, development of early maturing maize cultivars that remain productive under low N fertilizer farming system, consistent with the farmers' technologies is therefore a prerequisite to improving adoption of new varieties without increasing production cost in this agro-ecology. The objectives of this research therefore was to assess the performance of ten early OPVs and their F₁ hybrids under low N fertilizer regimes with the view to identify those that could be cultivated either as varieties *per se* under the farmer's practices or those that could be introgressed with other released cultivars to develop low N-tolerant varieties in the SGS ecology of Nigeria.

Materials and Methods

Field experiments were conducted at the Teaching and Research farm of the University of Ilorin (Latitude 80° 29'N, Longitude 40° 35'E and annual rainfall of 945 mm) in the SGS of Nigeria. The soil is classified as Typic paleustalf (United State Department of Agriculture, USDA soil taxonomy). Composite soil samples were collected randomly from the trial site at the depth of 0-15 cm with an auger prior to planting in 2005 and 2006 respectively. The samples were transferred to the laboratory for physico-chemical analysis. The collected samples were air-dried and passed through 2 mm sieve to remove large particles, debris and stones. The sieved samples were analyzed for pH in 1:1 soil to water ratio using the Coleman's pH meter. Organic carbon was determined by Walkley and Black procedure (Nelson and Somers 1992). Total Nitrogen was determined by the micro Kjeldahl method (Bremner 1965), while available phosphorus was extracted by Bray's P₁ method (Bray and Kurtz 1945) and read by the Atomic Absorption Spectrometer. Potassium was evaluated calorimetrically using molybdenum blue. Calcium and magnesium were estimated by Versenate ethylenediaminetetraacetic acid (EDTA) titration method (Juo 1979), while exchangeable potassium and sodium were analyzed using flame photometry (Juo 1979). The exchangeable cations were measured on extracts obtained after leaching soil samples with neutral normal ammonium acetate solution (IITA, 1979). Rainfall distribution data for the year 2005 and 2006 were also collected at the Lower Niger River Basin Development Authority, Ilorin, Nigeria.

The genetic materials used were ten OPVs of maize, selected for grain yield and adaptation to abiotic (drought) and biotic (stalk rot, striga and downy mildew) stress factors. They were early to medium maturing white cultivars with maturity period of 90 to 100 days. The cultivars were obtained from the International Institute of Tropical Agriculture (IITA), Ibadan. The origin, characteristic features and ecological adaptation of ten open pollinated maize varieties are shown in Table 1. The ten varieties were crossed in a half diallel to generate 45 F₁ hybrids during 2004 and 2005 cropping seasons. Planting were carried out during the growing seasons on 20th July, 2005 in the first year and 2nd July, 2006 in the second year. The resultant hybrids were harvested, processed and stored in the cold room prior to field evaluation.

It is worthwhile to mention that the ultimate goal is not diallel analysis in which using pure lines as parents is necessary. The trials were set up in a split plot arrangement with N-level as main plot and the genotypes as sub-plot. Each plot within N level was four-row, laid out in a randomized complete block design (RCBD) of four replications at spacing of 0.75 m between and 0.50 m within the rows. Each hill was planted to three seeds but was later thinned to two plants per hill to give a plant population of approximately 53,333 per hectare. Plantings were done on 28th July in 2005 and 22nd July in 2006. Nitrogen fertilizer in form of granular NPK 20-10-10 was applied at 0, 45, 90 kg N ha⁻¹. Phosphorus and potassium were obtained from single super phosphate (SSP) and murate of potash (MOP) respectively. The required quantities for each level of N were computed from the three combinations. All the SSP and MOP fertilizers were applied once at planting which included 60% of the NPK. The remaining 40% was applied at 6 weeks after planting (WAP). The fertilizers were applied by spot placement.

Data were collected from the two middle rows in each plot. Agronomic parameters recorded in each cropping seasons were seedling establishment count; plants and ear heights were measured as the distance from the ground level to the flag leaf and the node bearing the uppermost ear respectively using measuring tape. Days to 50% tasselling and silking were taken as the date when 50% of the plants in a plot had tasselled and extrude silk respectively. Anthesis-silking interval was estimated as the difference between days to pollen shed and silking. Moisture at harvest was determined by collection of 500 grain samples from each plot. The samples were first weighed to obtain initial weight, followed by drying to a constant weight in the oven at 80°C, and the difference between the two weights were recorded as the moisture at harvest. Grain yield was obtained from ear weight per plot (assuming 80% shelling percentage) and converted to tonnes per hectare after adjusting to 12% moisture content. Nitrogen use efficiency (NUE) was estimated as grain yield per unit of available soil N as
$$NUE = \frac{\text{Grain yield (g/plant) at N rate applied} - \text{grain yield at 0 kg ha}^{-1}}{N \text{ applied (gNf)}}$$
, where gNf = amount of N applied.

The data were subjected to analysis of variance and pertinent means were separated using Least Significant Difference (LSD) according to Gomez and Gomez (1984). The phenotypic, genotypic and environmental correlation coefficients were estimated from the mean squares and mean across products as suggested by Mode and Robinson (1959). Genstat Statistical Package (Genstat 1995) was used to compute simple linear correlation. Analysis of variance (ANOVA) on individual site basis was first computed before a combined ANOVA across sites.

Table 1. Origin, genetic background, breeding emphasis and ecological adaptation of the maize parent

Genotypes	Origin and genetic background	Breeding emphasis	Ecological adaptation
Acr 90 Pool 16-Dt	Early white dent CIMMYT cultivar, derived from crosses among large numbers of early late white flint materials from Mexico, the Caribbean area, Central and South America. Selected for drought tolerance.	Stalk rot, <i>Striga</i> and drought tolerance.	Forest and savannah.
Tze Comp 4-Dmr Srbc2	Early maturing white and semi dent cultivar, derived from diverse sources of early mid-altitude germplasm, intermated with TZESR-W and DMR-ESRW.	Yield and <i>Striga</i> tolerance .	Forest and savannah.
Tze Comp4 C2	Early maturing white and semi dent cultivar, derived from diverse sources of early mid-altitude germplasm, intermated with EV 8430-SR and IK 8149 SR. It has synchronous male and female flowering, lower plant height and small tassel size.	Yield and <i>Striga</i> tolerance.	Forest and savannah.
Acr 97 Tze Comp3 C4	Early white flint dent cultivar, derived from early mid-altitude germplasm with EV 8430-SR, DMR-ESRW, TZESR-W and IK 8149 SR intermated.	Yield, downy mildew and <i>Striga</i> tolerance.	Forest and savannah.
Hei 97 Tze Comp3 C4	Early white flint dent cultivar, derived from early mid-altitude germplasm with EV 8430-SR, DMR-ESRW, TZESR-W and IK 8149 SR intermated.	Yield, downy mildew and <i>Striga</i> tolerance.	Forest and savannah.
Acr 94 Tze Comp5	Early white flint dent cultivar, derived from early mid-altitude germplasm with <i>Striga</i> intermated with TZE-WC3.	<i>Striga</i> tolerance.	Savannah.
Tze Comp3 Dt	Early white flint dent cultivar, derived from diverse sources of early mid-altitude germplasm with drought tolerant cultivars, intermated with TZESR-W and DMR-ESRW.	Drought tolerance.	Forest and savannah
Tze Comp3 C2	Early white and flint dent cultivar, derived from diverse sources of early mid-altitude germplasm, produced by intermating TZESR-W and DMR-ESRW.	Downy mildew and <i>Striga</i> tolerance.	Forest and savannah.
Ak 95 Dmr-Esw	Early maturing and flint dent cultivar, developed from intermating diverse sources of early mid-altitude germplasm, produced by intermating DMR sources from Philippines with TZB, TZBP, TZSR and tropical late. Selected for earliness.	Downy mildew and <i>Striga</i> tolerance.	Forest.
Tze Msr-W	Early white semi dent cultivar, derived from early mid-altitude germplasm, developed from intermating local and early cultivars with TZSR.	Yield and <i>Striga</i> tolerance.	Forest and savannah.

Source: IITA Archival Report 1988-1992

Results

Soil Analysis: Soil samples were collected from the trial site before cropping in 2005 and 2006 were analyzed in the laboratory for selected physical and chemical properties (Table 2). The soil texture was loamy sand with similar values of soil properties in both years. The soil had a pH of 5.4 which is moderately acidic. The soil organic matter and total nitrogen was low, while the soil available Phosphate and the exchangeable cations (K, Na, Ca and Mg) were moderate for maize production in this area. However, rainfall distribution data for the year 2009 and 2010 were also collected (Fig. 1).

Table 2. Selected physical and chemical characteristics of the soil before growing maize in 2005 and 2006.

Physical characteristics	2005	2006
Texture	Loamy sand	Loamy sand
pH 1:1 (H ₂ O)	5.4	5.30
Sand %	84.1	84.0
Clay %	8.02	8.01
Silt %	6.42	6.40
Chemical characteristics		
Exchangeable Ca ²⁺ (C. mol kg ⁻¹)	1.12	1.10
Exchangeable Mg ²⁺ (C. mol kg ⁻¹)	1.62	1.61
Exchangeable Na ⁺ (C. mol kg ⁻¹)	0.19	0.18
Exchangeable K ⁺ (C. mol kg ⁻¹)	0.01	0.01
Total acidity H ⁺ (C. mol kg ⁻¹)	0.05	0.04
Cation exch. Cap. (C. mol kg ⁻¹)	2.83	2.82
% Organic Carbon	0.24	0.26
% Soil organic matter	1.03	1.04
% Total Nitrogen	0.24	0.26
Available Phosphate (mg kg ⁻¹)	20.31	20.30




Fig. 1. Rainfall distribution pattern for Ilorin in 2005 and 2006

Table 3. Maize grain yield, NUE and agronomic characters in open pollinated maize varieties and their F₁ hybrids as affected by N fertilizer rates in 2005 and 2006 at Ilorin, Nigeria.

Treatment	Grain Yield (t ha ⁻¹)	NUE	Maize establishment count (EPC)	Days to 50% tasselling (TS)	Days to 50% silking (SK)	Plant height (cm)	Ear height (cm)
Open pollinated varieties							
Year							
2005	2.42	1.82	39	52	53	129	44
2006	2.10	1.06	50	50	51	113	40
LSD α 0.05	0.08**	0.66**	0.76**	0.75**	0.85*	3.42*	2.81)
N Levels (kg N ha ⁻¹)							
0	0.89	1.00	36	54	56	101	35
45	2.18	2.07	38	52	53	121	37
90	2.61	2.45	37	51	52	129	40
LSD α 0.05	0.67*	0.45*	ns	0.45*	0.91*	3.54*	2.75*
F ₁ Hybrids							
Year							
2005	3.71	2.57	37	53	54	129	44
2006	3.70	2.51	33	50	51	118	40
LSD α 0.05	ns	ns	0.43*	0.36*	0.97*	3.28*	2.81*
N Levels (kg N ha ⁻¹)							
0	1.35	1.0	36	54	56	110	39
45	2.68	2.14	37	51	54	122	40
90	3.30	2.71	36	50	52	130	41
LSD α 0.05	0.98*	0.51*	ns	0.38*	0.96*	3.11*	2.61*

*, **, Significant F test at 0.05 and 0.01 levels of probability respectively.

Table 4. Interactive effects of year x nitrogen for grain yield and NUE in open pollinated varieties and their selected F₁ hybrids in 2005 and 2006 at Ilorin, Nigeria.

Treatment (N Levels (kg N ha ⁻¹))	Open pollinated varieties		F ₁ hybrids grain yield (t ha ⁻¹)	
	2005	2006	2005	2006
0	1.09	0.88	1.35	1.34
45	2.23	2.13	2.71	2.65
90	2.68	2.54	3.56	2.45
LSD α 0.05	0.18**		0.12**	
NUE				
0	1.00	1.00	1.00	1.00
45	2.12	2.01	2.15	2.13
90	2.64	2.35	2.74	2.68
LSD α 0.05	0.65*		0.39*	

*, **, Significant F test at 0.05 and 0.01 levels of probability respectively

Table 5. Genotypic means for maize grain yield, NUE and agronomic parameters in 10 open pollinated varieties and their selected F₁ hybrids in 2005 and 2006 at Ilorin, Nigeria.

Genotype	Grain Yield (t ha ⁻¹)	NUE	Maize establishment count (EPC)	Days to 50% tasselling (TS)	Days to 50% silking (SK)	Plant height (cm)	Ear height (cm)
Open pollinated varieties							
Acr 90 Pool 16-Dt	2.33	2.34	35	51	53	124	44
Tze Comp 4-Dmr Srbc2	2.29	1.16	38	50	54	122	35
Tze Comp4 C2	2.24	1.53	34	51	53	114	41
Acr 97 Tze Comp3 C4	2.21	1.24	37	51	54	114	42
Hei 97 Tze Comp3 C4	2.18	1.34	39	50	53	114	39
Acr 94 Tze Comp5	2.24	1.33	37	51	54	118	38
Tze Comp3 Dt	2.38	2.28	40	49	53	135	53
Tze Comp3 C2	2.22	1.34	37	51	53	117	42
Ak 95 Dmr-Esrw	2.27	1.56	36	51	54	122	44
Tze Msr-W	2.24	1.13	39	50	54	128	41
LSD α 005	0.14*	0.34*	0.45*	0.87*	0.56*	6.04**	4.42**
F₁ Hybrids							
Acr 94 Tze Comp5 x Tze Comp4 C2	3.46	1.90	35	50	52	137	45
Acr 90 Pool 16-Dt x Tze Comp3 C2	3.48	1.46	38	50	53	129	51
Acr 97 Tze Comp3 C4 x Tze Comp3 Dt	3.34	1.28	34	51	54	128	52
Acr 90 Pool 16-Dt x Ak 95 Dmr-Esrw	3.48	1.98	37	49	53	137	49
Tze Comp3 Dt x Ak 95 Dmr-Esrw	3.43	1.51	39	50	53	129	44
Tze Comp 4-Dmr Srbc2 x Ak 95 Dmr-Esrw	3.38	1.36	37	51	54	135	40
Ak 95 Dmr-Esrw x Tze Msr-W	3.44	1.81	40	49	53	128	43
Tze Comp3 Dt x Tze Comp3 C2	3.36	1.56	37	50	54	134	50
Acr 94 Tze Comp5 x Tze Comp3 Dt	3.34	1.48	36	50	53	126	53
Acr 90 Pool 16-Dt x Tze Comp3 Dt	3.48	1.90	39	49	52	128	52
Acr 94 Tze Comp5 x Tze Msr-W	3.38	1.46	35	50	52	127	49
Tze Comp 4-Dmr Srbc2 x Tze Msr-W	3.43	1.58	38	51	53	129	44
Tze Comp4 C2 x Tze Comp3 Dt	3.48	1.98	34	50	52	135	40
Tze Comp4 C2 x Tze Comp3 C2	3.43	1.98	36	51	54	128	43
LSD α 005	0.17*	0.56*	0.67*	0.89*	0.66*	5.98**	4.30**

*, **, Significant F test at 0.05 and 0.01 levels of probability respectively

Effect of cropping season on open pollinated varieties and their F₁ hybrids performance: The rainfall in 2006 growing season was more favourable for the expression of all studied characters among the genotypes than 2005 (Fig. 1). Each of the cropping seasons had significant effect on all the characters measured in the open pollinated varieties and hybrids, except for nitrogen use efficiency (NUE) and grain yield in the hybrids (Table 3). The plants were significantly taller, with higher ear placement, subsequently higher grain yield, more efficient utilization of N and higher anthesis-silking interval. However, expressions of these traits in the hybrids were relatively higher than the OPVs.

Effect of N fertilizer rates on grain yield, NUE and other agronomic characters: Varying the N-fertilizer rate significantly affected the expression of the characters in both OPVs and hybrids (Table 3). Grain yield increased with increasing in N-dosage in the OPVs and hybrids but was higher in the hybrids relative to OPVs. The NUE of hybrids and OPVs increased significantly by increasing the N-level from 0 to 45 kg N ha⁻¹. Though the NUE showed consistent trend in both groups from 45 to 90 kg N ha⁻¹, the increment was significant only for hybrids. The grain yield increased significantly from 0.89 to 2.18 t ha⁻¹ and 1.35 to 2.68 t ha⁻¹ for OPVs and hybrids respectively by increasing N-application from 0 to 45 kg N ha⁻¹. The total increase in grain yield observed was 1.72 t ha⁻¹ and 1.95 t ha⁻¹ for OPVs and hybrids respectively on application of 90 kg ha⁻¹ over no N-application. However, the increase in grain yield was not statistically significant in either case. Days to 50% tasselling and silking decreased with increasing N application rate, while plant and ear heights increased significantly with increasing N application rate in both groups.

Interactive effects of year x nitrogen on grain yield and NUE: The first order interaction of year x nitrogen was significant for both grain yield and NUE in the OPVs and hybrids (Table 4). Significantly, higher grain yield was obtained at 90 kg N ha⁻¹ in 2005 than 2006 in both genotypes. Grain yield increased significantly at every N level between 0 and 90 kg N ha⁻¹ with a difference of 1.59 and 1.66 t ha⁻¹ in OPVs and 2.22 and 2.11 t ha⁻¹ in the hybrids at the first and second years respectively. However, NUE was best at 90 kg N ha⁻¹ regardless of the growing season among the genotypes.

Genotypic performance for grain yield, NUE and other agronomic characters: The hybrids differed significantly for all the traits and showed superiority with respect to grain yield than their respective OPVs (Table 5). Although, there was no significant genotype x N fertilizer effects for grain yield and NUE in this study, but most of the hybrids appeared to exhibit capacity for utilizing N better than their respective parents. Two drought tolerant varieties (Acr 90 Pool 16-Dt and Tze Comp3 Dt) not only had the highest grain yield and NUE among OPVs, but also specifically combined well with each other and some cultivars. While Acr 90 Pool 16-Dt favourable combined with Tze Comp3 Dt, Tze Comp3 C2 and Ak 95 Dmr-Esrw for these characters, var. Tze Comp3 Dt combined well with Tze Comp4 C2. The genotypes were also taller with higher ear placement, increase in days to 50% tasselling and silking. However, correlation of grain with other traits among the parents and hybrids showed that grain yield correlated positively and significantly with growth characteristics measured except days to 50% silking (Table 6). NUE correlated positively and significantly with all agronomic characters except days to 50% silking in the OPVs.

Table 6. Simple linear correlation coefficient of grain yield, NUE and agronomic parameters in 10 open pollinated maize varieties and 45 F₁ hybrids in 2005 and 2006 at Ilorin, Nigeria.

Parameters	Open pollinated varieties		F ₁ Hybrids	
	Grain Yield (t ha ⁻¹)	NUE	Grain Yield (t ha ⁻¹)	NUE
Maize establishment count	0.320**	0.315**	0.411**	0.325**
Days to 50% tasselling	0.235**	0.261**	0.387**	0.341**
Days to 50% silking	0.042	-0.016	0.009	-0.076
Plant height	0.147**	0.140**	0.345**	0.134
Ear height	0.235**	0.125*	0.140*	0.110*
NUE	0.521**	-	0.242**	

*, **, Significant F test at 0.05 and 0.01 levels of probability respectively

Table 7. Estimates of genotypic and phenotypic variances, coefficients of genetic variability and genetic advance in agronomic characters among open pollinated maize varieties and their F₁ hybrids as affected by N fertilizer rates in 2005 and 2006 at Ilorin, Nigeria.

Character	δ^2g	δ^2p	GCV	PCV	GA (%)
Establishment count	3.6	17.4	52.4	118.8	12.2
Days to 50% tasselling	32.7	156.3	74.1	159.8	27.0
Days to 50% silking	10.5	317.2	16.8	254.6	11.9
Plant height	608.6	1062.5	129.6	517.5	1162.8
Ear height	121.4	1054.6	128.7	213.6	175.8
Grain yield	3.2	3.5	71.6	75.3	95.0

δ^2g = Genotypic variance, δ^2p = Phenotypic variance, GCV= Genotypic coefficient of variation, PCV= Phenotypic coefficient of variation, GA= Genetic advance

Genetic gains and variances for grain yield and other agronomic characters: Estimates of genotypic and phenotypic variances as well as coefficients of genotypic and phenotypic variations, and genetic advance are presented in Table 7. The value for genotypic variances ranged from 3.2 for grain yield to 608.6 for plant height, while the phenotypic variances ranged from 17.4 for establishment count to 1062.5 for plant height. The coefficients of variations also showed that the phenotypic coefficients of variations (PCV) were higher than the genotypic coefficients of variations (GCV) for all the parameters measured. Plant height had the highest GCV (129.6) and PCV (252.9). Similarly, highest genetic gain of 1162.8% was recorded for plant height and the least genetic gain of 11.9 % for days to 50% silking.

Discussion

The release of improved stress-tolerant maize varieties and hybrids has sparked optimism for increased maize productivity in Nigeria. These cultivars have the potential to provide farmers with opportunities to overcome the challenges to maize production. However, efficiency of plants in making use of soil nutrients within their environment is expected to be optimal with adequate soil moisture regime. Although rainfall distribution throughout the duration of this study appears favourable for crop growth and expression of studied traits in both OPVs and the F_1 hybrids, the second year precipitation was more beneficial (Fig. 1). However, the results from these experiments showed that hybrid maize efficiently utilized nitrogen better than OPVs as per earlier reports in Nigeria (Akintoye *et al.* 1999, Kogbe and Adediran 2003). In both years, hybrids gave higher yield than the OPVs even at low soil N status. Consequently, maize yield increased with increasing nitrogen availability and this is in agreement with previous report of Kogbe and Adediran (2003). All the genotypes responded adequately to N application even as low as 45 kg N ha⁻¹. Cultivars identified as less responsive to applied N sometimes perform better at low N than do N-responsive hybrids or OPVs (Pollmer *et al.* 1979). The low N tolerant cultivars are superior in the utilization of available N, either due to enhanced uptake capacity or because of more efficient use of absorbed N in grain production (Lafitte and Edmeades 1994, Jafartayari *et al.* 2012). However, number of days recorded in 2006 especially at anthesis was less than that recorded in 2005, which facilitated optimum utilization of available soil nutrients. This probably explains the higher grain yield and superiority in N utilization observed in the OPVs in 2005 relative to 2006 (Table 3) and also the significant year x nitrogen interaction for this character in both parents and hybrids (Table 4). Values for NUE, plant and ear heights were higher in 2005 compared to 2006 in both OPVs and hybrids. This observation tends to underscore the importance of moisture availability especially at anthesis in the accumulation of photo assimilates as well as its translocation for grain production in maize.

Grain yield and NUE increased in a consistent manner with increasing N rate in both parents and hybrids (Table 3). This is in line with previous reports (McCullough *et al.* 1994, Uhart and Andrade 1995) that variation in N supply affects both growth and development of maize plants. The capacity to utilize N as measured by NUE and consequently grain yield in this study was optimum at 45 kg N ha⁻¹ in both varieties and hybrids which were similar to the result obtained by Elemo (1997) in the northern Guinea savannah of Nigeria in which the author reported a yield increase of 200% over the control at 40 kg N ha⁻¹ in two maize varieties of different maturity period. This result is also in support with earlier findings (Lafitte and Banziger 1994, Namakka *et al.* 2008, Adekayode and Ogunkoya 2010, Gagnon and Ziadi 2010, Jafartayari *et al.* 2012) on increase in grain yield and NUE at low N levels. Kogbe and Adediran (2003) similarly observed that maize varieties responded adequately well to nitrogen application in the southern Guinea savanna of Nigeria and gave over 3 t ha⁻¹ grain yield by applying as low as 50 kg N ha⁻¹. They suggested that this could erase the fear that without high N input, hybrid maize would not give some appreciable returns. However, there was no significant genotype x N fertilizer effects for grain yield and NUE in this study with similar to the finding of Ma *et al.* (1999), but contradicts- the findings of Bertin and Gallais (2000) who observed that maize differed in their capacity for N uptake, assimilation and redistribution of N from vegetative to reproductive tissues. The fact that the OPVs used in this study are of the same maturing group may be responsible for the difference in the two results.

High phenotypic coefficients of variations values observed in both OPVs and hybrids may indicate significant year x nitrogen interactions for all the characters measured. This therefore affirms that the populations used in this study showed greater range of genetic variability under different N fertilizer rates. Genetic differences have been observed among maize genotypes (inbred lines, hybrids and OPVs) in response to N fertilizer application and efficient utilization of absorbed N for grain yield (Banziger *et al.* 1999, Kling *et al.* 1997, Abdel Rahman *et al.* 2002). The high genetic gain of plant and ear heights may indicate that these traits could be selected for genetic improvement of maize under different N regimes, as earlier suggested (Castleberry *et al.*

1984, Yusuf 2010). Plant height and days to 50% silking that had highest and lowest genetic gains respectively signify broader genetic base for plant height and narrower genetic base for days to 50% silking among the genotypes. Though low genetic gain was recorded for days to 50% silking and days for establishment count, there is a significant variability in these traits and hence may also be useful in maize breeding programme. Grain yield correlated positively with most of the parameters measured and this has also been reported by Tenaw (2000).

Conclusion

Two drought tolerant varieties (Acr 90 Pool 16-Dt and Tze Comp3 Dt) that combined well with specific cultivars for grain yield and NUE probably have gene pools for low N-tolerance. Therefore, these varieties can further be tried under farmers' growing conditions or could be hybridized for development of inbred lines in the development of N stress-tolerance varieties for cultivation in the SGS ecology.

References

- Abdel-Rahman AM, Magboul EL, Abdelatif EN. 2002. Effects of sowing date and cultivar on the yield and yield components of maize in northern Sudan.. In: Friesen DK, Palmer AFE (ed.) *Integrated approaches to higher maize productivity in the new millennium: Proc 7th Eastern Southern Africa Regional Maize Conf, Nairobi, Kenya.* 5–11 Feb. 2001. CIMMYT and Kenya Agric Res Inst, Nairobi, pp. 295–298.
- Adekayode FO, Ogunkoya MO. 2010. Effect of quantity and placement distances of inorganic 15-15-15 fertilizer in improving soil fertility status and the performance and yield of maize in a tropical rain forest zone of Nigeria. *J Soil Sci Environ Manag* 1(7), 155-163.
- Akintoye HA, Kling JG, Lucas EO. 1999. N use efficiency of single, double and synthetic maize lines grown at 4 N levels in three ecological zones of West Africa. *Field Crop Res* 60, 189-199. [http://dx.doi.org/10.1016/S0378-4290\(98\)00122-1](http://dx.doi.org/10.1016/S0378-4290(98)00122-1)
- Banziger M, Lafitte HR, Edemeades GO, Betran FJ, Berk DL, Elings A. 1999. Recent advances in breeding for tolerance to low nitrogen in tropical maize, pp. 21-34. In: Badu-Apraku B, Fakorede MAB, Ouedraogo M, Quin FM. (eds). *Proceedings of a Regional maize workshop*, 21-25 April, 1997, IITA Cotonou, Benin Republic.
- Bello OB, Abdulmalik SY, Afolabi MS, Ige SA. 2010. Correlation and path coefficient analysis of yield and agronomic characters among open pollinated maize varieties and their F₁ hybrids in a diallel cross. *Afr J Biotechnol* 9(18), 2633-2639.
- Bello OB. 2008. *Combining ability for maize grain yield in striga endemic and non-endemic environments of the southern Guinea savanna, Nigeria.* Ph.D. Thesis. The University of Ilorin, Ilorin, Nigeria. pp 86.
- Bertin P, Gallais A. 2000. Physiological and genetic basis of nitrogen use efficiency in maize. I. Agrophysiological results. *Maydica* 45, 53–66.
- Bray RH, Kurtz LT. 1945. Determination of total organic and available forms of phosphorus in soils. *Soil Sci* 59, 39-45. <http://dx.doi.org/10.1097/00010694-194501000-00006>
- Bremner JM. 1965. Total nitrogen. I methods of soil analysis. II chemical and microbiological properties. In: Black CA, Evans DD, White JL, Ensminger LE, Clerk FE, Dinauer RC (eds) *Agronomy Monograph*, 9, American Society of Agronomy Madison, Wisconsin USA. pp 26.
- Carsky RJ, Iwuafor ENO. 1999. Contribution of soil fertility research and maintenance to improved maize production and productivity in sub-Saharan Africa. In: B. Badu-Apraku et al. (eds). *Strategy for sustainable maize production in West and Central Africa.* Proc.Regional Maize Workshop, IITA-Cotonou, Benin Republic. 21–25 Apr. 1997. West and Central Africa Collaborative Maize Res. Network, Int. Inst. of Tropical Agric., Ibadan, Nigeria.
- Castleberry RM, Crum CW, Krull CF. 1984. Genetic yield improvement of U.S. maize cultivars under varying fertility and climatic environments. *Crop Sci* 24, 33-36. <http://dx.doi.org/10.2135/cropsci1984.0011183X002400010008x>
- Elemo KA. 1997. Extra-early and early maize varieties as affected by rate and time of nitrogen fertilization. In: Contribution of food self-sufficiency: Maize research and development in West and Central Africa. *Proceedings of a regional maize workshop*, IITA, Cotonou, Benin Republic. Pp. 241-246.
- Fakorede MAB, Badu-Apraku B, Coulibaly O, Fajemisin JM. 2001. Impact challenges and prospect of maize research and development in west and central Africa. *Proceedings of a regional maize workshop*, 4-7 May, 1999, WECAMAN/IITA Cotonou, Benin Republic. pp. 45.
- Gagnon B, Ziadi N. 2010. Grain corn and soil nitrogen responses to side-dress nitrogen sources and applications. *Agron J* 102, 1014-1022. <http://dx.doi.org/10.2134/agronj2010.0011>
- Genstat. 1995. Genstat 5 release 3.2. *Reference manual supplement*. Clarendon Press, Oxford, U.K.
- Gomez AG, Gomez AA. 1984. *Statistical procedures for Agricultural research*. IRRRI Book, 2nd edition, John Wiley & Sons Inc. N. Y.

- IITA (International Institute of Tropical Agriculture) 1989. Automated and semi-automated methods for soil and plant analysis. Manual series No. 7, IITA, pp. 726, Ibadan, Nigeria.
- Jafartayari D, Naderidarbahghsahi M, Javanmard H. 2012. Source limitation, nitrogen topdressing and seed yield relationship in different cultivars of maize (*Zea mays* L.). *Int J Agri Sci* 2(5), 408-413.
- Juo ASR, Mano A. 1996. Chemical dynamics in slash-and-burn Agriculture. *Agri Ecosys Environ* 58, 9-18. [http://dx.doi.org/10.1016/0167-8809\(95\)00656-7](http://dx.doi.org/10.1016/0167-8809(95)00656-7)
- Juo ASR. 1979. *Methods of Plant and Soil Analysis*. Manual Series No.1. International institute for Tropical Agriculture, Ibadan, Nigeria, 65pp.
- Kling JG, Heuberger HT, Oikeh SO, Akintoye HA, Horst WJ. 1997. Potential for developing nitrogen-use efficient maize for low-input Agricultural systems in the moist savannas of Africa. pp. 490-501. In: Edemeades GO, Banziger M, Mickelson HR, Pena-valdiva CB (eds). *Developing drought and low-N tolerant maize*. Proceedings of a symposium. March, 25-29, 1996, CIMMYT, E. I. Batan, Mexico, Mexico D. F.
- Kogbe JOS, Adediran JA. 2003. Influence of nitrogen, phosphorus and potassium application on the yield of maize in the savanna zone of Nigeria. *Afr J Biotechnol* 2 (10), 345-349.
- Lafitte HR, Banziger M. 1994. Strategies that improve maize varietal performance under low N; Progress due to selection. pp. 57-63. In: Jewell DC, Waddington SH, Ranson JK, Pixley KV (eds). *Maize research for stress environment*. Proc. of the fourth Eastern and Southern Africa regional maize conference, 28th March – 1st April, 1994, Harare, Zimbabwe.
- Lafitte HR, Edmeades GO. 1994. Improvements of tolerance to low soil nitrogen in tropical maize. I Selection criteria. *Field Crop Res* 39, 1-14. [http://dx.doi.org/10.1016/0378-4290\(94\)90066-3](http://dx.doi.org/10.1016/0378-4290(94)90066-3)
- Ma BL, Lianne MD, Edward GG. 1999. Soil nitrogen amendment effects on nitrogen uptake and grain yield of maize. *Agron J* 91, 650-656. <http://dx.doi.org/10.2134/agronj1999.914650x>
- McCullough DE, Girardin P, Mihajlovic M, Aguilera A, Tollenaar M. 1994. Influence of N supply on development and dry matter accumulation of an old and new maize hybrid. *Can J Plant Sci* 74, 471–477. <http://dx.doi.org/10.4141/cjps94-087>
- Mode CJ, Robinson HF. 1959. Pleiotropism and the genetic variance and covariance. *Biometrics* 15, 518-537. <http://dx.doi.org/10.2307/2527650>
- Namakka AI, Abubakar U, Sadik AI, Sharifai AI, Hassas AH. 2008. Effect of sowing date and nitrogen level on yield and yield components of two extra early maize varieties (*Zea mays* L.) in Sudan savanna of Nigeria. *ARPJ Agri Biol Sci* 3(2), 46-51.
- Nelson DW, Somers LE. 1992. Total carbon, total organic carbon and organic matter. pp. 539-580. In: Miller et al. (ed). *Methods of soil analysis*, Part 2, 2nd ed, Agronomy Monograph, 27, ASA, Madison, W. I.
- Pollmer WG, Eberhard D, Klein D, Dhillon BS. 1979. Genetic control of nitrogen uptake and translocation in maize. *Crop Sci* 19, 82-86. <http://dx.doi.org/10.2135/cropsci1979.0011183X001900010019x>
- Sanginga N, Okogun JA, Vanlauwe B, Diels J, Carsky RJ, Dashiell K. 2001. *Nitrogen contribution of promiscuous soybeans in maize based cropping systems*. SSSA Special Publication 58, Madison, USA. pp. 157-177.
- Smith ME, Miles CA, Beem JV. 1994. Genetic improvement of maize for nitrogen use efficiency in eastern and southern Africa. In: Jewell DC, Waddington SH, Ranson JK, Pixley KV (eds). *Maize research for stress environment*. Proc. of the fourth Eastern and Southern Africa Regional Maize Conference, 28/3-1/4 1994, Harare, Zimbabwe. Pp. 39-43.
- Tenaw W. 2000. Effect of nitrogen fertilizer rates and plant density on grain yield of maize. *Afr Crop Sci J* 8(3) 273-282.
- Uhart SA, Andrade FH. 1995. Nitrogen deficiency in maize: I. Effects on crop growth, development, dry matter partitioning, and kernel set. *Crop Sci* 35, 1376–1383. <http://dx.doi.org/10.2135/cropsci1995.0011183X003500050020x>
- Undie UL, Uwah DF, Attoe EE. 2012. Growth and development of late season maize/soybean intercropping in response to nitrogen and crop arrangement in the forest agro-ecology of South Southern Nigeria. *Int J Agric Res* 7 (1), 1-16. <http://dx.doi.org/10.3923/ijar.2012.1.16>
- Yusuf AA, Iwuafor ENO, Abaidoo RC, Olufajo OO, Sanginga N. 2009. Effect of crop rotation and nitrogen fertilization on yield and nitrogen efficiency in maize in the northern Guinea savanna of Nigeria, *Afr J Agric Res* 4 (10), 913-921.
- Yusuf AA, Janssen BH, Chude VO. 2003. Substitution rates of N, P and K in farmyard manure on an Alfisol in northern Guinea savanna of Nigeria. *J Agric Environ Int Dev* 97(12), 93-105.
- Yusuf M. 2010. Genetic variability and correlation in single cross hybrids of quality protein maize (*Zea Mays* L.). *Afr J Food Agric Nutr Dev* 10(2), 2166-2175.